



UNPRINTED AREA FOR DATE & LOT NUMBER -  
UNVARNISHED

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADS/BODY TEXT	8 pt
• LEADINGS	6.5 pt
• # OF CHARACTERS PER INCH	<50
• BULLETS	5 pt
• SPACE AFTER BULLETED SECTION	2 ems
• BARLINES, HAIRLINES	1.5 pt, 5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

100%



3/16" X 3/16" DATA MATRIX CODE (ECC200), PRINTED IN BLACK OR DARK BLUE ON UNPRINTED WHITE BACKGROUND WITH A MINIMUM OF 1/8" QUIET ZONE ALL 4 SIDES.

UNWARNISHED UNPRINTED AREA FOR LOT & EXP.

200%

**See box for important warnings and save for future use.**

**Uses:** temporarily relieves these symptoms associated with hay fever or other upper respiratory allergies, and the common cold: runny nose, sneezing, itchy, watery eyes, itching of the nose and throat, nasal congestion, headache, fever, minor aches and pains, and sinus pressure.

**Warnings:** **Allergy alert:** ibuprofen may cause a severe allergic reaction which may include: hives, facial swelling, asthma (wheezing), shock.

**Stomach bleeding warning:** Taking more than recommended may cause stomach bleeding.

**Do not use** if the child has ever had an allergic reaction to any pain reliever/fever reducer, nasal decongestant, or antihistamine. **Stop use and ask a doctor** if an allergic reaction occurs. Seek medical help right away. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Children's Advil Allergy Sinus**  
*Suspension*

Pain Reliever/Fever Reducer  
Nasal Decongestant/Antihistamine

**Active ingredients**  
(In each 5 mL teaspoon)

Chlorpheniramine maleate 1 mg.....	Antihistamine
Ibuprofen 100 mg.....	Pain reliever/fever reducer
Pseudoephedrine HCl 15 mg.....	Nasal decongestant

**Berry Flavored Liquid**  
Alcohol Free 1 FL OZ (31 mL)

**Do Not Use if breakable ring on bottle cap is separated or missing.**

**Directions:** **do not give more than directed**

- **shake well before using**
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- if needed, repeat dose every 4 hours while symptoms persist.
- do not use more than 4 times a day
- measure only with dosing cup provided

**Dosing Chart**

Weight (lb)	Age (yrs)	Dose (teaspoons)
under 48	under 6	ask a doctor
48 – 95	6 – 11	2 teaspoons

Wyeth Consumer Healthcare  
Madison, NJ 07940 U.S.A. U0183-10-1FDA05



128 CODE AREA- SUBSET C 10 MIL DENSITY WITH 4 OR 6 DIGITS-UNVARNISHED

UNPRINTED AREA FOR DATE & LOT NUMBER - UNVARNISHED

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TRIM	9 pt
• DRUG FACTS CONT INLUED	8 pt
• HEAD PAGES	8 pt
• SUBHEAD BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<30
• BULLETS	5 pt
• SPACE AFTER BULLETED SECTION	2 units
• BARLINES -HARLINES	1.5 pt, 5 pt
• SPACE BETWEEN HARLINES AND BOX END	2 spaces

100%

**See box for important warnings and save for future use.**

**Active ingredients (in each 5 mL teaspoon)**

Chlorpheniramine maleate 1 mg	Antihistamine
Ibuprofen 100 mg	Pain reliever/Fever reducer
Pseudoephedrine HCl 15 mg	Nasal decongestant

**Uses:** temporarily relieves these symptoms associated with hay fever or other upper respiratory allergies, and the common cold: runny nose, sneezing, itchy, watery eyes, itching of the nose and throat, nasal congestion, headache, fever, minor aches and pains, and sinus pressure.

**Warnings:**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** Taking more than recommended may cause stomach bleeding.

**Do not use** if the child has ever had an allergic reaction to any pain reliever/fever reducer, nasal decongestant, or antihistamine.

**Stop use and ask a doctor** if an allergic reaction occurs. Seek medical help right away.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Do Not Use If Breakable Ring on Bottle Cap Is Separated or Missing.**

**Children's Advil<sup>®</sup> Allergy Sinus Suspension**

Pain Reliever/Fever Reducer  
Nasal Decongestant/Antihistamine

**Berry Flavored Liquid**  
Alcohol Free 4 FL OZ (120 mL)

**Directions:**

- do not give more than directed
- shake well before using
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- if needed, repeat dose every 6 hours while symptoms persist
- do not use more than 4 times a day
- measure only with dosing cup provided

**Dosing Chart:**

Weight (lb)	Age (yrs)	Dose (teaspoons)
under 45	under 6	ask a doctor
45-95	6-11	2 teaspoons

**Other information:**

- each 5 mL teaspoon contains: sodium 10 mg
- store at 20-25°C (68-77°F)

**Questions or comments?** call weekdays from 8AM to 5PM EST at 1-800-84-ADWL or ask your pharmacist, doctor or healthcare professional.

Wyeth Consumer Healthcare, Madison, NJ 07940 U.S.A. U0183-20-1

3/16" X 3/16" DATA MATRIX CODE (01832001), PRINTED IN BLACK OR DARK BLUE ON UNPRINTED WHITE BACKGROUND WITH A MINIMUM OF 1/8" QUIET ZONE ALL 4 SIDES.

UNVARNISHED UNPRINTED AREA FOR LOT & EXP.

150%

**See box for important warnings and save for future use.**

**Active ingredients (in each 5 mL teaspoon)**

Chlorpheniramine maleate 1 mg	Antihistamine
Ibuprofen 100 mg	Pain reliever/Fever reducer
Pseudoephedrine HCl 15 mg	Nasal decongestant

**Uses:** temporarily relieves these symptoms associated with hay fever or other upper respiratory allergies, and the common cold: runny nose, sneezing, itchy, watery eyes, itching of the nose and throat, nasal congestion, headache, fever, minor aches and pains, and sinus pressure.

**Warnings:**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** Taking more than recommended may cause stomach bleeding.

**Do not use** if the child has ever had an allergic reaction to any pain reliever/fever reducer, nasal decongestant, or antihistamine.

**Stop use and ask a doctor** if an allergic reaction occurs. Seek medical help right away.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Do Not Use If Breakable Ring on Bottle Cap Is Separated or Missing.**

**Children's Advil<sup>®</sup> Allergy Sinus Suspension**

Pain Reliever/Fever Reducer  
Nasal Decongestant/Antihistamine

**Berry Flavored Liquid**  
Alcohol Free 4 FL OZ (120 mL)

**Directions:**

- do not give more than directed
- shake well before using
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- if needed, repeat dose every 6 hours while symptoms persist
- do not use more than 4 times a day
- measure only with dosing cup provided

**Dosing Chart:**

Weight (lb)	Age (yrs)	Dose (teaspoons)
under 45	under 6	ask a doctor
45-95	6-11	2 teaspoons

**Other information:**

- each 5 mL teaspoon contains: sodium 10 mg
- store at 20-25°C (68-77°F)

**Questions or comments?** call weekdays from 8AM to 5PM EST at 1-800-84-ADWL or ask your pharmacist, doctor or healthcare professional.

Wyeth Consumer Healthcare, Madison, NJ 07940 U.S.A. U0183-20-1

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	6 pt
• SUB-HEADS/BODY TEXT	6 pt
• LEADING	6.5 pt
• BULLETS	5 pt
• SPACE AFTER BULLETED SECTION	2 ems
• BOX LINE	.5pt
• SMALLEST @, TM etc.	4.5 pt



DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUB-HEADS/BODY TEXT	8 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<30
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ARTS
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

CONFIDENTIAL





150%



DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	6 pt
• SUBHEADS/BODY TEXT	6 pt
• LEADING	6.5 pt
• BULLETS	5 pt
• SPACE AFTER BULLETED SECTION	2 lines
• BOX LINE	5 pt
• SMALLEST ®, TM etc.	5.75 pt