

A light blue map of the Great Lakes region is centered on a dark blue background. The map shows the outlines of the five Great Lakes (Superior, Michigan, Huron, Erie, and Ontario) and the surrounding landmasses of North America and Europe. The text is overlaid on this map.

The Integrated Atmospheric Deposition Network

**Great Lakes Binational Toxics Strategy (GLBTS)
B(a)P / HCB Workgroup Meeting
December 12, 2007**

Todd Nettesheim

**Great Lakes National Program Office
U.S. Environmental Protection Agency**



Acknowledgements


- Most data slides from Ron Hites, Ilora Basu, Marta Venier, Lingyan Zhu, Eunha Hoh, and Ping “Sunny” Sun of Indiana University (IADN grantee)
- Binational IADN Steering Committee
- Tom Dann of Environment Canada for NAPS data
- And of course Melissa Hulting



Integrated Atmospheric Deposition Network (IADN)

- Joint EPA-Environment Canada project
 - Grantee is Ron Hites at Indiana University
- In operation since 1990 (GLWQA and CAAA of 1990)
- Measure PBTs in air and precipitation at 15 sites around the Great Lakes (U.S. runs five)
- Goals:
 - Determine atmospheric loadings
 - Look at trends in concentrations
 - Use data to measure progress

Integrated Atmospheric Deposition Network Monitoring Stations






National Air Pollution Surveillance (NAPS) Program

- The NAPS program is a cooperative federal-provincial-territorial network of over 800 ambient air quality monitoring instruments across Canada, mostly in urban centres.
- Substances measured:
 - PAH
 - PCDD/PCDF
 - Hexachlorobenzene (HCB)
 - Pentachlorophenol (PCP)
 - Octachlorostyrene (OCS)
 - Nitro-PAH - C13 to C22, 24 species including dinitropyrenes
 - Dioxin like PCBs
 - Metals - Hg, Cd, Pb
 - VOC - 1,4-dichlorobenzene
- <http://www.etc-cte.ec.gc.ca/NAPS/>



NAPS locations





Why air?

- Primary pathway for input to the Lakes
- Air concentrations respond rapidly to changes in emissions







HCB




Slow decrease in HCB



Annual Variation in Hexachlorobenzene Concentrations (ng/m³) at Ontario Sites (1997-2006)




Ma et al. 2004a. How Do Climate Fluctuations Affect Persistent Organic Pollutant Distribution in North America? Evidence from a Decade of Air Monitoring. *Enviro. Sci. Technol.* 38 (9): 2538 –2543.






Hexachlorobenzene


Vapor-phase conc. (pg/m³)




Particle-phase conc. (pg/m³)




Precip. conc. (pg/L)




Half-lives (years)



Max to min ratios



Maximum dates






Half-Lives for gas-phase HCB are decreasing.....

Data through....	2001	2002	2003
Eagle Harbor (Superior)	29	23.7	18
S.B. Dunes (Michigan)	15	15.7	12
Sturgeon Pt. (Erie)	18	17.1	15
IIT (Chicago)		19.4	8.3

Buehler et al. 2004

Sun et al. 2006

Most likely, increases in HCB in the late 1990s/early 2000s lengthened half-lives calculated previously.




A world map with a dark blue background and light blue landmasses. The text "Benzo(a)pyrene" is centered over the map in a bold, yellow font with a black outline. The map shows the outlines of continents and major islands, including North America, South America, Europe, Africa, Asia, and Australia.

Benzo(a)pyrene




Urban PAH levels are 10-100x higher in Chicago than in rural areas






Rural BaP concentrations are about 10x lower than urban



Benzo(a)pyrene Concentrations (ng/m³) 2005-2006
(Mean, 10th and 90th percentiles)


PAHs and PCBs are now largely an urban problem







B(a)P Particle Phase

- ◆ Superior
- Michigan
- ▲ Huron
- Erie
- Ontario








Concentration (pg/m³)





PAHs in precipitation are generally decreasing at Chicago





Some evidence of decreasing BaP concentrations on particles.....

but are levels increasing at Sleeping Bear Dunes?


Site	Precipitation	Particles
Eagle Harbor (Superior)	ns	ns
S.B. Dunes (Michigan)	ns	-11 (ms)
Burnt Island (Huron)	ns	11 (ms)
Sturgeon Pt. (Erie)	ns	4.8
Pt. Petre (Ontario)	ns	11 (ms)
IIT-Chicago	2.4	9.6 (ms)

But decreases in low MW
PAHs at Cdn sites

BaP half-lives for data through 2003




BaP higher in winter



Seasonal Variation of B(a)P (ng/m³) at Ontario Sites (1996-2006)




NAPS sites - Small or no decrease



Trend in Benzo(a)pyrene Concentrations (ng/m³) (1990-2006)




Possible decrease in *mean* in major source areas?



Benzo(a)pyrene Concentrations (ng/m^3) at Jonquière

Urban Impact for benzo(a)pyrene and other PAHs Significant: Chicago load increases Lake Michigan load by about 50%





Conclusions: HCB

- Levels are decreasing slowly again after increases in the late 90s
- Little spatial variation in concentrations



Conclusions: BaP/PAHs

- Rural sites:
 - In general, no trend over time for BaP and other PAHs in precipitation, except:
 - Decreases in low MW PAHs in precip seen only at Canadian IADN sites
 - Some evidence of decreases in the particle phase, except possible increase at Sleeping Bear Dunes (L Michigan)
- Chicago
 - PAHs in precipitation at Chicago are decreasing with a half-life of 2-5 years (2.4 years for BaP)
 - Half-life for BaP on particles is 9.6 years (marginally significant)
- Source reductions may be having an impact in urban/source areas
- Retene increasing at Brule River, Eagle Harbor and Sleeping Bear Dunes (due to wood burning?)



IADN Resources

- Main website: Station and other info, data request
 - ◆ <http://www.msc.ec.gc.ca/iadn/>
- U.S. IADN Information Page: Links to reports, Resource Page with SOPs, etc.
 - ◆ http://www.epa.gov/glnpo/monitoring/air_new_design/iadn_info.html
- U.S. IADN Government Reporting Indicator Page
 - ◆ <http://www.epa.gov/glnpo/glindicators/air/airb.html>
- Todd Nettesheim (EPA), Liisa Jantunen (EC)



Todd Nettesheim
U.S. EPA GLNPO
nettesheim.todd@epa.gov
312.353.9153