

Laboratory Astrophysics at the LLNL Electron Beam Ion Traps EBIT-I & EBIT-II

G.V. Brown, K.R. Boyce, R.L. Kelley, F.S. Porter, C.K. Stahle, A.E. Szymkowiak, W. Tillotson, P. Beiersdorfer, H. Chen, M.J. May, D. Thorn, E. Behar, M.F. Gu, S.M. Kahn

This article was submitted to NASA Laboratory Astrophysics Workshop, Moffett Field, CA, May 1-3, 2002

June 18, 2002

U.S. Department of Energy

Lawrence
Livermore
National
Laboratory

DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

This is a preprint of a paper intended for publication in a journal or proceedings. Since changes may be made before publication, this preprint is made available with the understanding that it will not be cited or reproduced without the permission of the author.

This work was performed under the auspices of the United States Department of Energy by the University of California, Lawrence Livermore National Laboratory under contract No. W-7405-Eng-48.

This report has been reproduced directly from the best available copy.

Available electronically at <http://www.doc.gov/bridge>

Available for a processing fee to U.S. Department of Energy
And its contractors in paper from
U.S. Department of Energy
Office of Scientific and Technical Information
P.O. Box 62
Oak Ridge, TN 37831-0062
Telephone: (865) 576-8401
Facsimile: (865) 576-5728
E-mail: reports@adonis.osti.gov

Available for the sale to the public from
U.S. Department of Commerce
National Technical Information Service
5285 Port Royal Road
Springfield, VA 22161
Telephone: (800) 553-6847
Facsimile: (703) 605-6900
E-mail: orders@ntis.fedworld.gov
Online ordering: <http://www.ntis.gov/ordering.htm>

OR

Lawrence Livermore National Laboratory
Technical Information Department's Digital Library
<http://www.llnl.gov/tid/Library.html>

Laboratory Astrophysics at the LLNL electron beam ion traps EBIT-I & EBIT-II

G. V. Brown, K. R. Boyce, R. L. Kelley, F. S. Porter, C. K. Stahle, A.E. Szymkowiak, and
W. Tillotson

NASA Goddard Space Flight Center, Greenbelt, MD 20771

P. Beiersdorfer, H. Chen, M. J. May, D. Thorn

*Physics and Advanced Technology, Lawrence Livermore National Laboratory, 7000 East
Avenue, Livermore, CA 94551*

E. Behar, M. F. Gu, S. M. Kahn

Columbia Astrophysics Laboratory, Columbia University, New York, NY 10027

ABSTRACT

In order to provide a complete, accurate set of atomic data for interpreting spectra provided by missions such as *XMM-Newton*, the *Chandra X-Ray Observatory*, and *Astro-E2*, we have harnessed the Lawrence Livermore National Laboratory's electron beam ion traps EBIT-I, EBIT-II, and Super-EBIT for laboratory astrophysics. In support of this work we have developed a number of unique techniques, including the ability to experimentally simulate a Maxwellian distribution of electron energies and measuring low-energy charge exchange cross sections using the "magnetic trapping mode". We have also built and operated a full suite of spectrometers spanning the 1–7000 Å wavelength band, the most recent being the NASA/Goddard Space Flight Center's Astro-E 6 X 6 engineering spare microcalorimeter array. Results of our efforts include a complete list of wavelengths of the Fe L-shell transitions, measurements of absolute and relative cross sections for direct impact, dielectronic, and resonance excitation, and measurements of low energy charge transfer reactions. A brief overview of the LLNL ebit facility, its capabilities, and some results will be discussed.

1. Introduction

Invented at the Lawrence Livermore National Laboratory (LLNL), the electron beam ion trap (Levine et al. 1988) has proven to be a useful tool for laboratory astrophysics.

The success of the LLNL ebit, EBIT-I, led to the second ebit, EBIT-II, completed at LLNL in 1990. EBIT-I was then modified to operate at higher electron beam energies and was renamed Super-EBIT. Later several other ebits were built (Silver et al. 1994; Gillaspy 1997; Biedermann et al. 1997; Curell et al. 1996; López-Urrutia et al. 2001). All of these ebits are based on the Livermore EBIT-I design, but none is an exact replica, making the operating parameters and capabilities of each device distinct.

Over the last decade the LLNL ebits have been harnessed as tools for laboratory astrophysics in support of the *Chandra*, *XMM-Newton*, and *Astro-E2* x-ray observatories. During this time several collaborations have been developed with astrophysical science programs, such as the Columbia Astrophysics Laboratory and the NASA/Goddard Space Flight Center. Our program has included the development of a plethora of instruments and methods for investigating astrophysically relevant problems.




Fig. 1.— Measured cross section as a function of electron impact energy for the $3p_{3/2} \rightarrow 2s_{1/2}$ Fe XXIV line at 10.6 \AA (Gu et al. 1999) labeled Li6. Notice the contribution below threshold from dielectronic recombination and the enhancement near threshold from resonance excitation.

2. Operating parameters and Results

EBIT-I is the current electron beam ion trap operating at LLNL. A detailed description can be found in (Beiersdorfer et al. 2000). EBIT-I operates at an electron density of $\sim 5 \times 10^{11} \text{ cm}^{-3}$ and has a mono-energetic electron beam energy that can be adjusted from $\sim 100 \text{ eV}$ to $100,000 \text{ eV}$ for a given experiment. It has five ports that observe the trap directly. For these ports we choose from over 20 spectrometers covering a range of $1\text{--}7000 \text{ \AA}$. Typically, an

experiment consists of a set of vacuum crystal spectrometers (occupying two ports) covering the 10–18 Å region continuously with 1 eV resolution; a flat-field grating spectrometer that covers the 10–50 Å region; a curved-crystal spectrometer for K-shell measurements, for example, of Fe XXV; and, the NASA/GSFC spare Astro-E spectrometer, the XRS. The XRS is a 6 X 6 square microcalorimeter array consisting of 32 active channels with an area of $\sim 13 \text{ mm}^2$ (Porter et al. 2001). Because of the high gain stability of all 32 active channels, we are able to accumulate data continuously without significant gain drift for over 12 hours.




Fig. 2.— A spectrum recorded by the NASA/GSFC microcalorimeter array. This spectrum includes the weak emission from radiative recombination (RR). The measurement of the RR emission, even with a 32 pixel array, requires long integration times.

In order to address problems encountered in the analysis of astrophysical spectra, we have developed several advanced operating techniques for EBIT-I. For example, to measure contributions from processes such as dielectronic recombination and resonance excitation, the e^- beam is swept linearly from below excitation threshold to above threshold while the spectrum is recorded as a function of beam energy. An example of one of these experiments is given in figure 1. With the addition of the NASA/GSFC XRS, we can provide absolute excitation cross sections by normalizing to radiative recombination (Chen et al. 2002) (figure 2). Other results and measuring techniques include the development of a Maxwellian temperature simulator making it possible to measure spectra in thermal plasmas at different temperatures we measured the wavelengths and identified all the Fe L-shell x-ray emission from Fe XVII–XXIV (Brown et al. 2002, 1998); we resolved a long standing problem with

relative line intensities (Brown et al. 2001). Using the magnetic trapping mode (Beiersdorfer et al. 1994), where the electron beam is off and the ions are trapped by the magnetic field of EBIT-I, we are able to measure charge exchange recombination cross sections (see Beiersdorfer et al. these proceedings). The complete list of results is too numerous to list here¹.

In support of present and future x-ray missions we will continue to study astrophysically relevant radiation processes and to develop new methods and instrumentation. This includes building and installing a new dedicated EBIT Calorimeter Spectrometer (ECS). The cryogenic and refrigerator lifetime of the ECS will see a dramatic improvement, increasing the data acquisition - cryogenic Dewar refill cycle. The ECS, in addition to the regular upgrades to our spectroscopic equipment, will improve our ability to address astrophysically relevant problems.

Work by the University of California, LLNL was performed under Contract No. W-7405-Eng-48 and supported by NASA SARA P.O. No. S-03958G.

REFERENCES

- Biedermann, C., et al. 1997, *Phys. Rev. A*, 56, R2522
- Beiersdorfer, P., et al. 2000, in *Atomic Data Needs for X-ray Astronomy*, ed. M. A. Bautista, T. R. Kallman, & A. K. Pradhan, NASA Goddard Space Flight Center, Greenbelt, MD 20771, 103-116
- Beiersdorfer, P., et al. et al. 1994, *Rapid Communications in Mass Spectrometry*, 8, 141
- Brown, G. V., et al. 2002, *Astrophys. J. Supp.*, 140(2), 589
- Brown, G. V., et al. 1998, *Astrophys. J.*, 502, 1015
- Brown, G. V., et al. 2001, *Astrophys. J. Lett.*, 557, L75-L78
- Chen, H., et al. 2002, *Astrophys. J. Lett.*, 567, L169
- Curell, F. J., et al. 1996, *J. Phys. Soc. Japan*, 65, 3186
- Gillaspy, J. 1997, *Phys. Scr.*, T71, 99
- Gu, M. F., et al. 1999, *Astrophys. J.*, 518, 1002
- Levine, M., Marrs, R., Knapp, D., & Schneider, M. 1988, *Phys. Scr.*, T22, 157
- López-Urrutia, J. C., et al. 2001, T92, 110-112
- Porter, F. S., et al. 2000, *Proc. SPIE*, 4140, 407

¹Visit <http://www-phys.llnl.gov/Research/EBIT/> and <http://homepage.mac.com/ebit> for a more complete list of measurements conducted at the LLNL EBIT facility

Silver, J., et al. 1994, Rev. Sci. Instrum., 65, 1072