

NDC 63824-056-82

Mucinex^{DM}

600 mg guaifenesin and
30 mg dextromethorphan HBr
extended-release bi-layer tablets

**EXPECTORANT AND
COUGH SUPPRESSANT**

2 EXTENDED-RELEASE BI-LAYER TABLETS

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: one or two tablets every 12 hours; not more than 4 tablets in 24 hours
- children under 12 years of age: do not use

**PHYSICIAN
SAMPLE-
NOT TO BE SOLD**

Adams[®]
LABORATORIES, INC.

Fort Worth, Texas 76155
US Pat. 6,372,252 B1
3105602 030904

3-6382405682-1

4 1/4"

15/16"

MUCINEX DM 600/30 mg (2 count) LABEL

Heading:
8 pt. Helvetica Bold
Italic, left justified

Bullets:
5 pt. solid square
Zapf Dingbats

Directions
■ do not crush, chew, or break tablet
■ take with a full glass of water
■ this product can be administered without regard for timing of meals
■ adults and children 12 years and older: one or two tablets every 12 hours; not more than 4 tablets in 24 hours
■ children under 12 years of age: do not use

Body text:
6 pt. Helvetica Regular
with 6.5 pt. leading,
left justified

2.5 pt. box barline

NDC 63824-056-20

Mucinex[®]DM

600 mg guaifenesin and
30 mg dextromethorphan HBr
extended-release bi-layer tablets

**EXPECTORANT AND
COUGH SUPPRESSANT**



20 EXTENDED-RELEASE BI-LAYER TABLETS

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: one or two tablets every 12 hours; not more than 4 tablets in 24 hours
- children under 12 years of age: do not use

US Pat. 6,372,252 B1

Adams[®]
LABORATORIES, INC. Fort Worth, Texas 76155

3-63824-056-20-3

*Please visit our Web site:
www.mucinex.com
or call 1-866-MUCINEX
(1-866-682-4639)*

3005620
030904

© 2004 Adams Laboratories, Inc.

5 1/4"

2"

MUCINEX DM 600/30 mg (20 count) LABEL

Heading:
8 pt. Helvetica Bold
Italic, left justified

Bullets:
5 pt. solid square
Zapf Dingbats

-Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: one or two tablets every 12 hours; not more than 4 tablets in 24 hours
- children under 12 years of age: do not use

Body text:
6 pt. Helvetica Regular
with 6.5 pt. leading,
left justified


2.5 pt. box barline

NDC 63824-056-40

Mucinex[®]DM

600 mg guaifenesin and
30 mg dextromethorphan HBr
extended-release bi-layer tablets

**EXPECTORANT AND
COUGH SUPPRESSANT**



40 EXTENDED-RELEASE BI-LAYER TABLETS

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: one or two tablets every 12 hours; not more than 4 tablets in 24 hours
- children under 12 years of age: do not use

US Pat. 6,372,252 B1

Adams[®]
LABORATORIES, INC. Fort Worth, Texas 76155

3-63824-056-40-1

*Please visit our Web site:
www.mucinex.com
or call 1-866-MUCINEX
(1-866-682-4639)*

3005640
030904

© 2004 Adams Laboratories, Inc.

5 1/4"

2"

MUCINEX DM 600/30 mg (40 count) LABEL

Heading:
8 pt. Helvetica Bold
Italic, left justified

Bullets:
5 pt. solid square
Zapf Dingbats

-Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: one or two tablets every 12 hours; not more than 4 tablets in 24 hours
- children under 12 years of age: do not use

Body text:
6 pt. Helvetica Regular
with 6.5 pt. leading,
left justified

2.5 pt. box barline

NDC 63824-037-82

Mucinex^{DM}

1200 mg guaifenesin and
60 mg dextromethorphan HBr
extended-release bi-layer tablets

**EXPECTORANT AND
COUGH SUPPRESSANT**

2 EXTENDED-RELEASE BI-LAYER TABLETS

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: one tablet every 12 hours; not more than 2 tablets in 24 hours
- children under 12 years of age: do not use

**PHYSICIAN
SAMPLE-
NOT TO BE SOLD**

Adams[®]
LABORATORIES, INC.

Fort Worth, Texas 76155
US Pat. 6,372,252 B1
3103702 030904

3-6382403782-0

15/16"

4 1/4"

MUCINEX DM 1200/60 mg (2 count) LABEL

Heading:
8 pt. Helvetica Bold
Italic, left justified

Bullets:
5 pt. solid square
Zapf Dingbats

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: one tablet every 12 hours; not more than 2 tablets in 24 hours
- children under 12 years of age: do not use

Body text:
6 pt. Helvetica Regular
with 6.5 pt. leading,
left justified

2.5 pt. box barline

NDC 63824-037-01

Mucinex^{DM}

1200 mg guaifenesin and
60 mg dextromethorphan HBr
extended-release bi-layer tablets

**EXPECTORANT AND
COUGH SUPPRESSANT**



10 EXTENDED-RELEASE BI-LAYER TABLETS

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: one tablet every 12 hours; not more than 2 tablets in 24 hours
- children under 12 years of age: do not use

US Pat. 6,372,252 B1

Adams[®]
LABORATORIES, INC. Fort Worth, Texas 76155

3-63824-037-01-1

*Please visit our Web site:
www.mucinex.com
or call 1-866-MUCINEX
(1-866-682-4639)*

3003701
030904

© 2004 Adams Laboratories, Inc.

5 1/4"

2"

MUCINEX DM 1200/60 mg (10 count) LABEL

Heading:
8 pt. Helvetica Bold
Italic, left justified

Bullets:
5 pt. solid square
Zapf Dingbats

-Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: one tablet every 12 hours; not more than 2 tablets in 24 hours
- children under 12 years of age: do not use

Body text:
6 pt. Helvetica Regular
with 6.5 pt. leading,
left justified

2.5 pt. box barline

NDC 63824-037-20

Mucinex^{DM}

1200 mg guaifenesin and
60 mg dextromethorphan HBr
extended-release bi-layer tablets

**EXPECTORANT AND
COUGH SUPPRESSANT**



20 EXTENDED-RELEASE BI-LAYER TABLETS

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: one tablet every 12 hours; not more than 2 tablets in 24 hours
- children under 12 years of age: do not use

US Pat. 6,372,252 B1

Adams[®]
LABORATORIES, INC. Fort Worth, Texas 76155

3-63824-037-20-2

*Please visit our Web site:
www.mucinex.com
or call 1-866-MUCINEX
(1-866-682-4639)*

3003720
030904

© 2004 Adams Laboratories, Inc.

5 1/4"

2"

MUCINEX DM 1200/60 mg (20 count) LABEL

Heading:
8 pt. Helvetica Bold
Italic, left justified

Bullets:
5 pt. solid square
Zapf Dingbats


-Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: one tablet every 12 hours; not more than 2 tablets in 24 hours
- children under 12 years of age: do not use

Body text:
6 pt. Helvetica Regular
with 6.5 pt. leading,
left justified

2.5 pt. box barline

FOR LAYOUT PURPOSES ONLY



MUCINEX DM 600/30 mg SAMPLE TRAY

Title:
14 pt. Helvetica Bold
Italic, left justified

Headings:
8 pt. Helvetica Bold
Italic, left justified

Bullets:
5 pt. solid square
Zapf Dingbats

Subheadings:
6 pt. Helvetica Bold
left justified

Graphic leading
to next panel

<p>Drug Facts</p> <p>Active ingredients (in each extended-release bi-layer tablet)</p> <table border="0"> <tr> <td>Dextromethorphan HBr 30 mg.....</td> <td>Cough suppressant</td> </tr> <tr> <td>Guaifenesin 600 mg.....</td> <td>Expectorant</td> </tr> </table>		Dextromethorphan HBr 30 mg.....	Cough suppressant	Guaifenesin 600 mg.....	Expectorant
Dextromethorphan HBr 30 mg.....	Cough suppressant				
Guaifenesin 600 mg.....	Expectorant				
<p>Uses</p> <ul style="list-style-type: none"> ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive ■ temporarily relieves: <ul style="list-style-type: none"> ■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants ■ the intensity of coughing ■ the impulse to cough to help you get to sleep 					
<p>Warnings</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ for children under 12 years of age ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product. 					
<p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ cough accompanied by too much phlegm (mucus) 					
<p>Drug Facts (continued)</p> <p>When using this product</p> <ul style="list-style-type: none"> ■ do not use more than directed <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness. <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>					
<p>Directions</p> <ul style="list-style-type: none"> ■ do not crush, chew, or break tablet ■ take with a full glass of water ■ this product can be administered without regard for timing of meals ■ adults and children 12 years and older: one or two tablets every 12 hours; not more than 4 tablets in 24 hours ■ children under 12 years of age: do not use 					
<p>Other information</p> <ul style="list-style-type: none"> ■ tamper evident: do not use if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing ■ store at 20-25°C (68-77°F) ■ see bottom of bottle for lot code and expiration date 					
<p>Inactive ingredients carbomer 934P, NF; D&C yellow #10 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF</p>					

Right justified

Title for continued panel:
8 pt. Helvetica Bold Italic

8 pt. Helvetica Regular

0.5 pt. hairline

2.5 pt. barline

Body text:
6 pt. Helvetica Regular
with 6.5 pt. leading,
left justified

2.5 pt. box barline

1 3/4"
3 1/4"

20 EXTENDED-RELEASE BI-LAYER TABLETS

Mucinex[®]DM
NEW
600 mg guaifenesin and
30 mg dextromethorphan HBr
extended-release bi-layer tablets
EXPECTORANT AND
COUGH SUPPRESSANT

Drug Facts (continued)

When using this product
■ do not use more than directed
Stop use and ask a doctor if
■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
■ do not crush, chew, or break tablet
■ take with a full glass of water
■ this product can be administered without regard for timing of meals
■ adults and children 12 years and older: one or two tablets every 12 hours; not more than 4 tablets in 24 hours
■ children under 12 years of age: do not use

Other information
■ tamper evident: do not use if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing
■ store at 20-25°C (68-77°F)
■ see bottom of bottle for lot code and expiration date


Inactive ingredients carbomer 934P, NF; D&C yellow #10 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

NEW
Mucinex[®]DM
600 mg guaifenesin and
30 mg dextromethorphan HBr
extended-release bi-layer tablets
EXPECTORANT AND
COUGH SUPPRESSANT

Please visit our Web site:
www.mucinex.com

Adams[®]
LABORATORIES, INC.
Fort Worth, Texas 76155
© 2004 Adams Laboratories, Inc.
US Pat. 6,372,252 B1

NEW NDC 63824-056-20
Mucinex[®]DM
600 mg guaifenesin and
30 mg dextromethorphan HBr
extended-release bi-layer tablets
EXPECTORANT AND
COUGH SUPPRESSANT



20 EXTENDED-RELEASE BI-LAYER TABLETS

3-63824-056-20-3

Drug Facts

Active ingredients **Purpose**
(in each extended-release bi-layer tablet)
Dextromethorphan HBr 30 mg Cough suppressant
Guaifenesin 600 mg Expectorant

Uses ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
■ temporarily relieves:
■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
■ the intensity of coughing
■ the impulse to cough to help you get to sleep

Warnings
Do not use ■ for children under 12 years of age
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
■ cough accompanied by too much phlegm (mucus) ▶

MUCINEX DM 600/30 mg (20 count) CARTON

Title:
14 pt. Helvetica Bold
Italic, left justified

Headings:
8 pt. Helvetica Bold
Italic, left justified

Bullets:
5 pt. solid square
Zapf Dingbats

Subheadings:
6 pt. Helvetica Bold
left justified

Graphic leading
to next panel

Drug Facts	
Active ingredients (in each extended-release bi-layer tablet)	Purpose
Dextromethorphan HBr 30 mg..... Cough suppressant Guaifenesin 600 mg..... Expectorant	
Uses	■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
	■ temporarily relieves:
	■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
	■ the intensity of coughing
	■ the impulse to cough to help you get to sleep
Warnings	
Do not use	■ for children under 12 years of age
	■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have	
	■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
	■ cough accompanied by too much phlegm (mucus)

Right justified

Drug Facts (continued)
When using this product
■ do not use more than directed
Stop use and ask a doctor if
■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
■ do not crush, chew, or break tablet
■ take with a full glass of water
■ this product can be administered without regard for timing of meals
■ adults and children 12 years and older: one or two tablets every 12 hours; not more than 4 tablets in 24 hours
■ children under 12 years of age: do not use
Other information
■ tamper evident: do not use if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing
■ store at 20-25°C (68-77°F)
■ see bottom of bottle for lot code and expiration date
Inactive ingredients carbomer 934P, NF; D&C yellow #10 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

Title for continued panel:
8 pt. Helvetica Bold Italic

8 pt. Helvetica Regular

0.5 pt. hairline

2.5 pt. barline

Body text:
6 pt. Helvetica Regular
with 6.5 pt. leading,
left justified

2.5 pt. box barline

1 3/4"

3 1/4"

40 EXTENDED-RELEASE BI-LAYER TABLETS

Mucinex[®]DM
NEW
600 mg guaifenesin and
30 mg dextromethorphan HBr
extended-release bi-layer tablets
EXPECTORANT AND
COUGH SUPPRESSANT

Drug Facts (continued)

When using this product

- do not use more than directed

Stop use and ask a doctor if

- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: one or two tablets every 12 hours; not more than 4 tablets in 24 hours
- children under 12 years of age: do not use

Other information

- tamper evident: do not use if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing
- store at 20-25°C (68-77°F)
- see bottom of bottle for lot code and expiration date

Inactive ingredients carbomer 934P, NF; D&C yellow #10 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

NEW
Mucinex[®]DM
600 mg guaifenesin and
30 mg dextromethorphan HBr
extended-release bi-layer tablets
EXPECTORANT AND
COUGH SUPPRESSANT


Please visit our Web site:
www.mucinex.com

Adams[®]
LABORATORIES, INC.
Fort Worth, Texas 76155
© 2004 Adams Laboratories, Inc.
US Pat. 6,372,252 B1

030904
2005640

1 3/4"

NEW NDC 63824-056-40
Mucinex[®]DM
600 mg guaifenesin and
30 mg dextromethorphan HBr
extended-release bi-layer tablets
EXPECTORANT AND
COUGH SUPPRESSANT



40 EXTENDED-RELEASE BI-LAYER TABLETS

Drug Facts

Active ingredients **Purpose**
(in each extended-release bi-layer tablet)
Dextromethorphan HBr 30 mg Cough suppressant
Guaifenesin 600 mg Expectorant

Uses ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help you get to sleep

Warnings

Do not use ■ for children under 12 years of age

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus) ▶

3-63824-056-40-1

MUCINEX DM 600/30 mg (40 count) CARTON

Title:
14 pt. Helvetica Bold
Italic, left justified

Headings:
8 pt. Helvetica Bold
Italic, left justified

Bullets:
5 pt. solid square
Zapf Dingbats

Subheadings:
6 pt. Helvetica Bold
left justified

Graphic leading
to next panel

Right justified

Drug Facts	
Active ingredients <i>(in each extended-release bi-layer tablet)</i>	Purpose
Dextromethorphan HBr 30 mg.....	Cough suppressant
Guaifenesin 600 mg.....	Expectorant
Uses	
<ul style="list-style-type: none"> ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive ■ temporarily relieves: <ul style="list-style-type: none"> ■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants ■ the intensity of coughing ■ the impulse to cough to help you get to sleep 	
Warnings	
<p>Do not use</p> <ul style="list-style-type: none"> ■ for children under 12 years of age ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product. 	
<p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ cough accompanied by too much phlegm (mucus) 	

Title for continued panel:
8 pt. Helvetica Bold Italic

Drug Facts (continued)
When using this product
<ul style="list-style-type: none"> ■ do not use more than directed
Stop use and ask a doctor if
<ul style="list-style-type: none"> ■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
<ul style="list-style-type: none"> ■ do not crush, chew, or break tablet ■ take with a full glass of water ■ this product can be administered without regard for timing of meals ■ adults and children 12 years and older: one or two tablets every 12 hours; not more than 4 tablets in 24 hours ■ children under 12 years of age: do not use
Other information
<ul style="list-style-type: none"> ■ tamper evident: do not use if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing ■ store at 20-25°C (68-77°F) ■ see bottom of bottle for lot code and expiration date
Inactive ingredients carbomer 934P, NF; D&C yellow #10 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

Title for continued panel:
8 pt. Helvetica Bold Italic

8 pt. Helvetica Regular


0.5 pt. hairline

2.5 pt. barline

Body text:
6 pt. Helvetica Regular
with 6.5 pt. leading,
left justified

2.5 pt. box barline

FOR LAYOUT PURPOSES ONLY



MUCINEX DM 1200/60 mg SAMPLE TRAY

Title:
14 pt. Helvetica Bold
Italic, left justified

Headings:
8 pt. Helvetica Bold
Italic, left justified

Bullets:
5 pt. solid square
Zapf Dingbats

Subheadings:
6 pt. Helvetica Bold
left justified

Graphic leading
to next panel

<p><i>Drug Facts</i></p> <p>Active ingredients (in each extended-release bi-layer tablet)</p> <p>Dextromethorphan HBr 60 mg Cough suppressant Guaifenesin 1200 mg Expectorant</p> <p>Purpose</p> <p>Uses ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive</p> <p>■ temporarily relieves:</p> <ul style="list-style-type: none"> ■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants ■ the intensity of coughing ■ the impulse to cough to help you get to sleep <p>Warnings</p> <p>Do not use ■ for children under 12 years of age</p> <ul style="list-style-type: none"> ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product. <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ cough accompanied by too much phlegm (mucus) 		<p><i>Drug Facts (continued)</i></p> <p>When using this product</p> <ul style="list-style-type: none"> ■ do not use more than directed <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness. <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ do not crush, chew, or break tablet ■ take with a full glass of water ■ this product can be administered without regard for timing of meals ■ adults and children 12 years and older: one tablet every 12 hours; not more than 2 tablets in 24 hours ■ children under 12 years of age: do not use <p>Other information</p> <ul style="list-style-type: none"> ■ tamper evident: do not use if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing ■ store at 20-25°C (68-77°F) ■ see bottom of bottle for lot code and expiration date <p>Inactive ingredients carboxymethylcellulose, NF; FD&C blue #1 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF</p>
--	--	---

Right justified

Title for continued panel:
8 pt. Helvetica Bold Italic

8 pt. Helvetica Regular

0.5 pt. hairline

2.5 pt. barline

6 pt. Helvetica Regular
with 6.5 pt. leading,
left justified

2.5 pt. box barline

1 3/4"
3 1/4"

10 EXTENDED-RELEASE
BI-LAYER TABLETS

Mucinex[®]DM
NEW
1200 mg guaifenesin and
60 mg dextromethorphan HBr
extended-release bi-layer tablets
EXPECTORANT AND
COUGH SUPPRESSANT

Drug Facts (continued)
When using this product
■ do not use more than directed
Stop use and ask a doctor if
■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
■ do not crush, chew, or break tablet
■ take with a full glass of water
■ this product can be administered without regard for timing of meals
■ adults and children 12 years and older: one tablet every 12 hours; not more than 2 tablets in 24 hours
■ children under 12 years of age: do not use

Other information
■ tamper evident: do not use if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing
■ store at 20-25°C (68-77°F)
■ see bottom of bottle for lot code and expiration date


Inactive ingredients carbomer 934P, NF; FD&C blue #1 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

NEW
Mucinex[®]DM
1200 mg guaifenesin and
60 mg dextromethorphan HBr
extended-release bi-layer tablets
EXPECTORANT AND
COUGH SUPPRESSANT

Please visit our Web site:
www.mucinex.com

Adams[®]
LABORATORIES, INC.
Fort Worth, Texas 76155
© 2004 Adams Laboratories, Inc.
US Pat. 6,372,252 B1

NEW NDC 63824-037-01
Mucinex[®]DM
1200 mg guaifenesin and
60 mg dextromethorphan HBr
extended-release bi-layer tablets
EXPECTORANT AND
COUGH SUPPRESSANT



10 EXTENDED-RELEASE
BI-LAYER TABLETS

3-63824-037-01-1

Drug Facts
Active ingredients **Purpose**
(in each extended-release bi-layer tablet)
Dextromethorphan HBr 60 mg Cough suppressant
Guaifenesin 1200 mg Expectorant

Uses ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
■ temporarily relieves:
■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
■ the intensity of coughing
■ the impulse to cough to help you get to sleep

Warnings
Do not use ■ for children under 12 years of age
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
■ cough accompanied by too much phlegm (mucus) ▶

MUCINEX DM 1200/60 mg (10 count) CARTON

Title:
14 pt. Helvetica Bold
Italic, left justified

Headings:
8 pt. Helvetica Bold
Italic, left justified

Bullets:
5 pt. solid square
Zapf Dingbats

Subheadings:
6 pt. Helvetica Bold
left justified

Graphic leading
to next panel

Drug Facts

Active ingredients (in each extended-release bi-layer tablet)	Purpose
Dextromethorphan HBr 60 mg.....	Cough suppressant
Guaifenesin 1200 mg.....	Expectorant

Uses ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help you get to sleep

Warnings

Do not use ■ for children under 12 years of age

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

Right justified

Title for continued panel:
8 pt. Helvetica Bold Italic

Drug Facts (continued)

When using this product

- do not use more than directed

Stop use and ask a doctor if

- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: one tablet every 12 hours; not more than 2 tablets in 24 hours
- children under 12 years of age: do not use

Other information

- tamper evident: do not use if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing
- store at 20-25°C (68-77°F)
- see bottom of bottle for lot code and expiration date

Inactive ingredients carbomer 934P, NF; FD&C blue #1 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

8 pt. Helvetica Regular

0.5 pt. hairline

2.5 pt. barline

Body text:
6 pt. Helvetica Regular
with 6.5 pt. leading,
left justified

2.5 pt. box barline

1 3/4"
3 1/4"

20 EXTENDED-RELEASE BI-LAYER TABLETS

Mucinex[®]DM
NEW

1200 mg guaifenesin and
60 mg dextromethorphan HBr
extended-release bi-layer tablets

**EXPECTORANT AND
COUGH SUPPRESSANT**

Drug Facts (continued)

When using this product

- do not use more than directed

Stop use and ask a doctor if

- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: one tablet every 12 hours; not more than 2 tablets in 24 hours
- children under 12 years of age: do not use

Other information

- tamper evident: do not use if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing
- store at 20-25°C (68-77°F)
- see bottom of bottle for lot code and expiration date

Inactive ingredients carbomer 934P, NF; FD&C blue #1 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

NEW

Mucinex[®]DM

1200 mg guaifenesin and
60 mg dextromethorphan HBr
extended-release bi-layer tablets

**EXPECTORANT AND
COUGH SUPPRESSANT**

Please visit our Web site:
www.mucinex.com

Adams[®]
LABORATORIES, INC.
Fort Worth, Texas 76155
© 2004 Adams Laboratories, Inc.
US Pat. 6,372,252 B1

030904
2003720

NEW NDC 63824-037-20

Mucinex[®]DM

1200 mg guaifenesin and
60 mg dextromethorphan HBr
extended-release bi-layer tablets

**EXPECTORANT AND
COUGH SUPPRESSANT**



20 EXTENDED-RELEASE BI-LAYER TABLETS

Drug Facts

Active ingredients **Purpose**
(in each extended-release bi-layer tablet)

Dextromethorphan HBr 60 mg Cough suppressant
Guaifenesin 1200 mg Expectorant

Uses ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help you get to sleep

Warnings

Do not use ■ for children under 12 years of age

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus) ▶

3-63824-037-20-2

MUCINEX DM 1200/60 mg (20 count) CARTON

Title:
14 pt. Helvetica Bold
Italic, left justified

Headings:
8 pt. Helvetica Bold
Italic, left justified

Bullets:
5 pt. solid square
Zapf Dingbats

Subheadings:
6 pt. Helvetica Bold
left justified

Graphic leading
to next panel

Drug Facts

Active ingredients (in each extended-release bi-layer tablet)	Purpose
Dextromethorphan HBr 60 mg.....	Cough suppressant
Guaifenesin 1200 mg.....	Expectorant

Uses ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help you get to sleep

Warnings

Do not use ■ for children under 12 years of age

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

Right justified

Title for continued panel:
8 pt. Helvetica Bold Italic

Drug Facts (continued)

When using this product

- do not use more than directed

Stop use and ask a doctor if

- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: one tablet every 12 hours; not more than 2 tablets in 24 hours
- children under 12 years of age: do not use

Other information

- tamper evident: do not use if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing
- store at 20-25°C (68-77°F)
- see bottom of bottle for lot code and expiration date

Inactive ingredients carbomer 934P, NF; FD&C blue #1 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

8 pt. Helvetica Regular

0.5 pt. hairline

2.5 pt. barline

Body text:
6 pt. Helvetica Regular
with 6.5 pt. leading,
left justified

2.5 pt. box barline