Drug Facts (continued)

Directions

Surgical Hand Antiseptic

. For the first scrub of the day or when hands are visibly soiled it is recommended to use an impregnated surgical scrub brush Clean under nalls with a nall orck Natis should be maintained with a 1 millimeter free edge • Apply to clean, dry hands and nails Dispense a half palmful (approximately 3.0 mL) of solution in one hand \cdot Dip the lingers of the opposite hand into the solution and work in under the nails \cdot Spread the remaining solution over the hand and up to just above the elbow Rub thoroughly for 1 minute or until dry, paying particular attention to the nails, cutricles and interdigital spaces • Using another half palmful (approximately 3 0 mL), repeat with the other hand • Dispense another half palmful (approximately 3 0 mL) of solution into either hand and reapply to all aspects of both hands up the wrists Rub thoroughly for 1 minute or until dry . Rinse thoroughly with water Healthcare Personnel Hand Antiseptic

 Apply to clean dry hands and nails • Dispense a palmful (approximately 5 0 mL) into the palm of one hand • Spread the solution evenly to cover both hands up to the wrists, paying particular attention to the nails cuticles and interdigital spaces Rub hands for 30 seconds or until dry Rinse thoroughly with water

Other information avoid excessive heat 40°C (104°F) • discard container when empty suitable for use with CHG Products

Inactive ingredients behentrimonium methosulfate, benzalkonium chloride, benzethonium Alonde, editors and existing to ensure the state of th

Questions? 1-800-453-4538 Monday to Friday. 8 a m to 5 p m MST

⊗BD E-Z Care[™]

Surgical and Healthcare Personnel Hand Antiseptic (70% Ethyl Alcohol w/w)

Brushless Antiseptic with Moisturizers Suitable for use with CHG products

REF 372403 • NDC 17271-240-30

Foreign Patents and Patents Pending 32 FL. OZ (946 ml)



Drug Facts

Active ingredient Ethyl alcohol 70% w/w	Purpose Antiseptic
Uses - Surgical hand antiseptic. Significantly reduces the number of microon nands and forearms prior to surgery or patient care - Healthcare personnel hand antiseptic. Handwash to help reduce bacteria th cause disease	
Warnings Flammable: keep away from fire or flame. For external use only.	
Do not use if redness or irritation occurs	
When using this product avoid contact with eyes	
Stop use and ask a doctor if severe adverse reactions occur	
Keep out of reach of children If swallowed or gets in eyes, get medical help or Control Center immediately	contact a Poison

To Instail

1 Remove product by tearing shrink wrap from the bottom

2 Remove bottle cap and apply dispensing cap 3 Slide dispenser into bracket, inserting the open air inlet on to proper receptacle

To Remove:

1 Hold bracket against the wall and pull the empty dispenser straight forward

 Discard the entire dispenser
Container is single use only of Discard when empty.
BD, BD Logo and all trademarks are property of Becton, Dickinson and Company @2002 BD 8011604 D14492 4 A(6-03) Becton, Dickinson and Company, Franklin Lakes NJ 07417 Made in USA

Drug Facts

Active ingredient		 		Purpose
Ethyl alcohol 70% w/w				. Antiseptic
	المربوبي تتخفي فستعافذ فست		***	

Uses • Surgical hand antiseptic: Significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care • Healthcare personnel hand antiseptic: Handwash to help reduce bacteria that potentially can cause disease

Warnings

Flammable: keep away from fire or flame For external use only

Do not use if redness or irritation occurs

When using this product avoid contact with eyes

Stop use and ask a doctor if severe adverse reactions occur

Keep out of reach of children. If swallowed or gets in eyes, get medical help or contact a Poison Control Center immediately

Directions • Surgical Hand Antiseptic • Apply to clean, dry hands and nails Clean under nails with a nail pick Nails should be maintained with a 1-millimeter free edge • Dispense a half palmful (approximately 3 0 mL) of solution in one hand • Dip the lingers of the opposite hand into the solution and work in under the nails. • Spread the remaining solution over the hand and up to just above the elbow. Rub thoroughly for 30 seconds or until dry, paying particular attention to the nails, cuticles and interdigital spaces • Using another half palmful (approximately 3 0 mL) of solution into either hand and other hand • Dispense another half palmful (approximately 3 0 mL) of solution into either hand and

Drug Facts (continued)

reapply to all aspects of both hands up the wrists Rub thoroughly for 30 seconds or until dry Healthcare Personnel Hand Antiseptic • Apply to clean, dry hands and nails • Dispense a half palmful (approximately 3 0 mL) into the palm of one hand • Spread the solution evenly to cover both hands up to the wrists, paying particular attention to the nails, cuticles and interdigital spaces Rub hands for 15 seconds or until dry.

Other Information avoid excessive heat 40°C (104°F) • discard container when empty • suitable for use with CHG Products

Inactive ingredients behentrimonium methosulfate, benzalkonium chloride, benzethonium chloride, cetearyl alcohol, chloritexidine gluconate, dimethicone, docosanol, fragrance, glycerin, isopropyl patimitate, methylparaben, mineral oil, PEG-10 behenyl ether, PEG-40 castor oil, propylparaben, stearalkonium chloride, USP water.

Questions? 1-800-453-4538 Monday to Friday, 8 a m to 5 p m MST

Becton, Dickinson and Company, Franklin Lakes NJ 07417 BD, BD Logo and all trademarks are property of Becton, Dickinson and Company @2003 BD

byo and an trademarks are property of becton, dickinson and company @2003 8D

8007157, D14755-3 A(6-03)

Drug Facts (continued)

Directions

Surgical Hand Antiseptic

 Apply to clean, dry hands and nais. Clean under naits with a nail pick. Nais should be maintained with a 1-mil/meter free dage * 0)sponse a hall paimful (approximately 30 mL) of solution in one hand * Dip the ingers of the opposite hand into the solution and work in under the nais. Spread the remaining solution over the hand and up to just above the ellow. Rub thoroughly for 30 seconds or unit dry, paying particular attention to the nais: culcles and interrigidial spaces. Using another half palmful (approximately 30 mL), repeat with the other hand • Dispense another half palmful (approximately 30 mL) of solution into either hand and respity to all aspects of both thands to the roughly to 30 mL of solution response or unit dry.

Healthcare Personnel Hand Antiseptic • Apply to clean, dry hands and nais • Dispense a half palmful (approximately 30 mL) into the paim of one hand • Spread the solution evenly to cover both hands up to the wrists paying particular attention to the nails cuticles and interdigital spaces. Rub hands for 15 seconds or until dry.

Other Information avoid excessive heat 40°C (104°F) • discard container when empty • suitable for use with CHG Products

Inactive ingredients behavirmonium methosulfate, benzelkonium chloride benzelhorium chloride cetearyl alcohol, chlorhexidine gluconate, dimethicone docosanol, fragrance, glycerin, isopropyl palmitate methylpataben, mineral oli, PEG-10 behenyl ether, PEG-40 castor oli provplaraben, stearalkonium chloride, USP water

Questions? 1-800-453-4538 Monday to Friday, 8 a m to 5 p m MST

SD E-Z Care™

Rinseless, Brushless Antiseptic

REE 372408 * NOG * 271 240 80 *****

Surgical and Healthcare Personnel Hand Antiseptic with Moisturizers 70% Ethyl Atonal www Suitable for use with CHG products

Drug Facts

Active ingredient Ethyl alcohol 70% w/w	Purpose Antiseptic
USES - Surgical hand antiseptic: Significantly reduces the number of hands and forearms prior to surgery or patient care - Healthcare personnel hand antiseptic: Handwash to help reduce ba cause disease	
Warnings Flammable: keep away from fire or flame For external use only.	1111
Do not use if redness or irritation occurs	
When using this product avoid contact with eyes	
Stop use and ask a doctor if severe adverse reactions occur	
Keep out of reach of children If swallowed or gets in eyes, get medica Control Center immediately	I help or contact a Poison
e fremili	



