

PATIENT INFORMATION

Premarin[®]

(conjugated estrogens tablets, USP)

Read this PATIENT INFORMATION before you start taking Premarin and read what you get each time you refill your Premarin prescription. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or your treatment.

What is the most important information I should know about Premarin (an estrogen mixture)?

- Estrogens increase the chance of getting cancer of the uterus.

Report any unusual vaginal bleeding right away while you are taking Premarin. Vaginal bleeding after menopause may be a warning sign of cancer of the uterus (womb). Your healthcare provider should check any unusual vaginal bleeding to find out the cause.

- Do not use estrogens with or without progestins to prevent heart disease, heart attacks, strokes, or dementia.

Using estrogens, with or without progestins, may increase your chance of getting heart attacks, strokes, breast cancer, and blood clots. Using estrogens, with or without progestins, may increase your chance of getting dementia, based on a study of women age 65 years or older. You and your healthcare provider should talk regularly about whether you still need treatment with Premarin.

What is Premarin?

Premarin is a medicine that contains a mixture of estrogen hormones.

Premarin is used after menopause to:

- **Reduce moderate to severe hot flashes.** Estrogens are hormones made by a woman's ovaries. The ovaries normally stop making estrogens when a woman is between 45 and 55 years old. This drop in body estrogen levels causes the “change of life” or menopause (the end of monthly menstrual periods). Sometimes both ovaries are removed during an operation before natural menopause takes place. The sudden drop in estrogen levels causes “surgical menopause.”

When the estrogen levels begin dropping, some women get very uncomfortable symptoms, such as feelings of warmth in the face, neck, and chest, or sudden strong feelings of heat and sweating (“hot flashes” or “hot flushes”). In some women the symptoms are mild, and they will not need to take estrogens. In other women, symptoms can be more severe. You and your healthcare provider should talk regularly about whether you still need treatment with Premarin.

- **Treat moderate to severe dryness, itching, and burning, in and around the vagina.** You and your healthcare provider should talk regularly about whether you still need treatment with Premarin to control these problems. If you use Premarin only to treat your dryness, itching, and burning in and around your vagina, talk with your healthcare provider about whether a topical vaginal product would be better for you.
- **Help reduce your chances of getting osteoporosis (thin weak bones).** Osteoporosis from menopause is a thinning of the bones that makes them weaker and easier to break. If you use Premarin only to prevent osteoporosis due to menopause, talk with your healthcare provider about whether a different treatment or medicine without estrogens might be better for you. You and your healthcare provider should talk regularly about whether you should continue with Premarin.

Weight-bearing exercise, like walking or running, and taking calcium and vitamin D supplements may also lower your chances for getting postmenopausal osteoporosis. It is important to talk about exercise and supplements with your healthcare provider before starting them.

Premarin is also used to:

- **Treat certain conditions in women before menopause if their ovaries do not make enough estrogen naturally.**
- **Ease symptoms of certain cancers that have spread through the body, in men and women.**

Who should not take Premarin?

Do not start taking Premarin if you:

- **Have unusual vaginal bleeding.**
- **Currently have or have had certain cancers.** Estrogens may increase the chance of getting certain types of cancers, including cancer of the breast or uterus. If you have or have had cancer, talk with your healthcare provider about whether you should take Premarin.
- **Had a stroke or heart attack in the past year.**
- **Currently have or have had blood clots.**
- **Currently have or have had liver problems.**
- **Are allergic to Premarin tablets or any of its ingredients.** See the list of ingredients in Premarin at the end of this leaflet.
- **Think you may be pregnant.**

Tell your healthcare provider:

- **If you are breast feeding.** The hormones in Premarin can pass into your milk.
- **About all of your medical problems.** Your healthcare provider may need to check you more carefully if you have certain conditions, such as asthma (wheezing), epilepsy (seizures), migraine, endometriosis, lupus, problems with your heart, liver, thyroid, kidneys, or have high calcium levels in your blood.
- **About all the medicines you take,** including prescription and nonprescription medicines, vitamins, and herbal supplements. Some medicines may affect how Premarin works. Premarin may also affect how your other medicines work.
- **If you are going to have surgery or will be on bedrest.** You may need to stop taking estrogens.

How should I take Premarin?

- Take one Premarin tablet at the same time each day.
- If you miss a dose, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your normal schedule. Do not take 2 doses at the same time.
- Estrogens should be used at the lowest dose possible for your treatment only as long as needed. You and your healthcare provider should talk regularly (for example, every 3 to 6 months) about the dose you are taking and whether you still need treatment with Premarin.

What are the possible side effects of Premarin?

Side effects are grouped by how serious they are and how often they happen when you are treated.

Serious but less common side effects include:

- Breast cancer
- Cancer of the uterus
- Stroke
- Heart attack
- Blood clots
- Dementia
- Gallbladder disease
- Ovarian cancer

- High blood pressure
- Liver problems
- High blood sugar
- Enlargement of benign tumors of the uterus (“fibroids”)

Some of the warning signs of these serious side effects include:

- Breast lumps
- Unusual vaginal bleeding
- Dizziness and faintness
- Changes in speech
- Severe headaches
- Chest pain
- Shortness of breath
- Pains in your legs
- Changes in vision
- Vomiting
- Yellowing of the skin, eyes or nail beds

Call your healthcare provider right away if you get any of these warning signs, or any other unusual symptoms that concern you.

Less serious but common side effects include:

- Headache
- Breast pain
- Irregular vaginal bleeding or spotting
- Stomach/abdominal cramps, bloating
- Nausea and vomiting
- Hair loss
- Fluid retention

- Vaginal yeast infection

These are not all the possible side effects of Premarin. For more information, ask your healthcare provider or pharmacist.

What can I do to lower my chances of getting a serious side effect with Premarin?

- Talk with your healthcare provider regularly about whether you should continue taking Premarin.
- If you have a uterus, talk to your healthcare provider about whether the addition of a progestin (a different prescribed hormone medication) is right for you. The addition of a progestin is generally recommended for women with a uterus to reduce the chance of getting cancer of the uterus.
- See your healthcare provider right away if you get vaginal bleeding while taking Premarin.
- Have a breast exam and mammogram (breast X-ray) every year unless your healthcare provider tells you something else. If members of your family have had breast cancer or if you have ever had breast lumps or an abnormal mammogram, you may need to have breast exams more often.
- If you have high blood pressure, high cholesterol (fat in the blood), diabetes, are overweight, or if you use tobacco, you may have higher chances for getting heart disease. Ask your healthcare provider for ways to lower your chances for getting heart disease.

General information about the safe and effective use of Premarin

Medicines are sometimes prescribed for conditions that are not mentioned in patient information leaflets. Do not take Premarin for conditions for which it was not prescribed. Do not give Premarin to other people, even if they have the same symptoms you have. It may harm them.

Keep Premarin out of the reach of children.

This leaflet provides a summary of the most important information about Premarin. If you would like more information, talk with your healthcare provider or pharmacist. You can ask for information about Premarin that is written for health professionals. You can get more information by calling the toll free number 800-934-5556.

What are the ingredients in Premarin?

Premarin contains a mixture of conjugated estrogens, which are a mixture of sodium estrone sulfate and sodium equilin sulfate and other components including sodium sulfate conjugates, 17 α -dihydroequilin, 17 α -estradiol, and 17 β -dihydroequilin.

Premarin 0.3 mg, 0.45 mg, 0.625 mg, 0.9 mg, and 1.25 mg tablets also contain the following inactive ingredients: calcium phosphate tribasic, hydroxypropyl cellulose, microcrystalline cellulose, powdered cellulose, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, sucrose and titanium dioxide.

The tablets come in different strengths and each strength tablet is a different color. The color ingredients are:

— 0.3 mg tablet (green color): D&C Yellow No. 10 and FD&C Blue No. 2.

— 0.45 mg tablet (blue color): FD&C Blue No. 2.

— 0.625 mg tablet (maroon color): FD&C Blue No. 2 and FD&C Red No. 40.

— 0.9 mg tablet (white color): D&C Red No. 30 and D&C Red No. 7.

— 1.25 mg tablet (yellow color): black iron oxide, D&C Yellow No. 10, and FD&C Yellow No. 6.

The appearance of these tablets is a trademark of Wyeth Pharmaceuticals.



This product's label may have been updated. For current package insert and further product information, please visit www.wyeth.com or call our medical communications department toll-free at 1-800-934-5556.



Wyeth[®]

Wyeth Pharmaceuticals Inc.
Philadelphia, PA 19101

W10405C021
ET01
Rev March 2008