

to obtain uniform distribution of sperm, and the finger bowl set aside for 5 hours, half submerged in running sea water. At the expiration of this period a sample was taken for examination and cleaved and uncleaved eggs were counted. In each case 300 eggs of the sample were examined. All tests were made in water of 31 to 32‰ salinity and 20.8° to 21.4° C. During the first 4 hours of aging the percentages of cleaved eggs declined from 90 to 70. After 5 or 6 hours the percentages dropped to 60. Then the fertilizability decreased to about 20 percent in 10 hours, and only a few eggs cleaved normally after 12 and 24 hours of aging.

It is common knowledge among embryologists that the fertilizing power of spermatozoa is not decreased if sperm is kept at 10° to 12° C. in a concentrated suspension in a tightly closed container. This is also true for oyster sperm. Its fertilizing power is affected by dilution and increased temperature. At room temperature in a dilute suspension, the spermatozoa lose their fertilizing ability within 4 to 5 hours. However, in a concentrated suspension, protected from evaporation, and stored in a refrigerator at about 10° C. the sperm remains active and retains its full fertilizing power for 24 hours and possibly longer.

The effect of cold storage on the fertilizability of eggs is not known. On several occasions ripe females with intact shells were kept for 3 to 4 days in a refrigerator (about 10° C.) and after that were successfully used in spawning experiments. The effect of cold storage on eggs of the excised ovary has not been studied.

#### POLARITY OF EGG

The polarity of all molluscan eggs apparently is determined while they are still attached to the wall of the ovarian follicle. Presumably the side on which the food reaches the growing ovocyte becomes the vegetative pole of the mature egg (Raven, 1958).

The metabolic gradient along the egg axis is indicated by a concentration of cytochrome oxidase which Kobayashi (1959) detected with M-Nadi reaction; the activity of the enzyme was observed using Gräff's modification of this method. (The reader not familiar with the reaction and its significance in cytochemical research is referred to the publications of Danielli (1958), Deane, Barnett, and Seligman (1960), and to a review by

Novikoff (1961a, p. 308).) In brief, the localization of oxidative enzymes and their presence in mitochondria can be determined by the staining reactions. The rate of respiration of eggs of *C. virginica* increases with fertilization by a factor of 1.4 (Ballentine, 1940) but in the eggs of the Sydney rock oyster, *C. commercialis*, the rate of respiration increases only at the onset of the first cleavage (Cleland, 1950).

#### CLEAVAGE

The spermatozoon may enter the oyster egg anywhere. Its path inside the egg cytoplasm toward the nucleus has not been described, and cytological details of the process leading to the fusion of the female and male pronuclei have not been studied. It is probable that the major features of these events are not different from those found in other mollusks. Maturation divisions occur in the oyster egg after the elevation of the fertilization membrane. The germinal vesicle (ovocyte nucleus) breaks down and moves toward the egg's periphery. At temperatures of 22° to 24° C. the first polar body is formed within 25 to 50 minutes after the addition of sperm. The reduction of the number of chromosomes probably takes place during the first meiotic division. This is to a certain extent corroborated by an examination of fertilized eggs of *C. virginica* stained in toto with Feulgen reagent or with acetic orcein. Unfortunately the results are not consistent enough to draw a final conclusion and the question remains unanswered, awaiting a complete cytological study.

The second polar body is formed shortly after the first, within 45 to 70 minutes after fertilization (at 22.5° to 24° C.). The two polar bodies remain attached to the surface of the egg (fig. 319) until the completion of cleavage and emergence of the trochophore.

The first cleavage following the formation of the second polar body divides the egg meridionally into two unequal cells designated as AB and CD (fig. 320). The inequality of the blastomeres is due to the occurrence of a polar lobe. Because the egg appears to consist of three cells this stage received the name trefoil.

The plane of the second division, also meridional, is at a right angle to the first. Both blastomeres divide synchronously and separate into the four quadrants. In fig. 321, drawn from a photograph of a cleaving egg taken from the animal pole, the position of the spindles indicates the plane of new




FIGURE 319.—Photomicrograph of a live fertilized egg of *C. virginica* after the formation of two polar bodies (top of egg). High-phase oil immersion lens.




FIGURE 321.—Beginning of second cleavage of the egg. Viewed from the animal pole. Whole mount, Kahle, Feulgen stain.




FIGURE 320.—First cleavage division of the egg of *C. virginica* 70 minutes after fertilization. Blastomere AB (left) and CD (right). Polar body on top.




FIGURE 322.—Section of an egg of *C. virginica* at the second cleavage. Beginning of anaphase. Kahle, Heidenhain iron-hematoxylin.

furrows which will intersect the egg into four cells, A, B, C, and D. At this stage the mitotic figures are fairly large and the chromosomes are in a favorable position for examination. In the best sectioned preparations of the cleaving egg, eight daughter chromosomes were counted at the beginning of anaphase (fig. 322). It would be, however, premature to state that the diploid number of chromosomes in *C. virginica* is 8 because on other preparations 7, 9, and 10 were counted.

The third division of each quadrant cuts the cells in the equatorial plane and separates the

first quartet of micromeres, small cells at the animal pole, from the macromeres, or larger cells at the vegetal pole (fig. 323).

At the fourth and fifth cleavages, resulting in 16- and 32-cell stages, the micromeres overgrow the macromeres. Only one of the macromeres is visible in the figure 324 and two in 325, which show a side view of an oyster egg at these two stages of development.

Cell lineage, or tracing the developmental history of the cleavage blastomeres through to




FIGURE 323.—Third division of fertilized egg of *C. virginica*; side view. Separation of micromeres (on top) from macromeres (bottom). Only one macromere is seen in the plane of view. Live egg.




FIGURE 325.—Fifth cleavage of egg of *C. virginica* and the formation of the third quartet of micromeres (32-cell stage). Side view. Drawn from photomicrograph of live cell.




FIGURE 324.—Fourth cleavage of egg of *C. virginica* and the formation of 8 micromeres (2d quartet). Side view. Drawn from photomicrograph of live cell.

their ultimate fates as parts of the larva or adult, was first described by Whitman (1878) for the egg of *Clepsine*, and followed by Wilson (1892) for the egg of *Nereis*. The works of Lillie (1895) on the development of Unionidae, Conklin (1897, 1908) on *Crepidula* and *Fulgur*, Meisenheimer

(1901a, 1901b) on *Dreissensia* and *Cyclas*, and Wilson (1904a, 1904b) on *Dentalium* and *Patella* constitute major contributions to the embryology of mollusks.

The nomenclature of the cleavage blastomeres, as developed by Wilson (1892), was progressively modified by Conklin (1897), Mead (1897), and Child (1900); the present system is based largely on the work of Robert (1902) on the development of the *Trochus* egg. The system is a combination of letters and numbers by which the blastomeres are identified.

The first four cells or macromeres are designated as A, B, C, and D; in the majority of cases studied D is the largest of the four and is situated at the side which will develop into the posterior portion of the embryo. When the first four blastomeres divide, their daughter cells are denominated 1a and 1A, 1b and 1B, 1c and 1C, and 1d and 1D, the small letters in each case referring to the micromere and the capital letter to the macromere.

In successive divisions 1A divides into 2a and 2A, 1B into 2b and 2B, and so on. When the micromeres divide, 1a is divided into 1a<sup>1</sup> and 1a<sup>2</sup>, the superscript 1 denoting the daughter cell which is nearest to the animal pole and superscript 2 the one nearer to the vegetal pole. The nomenclature is capable of indefinite expansion, but certain confusion arises when the two daughter cells resulting

from the division of one cell lie at an equal distance from the pole. In this case the letters r for right and l for left are used. The practice is, however, not generally followed.

Descriptions of various types of cleavage can be found in volume 1 of MacBride (1914). The equal and unequal cleavages in the spirally cleaving eggs of annelids and molluscs are discussed by Costello (1955) in chapter 2 of Willier, Weiss, and Hamburger (1955).

During division the micromeres of a quartet, viewed from the animal pole, become slightly displaced because the spindles of the dividing cells (not shown in fig. 324 or 325) occupy an oblique position with respect to the egg's axis. At the following divisions the plane of separation of daughter cells is oriented approximately at a right angle to the preceding divisions. The pattern of such cleavage is called spiral. It gives rise to an irregular morula (sterroblastula according to Korschelt and Heider, 1895, from the Greek "sterros" meaning firm) found in annelids (*Nereis*), in some bivalves (Ostreidae, *Teredo*), and in gastropods (*Crepidula*, *Fulgur*, *Nassa*), and others. In all cases the sterroblastula arises from an unequal cleavage during which the micromeres overlie the macromeres, and at each division are slightly displaced to the right (dextrotropic cleavage) or to the left (laetotropic cleavage). Sometimes, as in the case of *Dreissensia*, the second dextrotropic cleavage is followed by a third dextrotropic cleavage after which the normal alternating course is established (Meisenheimer, 1901a). In the case of oyster eggs, as shown by Fujita (1929) for *C. gigas*, the cleavage is laetotropic.

The multicellular stages of a *C. virginica* egg are reached in the course of the sixth and ensuing cleavages (figs. 326 and 327) during which the micromeres divide much more rapidly than the macromeres, become progressively smaller and overgrow the vegetal pole. Approximately at this stage the sterroblastula of an oyster is formed.

Gastrulation begins with epibolic extension of the micromeres. At 22° to 24° C. the stage shown in fig. 328 is reached within 4 to 6 hours.

The cell lineage of *C. virginica* has not been studied; the stages of development of eggs of the species shown in figures 320 to 328 are similar to those previously described by Brooks (1898); Horst (1882) for *O. edulis*; Seno (1929) for *O. denselamellosa*; Hori (1933) for *O. lurida*; Yasugi (1938) for *O. spinosa*, and *C. gigas*. Yasugi




FIGURE 326.—Formation of sterroblastula in *C. virginica* egg; of the four macromeres only two are visible from the side. Micromeres begin to overgrow the vegetal pole. Drawn from photomicrograph of live egg.




FIGURE 327.—Advanced sterroblastula stage of *C. virginica*. One of the macromeres has divided into two daughter cells of equal size. Drawn from photomicrograph of live egg.




FIGURE 328.—Early stage of gastrulation in the egg of *C. virginica*. Seen at a section of an egg cut along its axis. Heidenhain, iron-hematoxylin.

found that equal cleavage can be induced artificially in eggs of the Japanese oyster by centrifuging for 2 minutes at 1,500 r.p.m. and at the centrifuge radius of 14 cm.

Fujita (1929) gives a brief account of the cell lineage of the eggs of *C. gigas* and states that the mode of cleavage of this species is identical to that of *C. virginica*. The main features described by him are as follows. The first polar body in the fertilized egg of *C. gigas* appears 15 minutes after insemination. At the two-cell stage (fig. 329) the two blastomeres of unequal size, AB and CD, are separated along the meridional plane. Their position corresponds to the anterior (Ant.) and posterior (Pst.) ends of the embryo. The second division, also meridional, separates the four blastomeres A, B, C, and D (fig. 329b).

A and B represent the anterior, and C and D the posterior halves of the embryo, while B and C form its left and A and D its right halves (fig. 329b). The ensuing cleavage starts with the blastomere B and proceeds in laeotropic order to C, D, and A; the resulting daughter cells, the micromeres  $a_1$ ,  $b_1$ ,  $c_1$ , and  $d_1$ , retain the shape of the mother cells but are smaller. The macromere D and micromere  $d_1$  are respectively the largest. The four daughter cells  $a_1$  through  $d_1$  form the

first quartet of micromeres located between the macromeres on the dorsal side of the embryo.

The 12-cell stage is initiated by the division of the macromere D; the ensuing larger cell  $d_2$  (fig. 329c) is generally known as the first somatoblast X. (In the system of nomenclature used by American and European embryologists (see p. 346) the 1D cell gives rise to 2d and 2D and the 2d is the X cell.)

The cleavage is continued laeotropically, and the second generations of micromeres  $a_2$ ,  $b_2$ , and  $c_2$  are smaller than the first macromeres. They lie on the outside of the macromeres. The third cleavage of macromeres A, B, and C continues in laeotropic order and results in the micromeres  $a_3$ ,  $b_3$ , and  $c_3$ ; they are larger than other micromeres. After the third cleavage the macromeres make no further contribution to the formation of micromeres and in the course of development become the entoderm. The first somatoblast (X cell) gives rise to many organs of ectodermal origin. At the 18-cell stage of the embryo the position of cell X and its first divisions mark the beginning of the transition from spiral to bilateral symmetry (fig. 329d).

The mesoderm begins to form at about the 32-cell stage with the appearance of cell 4d, the second somatoblast, also designated as cell M. In bivalves the cell M remains at the surface for a long time, then divides into the two mesodermal teloblasts which sink into blastocoel (Raven, 1958, p. 117). The formation of mesoderm in *C. gigas* has not been followed in detail, but as a rule the mesoderm bands in bivalves remain rather rudimentary (Raven, 1958). Fujita states that the establishment of the three germinal layers in *C. gigas* is completed at the 30-cell stage (fig. 329 e and f).

The gastrula stage is reached in 4 to 6 hours. The cell lineage of *C. gigas* is generally comparable to that described for other bivalves (see: Raven, 1958, p. 70), but for details the reader should consult Fujita's (1929) original text and his drawings.

In about 4 to 6 hours after fertilization, an egg of *C. virginica* reaches the stage (fig. 330) when a few large cilia become visible at the vegetal pole, the oval-shaped body is covered with very




FIGURE 329.—Several stages of development of the egg of *C. gigas*. Redrawn from Fujita, 1929. a—Two-cell stage, Ant.—anterior, Pst.—posterior ends; b—Four-cell stage, formation of blastomeres A, B, C, and D; c—12-cell stage and the formation of the first somatoblast (cell X), viewed from the animal pole; d—embryo viewed from vegetative pole after the formation of the mesomere M; e—cleavage of mesomere M and the first somatoblast X, posterior view optical section; f—advanced stage of development showing the arrangement of the mesomeres M, M, and the somatoblasts, X, X, X<sub>1</sub>, X<sub>1</sub>, posterior view optical section. Cleavage nomenclature as given by Fujita.




FIGURE 330.—Larva of *C. virginica* ready to hatch. Drawn from photomicrograph of live larva.

fine ciliation, and two polar bodies still remain attached to the animal pole. The beating of the cilia is not coordinated at this stage, and the movements of the larva are irregular and spasmodic. A few minutes later a girdle of powerful cilia is formed, the polar bodies are lost, and the larva begins to swim upward (fig. 331). In a finger bowl containing cleaving eggs, the newly hatched larvae appear as white columns rising from the layer of fertilized eggs on the bottom of the container (fig. 332). The larvae can be pipetted off easily, transferred into larger containers and provided with suitable food.

The time required to complete the development of an oyster egg varies, depending on condition of eggs, temperature, salinity, oxygenation of water, and other environmental factors. Records of three sets of observations made in the Woods




FIGURE 331.—Larva of *C. virginica* at the time of its emergence 6 to 6½ hours after fertilization. Drawn from a photomicrograph of a live larva.

Hole laboratory at room temperatures varying from 22.5° to 24.5° C. and salinity of water of 32.2 ‰ are given in table 37. To obtain records of rates of development at different temperatures, several hundred artificially fertilized eggs were placed in each Syracuse dish filled with fresh sea water and covered to prevent evaporation. The debris was removed, and the water contained no unfertilized or cytolized eggs.

TABLE 37.—Observations on the time required for artificially fertilized eggs of *C. virginica* to reach trochophore stage

All observations were made at Woods Hole in July at room temperatures varying from 23° to 25° C. The time required to reach different stages varied in different groups of eggs. The observations are arranged in two groups: A and B, which differ primarily in the duration of time required to reach rotating blastula and trochophore stages.

STAGES		
Stage of development	A	B
Fertilization membrane.....	5 min.....	10 to 25 min.
First polar body.....	40 min.....	25 to 52 min.
Second polar body.....	1 hr. 10 min.....	40 to 65 min.
First cleavage.....	1 hr. 12 min.....	45 min.
Second cleavage.....	.....	52 to 120 min.
Third cleavage.....	2 hr. 10 min.....	55 to 195 min.
Morula stage.....	.....	135 min.
Rotating blastula.....	4 hr.....	6 hr. 30 min.
Trochophore.....	5 hr.....	8 to 9 hr.




FIGURE 332.—The emergence of larvae of *C. virginica* from fertilized eggs kept in a finger bowl. The free-swimming larvae form columns, which tend to disperse at the surface. Drawn from life.

## BIBLIOGRAPHY

- AFZELIUS, BJÖRN A.  
1955. The fine structure of the sea urchin spermatozoa as revealed by the electron microscope. *Zeitschrift für Zellforschung und mikroskopische Anatomie*, Band 42, pp. 134-148.
- ALLEN, ROBERT DAY.  
1953. Fertilization and artificial activation in the egg of the surf-clam, *Spisula solidissima*. *Biological Bulletin*, vol. 105, No. 2, pp. 213-239.
- AMEMIYA, IKUSAKU.  
1926. Notes on experiments on the early developmental stages of the Portuguese, American and English native oysters, with special reference to the effect of varying salinity. *Journal of the Marine Biological Association of the United Kingdom*, vol. 14, No. 1, pp. 161-175.
- BALLENTINE, ROBERT.  
1940. Analysis of the changes in respiratory activity accompanying the fertilization of marine eggs. *Journal of Cellular and Comparative Physiology*, vol. 15, No. 2, pp. 217-232.
- BERG, WILLIAM E.  
1950. Lytic effects of sperm extracts on the eggs of *Mytilus edulis*. *Biological Bulletin*, vol. 98, No. 2, pp. 128-138.
- BRACHET, JEAN.  
1944. *Embryologie chimique*. Masson et Cie, Paris, 509 pp.  
1960. *The biochemistry of development*. Pergamon Press, New York, 320 pp.
- BRAMBELL, F. W. ROGERS.  
1924. The nature and origin of yolk. *Experimental studies of the oöcytes of Helix aspersa and Patella vulgata*. *British Journal of Experimental Biology*, vol. 1, No. 4, pp. 501-517.
- BRETSCHNEIDER, L. H., and CHR. P. RAVEN.  
1954. Structural and topochemical changes in the egg cells of *Limnaea stagnalis* L. during oogenesis. *Archives Néerlandaises de Zoologie*, tome 10, livraison 1 (1951), pp. 1-31.
- BROOKS, W. K.  
1880. Development of the American oyster (*Ostrea virginica* List.). *Studies from the Biological Laboratory, Johns Hopkins University, Baltimore, Md.*, No. 4 (vol. 1), pp. 1-81.  
1898. Embryonic development. *In* H. F. Moore, *Oysters and methods of oyster-culture*, pp. 270-272. U.S. Commission of Fish and Fisheries, Part 23, Report of the Commissioner for the year ending June 30, 1897.
- CHILD, CHARLES MANNING.  
1900. The early development of *Arenicola* and *Sternaspis*. *Archiv für Entwicklungsmechanik der Organismen*, Band 9, pp. 587-723.
- CLELAND, K. W.  
1947. Some observations on the cytology of oogenesis in the Sydney Rock oyster (*Ostrea commercialis* I. and R.). *Proceedings of the Linnean Society of New South Wales*, vol. 72, pp. 159-182.
1950. Respiration and cell division in developing oyster eggs. *Proceedings of the Linnean Society of New South Wales*, vol. 75, pp. 282-295.
1951. The enzymatic architecture of the unfertilized oyster egg. *Australian Journal of Experimental Biology and Medical Science*, vol. 29, part 1, pp. 35-45.
- COE, WESLEY R.  
1931. Spermatogenesis in the California oyster (*Ostrea lurida*). *Biological Bulletin*, vol. 61, No. 3, pp. 309-315.  
1932a. Sexual phases in the American oyster (*Ostrea virginica*). *Biological Bulletin*, vol. 63, No. 3, pp. 419-441.  
1932b. Development of the gonads and the sequence of the sexual phases in the California oyster (*Ostrea lurida*). *Bulletin of the Scripps Institution of Oceanography of the University of California*, Technical Series, vol. 3, No. 6, pp. 119-144.  
1934. Alternation of sexuality in oysters. *American Naturalist*, vol. 68, No. 716, pp. 236-251.  
1936. Environment and sex in the oviparous oyster *Ostrea virginica*. *Biological Bulletin*, vol. 71, No. 2, pp. 353-359.  
1938. Conditions influencing change of sex in mollusks of the genus *Crepidula*. *Journal of Experimental Zoology*, vol. 77, No. 3, pp. 401-424.
- COLWIN, ARTHUR L., and LAURA HUNTER COLWIN.  
1955. Sperm entry and the acrosome filament (*Holothuria atra* and *Asterias amurensis*). *Journal of Morphology*, vol. 97, No. 3, pp. 543-567.  
1957. Morphology of fertilization: Acrosome filament formation and sperm entry. *In* Albert Tyler, R. C. von Borstel, and Charles B. Metz (editors), *The beginnings of embryonic development*, pp. 135-168. American Association for the Advancement of Science, Publication No. 48, Washington, D.C.  
1960. Egg membrane lytic activity of sperm extract and its significance in relation to sperm entry in *Hydroides hexagonus* (Annelida). *Journal of Biophysical and Biochemical Cytology*, vol. 7, No. 2, pp. 321-328.  
1961a. Changes in the spermatozoon during fertilization in *Hydroides hexagonus* (Annelida). II. Incorporation with the egg. *Journal of Biophysical and Biochemical Cytology*, vol. 10, No. 2, pp. 255-274.  
1961b. Fine structure of the spermatozoon of *Hydroides hexagonus* (Annelida), with special reference to the acrosomal region. *Journal of Biophysical and Biochemical Cytology*, vol. 10, No. 2, pp. 211-230.
- COLWIN, LAURA HUNTER, and ARTHUR L. COLWIN.  
1956. The acrosome filament and sperm entry in *Thyone briareus* (Holothuria) and *Asterias*. *Biological Bulletin*, vol. 110, No. 3, pp. 243-257.  
1960. Formation of sperm entry holes in the vitelline membrane of *Hydroides hexagonus* (Annelida) and evidence of their lytic origin. *Journal of Biophysical and Biochemical Cytology*, vol. 7, No. 2, pp. 315-320.



- COLWIN, LAURA HUNTER, and AUTHUR L. COLWIN.  
1961. Changes in the spermatozoon during fertilization in *Hydroides hexagonus* (Annelida). I. Passage of the acrosomal region through the vitelline membrane. *Journal of Biophysical and Biochemical Cytology*, vol. 10, No. 2, pp. 231-254.
- COKLIN, EDWIN GRANT.  
1897. The embryology of *Crepidula*. A contribution to the cell lineage and early development of some marine gasteropods. *Journal of Morphology*, vol. 13, No. 1, pp. 1-226.  
1908. The embryology of *Fulgur*: a study of the influence of yolk on development. *Proceedings of the Academy of Natural Sciences of Philadelphia*, vol. 59, 1907, pp. 320-359.
- COSTELLO, DONALD PAUL.  
1939. The volumes occupied by the formed cytoplasmic components in marine eggs. *Physiological Zoology*, vol. 12, No. 1, pp. 13-21.  
1955. Cleavage, blastulation and gastrulation. In Benjamin H. Willier, Paul A. Weiss, and Viktor Hamburger (editors), *Analysis of development*, sec. 5, Embryogenesis: preparatory phases, ch. 2, pp. 213-229. W. B. Saunders Company, Philadelphia, Pa.
- DALCO, A. M.  
1960. Germinal organisation and induction phenomena. In W. W. Nowinski (editor), *Fundamental aspects of normal and malignant growth*, ch. 4, pp. 305-494. Elsevier, Amsterdam.
- DAN, JEAN C., and SEIJI K. WADA.  
1955. Studies on the acrosome. IV. The acrosome reaction in some bivalve spermatozoa. *Biological Bulletin*, vol. 109, No. 1, pp. 40-55.
- DANIELLI, J. F.  
1958. *General cytochemical methods*. Vol. 1. Academic Press, New York, 471 pp.
- DAVAINE, C.  
1853. Recherches sur la génération des huîtres. *Comptes Rendus des Séances et Mémoires de la Société de Biologie*, tome 4, série 1, année 1852, pp. 297-339.
- DEANE, H. W., R. J. BARNETT and A. M. SELIGMAN  
1960. Histochemical methods for the demonstration of enzymatic activity. In Walther Graumann and Karlheinz Neumann (editors), *Handbuch der Histochemie*, Vol. 7, Enzymes, pt. 1, 202 pp. Gustav Fisher, Stuttgart.
- DEROBERTIS, E. D. P., W. W. NOWINSKI, and FRANCISCO A. SAEZ.  
1960. *General cytology*. 3d ed. W. B. Saunders Company, Philadelphia, Pa., 555 pp.
- DREW, GILMAN A.  
1900. *Yoldia limatula*. Memoirs from the Biological Laboratory of the Johns Hopkins University, IV, Selected morphological monographs, 3. The Johns Hopkins Press, Baltimore, Md., 37 pp.  
1906. The habits, anatomy, and embryology of the giant scallop (*Pecten tenuicostatus* Mighels). *The University of Maine Studies*, Orono, Maine, No. 6, 71 pp.
- FRANZÉN, ÅKE.  
1956. Comparative morphological investigations into the spermiogenesis among mollusca. *Zoologiska Bidrag Från Uppsala*, Band 30, pp. 399-456.
- FUJITA, TSUNENOBU.  
1929. On the early development of the common Japanese oyster. *Japanese Journal of Zoology*, Transactions and Abstracts, vol. 2, No. 3, pp. 353-358.
- GALTSOFF, PAUL S., and D. E. PHILPOTT.  
1960. Ultrastructure of the spermatozoon of the oyster, *Crassostrea virginica*. *Journal of Ultrastructure Research*, vol. 3, No. 3, pp. 241-253.
- GATENBY, J. BRONTÉ.  
1919. The cytoplasmic inclusions of the germ-cells. Part V. The gametogenesis and early development of *Limnaea stagnalis* (L.), with special reference to the Golgi apparatus and the mitochondria. *Quarterly Journal of Microscopical Science*, vol. 63, No. 252, pp. 445-491.
- GATENBY, J. BRONTÉ, and J. H. WOODGER.  
1920. On the relationship between the formation of yolk and the mitochondria and Golgi apparatus during oögenesis. *Journal of the Royal Microscopical Society for the year 1920*, pp. 129-156.
- GOMORI, GEORGE.  
1952. *Microscopic histochemistry—principles and practice*. University of Chicago Press, Chicago, Ill., 273 pp.
- HAGSTRÖM, B. E.  
1956a. Studies on the fertilization of jelly-free sea urchin eggs. *Experimental Cell Research*, vol. 10, No. 1, pp. 24-28.  
1956b. The effect of removal of the jelly coat on fertilization in sea urchins. *Experimental Cell Research*, vol. 10, No. 3, pp. 740-743.  
1956c. The influence of the jelly coat in situ and in solution on cross fertilization in sea urchins. *Experimental Cell Research*, vol. 11, No. 2, pp. 306-316.  
1956d. Further studies on cross fertilization in sea urchins. *Experimental Cell Research*, vol. 11, No. 2, pp. 507-510.
- HARVEY, ETHEL BROWNE.  
1956. *The American Arbacia and other sea urchins*. Princeton University Press, Princeton, N.J., 298 pp.
- HATSCHKE, B.  
1881. Ueber Entwicklungsgeschichte von *Teredo*. *Arbeiten aus dem Zoologischen Institute der Universität Wien und der Zoologischen Station in Triest*, tome 3, pp. 1-44.
- HORI, JUZO.  
1933. On the development of the Olympia oyster, *Ostrea lurida* Carpenter, transplanted from United States to Japan. *Bulletin of the Japanese Society of Scientific Fisheries*, vol. 1, No. 6, pp. 269-276.
- HORST, R.  
1882. On the development of the European oyster (*Ostrea edulis* L.). *Quarterly Journal of Microscopical Science*, vol. 22, pp. 341-346.

- IMMERS, J., and E. VASSEUR.  
1949. Comparative studies on the coagulation process with heparin and sea-urchin fertilizin. *Experimentia*, vol. 5, fascicule 3, pp. 124-125.
- JACKSON, ROBERT TRACY.  
1888. The development of the oyster with remarks on allied genera. *Proceedings of the Boston Society of Natural History*, vol. 23, pp. 531-556.  
1890. Phylogeny of the Pelecypoda. The Aviculidae and their allies. *Memoirs of the Boston Society of Natural History*, vol. 4, No. 8, pp. 277-400.
- KELLY, JOHN W.  
1954. Metachromasy in the eggs of fifteen lower animals. *Protoplasma*, Band 43, pp. 329-346.  
1956. The metachromatic reaction. *Protoplasmatologia*, Handbuch der Protoplasmaforschung, Band 2, D 2, 98 pp.
- KOBAYASHI, HARUO.  
1959. Bipolarity of egg of oyster (*Gryphaea gigas* (Thunberg)). *Cytologia*, vol. 24, No. 2, pp. 237-243.  
1960. Cytochemical studies on the second gradient axis of fertilized oyster egg. *Japanese Journal of Zoology*, vol. 12, No. 4, pp. 497-506.
- KORSCHULT, E., and K. HEIDER.  
1895. Text-book of the embryology of invertebrates. Part 1. Porifera, Cnidaria, Ctenophora, Vermes, Enteropneusta, Echinodermata. MacMillan and Company, New York, 484 pp. Translated from the German by Edward L. Mark and W. McM. Woodworth.
- KOSTANECKI, K.  
1904. Cytologische Studien an künstlich parthenogenetisch sich entwickelnden Eiern von *Mactra*. *Archiv für mikroskopische Anatomie und Entwicklungsgeschichte*, Band 64, pp. 1-98.  
1908. Zur Morphologie der künstlichen parthenogenetischen Entwicklung bei *Mactra*. Zugleich ein Beitrag zur Kenntnis der veilpoligen mitose. *Archiv für mikroskopische Anatomie und Entwicklungsgeschichte*. Band 72, pp. 327-352.
- LENHOSSÉK, M. VON.  
1898. Untersuchungen über Spermatogenese. *Archiv für mikroskopische Anatomie und Entwicklungsgeschichte*, Band 51, pp. 215-318.
- LILLIE, FRANK R.  
1895. The embryology of the Unionidae. A study in cell-lineage. *Journal of Morphology*, vol. 10, No. 1, pp. 1-100.  
1919. *Problems of fertilization*. University of Chicago Press, Chicago, Ill., 278 pp.
- MACBRIDE, ERNEST WILLIAM.  
1914. Invertebrata. In Walter Heape (editor), *Text-book of embryology*, vol. 1, 692 pp. Macmillan and Company, London.
- MEAD, A. D.  
1897. The early development of marine annelids. *Journal of Morphology*, vol. 13, No. 2, pp. 227-326.
- MEISENHEIMER, JOHANNES.  
1897. Entwicklungsgeschichte von *Limax maximus* L. Theil I. Furchung und Keimblätterbildung. *Zeitschrift für wissenschaftliche Zoologie*, Band 69, pp. 417-428.
- 1901a. Entwicklungsgeschichte von *Dreissensia polymorpha* Pall. *Zeitschrift für wissenschaftliche Zoologie*, Band 69, pp. 1-137.  
1901b. Die Entwicklung von Herz, Perikard, Niere und Genitalzellen bei *Cyclas* im Verhältnis zu den übrigen Mollusken. *Zeitschrift für wissenschaftliche Zoologie*, Band 69, pp. 417-428.
- MULNARD, JACQUES.  
1958. La métachromasie in vivo et son analyse cytochimique dans l'oeuf de l'Annélide *Chaetopterus pergamentaceus*. *Archives de Biologie*, tome 69, pp. 645-685.
- MULNARD, JACQUES, WALTER AUCLAIR, and DOUGLAS MARSLAND.  
1959. Metachromasia observed in the living eggs of *Arbacia punctulata* and its cytochemical analysis. *Journal of Embryology and Experimental Morphology*, vol. 7, part 2, pp. 223-240.
- NOVIKOFF, ALEX B.  
1961a. Mitochondria (Chondriosomes). In Jean Brachet and Alfred E. Mirsky (editors), *The cell*, vol. 2, ch. 5, pp. 299-421. Academic Press, New York.  
1961b. Lysosomes and related particles. In Jean Brachet and Alfred E. Mirsky (editors), *The cell*, vol. 2, ch. 6, pp. 423-488. Academic Press, New York.
- PASTEELS, J. J., and J. MULNARD.  
1957. La Métachromasie in vivo au bleu de toluidine et son analyse cytochimique dans les oeufs de *Barnea candida*, *Gryphaea angulata* (Lamellibranches) et de *Psammecchinus miliaris*. *Archives de Biologie*, tome 68, pp. 115-163.
- PHILPOTT, DELBERT E.  
1955. A simple and economical microtome for ultrathin sectioning. *Experimental Medicine and Surgery*, vol. 13, No. 2, pp. 189-192.
- RAI, HARDIT SINGH.  
1930. On the origin of yolk in the egg of *Ostrea cucullata*. *Journal of the Royal Microscopical Society*, vol. 50, series 3, pp. 210-217.
- RAVEN, CHR. P.  
1958. *Morphogenesis: the analysis of molluscan development*. Pergamon Press, New York, 311 pp.
- REBHUN, LIONEL I.  
1960. Aster-associated particles in the cleavage of marine invertebrate eggs. *Annals of the New York Academy of Science*, vol. 90, art. 2, pp. 357-380.
- RETZIUS, GUSTAF.  
1904. Zur Kenntniss der Spermien der Evertebraten. *Biologische Untersuchungen*, Neue Folge, 11, pp. 1-32. Gustav Fischer, Jena.
- ROBERT, A.  
1902. Recherches sur le développement des troques. *Archives de Zoologie Expérimentale et Générale*, serie 3, tome 10, pp. 269-538.
- RUNNSTRÖM, JOHN.  
1952. The cell surface in relation to fertilization. *Symposia of the Society for Experimental Biology*, number 6, Structural aspects of cell physiology. Academic Press, New York, pp. 39-88.

- RUNNSTRÖM, J., B. E. HAGSTRÖM, and P. PERLMANN.  
1959. Fertilization. In Jean Brachet and Alfred E. Mirsky (editors), *The cell*, vol. 1, ch. 9, pp. 327-397. Academic Press, New York.
- SENŌ, HIDEKI.  
1929. A contribution to the knowledge of the development of *Ostrea denselamellosa* Lischke. *Journal of the Imperial Fisheries Institute*, vol. 24, No. 5, pp. 129-135.
- SENŌ, HIDEKI, JUZŌ HORI, and DAIJIRŌ KUSAKABE.  
1926. Effects of temperature and salinity on the development of the eggs of the common Japanese oyster, *Ostrea gigas* Thunberg. *Journal of the Imperial Fisheries Institute*, vol. 22, No. 2, pp. 41-47.
- SHAVER, JOHN R.  
1957. Some observations on cytoplasmic particles in early echinoderm development. In Albert Tyler, R. C. von Borstel, and Charles B. Metz (editors), *The beginnings of embryonic development*, pp. 263-290. American Association for the Advancement of Science, Publication No. 48, Washington, D.C.
- STAFFORD, J.  
1912. Supplementary observations on the development of the Canadian oyster. *American Naturalist*, vol. 46, No. 541, pp. 29-40.  
1913. The Canadian oyster, its development, environment and culture. Committee on Fisheries, Game and Fur-bearing Animals, Commission of Conservation, Canada. The Mortimer Company, Ltd., Ottawa, Canada, 159 pp.
- SWANSON, CARL P.  
1957. *Cytology and cytogenetics*. Prentice-Hall Company, Englewood Cliffs, N.J., 596 pp.
- TERAO, ARATA.  
1927. On the fertilizin of the oyster, *Ostrea circumpecta*. *Science Reports of the Tôhoku Imperial University*, series 4, Biology, vol. 2, No. 2, pp. 127-132.
- THOMAS, LYEELL J., JR.  
1951. A blood anti-coagulant from surf clams. *Biological Bulletin*, vol. 101, No. 2, pp. 230-231.  
1954. The localization of heparin-like blood anti-coagulant substances in the tissues of *Spisula solidissima*. *Biological Bulletin*, vol. 106, No. 1, pp. 129-138.
- TYLER, ALBERT.  
1939. Extraction of an egg membrane-lysin from sperm of the giant keyhole limpet (*Megathura crenulata*). *Proceedings of the National Academy of Sciences of the United States*, vol. 25, pp. 317-323.  
1948. Fertilization and immunity. *Physiological Reviews*, vol. 28, No. 2, pp. 180-219.  
1949. Properties of fertilizin and related substances of eggs and sperm of marine animals. *American Naturalist*, vol. 83, No. 11, pp. 195-219.
- VASSEUR, ERIK.  
1948a. The sulphuric acid content of the egg coat of the sea urchin, *Strongylocentrotus droebachiensis* Müll. *Arkiv för Kemi, Mineralogi och Geologi*, Band 25, B, No. 6, pp. 1-2.  
1948b. Chemical studies on the jelly coat of the sea-urchin egg. *Acta Chemica Scandinavica*, vol. 2, pp. 900-913.
- WADA, S. K., J. R. COLLIER, and J. C. DAN.  
1956. Studies on the acrosome. V. An egg-membrane lysin from the acrosomes of *Mytilus edulis* spermatozoa. *Experimental Cell Research*, vol. 10, No. 1, pp. 168-180.
- WHITMAN, CHARLES OTIS.  
1878. The embryology of *Clepsine*. *Quarterly Journal of Microscopical Science*, vol. 18, new series, No. 71, *Memoirs*, pp. 215-314.
- WILLIER, BENJAMIN, PAUL A. WEISS, and VIKTOR HAMBURGER.  
1955. *Analysis of development*. W. B. Saunders Company, Philadelphia, Pa., 735 pp.
- WILSON, EDMUND B.  
1892. The cell-lineage of *Nereis*. A contribution to the cytogeny of the annelid body. *Journal of Morphology*, vol. 6, No. 3, pp. 361-480.  
1904a. Experimental studies on germinal localization. I. The germ-regions in the egg of *Dentalium*. *Journal of Experimental Zoölogy*, vol. 1, No. 1, pp. 1-72.  
1904b. Experimental studies on germinal localization. II. Experiments on the cleavage-mosaic in *Patella* and *Dentalium*. *Journal of Experimental Zoölogy*, vol. 1, No. 2, pp. 197-268.  
1928. *The cell in development and heredity*. 3d ed. Macmillan Company, New York, 1232 pp.
- WOODS, FARRIS H.  
1932. Keimbahn determinants and continuity of the germ cells in *Sphaerium striatinum* (Lam.). *Journal of Morphology*, vol. 53, No. 2, pp. 345-365.
- WORLEY, LEONARD G.  
1944. Studies of the vitally stained Golgi apparatus. II. Yolk formation and pigment concentration in the mussel *Mytilus californianus* Conrad. *Journal of Morphology*, vol. 75, No. 1, pp. 77-101.
- YASUGI, RYUICHI.  
1938. On the mode of cleavage of the eggs of the oysters *Ostrea spinosa* and *O. gigas* under experimental conditions (a preliminary note). *Annotationes Zoologicae Japonenses*, vol. 17, pp. 295-300.
- ZIEGLER, H. ERNST.  
1885. Die Entwicklung von *Cyclas cornea* Lam. (*Sphaerium corneum* L.). *Zeitschrift für wissenschaftliche Zoologie*, Band 41, pp. 525-569.