Do not use if printed foil seal under bottle cap is open or torn.

Active ingredient (in each gelcap) **Purpose**
Famotidine 10 mg ....Acid reducer

Uses • relieves heartburn associated with acid indigestion and sour stomach
• prevents heartburn associated with acid indigestion and sour stomach
brought on by eating or drinking certain food and beverages

Warnings **Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers

LOT

EXP.

XXXXXXX

Do not use
• if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
• with other acid reducers

ORIGINAL STRENGTH

NDC 16837-856-61

Pepcid.®
Famotidine Tablets 10mg   *Acid Reducer*
AC
Just One Tablet

Prevents & Relieves Heartburn
Due to Acid Indigestion

60 GELCAPS
(GELATIN COATED, CAPSULE SHAPED TABLETS)

Ask a doctor before use if you have • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with **lightheadedness, sweating, or dizziness** • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness • frequent **chest pain** • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain. **Stop use and ask a doctor if** • your heartburn continues or worsens • you need to take this product for more than 14 days. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions** • adults and children 12 years and over: • to **relieve** symptoms, swallow 1 gelcap with a glass of water. Do not chew. • to **prevent** symptoms, swallow 1 gelcap with a glass of water at any time from **15 to 60 minutes before** eating food or drinking beverages that cause heartburn • do not use more than 2 gelcaps in 24 hours • children under 12 years: ask a doctor

Dist. by: *Johnson&Johnson* • MERCK
CONSUMER PHARMACEUTICALS CO., FORT WASHINGTON, PA 19034 USA
®Registered trademark of Merck & Co., Inc.

150% OF SIZE

Labeling Format Information:

Fonts: Universe Cond regular, bold and bold italic.

Drug Facts:	NA	**Leading**:	4.5 pt
Header:	NA	**Bullets**:	4.5 pt
Subheader:	4.5 pt	**Barlines**:	NA
Body Text:	4.5 pt	**Hairlines**:	NA
Drug Facts (continued):	NA		

Horizontal Scale: 75-80%	Average Kerning: -15

LABEL PEPCID GC 60 (CLASS LABELING)   **3004920 C-2**
OCTOBER 12, 2004

Do not use if printed foil seal under bottle cap is open or torn.

WHAT YOU SHOULD KNOW ABOUT:

Pepcid® AC
Famotidine Tablets 10mg/ Acid Reducer
Gelcaps

PACKAGE INSERT

A non-prescription stomach medicine

PEPCID AC Gelcaps contain a medicine, different from antacids, that doctors have prescribed for years to treat acid-related problems in millions of people.

Product Benefits:

Shown actual size

- **1 gelcap** relieves heartburn due to acid indigestion.
- Pepcid AC prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods or beverages.
- It contains famotidine, a prescription-proven medicine.

How to use PEPCID AC Gelcaps

Adults and children 12 years and over:

Use PEPCID AC Gelcaps to relieve or prevent heartburn due to acid indigestion and sour stomach.

TO **RELIEVE** SYMPTOMS
Swallow 1 gelcap with a glass of water (do not chew).

TO **PREVENT** SYMPTOMS
Swallow 1 gelcap with a glass of water at any time from **15 to 60 minutes before** eating food or drinking beverages that cause heartburn.

PEPCID AC can be used up to twice daily (up to 2 gelcaps in 24 hours).

Excess acid: a burning problem

It is normal for the stomach to produce acid, especially after consuming food and beverages. However, acid in the wrong place (the esophagus), or too much acid, can cause burning pain and discomfort that interfere with everyday activities.

Heartburn— Caused by acid in the esophagus

A valve-like muscle called the lower esophageal sphincter (LES) is relaxed in an open position

Burning pain/ discomfort

Excess acid moves up into esophagus

Tips for Managing Heartburn
- Do not lie flat or bend over soon after eating.
- Do not eat late at night, or just before bedtime.
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, and even some fruits and vegetables.
- Eat slowly and do not eat big meals.
- If you are overweight, lose weight.
- If you smoke, quit smoking.
- Raise the head of your bed.
- Wear loose fitting clothing around your stomach.

Excellent safety record

The ingredient in PEPCID AC, famotidine, has been prescribed by doctors for years to treat millions of patients safely and effectively. The active ingredient in PEPCID AC has been taken safely with many frequently prescribed medications.

Know when to see your doctor

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use
- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools.
These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Stop use and ask a doctor if
- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Questions or comments?

For more information and a guide of how to help your heartburn, please call 1-800-755-4008 (English) or 1-888-466-8746 (Spanish), www.pepcidac.com

Distributed by:
Johnson&Johnson • MERCK
CONSUMER PHARMACEUTICALS CO.
FORT WASHINGTON, PA 19034 USA
® Registered trademark of Merck & Co., Inc.
XXXXXXX

VALUABLE MAIL-IN OFFER

Save $3.00 on any Pepcid® AC® or Pepcid® Complete® product with purchase of 25 count or larger

Here's how you can get $3.00 off your next purchase:
BUY: Any Pepcid AC or Pepcid Complete Package 25 count or Larger.
SEND: This completed certificate, UPC from the Pepcid AC or Pepcid Complete package, plus dated cash register receipt with purchase price circled.
RECEIVE: $3.00 Coupon by mail.
SEND TO: Pepcid AC / Pepcid Complete $3 Offer 780
P.O. Box 6607
Douglas, AZ 85655-6607
XXXXXXX
NOTE: Offer good only in U.S.A. and APO/FPO addresses. This request form may not be mechanically reproduced. LIMIT TWO OFFERS PER FAMILY AND/OR ADDRESS, PER YEAR. No group or organization requests will be honored. Your offer rights may not be transferred or assigned. Offer void where prohibited or taxed. You should expect to receive your coupon in 6-8 weeks. Cannot be combined with any other offers.

☐ Yes, I'd like to receive future offers and information from McNeil Consumer & Specialty Pharmaceuticals and McNeil Nutritionals, Divisions of McNeil-PPC, Inc. and Johnson & Johnson • Merck Consumer Pharmaceuticals Co.
℗ Johnson & Johnson • Merck 2004

Please circle one: Dr. Mr. Mrs. Miss.
Name (please print)
Address   Apt.
City   State   Zip
E-mail
Age: ☐ under 35 ☐ 35-49 ☐ 50-54 ☐ 55-64 ☐ 65 yrs+

Labeling Format Information:

Fonts: Universe regular and bold

Drug Facts:	NA	**Leading:**	6.5 pt
Header:	NA	**Bullets:**	5 pt
Subheader:	6 pt	**Barlines:**	NA
Body Text:	6 pt	**Hairlines:**	NA
Drug Facts (continued):	NA		

Horizontal Scale: 100%	Average Kerning: 0

INSERT PEPCID GC 60 BOTTLE CODES (CLASS LABELING)   3004920 D-2
OCTOBER 1, 2004