

Do not use if printed foil seal under bottle cap is open or torn.

Active ingredient (in each tablet) **Purpose**
Famotidine 10 mg Acid reducer

Uses • relieves heartburn associated with acid indigestion and sour stomach
• prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings **Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers

LOT
[]

EXP.
[]

XXXXXXXXXX

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools.

These may be signs of a serious condition. See your doctor.

- with other acid reducers

ORIGINAL STRENGTH

NDC 16837-872-60



Pepcid[®] AC

Famotidine Tablets 10mg Acid Reducer

Just One Tablet

Prevents & Relieves Heartburn Due to Acid Indigestion

60 TABLETS

Ask a doctor before use if you have • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with **lightheadedness, sweating, or dizziness** • chest pain or shoulder pain with shortness of breath, sweating; pain spreading to arms, neck or shoulders; or lightheadedness

- frequent **chest pain** • frequent wheezing, particularly with heartburn
- unexplained weight loss • nausea or vomiting • stomach pain. **Stop use and ask a doctor if** • your heartburn continues or worsens • you need to take this product for more than 14 days. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions**
- adults and children 12 years and over: • **to relieve symptoms**, swallow 1 tablet with a glass of water. Do not chew. • **to prevent symptoms**, swallow 1 tablet with a glass of water at any time from **15 to 60 minutes before** eating food or drinking beverages that cause heartburn • do not use more than 2 tablets in 24 hours • children under 12 years: ask a doctor

Dist. by: **Johnson & Johnson** • MERCK
CONSUMER PHARMACEUTICALS CO., FORT WASHINGTON, PA 19034 USA
©Registered trademark of Merck & Co., Inc.

150% OF SIZE

Do not use if printed foil seal under bottle cap is open or torn.

Active ingredient (in each tablet) **Purpose**
Famotidine 10 mg Acid reducer

Uses • relieves heartburn associated with acid indigestion and sour stomach
• prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings **Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers

LOT
[]

EXP.
[]

XXXXXXXXXX

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools.

These may be signs of a serious condition. See your doctor.

- with other acid reducers

ORIGINAL STRENGTH

NDC 16837-872-60



Pepcid[®] AC

Famotidine Tablets 10mg Acid Reducer

Just One Tablet

Prevents & Relieves Heartburn Due to Acid Indigestion

60 TABLETS


Ask a doctor before use if you have • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with **lightheadedness, sweating, or dizziness** • chest pain or shoulder pain with shortness of breath, sweating; pain spreading to arms, neck or shoulders; or lightheadedness

- frequent **chest pain** • frequent wheezing, particularly with heartburn
- unexplained weight loss • nausea or vomiting • stomach pain. **Stop use and ask a doctor if** • your heartburn continues or worsens • you need to take this product for more than 14 days. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions**
- adults and children 12 years and over: • **to relieve symptoms**, swallow 1 tablet with a glass of water. Do not chew. • **to prevent symptoms**, swallow 1 tablet with a glass of water at any time from **15 to 60 minutes before** eating food or drinking beverages that cause heartburn • do not use more than 2 tablets in 24 hours • children under 12 years: ask a doctor

Dist. by: **Johnson & Johnson** • MERCK
CONSUMER PHARMACEUTICALS CO., FORT WASHINGTON, PA 19034 USA
©Registered trademark of Merck & Co., Inc.

Labeling Format Information:			
Fonts: Universe Cond regular, bold and bold italic.			
Drug Facts:	NA	Leading:	4.5 pt
Header:	NA	Bullets:	4.5 pt
Subheader:	4.5 pt	Barlines:	NA
Body Text:	4.5 pt	Hairlines:	NA
Drug Facts (continued):	NA		
Horizontal Scale: 75-80%		Average Kerning: -15	

LABEL PEPCID FCT 60 (CLASS LABELING) 3004907 C-4
OCTOBER 12, 2004



Do not use if printed foil seal under bottle cap is open or torn.

WHAT YOU SHOULD KNOW ABOUT:

Pepcid[®] AC
Famotidine Tablets 10mg/
Acid Reducer

PACKAGE INSERT
Tablets

A non-prescription stomach medicine

PEPCID AC contains a medicine, different from antacids, that doctors have prescribed for years to treat acid-related problems in millions of people.

- Product Benefits:**
- 1 tablet relieves heartburn due to acid indigestion.
 - Pepcid AC prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods or beverages.
 - It contains famotidine, a prescription-proven medicine.



Shown actual size

How to use PEPCID AC

Use PEPCID AC to relieve or prevent these symptoms:

- Heartburn
- Acid indigestion
- Sour stomach



TO RELIEVE SYMPTOMS

Swallow 1 tablet with a glass of water (do not chew).

PEPCID AC can be used up to twice daily (up to 2 tablets in 24 hours).



TO PREVENT SYMPTOMS


Swallow 1 tablet with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn.

Excess acid: a burning problem

It is normal for the stomach to produce acid, especially after consuming food and beverages. However, acid in the wrong place (the esophagus), or too much acid, can cause burning pain and discomfort that interfere with everyday activities.

Heartburn— Caused by acid in the esophagus

A valve-like muscle called the lower esophageal sphincter (LES) is relaxed in an open position



Tips for Managing Heartburn

- Do not lie flat or bend over soon after eating.
- Do not eat late at night, or just before bedtime.
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, and even some fruits and vegetables.
- Eat slowly and do not eat big meals.
- If you are overweight, lose weight.
- If you smoke, quit smoking.
- Raise the head of your bed.
- Wear loose fitting clothing around your stomach.

Age: under 35 35-49 50-54 55-64 65 yrs+

E-mail: _____
City: _____ State: _____ Zip: _____ Apt: _____
Name (please print): _____
Please circle one: Dr. Mr. Mrs. Miss.

Yes, I'd like to receive future offers and information from McNeil-PPC, Inc. and Johnson & Johnson Consumer & Specialty Pharmaceuticals and McNeil Nutritional, Divisions of McNeil-PPC, Inc. and Johnson & Johnson Consumer & Specialty Pharmaceuticals Co.

© Johnson & Johnson • Merck 2004
P.O. Box 6607
Douglas, AZ 85555-6607

SEND TO: Pepcid AC / Pepcid Complete \$3 Offer 780
FAMILY AND/OR ADDRESS, PER YEAR. No group or organization requests will be honored. Your rights may not be transferred or assigned. Offer void where prohibited or taxed. You should expect to receive your coupon in 8-9 weeks. Cannot be combined with any other offers.

NOTE: Offer good only in U.S.A. and APO/FPO addresses. This request form may not be mechanically reproduced. LIMIT TWO OFFERS PER

RECEIVE: \$3.00 Coupon by mail.
SEND: This completed certificate, UPC from the Pepcid AC or Pepcid Complete Package 25 count or Larger.
BUY: Any Pepcid AC or Pepcid Complete Package 25 count or Larger.
SEND: \$3.00 Coupon with purchase of Pepcid AC or Pepcid Complete Package, plus dated

Here's how you can get \$3.00 off your next purchase:
product with purchase of 25 count or larger
Save \$3.00 on any Pepcid[®] AC or Pepcid Complete[®]

VALUABLE MAIL-IN OFFER

© Registered trademark of Merck & Co., Inc.
FAMOTIDINE PHARMACEUTICALS CO.
FORT WASHINGTON, PA 19034 USA

1-888-466-8746 (Spanish), www.pepcidac.com
or 1-800-755-4008 (English)
Distributed by: **Johnson & Johnson • MERCK**

For more information and a guide of how to help your heartburn, please call 1-800-755-4008 (English) or 1-888-466-8746 (Spanish), www.pepcidac.com

Questions or comments?

Center right away.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
If pregnant or breast-feeding, ask a health professional before use.
• You need to take this product for more than 14 days
• Your heartburn continues or worsens
• Stop use and ask a doctor if:
• stomach pain
• nausea or vomiting
• unexplained weight loss
• frequent wheezing, particularly with heartburn
• frequent chest pain
• shoulders, or lightheadedness
• chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or heartburn with lightheadedness, sweating, or dizziness
• had heartburn over 3 months. This may be a sign of a more serious condition.
Ask a doctor before use if you have
• with other acid reducers
• if you have signs of a serious condition. See your doctor.
• if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools.

Do not use
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Know when to see your doctor

The ingredient in PEPCID AC, famotidine, has been prescribed by doctors for years to treat millions of patients safely and effectively. The active ingredient in PEPCID AC has been taken safely with many frequently prescribed medications.

Excellent safety record

Labeling Format Information:

Fonts:	Universe regular and bold	
Drug Facts:	NA	Leading: 6.5 pt
Header:	NA	Bullets: 5 pt
Subheader:	6 pt	Barlines: NA
Body Text:	6 pt	Hairlines: NA
Drug Facts (continued):	NA	
Horizontal Scale: 100%		Average Kerning: 0

INSERT PEPCID FCT BOTTLE CODES (CLASS LABELING) 3004907 D-3
OCTOBER 4, 2004