

# Nitropyrene: DNA Binding and Adduct Formation in Respiratory Tissues

by M. A. Jackson,\* L. C. King,† L. M. Ball,‡  
S. Ghayourmanesh,\*\* A. M. Jeffrey,\*\* and J. Lewtas†§

Binding of 1-nitro (<sup>14</sup>C)pyrene (NP) or its metabolites to cellular DNA and protein in cultures of rabbit alveolar macrophages, lung tissue, and tracheal tissue was examined. DNA binding in tracheal tissue (136 ± 18.3 pmole NP/mg DNA) was four to five times the levels measured in either lung tissue (38 ± 9.4 pmole NP/mg DNA) or macrophages (26 ± 7.5 pmole NP/mg DNA). Adduct analysis of DNA isolated from lung tissue incubated with 1-nitro[<sup>3</sup>H]pyrene *in vitro* resulted in the identification of 2 to 5% of the NP adducts as C8-deoxyguanosine 1-aminopyrene. NP was also bound to cellular protein in tracheal tissue and lung tissue, and at a lower level in macrophages. Cocultivation of the macrophages with lung and tracheal tissue decreased the DNA binding in tracheal tissue by 45%. Following intratracheal instillation of diesel particles (5 mg) vapor-coated with <sup>14</sup>C-NP (380 ppm, 0.085 μCi/mg) particles into rats, 5-8% of the radioactivity remained in the lungs after 20 hr. Most of the diesel particles were also deposited in the lung. Examination of DNA and protein binding in this tissue showed 5 to 12% of the pulmonary <sup>14</sup>C bound to protein and no detectable levels of <sup>14</sup>C bound to DNA.

## Introduction

Polycyclic aromatic hydrocarbons (PAHs) which pollute our atmosphere have been associated with the etiology of human cancer (1). Nitro-substituted PAHs (NO<sub>2</sub>-PAH) have been found in extracts from diesel particles (2-5) and ambient air particulates (6). Many of these NO<sub>2</sub>-PAH are mutagenic in mammalian cells (7-10) as well as in bacteria (8-12) and may, therefore, be a potential risk to human health.

Among these NO<sub>2</sub>-PAH, 1-nitropyrene (1-NP) has been found to be a potent bacterial mutagen (12) and potentially an animal carcinogen (13). This compound has been identified and quantitated in diesel particle extracts (3,4) and may be responsible for up to 30% of the mutagenicity of these extracts (4,14). The bacterial mutagenicity of 1-NP is believed to result from reduction to reactive electrophiles which bind to DNA (12,15).

Howard and co-workers (1) have identified one DNA adduct formed by *Salmonella typhimurium* from the reduction of 1-NP as an *N*-(deoxyguanosin-8-yl)-1-aminopyrene, i.e., C8-deoxyguanosine-1-aminopyrene adduct (C8-dG-AP). They have also shown that some mammalian enzymes are capable of reducing 1-NP to

form this same adduct. In addition, mammalian tissues are capable of oxidizing 1-NP both *in vivo* (16) and *in vitro* (17,18) to metabolites that are themselves mutagenic. We have shown in an earlier study (19) that rabbit pulmonary macrophages and tissues activated 1-NP to DNA-bound adducts.

In this work we evaluate DNA adducts resulting from the metabolism of 1-NP by mammalian cells and tissues to see whether and to what extent oxidative metabolism of 1-NP forms the C8-dG-AP adduct previously characterized as a product of reductive metabolism. Respiratory tissue has been selected for this evaluation because of potential for exposure to 1-NP associated with diesel particles. *In vivo* studies have shown that inhaled diesel particles are readily deposited in the respiratory tract (20,21) in particular, in the alveolar macrophages (21,22). Although many of the particles are rapidly cleared from the lung, a substantial portion remain in the lung for a long period (20,22), thus increasing exposure to compounds associated with the particles. It has also been shown that both 1-NP and mutagenic activity were lost from diesel particles after incubation with pulmonary alveolar macrophages *in vitro* (23). We have now used diesel particles vapor-coated with <sup>14</sup>C-labeled 1-NP to study *in vivo* the fate of 1-NP absorbed onto diesel particles and evaluate interactions with pulmonary macromolecules.

## Materials and Methods

### Chemicals

1-Nitro[4,5,9,10-<sup>14</sup>C]pyrene (<sup>14</sup>C-1NP; 60 mCi/mmol), 1-nitro[<sup>3</sup>H]pyrene (<sup>3</sup>H-1NP, 6 Ci/mmol) and unlabeled

\*Environmental Health Research and Testing, P. O. Box 12199, Research Triangle Park, NC 27709.

†Genetic Toxicology Division, HERTL, U. S. Environmental Protection Agency, Research Triangle Park, NC 27711.

‡Department of Environmental Sciences and Engineering, University of North Carolina, Chapel Hill, NC 27514.

\*\*Columbia University Health Science Center, New York, NY 10032.

§Author to whom correspondence and reprint requests should be addressed.

1-nitropyrene (1-NP) were synthesized by Midwest Research Institute (Kansas City, MO). Unlabeled 1-NP was added to  $^{14}\text{C}$ -1NP to adjust the specific activity to 10 mCi/mmol. Diesel particles vapor-coated with  $^{14}\text{C}$ -1NP were prepared at Battelle Columbus Laboratories (Columbus, OH), by vaporizing  $^{14}\text{C}$ -1NP at 275°C onto a stream of aerosolized diesel particles. The recovered particles contained 380 ppm nitropyrene and 195,000 dpm/mg particle. All chemicals used in the isolation and preparation of DNA for adduct analysis were purchased as follows: chloroform (spectro grade) from Burdick and Jackson (Muskegon, MI); isoamyl alcohol from Fisher Scientific (Raleigh, NC); phenol (99+%) from Aldrich Chemical Co. (Milwaukee, WI); and enzymes for DNA hydrolysis from Sigma Chemical Co. (St. Louis, MO) and Worthington Biochemical Corp. (Freehold, NJ). All other materials were purchased at the highest available grade of purity.

### **In Vitro Tissue Preparation and Incubation**

Rabbit alveolar macrophages, tracheas and lung tissue were obtained from New Zealand White male rabbits (1.5–2.0 kg; Dutchland Laboratory Animals, Denver, PA) following lung lavage (24) and perfusion as described previously (19).

Macrophages ( $0.5 \times 10^6$  cells/mL), lung tissue ( $31.0 \pm 4.0$  mg/mL), or tracheal tissue ( $14.0 \pm 0.4$  mg/mL) were each incubated separately with  $^{14}\text{C}$ -1NP ( $2 \mu\text{g/mL}$ ) at 37°C in a humidified atmosphere of 5%  $\text{CO}_2$  and air for 20 hr. The culture medium (M-199) contained 10% heat-deactivated fetal calf serum (FCS), penicillin (100 units/mL), streptomycin (100  $\mu\text{g/mL}$ ), kanamycin (100  $\mu\text{g/mL}$ ), and  $^{14}\text{C}$ -1NP (8.1  $\mu\text{M}$ ; 81.0 nCi/mL) dissolved in DMSO (final concentration 0.5%, Burdick and Jackson Laboratories, Inc., Muskegon, MI). Macrophages ( $0.5 \times 10^6$  cells/mL), lung tissue ( $29.0 \pm 2.0$  mg/mL), and tracheal tissue ( $14.0 \pm 1.0$  mg/mL) were also cocultured under the same exposure conditions. Each experiment was repeated three times.

After incubation, the attached macrophages were detached by trypsinization and recovered by centrifugation at 300g, then lysed with 25 mL lysing buffer (50 mM Tris-HCl, 10 mM EDTA, 1% SDS, pH 8.0). The lung or tracheal tissues were separated from the culture medium by filtration through nylon mesh, then homogenized in 20 mL of the lysing buffer. A full description of these procedures has been given by King et al. (19).

Lung tissue ( $29.0 \pm 2.0 \mu\text{g/mL}$ ) was also incubated with  $^3\text{H}$ -1NP ( $2 \mu\text{g/mL}$ ) as described above to provide modified DNA of higher specific radioactivity for identification of the adducts formed.

### **In Vivo Studies**

Male rats (300 g; Sprague-Dawley, Charles River CD-1, Wilmington, MO) received  $^{14}\text{C}$ -NP-coated diesel particles by intratracheal instillation of 5 mg of particles in

0.2 mL phosphate-buffered saline with 1% Tween 80 under light halothane anesthesia. The animals were maintained individually in metabolism cages (Nalgene Co., Rochester, NY) for collection of urine and feces and received food and water *ad libitum*.

After 24 hr each animal was anesthetized with ether and the thoracic cavity opened to expose the lungs. The lungs were removed *en bloc* following total exsanguination of the animal. All vascular and connective tissues were removed and the lung tissue was placed into a 50-mL tube on ice. The tissue was thoroughly minced, than an aliquot (1 g) was weighed out and homogenized in 10 mL lysing buffer.

### **Quantitation of DNA and Protein Binding**

DNA and protein were isolated according to an extraction scheme previously described (25) and illustrated in Figure 1. Briefly, protein was extracted from cellular nucleic acid in the lysates or homogenates with chloroform:isoamyl alcohol:phenol (CIP, 24:1:25) and precipitated with acetone. DNA was removed from the aqueous portion by precipitation with cold ethanol. Residual protein and RNA were removed by incubation with proteinase K (1 mg/mL) and NaOH, respectively. The radioactivity associated with the DNA was determined by liquid scintillation counting after acid precipitation onto glass fiber filters. Radioactivity associated with protein was determined by liquid scintillation counting of NaOH digested protein neutralized with HCl. Total DNA content was measured by the diphenylamine colorimetric assay (26) and total protein by the Lowry method (27).

### **Identification of DNA Adducts**

The isolated DNA was redissolved in water and precipitated with ethanol in presence of sodium acetate until at least 95% of the radioactivity was precipitable. The DNA (<1 mg/mL) was dissolved in 10 mM Tris-HCl–0.1 M NaCl–5 mM  $\text{MgCl}_2$  (pH 7.9) and was treated consecutively with 200 units of DNase I for 2 hr, 5 units of alkaline phosphatase for 2 hr, 2 units of phosphodiesterase I for 4 hr, 2 units phosphodiesterase II for 2 hr, and 5 units of alkaline phosphatase for 2 hr.

The resulting deoxyribonucleosides were then separated on a Sephadex LH-20 column. After application of the sample, the column was washed with water to remove unmodified deoxyribonucleosides. The modified adducts were then eluted with methanol and the solvent evaporated under reduced pressure. The sample was dissolved in 10  $\mu\text{L}$  of methanol containing unlabeled 1-nitropyrene-modified DNA adducts [prepared according to Howard et al. (1)] and was analyzed by HPLC on a Dupont Instruments Model 850 high-pressure liquid chromatograph fitted with a  $\text{C}_{15}$ - $\mu\text{Bondapak}$  column (Waters Associates) using a concave gradient (number 2) from 50 to 100% methanol in water at 50°C and a flow rate of 1 mL/min. The eluate was monitored




FIGURE 1. DNA isolation scheme. Protein was removed from cellular nucleic acid by two extractions of the cell lysates or tissue homogenates with chloroform:isoamyl alcohol:phenol (CIP, 24:1:25). Protein in the CIP portion was precipitated with acetone and DNA in the aqueous portion was precipitated with cold ethanol. After residual protein and RNA were hydrolyzed with proteinase K and NaOH, respectively, DNA was acid precipitated onto glass fiber filters and associated radioactivity measured in a liquid scintillation spectrometer. Samples used for identification of DNA adducts were not taken through the hydrolysis procedure, but were prepared directly as described in "Materials and Methods."

at 254 nm, and 1-min fractions were collected then assayed for radioactivity by liquid scintillation counting.

## Results

Binding levels of  $^{14}\text{C}$ -1NP metabolites to DNA and protein from respiratory tissues exposed *in vivo* are presented in Figure 2. Tracheal tissue exhibited the highest binding to DNA (Fig. 2A) and protein (Fig. 2B) in both the separate 20-hr cultures of isolated macrophages and lung and tracheal tissue and the cocultured system.

DNA from lung tissue incubated for 20 hr with  $^3\text{H}$ -1NP was hydrolyzed and the resulting deoxyribonucleosides prepared for identification of modified adducts by high-pressure liquid chromatography as described in "Materials and Methods." A chromatogram from this analysis is shown in Figure 3. From 2 to 5% of the total DNA adducts co-eluted with the synthetic C8-dG-AP adduct in fraction number 60 (retention time 60 min). However, the majority (over 90%) of the modified DNA




FIGURE 2. Binding of 1-nitro $^{14}\text{C}$ pyrene metabolites to DNA and protein. Cultures of rabbit alveolar macrophages ( $0.5 \times 10^6$  cells/mL), lung tissue 31.0 mg/mL, or tracheal tissue (14.0 mg/mL) were incubated 20 hr in tissue culture media containing  $^{14}\text{C}$ -1NP (8.1  $\mu\text{M}$ ; 81.0 nCi/mL). Cellular DNA (A) and protein (B) were isolated for analysis of binding as previously described. Similarly, rabbit alveolar macrophages ( $0.5 \times 10^6$  cells/mL), lung tissue 29.0 mg/mL and tracheal tissue (14.0 mg/mL) were co-cultured 20 hr in media containing  $^{14}\text{C}$ -1NP (8.1  $\mu\text{M}$ ; 81.0 nCi/mL). The cells and tissues were separated from one another as described and prepared for analysis of DNA and protein binding. Data are presented as the mean of three experiments  $\pm$  standard error of the mean.

was not retained on the Sephadex LH-20 column and therefore was not amenable to HPLC analysis. Of the material retained on the LH20 column, 30% or more eluted with the solvent front, and the other two peaks observed (35 and 94 min) did not correspond to any synthetic standard currently available.

$^{14}\text{C}$ -1NP vapor-coated onto diesel particles was administered to rats by intratracheal instillation to investigate the binding of 1-NP to respiratory tissue *in vivo* in an attempt to more closely model realistic exposure routes. Results are presented in Figure 4. After 24 hr, 6–8% of the total  $^{14}\text{C}$  dose as well as a major portion of the particles (based on visual inspection) remained in the lungs. Analysis of DNA and protein binding in the lung tissue showed that 5 to 13% of the residual pulmonary  $^{14}\text{C}$  (representing 0.6% of the total dose) was bound to protein, and no detectable  $^{14}\text{C}$  was bound to DNA.

## Discussion

Although macrophages and respiratory tissue are known to be able to metabolize PAH to reactive inter-




FIGURE 3. Chromatogram of lung DNA adducts from [ $^3\text{H}$ ]1-nitropyrene. This chromatogram represents the HPLC profile of the modified deoxyribonucleosides obtained from the hydrolysis of DNA isolated from lung tissue exposed *in vitro* to  $^3\text{H}$ -1NP (8.1  $\mu\text{M}$ ; 48  $\mu\text{Ci}/\text{mL}$ ). The major peak (fraction 60) is identified as the C8-deoxyguanosine-1-aminopyrene adduct and represents 2–5% of the total activity bound to DNA.

mediates that bind DNA (28), the metabolism and binding of  $\text{NO}_2$ -PAH have only recently begun to receive attention. We have previously presented data that clearly show that macrophages and tissues from the respiratory tract are capable of binding 1-NP or its metabolites to DNA and protein (19). We have now identified one adduct bound to DNA from lung tissue exposed to  $^3\text{H}$ -1NP.




FIGURE 4. Disposition of  $^{14}\text{C}$  from 1-nitro[ $^{14}\text{C}$ ]pyrene vapor-coated onto diesel particles. Rats were treated with  $^{14}\text{C}$ -1NP vapor-coated onto diesel particles by intratracheal instillation with the particles. Animals were maintained for 24 hr after dosing, then the lungs were removed and analyzed for total  $^{14}\text{C}$  content. DNA and protein were isolated and  $^{14}\text{C}$  bound to these macromolecules was measured as described in "Materials and Methods."

The *N*-(deoxyguanosin-8-yl)-1-aminopyrene (C8-dG-AP) adduct, which had previously been identified as the major DNA adduct formed from 1-NP by *Salmonella typhimurium* TA 1538 and by xanthine oxidase (1), was found to represent a small portion (2–5%) of the total DNA adducts formed by lung tissue. The majority of the NP-DNA adducts either did not respond to the digestive enzyme treatments, or yielded mononucleoside products that were not hydrophobic enough to be retained on the Sephadex LH-20 column. This is in marked contrast to the xanthine oxidase-catalyzed adducts, of which more than 80% were extractable and resolved into the C8-dG-AP adduct, and to the adducts formed in TA 1538, where the C8-dG-AP adduct was the only peak detected by HPLC (1). However, the presence of even small quantities of C8-dG-AP in mixtures of NP-DNA adducts from lung tissue is significant, in that it indicates that some portion of pulmonary metabolism and activation of 1-NP is by reduction of the nitro function, thus forming a C8-dG type of adduct that appears to be a characteristic of arylamines activated by *N*-hydroxylation (1).

Analysis of binding to DNA and protein from lung tissue exposed *in vivo* to  $^{14}\text{C}$ -1NP vapor-coated onto diesel particles showed that a very small portion of the total radioactivity instilled in the lungs was bound to protein and that there was no detectable DNA binding. This is significant, however, because less than 10% of the total activity administered remained in the lung tissue even though most of the particulate matter was still present. This indicates that not only is respiratory tissue able to metabolize 1-NP, but that it is capable of removing it from diesel particles *in vivo*, as had previously been demonstrated for pulmonary alveolar macrophages *in vitro* (23). The majority of NP originally present on the diesel particles is therefore fully available for distribution throughout the body and for further metabolism and possible activation.

The information we have presented here strongly suggests that additional studies need to be conducted on the metabolism and binding of the  $\text{NO}_2$  PAH found in diesel exhaust and urban air. More information is needed for defining the genotoxic burden of  $\text{NO}_2$  PAH in ambient air. Further studies are currently being conducted in our laboratories to elucidate better the DNA and protein binding of 1-NP to tracheal epithelial cells. In addition, new DNA hydrolysis techniques are being evaluated to increase total DNA adduct recovery and identification.

Part of the research described in this article is conducted under contract to the Health Effects Institute (HEI, #83-23, with the Columbia University Health Science Center), an organization that supports the conduct of independent research and is jointly funded by the United States Environmental Protection Agency (EPA) and automotive manufacturers. Publication here implies nothing about the view of the contents by HEI of its research sponsors. HEI's Health Review Committee may comment at any time and will evaluate the final report of the project.

The work described in this document has also been funded in part by the U. S. Environmental Protection Agency under Contract 68-

02-4031 with Environmental Research and Testing, Inc., under cooperative agreement CR-810849-010 with the Department of Environmental Sciences and Engineering, University of North Carolina, and under assistance agreement X808859 with HEI. The contents, however, do not reflect the views and policies of the Agency; nor does mention of trade names or commercial products constitute endorsement or recommendations for use.

## REFERENCES

- Howard, P. C., Heflich, R. H., Evans, F. E., and Beland, F. A. Formation of DNA adducts *in vitro* and in *Salmonella typhimurium* upon metabolic reduction of the environmental mutagen 1-nitropyrene. *Cancer Res.* 43: 2052-2058 (1983).
- Pitts, J. N., Jr., Lokensgard, D. M., Harger, W., Fisher, T. S., Mejia, V., Schuler, J. J., Scorziell, G. M., and Katzenstein, Y. A. Mutagens in diesel exhaust particulate: identification and direct activities of 6-nitrobenzo(a)pyrene, 9-nitroanthracene, 1-nitropyrene and 5H-phenanthro[4,5-b]pyran-5-one. *Mutat. Res.* 103: 241-249 (1982).
- Schuetzle, D., Riley, T. L., Prater, T. J., Harvey, T. M., and Hunt, D. F. Analysis of nitrated polycyclic hydrocarbons in diesel particulates. *Anal. Chem.* 54: 265-271 (1982).
- Nishioka, M. B., Peterson, B., and Lewtas, J. Comparison of nitro-PNA content and mutagenicity of diesel emissions. In: *Polynuclear Aromatic Hydrocarbons: Physical and Biological Chemistry* (M. Cook, A. J. Dennis, and G. L. Fisher, Eds.), Battelle Press, Columbus, 1982, pp. 603-613.
- Xu, X. B., Nachtman, J. P., Jin, A. L., Wel, E. T., Rappaport, S. M., and Burlingame, A. L. Isolation and identification of mutagenic nitro-PAH in diesel-exhaust particulates. *Anal. Chem. Acta* 136: 163-174 (1982).
- Tokiwa, H., Kitamori, S., Nakagawa, R., Horikawa, K., and Matenal, L. Demonstration of a powerful mutagenic dinitropyrene in airborne particulate matter. *Mutat. Res.* 121: 107-116 (1983).
- Cole, J., Arlett, C. F., Lowe, J., and Bridges, B. A. The mutagenic potency of 1,8-dinitropyrene in cultured mouse lymphoma cells. *Mutat. Res.* 93: 213-220 (1982).
- Campbell, J., Crumplin, G. C., Garner, R. C., Martin, C. N., and Rutter, A. Nitrated polycyclic aromatic hydrocarbons: potent bacterial mutagens and stimulators of DNA repair synthesis in cultured human cells. *Carcinogenesis* 2: 599-656 (1981).
- Lewtas, J. Mutagenic activity of diesel emissions. In: *Toxicological Effects of Emissions From Diesel Engines* (J. Lewtas, Ed.), Elsevier Science Publishing, New York, 1982, pp. 243-264.
- Nachtman, J. P., Wolff, S. Activity of nitro-polynuclear aromatic hydrocarbons in the sister chromatid exchange assay with and without metabolic activation. *Environ. Mutagen.* 4: 1-7 (1982).
- Rosenkranz, H. S., McCoy, E. C., Mermelstein, R., and Speck, W. T. A cautionary note on the use of nitroreductase-deficient strains of *Salmonella typhimurium* for detection of nitroarenes as mutagens in complex mixtures including exhaust. *Mutat. Res.* 91: 103-105 (1981).
- Mermelstein, R., Kiriazides, D. K., Butler, M., McCoy, E. C., and Rosenkranz, H. S. The extraordinary mutagenicity of nitropyrenes in bacteria. *Mutat. Res.* 89: 187-196 (1981).
- Hirose, M., Lee, M. S., Wang, C. Y., and King, C. M. Induction of rat mammary gland tumors by 1-nitropyrene, a recently-recognized environmental mutagen. *Cancer Res.* 44: 1158-1162 (1984).
- Salmeeen, I., Durisin, A. M., Prater, T. J., Riley, T., and Schuetzle, D. Contribution of 1-nitropyrene to direct-acting Ames assay mutagenicities of diesel particle extracts. *Mutat. Res.* 104: 17-23 (1982).
- Messlier, F., Lu, C., Andrews, P., McCarry, B. E., Quilliam, M. A., and McCalla, D. R. Metabolism of 1-nitropyrene and formation of DNA adducts in *Salmonella typhimurium*. *Carcinogenesis* 2: 1007-1011 (1981).
- Ball, L. M., Kohan, M. J., Inmon, J. P., Claxton, L. D., and Lewtas, J. Metabolism of 1-nitro[<sup>14</sup>C]pyrene *in vivo* in the rat and mutagenicity of urinary metabolites. *Carcinogenesis* 5: 1557-1564 (1984).
- El-Bayoumy, K., and Hecht, S. S. Identification and mutagenicity of metabolites of 1-nitropyrene formed by rat liver. *Cancer Res.* 43: 3132-3137 (1983).
- Ball, L. M., Kohan, M. J., Claxton, L. D., and Lewtas, J. Mutagenicity of derivatives and metabolites of 1-nitropyrene: activation by rat liver S9 and bacterial enzymes. *Mutat. Res.* 138: 113-125 (1984).
- King, L. C., Jackson, M., Ball, L. M., and Lewtas, J. Binding of 1-nitro[<sup>14</sup>C]pyrene to DNA and protein in cultured lung macrophages and respiratory tissues. *Cancer Letters* 19: 241-246 (1983).
- Chan, T. L., Lee, P. S., and Herring, W. E. Deposition and clearance of inhaled diesel exhaust particles in the respiratory tract of Fischer rats. *J. Appl. Toxicol.* 1: 77-82 (1981).
- White, H. J., and Garg, B. D. Early pulmonary response of the rat lung to inhalation of high concentration of diesel particles. *J. Appl. Toxicol.* 1: 104-110 (1981).
- Rudd, C. J., and Strom, K. A. Spectrophotometric method for the quantitation and diesel exhaust particles in guinea pig lung. *J. Appl. Toxicol.* 1: 83-87 (1981).
- King, L. C., Loud, K., Tejada, S. B., Kohan, M. J., and Lewtas, J. Evaluation of the release of mutagens and 1-nitropyrene from diesel particles in the presence of lung macrophage cells in culture. *Environ. Mutag.* 5: 577-588 (1983).
- Garrett, N. E., Campbell, J. A., Stack, H. F., Waters, M. D., and Lewtas, J. The utilization of the rabbit alveolar macrophage and Chinese hamster ovary cell for evaluation of the toxicity of particulate materials: 1. Model compounds and metal-coated fly ash. *Environ. Res.* 24: 345-365 (1981).
- Jackson, M. A., King, L. C., and Ball, L. M. A rapid technique for estimating DNA binding used to evaluate 1-nitropyrene adduct formation. *Drug Chem. Toxic.* 6: 549-562 (1983).
- Schneider, W. C. Determination of nucleic acids in tissues by pentose analysis. In: *Methods in Enzymology*, Vol. 3 (S. P. Colowick and N. O. Kaplan, Eds.), Academic Press, New York, 1957, p. 680.
- Lowry, O. H., Rosebrough, N. G., Farr, A. L., and Randall, R. J. Protein measurement with the Folin reagent. *J. Biol. Chem.* 193: 265-275 (1951).
- Autrup, H., WeFald, F. C., Jeffrey, A. M., Tate, H., Schwartz, R. D., Trump, B. F., and Harris, C. C. Metabolism of benzo(a)pyrene by cultured tracheobronchial tissues from mice, rats, hamsters, bovines and humans. *Int. J. Cancer* 25: 293-300 (1980).