

# Chelation of Cadmium

by Ole Andersen\*

The toxicity of cadmium is determined by chelation reactions: *in vivo*, Cd<sup>2+</sup> exists exclusively in coordination complexes with biological ligands, or with administered chelating agents.

The Cd<sup>2+</sup> ion has some soft character, but it is not a typical soft ion. It has a high degree of polarizability, and its complexes with soft ligands have predominantly covalent bond characteristics. Cd<sup>2+</sup> forms the most stable complexes with soft donor atoms (S >> N > O). The coordination stereochemistry of Cd<sup>2+</sup> is unusually varied, including coordination numbers from 2 to 8. Even though the Cd<sup>2+</sup> ion is a d<sup>10</sup> ion, disturbed coordination geometries are often seen. Generally, the stability of complexes increases with the number of coordination groups contributed by the ligand; consequently, complexes of Cd<sup>2+</sup> with polydentate ligands containing SH groups are very stable. Cd<sup>2+</sup> in metallothionein (MT) is coordinated with 4 thiolate groups, and the log stability constant is estimated to 25.5.

Complexes between Cd<sup>2+</sup> and low molecular weight monodentate or bidentate ligands, e.g., free amino acids (LMW-Cd), seem to exist very briefly, and Cd<sup>2+</sup> is rapidly bound to high molecular weight proteins, mainly serum albumin. These complexes (HMW-Cd) are rapidly scavenged from blood, mainly by the liver, and Cd<sup>2+</sup> is redistributed to MT. After about 1 day the Cd-MT complex (MT-Cd) almost exclusively accounts for the total retained dose of Cd<sup>2+</sup>, independent of the route of exposure. MT-Cd is slowly transferred to and accumulated in kidney cortex. The acute toxicity and interorgan distribution of parenterally administered Cd<sup>2+</sup> are strongly influenced by preceding MT induction, or decreased capacity for MT synthesis; however, the gastrointestinal (GI) uptake of Cd<sup>2+</sup> seems unaffected by preceding MT induction resulting in considerable capacity for Cd<sup>2+</sup> chelation in intestinal mucosa, and this finding indicates that endogenous MT is not involved in Cd<sup>2+</sup> absorption. The toxicity of parenterally administered Cd<sup>2+</sup> is strongly enhanced when administered as complexes with NTA or STPP, but it is much decreased when administered as a complex with EDTA. In chronic oral exposure the toxicity and GI uptake of Cd<sup>2+</sup> is not changed when Cd<sup>2+</sup> is administered as a complex with the detergent formula chelating agents NTA, EDTA and STPP. The uptake of Cd<sup>2+</sup> from ligated intestine *in vivo* was not affected by administration of Cd<sup>2+</sup> as complexes with CYS or GSH, but significantly reduced by complexation with EDTA or BAL. The acute toxicity of orally administered Cd<sup>2+</sup> is reduced when Cd<sup>2+</sup> is administered as a complex with EDTA. The renal toxicity and deposition of Cd<sup>2+</sup> is markedly increased when Cd<sup>2+</sup> is parenterally administered as MT-Cd. Orally administered Cd-MT is taken up by intestinal mucosa and transferred to kidneys, where MT is broken down. Perfusion of intestinal loops *in situ* with MT-Cd induces necrosis of intestinal mucosa, not observed with Cd<sup>2+</sup> alone.

During acute treatment of Cd<sup>2+</sup> poisoning the superior agents seem to be the polyamino-polycarboxylates, especially DTPA, which increases renal excretion. The time lapse between Cd intoxication and antidote administration is critical due to the rapid intracellular accumulation of Cd<sup>2+</sup>, mainly in liver. During chronic Cd<sup>2+</sup> exposure, BAL significantly increases biliary excretion of Cd<sup>2+</sup>, and part of the mobilized Cd<sup>2+</sup> is chelated out of MT.

## Introduction

The element cadmium (Cd) was discovered in 1817. Since then Cd has gained increasing technological importance, and the annual world

smelter production of Cd has increased from 10,000 short tons in 1958 to 18,780 in 1974. Cd is mainly obtained as a by-product from smelting zinc ore, where Cd often is present as an impurity (0.2-0.3%). Cd metal and compounds of Cd are used for electroplating (50% of world production), Cd-bearing alloys, Cd-Ni batteries, pigments, stabilizer in PVC plastics, and control rods in

\*Institute of Community Health, Department of Environmental Medicine, Odense University, J. B. Winsløvs Vej 19, DK-5000 Odense C, Denmark.

nuclear fission reactors. Human exposure to Cd has taken place in industries producing and utilizing Cd, but serious exposures have also occurred in the zinc industry, in operations (mainly welding) on Cd-plated items, in scrap remelting, and due to environmental pollution. Today Cd is ubiquitous in the biosphere. Due to the great concern regarding present and future human health aspects of Cd pollution, a number of reviews of chemical, environmental and biological effects of Cd have appeared (1-8).

Cd is very toxic to mammals both during acute and chronic exposure. In biological systems, soluble  $\text{Cd}^{2+}$  inevitably exists as coordination complexes with biological ligands, and the toxicity of  $\text{Cd}^{2+}$  is beyond doubt mediated through chelation reactions at the target. Although the exact mechanism of Cd toxicity has not been established in any organism, the modifying effect of metallothionein on Cd toxicity is widely recognized (9,10). Recent reviews of the chemistry of metals with special emphasis on biological effects have been prepared by Martell, Schubert and Nieboer (11-13), and two recent reviews (14,15) were devoted to the coordination chemistry of cadmium.

## Chemistry of Cd

A chemical description of the  $\text{Cd}^{2+}$  ion, in relation to those metal ions known to interplay with  $\text{Cd}^{2+}$  in biological systems, i.e.,  $\text{Cu}^{2+}$  and  $\text{Zn}^{2+}$ , as well as those donor groups available for complex formation in the biological systems, may contribute to the understanding of the ligand preference of  $\text{Cd}^{2+}$  during Cd intoxication, as well as the possibilities for induction of ligand exchange reactions during chelation therapy.

Together with Zn and Hg, Cd forms group IIB in the periodical table of the elements. Although a few compounds of  $\text{Cd}^+$  exist, Cd is usually found in valence +2, with outer electron configuration  $4d^{10}$ . The polarizability of the  $\text{Cd}^{2+}$  ion is rather high ( $\alpha = 1.8 \text{ \AA}^3$ ; 16), and coordination bonds formed with soft donors have a high degree of covalent character. Organic compounds of Cd ( $\text{R}_2\text{Cd}$  and  $\text{RCd}^+$ , R = aliphatic) exist (diethylcadmium is used in the production of tetraethyllead), but are unstable and decompose upon contact with water.

The stability of complexes formed between Cd and polar groups, in proteins (mainly those containing N, O or S), nucleic acids (the phosphate moieties or the bases) or added chelating agents, depend on energy changes associated with complex formation. The size of these energy changes depend on chemical features of the ligand (and

the metal ion, in this case  $\text{Cd}^{2+}$ ), the most important being stereochemical conditions at the ligand, "hardness/softness," the chelate effect and rate effects. These four factors of course interplay, and they cannot be considered independent of each other.

## Stereochemistry of $\text{Cd}^{2+}$ Complexes

Information about the stereochemistry of coordination complexes of  $\text{Cd}^{2+}$  with biological ligands in aqueous solution is of great importance for understanding the biological effects of  $\text{Cd}^{2+}$ . However, the largest part of information available concern the coordination chemistry of  $\text{Cd}^{2+}$  complexes in the solid state, examined by X-ray crystallographic methods. An extensive review of the stereochemistry of coordination compounds of Cd was recently published by Tuck (14). From the  $d^{10}$  state of the  $\text{Cd}^{2+}$  ion the coordination chemistry of  $\text{Cd}^{2+}$  would be expected to be dominated by examples of coordination to four donors of two electrons, according to the inert gas rule. Many examples of this are known. However, cadmium is also found in a large number of complexes with other coordination numbers ranging from 2 to 8, in many cases with the structure distorted from regular tetrahedral or octahedral. In Table 1 examples of coordination compounds of  $\text{Cd}^{2+}$  are given. It should be noted that, except for the aquo-cation of  $\text{Cd}^{2+}$ , the data pertain to nonaqueous conditions, and use of these data for extrapolation to biological systems is questionable.

The only known examples of two-coordinate  $\text{Cd}^{2+}$  species occur in gas phase. In aqueous phase such structures are hydrated. The  $\text{CdBr}_3^-$  complex has pyramidal structure in aqueous solution, indicating hydration. The only known example of tetrahedral  $\text{CdO}_4$  coordination is in crystalline  $\text{CdZn}(\text{PO}_4)_2$ .  $\text{CdN}_4$  kernels have not been definitely proven. However, a few examples of  $\text{CdC}_4$  are known, and  $\text{CdS}_4$  coordination is observed with several chelating agents with S-donor ligands, all with more or less distorted  $T_d$  structure. The stereochemistry of five-coordinated  $\text{Cd}^{2+}$  is varied, and many examples of distorted  $C_{4v}$  and  $D_{3h}$  symmetry have been reported. Examples of  $\text{CdO}_5$ ,  $\text{CdN}_5$  and  $\text{CdS}_5$  coordination are known, in most cases forming dimers or polymers.

The aquo-cation of  $\text{Cd}^{2+}$  is  $(\text{Cd}(\text{H}_2\text{O})_6)^{2+}$ . Several examples of  $\text{CdO}_6$  and  $\text{CdN}_6$  kernels have been reported. The highest number of S reported in hexacoordinate  $\text{Cd}^{2+}$  complexes is 4, in the kernels  $\text{CdO}_2\text{S}_4$  and  $\text{CdN}_2\text{S}_4$ , and  $\text{Cd}^{2+}$  complexes containing S in the kernel with coordination numbers higher than six have not been reported.

Table 1. Stereochemistry of coordination complexes of Cd<sup>2+</sup>.

<i>N</i> <sub>a</sub>	Compound	Coordin. Kernel	Stereochemistry	Reference
2	CdX <sub>2</sub> (gas phase), X = F, Cl, Br, I	—	<i>D</i> <sub>∞h</sub> symmetry	(17)
3	CdBr <sub>3</sub> - (non aq. sol.)	—	<i>D</i> <sub>3h</sub> symmetry	(18)
4	CdZn <sub>2</sub> (PO <sub>4</sub> ) <sub>2</sub>	CdO <sub>4</sub>	Distorted <i>T</i> <sub>d</sub> ; polymeric lattice	(19)
	Cd(CN) <sub>2</sub> ·en·2C <sub>6</sub> H <sub>6</sub>	CdC <sub>4</sub>	<i>T</i> <sub>d</sub> ; polymeric structure	(20)
	Cd(2-methyl thioxine) <sub>2</sub>	CdN <sub>2</sub> S <sub>2</sub>	Distorted <i>T</i> <sub>d</sub> ; monomer	(21)
	CdCl <sub>2</sub> thiourea <sub>2</sub>	CdCl <sub>2</sub> S <sub>2</sub>	Distorted <i>T</i> <sub>d</sub> ; monomer	(22)
	Cd(Et-xanthate) <sub>2</sub>	CdS <sub>4</sub>	Distorted <i>T</i> <sub>d</sub> ; each S from different xanthate	(23)
5	Cd <sub>2</sub> Zn(PO <sub>4</sub> ) <sub>2</sub>	CdO <sub>5</sub>	Trigonal pyramidal; polymeric lattice	(19)
	Cd(N,N-diethyldithio carbamic acid) <sub>2</sub>	CdS <sub>5</sub>	Distorted <i>C</i> <sub>4v</sub> ; dimer	(24)
	Cd(thioglycollate) <sub>2</sub>	CdS <sub>4</sub> O	Distorted <i>D</i> <sub>3h</sub> ; polymeric layer	(25)
6	[Cd(OH <sub>2</sub> ) <sub>6</sub> ] <sup>2+</sup> (aq. sol.)	CdO <sub>6</sub>	<i>O</i> <sub>h</sub>	(26)
	Cd(acetylacetonate) <sub>2</sub>	CdO <sub>6</sub>	Approx. <i>O</i> <sub>h</sub> ; parallel chain structures	(27)
	[Cd(imidazole) <sub>6</sub> ]CO <sub>3</sub> ·5H <sub>2</sub> O	CdN <sub>6</sub>	Approx. <i>O</i> <sub>h</sub>	(28)
	Cd(NCS) <sub>2</sub>	CdN <sub>2</sub> S <sub>4</sub>	Approx. <i>O</i> <sub>h</sub> ; three-dimensional polymer, NCS-linked	(29)
	Cd(HCOO) <sub>2</sub> (thiourea) <sub>2</sub>	CdO <sub>2</sub> S <sub>4</sub>	Approx. <i>O</i> <sub>h</sub> ; chain polymer, S-bridged	(30)
	Cd(L-methionate) <sub>2</sub>	CdO <sub>4</sub> N <sub>2</sub>	Distorted <i>O</i> <sub>h</sub>	(31)
	Cd(D-penicillamine)·H <sub>2</sub> O	CdO <sub>3</sub> S <sub>2</sub> N	Distorted <i>O</i> <sub>h</sub>	(32)
7	Cd(CH <sub>3</sub> COO) <sub>2</sub> ·2H <sub>2</sub> O	CdO <sub>7</sub>	Square-based trigonal cap	(33)
	Cd(malonate) <sub>2</sub> ·H <sub>2</sub> O	CdO <sub>7</sub>	Distorted pentagonal bipyramid	(34)
8	Cd(NO <sub>3</sub> ) <sub>2</sub> ·4H <sub>2</sub> O	CdO <sub>8</sub>	Bidentate NO <sub>3</sub> on each side of square planar Cd(OH <sub>2</sub> ) <sub>4</sub>	(35)
	Cd(maleate)·2H <sub>2</sub> O	CdO <sub>8</sub>	Distorted dodecahedron	(36)

<sup>a</sup>*N* = coordination number.

The stereochemistry of hexacoordinate Cd<sup>2+</sup> is dominated by distorted or approximately *O*<sub>h</sub> structure. Examples of heptacoordinate Cd<sup>2+</sup> complexes with 4–7 O in the kernel are known, mainly with distorted pentagonal bipyramidal structure. In octacoordinate complexes, Cd<sup>2+</sup> has been found in *O*<sub>8</sub> and *N*<sub>8</sub> kernels, mainly with distorted dodecahedron structure.

### Scales for “Hardness/Softness”

The “hardness/softness” (or class A or B character) of metal ions, and thus their behavior (= ligand preference) in aqueous solution, is determined by physical and chemical characteristics of the ion. The same, of course, holds true for the donor groups. Some discrepancy exists in the chemical literature about whether to consider Cd<sup>2+</sup> a “soft” ion or an “intermediate” ion. A little scrutiny on this subject may therefore be relevant.

Several authors have developed formal descriptors for tabulating donors and acceptors in complexation reactions into the three classes, “hard” or class A, “borderline” or “intermediate,” and “soft” or class B. This classification was originally suggested by Schwarzenbach (37) and Ahrland et al. (38) and further elaborated by Pearson (39),

mainly on the basis of differences in the stability of metal halide complexes in water. Thus, a typical hard metal ion has the donor atom preference sequence: F<sup>-</sup> > Cl<sup>-</sup> > Br<sup>-</sup> > I<sup>-</sup>, and a typical soft metal ion has the opposite preference sequence: I<sup>-</sup> > Br<sup>-</sup> > Cl<sup>-</sup> > F<sup>-</sup>. Based on this definition, Cd<sup>2+</sup> is a soft ion.

Several softness descriptors are listed in Table 2, and values for Cd<sup>2+</sup> and other divalent ions tabulated: The ionic index correlates well with ionic interactions and indicates the propensity of the ion to form ionic bonds (40,41). The ratio of the sum of ionization energies of an ion over the ionic index, i.e.,  $r\Sigma I/Z^2$  was used by Williams as a measure of softness (41). Williams and Hale (41) defined the softness parameter *Q* as

$$Q = \frac{\Delta H_{\text{ion(g)}}(\text{MF}_n) - \Delta H_{\text{ion(g)}}(\text{MI}_n)}{1/2 [(\Delta H_{\text{ion(g)}}(\text{MF}_n) + \Delta H_{\text{ion(g)}}(\text{MI}_n))]}$$

where  $\Delta H_{\text{ion(g)}}(\text{MF}_n)$  is the coordinate bond energy, i.e., the enthalpy change for the reaction



in the gas phase, and analogously for iodide. Pearson and Mawbey (42) defined a related softness parameter

$$\sigma_p = \frac{\Delta H_{\text{ion(g)}}(\text{MF}_n) - \Delta H_{\text{ion(g)}}(\text{MI}_n)}{\Delta H_{\text{ion(g)}}(\text{MF}_n)}$$

The softness of metals increases with decreasing  $Q$  and decreasing  $\sigma_p$ .

Klopman (43) defined the softness parameter  $\sigma_K$  as the difference (in eV) between the empty frontier orbital energy and the desolvation energy of a metal ion. Ahrland (44) defined a softness parameter,

$$\sigma_A = (I_n + \Delta H_n)/Z$$

i.e., the difference between the total ionization potential and the dehydration energy ( $-\Delta H_n$ ) divided by the ion's charge. The  $\sigma_A$  increase with increasing softness.

The covalent index,  $X_m^{2r}$ , where  $X_m$  is the electronegativity of the ions, was used by Nieboer and Richardson (13) to describe softness. The rationale is that  $X_m$  is related to the empty frontier orbital energy of a metal ion, and this parameter indicates the ability of the ion to accept electrons and form covalent bonds.  $X_m^{2r}$  increases with increasing softness.


In Table 2, several divalent metal ions are placed in order of average increasing softness, as based on the softness parameters. A reasonably good correlation between the different descriptors is demonstrated, although many different thermodynamic functions and experimental measurements have been used. The selection of exclusively divalent ions of course favors this agreement.

Generally, hard metal ions form stable complexes with hard donors, while soft metal ions form stable complexes with soft donors. However, softness itself favors formation of stable complexes. Thus, in a series of analogous coordination complexes between ions with the same charge and different ligands with the same number of

donor groups, the stability of the complexes depends mainly on the softness of the ion and the ligand. This is illustrated in Table 3, which gives the stability constants of complexes between several divalent metal ions and different bidentate ligands. Table 3 demonstrates the general trend that the stability constants increase downwards and towards the right. This may be explained by increasing softness of the metals downwards and of the chelating agents' donor groups towards the right, i.e., the coordination bonds formed have a high degree of ionic character in the upper left part of the table and a high degree of covalent character in the lower right part. Table 4 summarizes the most important biological ligands, grouped according to their metal preference.

## Thermodynamics of Complex Formation

The stability of coordination complexes can conveniently be described by thermodynamic functions. When  $n$  ligands form a coordination complex with a metal ion,



the ligational free energy,  $\Delta G_n$ , is related to the stability constant:

$$\beta_n = (MA_n)/(M)(A)^n$$

through  $\Delta G_n = -RT \ln \beta_n$ . The energy change is contributed by changes in enthalpy,  $\Delta H_n$  and entropy,  $\Delta S_n$ :

$$\Delta G_n = \Delta H_n - T\Delta S_n$$

$\Delta H$  mainly describes energy changes due to bonding, while  $T\Delta S$  mainly describes energy changes

Table 2. Values of softness parameters for divalent metal ions.

Hardness <sup>a</sup>	Metal ion	$\alpha$	$\frac{Z^2}{r}$	$\frac{rI}{Z^2}$	$Q$	$\sigma_p$	$\sigma_K$	$\sigma_A$	$X_m^{2r}$
H	Be <sup>2+</sup>	0.01	9.09	3.02	0.20	0.172	-3.75	1.2	0.85
H	Mg <sup>2+</sup>	0.10	4.87	—	0.21	0.167	-2.42	1.4	1.25
H	Ca <sup>2+</sup>	1.10	4.04	4.45	0.21	0.180	-2.33	0.9	1.00
I	Ni <sup>2+</sup>	1.00	7.79	3.30	—	0.126	-0.29	2.0	2.50
I	Fe <sup>2+</sup>	1.20	5.40	4.45	—	0.127	-0.69	2.1	2.60
I	Co <sup>2+</sup>	1.20	5.55	4.48	—	0.130	—	1.8	2.65
I	Mn <sup>2+</sup>	1.25	5.00	4.61	—	0.124	—	2.0	2.00
I	Zn <sup>2+</sup>	0.80	5.40	5.06	0.12	0.115	—	3.1	2.20
S	Cu <sup>2+</sup>	—	5.55	5.04	—	0.104	0.55	3.1	2.95
S	Cd <sup>2+</sup>	1.80	4.12	6.28	0.09	0.081	2.04	3.5	2.80
S	Pb <sup>2+</sup>	4.80	3.33	—	0.14	0.131	—	4.1	3.40
S	Hg <sup>2+</sup>	2.90	3.63	8.03	0.07	0.064	4.64	4.64	4.10
Reference		(16)	b	c	(41)	(45)	(45)	(45)	(13)

<sup>a</sup>H = hard; I = intermediate; S = soft.

<sup>b</sup>Calculated from the formal charge and the crystal ionic radius (46).

<sup>c</sup>Calculated from the ionic index and the total ionization potential (45).

Table 3. Equilibrium constants for complexes between divalent metals and simple bidentate ligands:  $M + L \rightarrow ML$ .

Hard-ness <sup>a</sup>	Metal ion	log $\beta$				
		Oxalate	Glycine	Ethylene-diamine	Mercapto-acetate	Mercapto-ethylamine
H	Be <sup>2+</sup>	4.08 <sup>b</sup>	—	—	—	—
H	Mg <sup>2+</sup>	2.76 <sup>b</sup>	2.22 <sup>b</sup>	0.37 <sup>c</sup>	—	2.30 <sup>b</sup>
H	Ca <sup>2+</sup>	3.00 <sup>c</sup>	1.39 <sup>d</sup>	—	—	2.21 <sup>b</sup>
I	Ni <sup>2+</sup>	5.16 <sup>d</sup>	5.78 <sup>b</sup>	7.35 <sup>b</sup>	(6.2) <sup>f</sup>	10.05 <sup>e</sup>
I	Fe <sup>2+</sup>	3.05 <sup>d</sup>	4.31 <sup>d</sup>	4.34 <sup>c</sup>	—	—
I	Co <sup>2+</sup>	3.84 <sup>b</sup>	4.64 <sup>b</sup>	5.96 <sup>c</sup>	5.84 <sup>b</sup>	7.68 <sup>e</sup>
I	Mn <sup>2+</sup>	3.2 <sup>b</sup>	2.80 <sup>b</sup>	2.67 <sup>b</sup>	4.38 <sup>b</sup>	—
I	Zn <sup>2+</sup>	3.88 <sup>b</sup>	4.96 <sup>b</sup>	5.7 <sup>b</sup>	7.86 <sup>b</sup>	9.90 <sup>b</sup>
S	Cu <sup>2+</sup>	4.84 <sup>b</sup>	8.15 <sup>b</sup>	10.54 <sup>b</sup>	—	(16)
S	Cd <sup>2+</sup>	3.89 <sup>d</sup>	4.22 <sup>b</sup>	5.45 <sup>b</sup>	—	10.97 <sup>b</sup>
S	Pb <sup>2+</sup>	4.00 <sup>b</sup>	5.47 <sup>d</sup>	7.00 <sup>b</sup>	(8.5) <sup>f</sup>	11.10 <sup>b</sup>
S	Hg <sup>2+</sup>	9.66 <sup>b</sup>	10.3 <sup>c</sup>	14.3 <sup>b</sup>	(24) <sup>f</sup>	—

<sup>a</sup>H = Hard, I = Intermediate, S = Soft. All values from Martell and Smith (47), except as noted.

<sup>b</sup>25°C, ionic strength 0.1.

<sup>c</sup>25°C, ionic strength 0.5.


<sup>d</sup>25°C.

<sup>e</sup>30°C, ionic strength, 1.0.

<sup>f</sup>Data of Jacobson and Turner (15).

related to the number of independent species, e.g., associated with changes in solvation. During complex formation with highly charged negative polydentate ligands such as the polyaminopolycarboxylates, the main contribution to the stability of the complex often is the increase in entropy due to loss of solvation, because of charge neutralization (Table 5). In the formation of complexes with uncharged ligands, e.g., the polyamines, the stability is generally due to decrease in enthalpy. The reason enthalpy changes are not totally outnumbered by entropy changes, e.g., during ligand selection in mixed chelation, is that the high dielectric constant of water efficiently screens electrostatic forces.

Chelating ligands form much more stable complexes than unidentate ligands: In series of analogous complexes with equal numbers of the same ligand, the stability generally increases with the number of rings formed. From the displacement reaction




where  $I_n$  is a polydentate ligand with the same donor atoms as A, Schwarzenbach (49) defined the chelate effect as the logarithm to the equilibrium constant for the displacement reaction. This entity can be related to the change in free energy. Assuming (what may not always be true, however) that the change in enthalpy associated with complex formation is independent on whether the coordination groups are unidentate or bound together in a polydentate ligand, the chelate effect should be due to increased entropy. A significant enthalpy effect is, however, often observed, since

Table 4. Biological ligands.

Ligands preferred by hard ions	Ligands preferred by intermediate or soft ions	Ligands preferred by soft ions
OH <sup>-</sup>	Cl <sup>-</sup>	R-SH
H <sub>2</sub> O	NH <sub>3</sub>	R-S-R
CO <sub>3</sub> <sup>2-</sup>	N <sub>2</sub>	R-S-S-R
SO <sub>4</sub> <sup>2-</sup>	O <sub>2</sub>	
R-O-SO <sub>3</sub> <sup>-</sup>	R-NH <sub>2</sub>	
NO <sub>3</sub> <sup>-</sup>	R <sub>2</sub> NH	
HPO <sub>4</sub> <sup>2-</sup>	R <sub>3</sub> N	
Mono- and diesters of phosphate	R = N-R	
R-OH	R-CO-N-R	
R-COOH		
R-CHO		
R-CO-R		
R-O-R		

Table 5. Effect of ring size on the chelate effect in the displacement reaction.



$n$	log $\beta^a$	$G$ , kcal/mole	$H$ , kcal/mole	$T/S$
2	7.2	-5.3	-1.8	3.5
3	2.4	-1.8	1.8	3.6
4	-1	0.7	4.4	3.7
5	-1.6	1.2	2.8	1.6

<sup>a</sup>Calculated from  $\Delta G$ , 20°C, 0.1 (48).

mutual repulsive forces between the donor groups are overcome by introducing them into one molecule. The chelate effect may be counteracted by adverse energy changes, due to necessary conformational changes during complex formation, especially when strained rings are formed. This is shown in Table 5:  $\Delta H$  more than counteracts the entropy contribution to stability as the ring size increases.

Table 6. Stability of complexes between Cd<sup>2+</sup> and polydentate ligands, M + L → ML.


Ligand	Donor groups	log β
Malonic acid	2	2.64 <sup>a</sup>
Citric acid	3	3.75 <sup>a</sup>
Ditartronic acid	4	5.44 <sup>a</sup>
Ethylenediamine (en)	2	5.45 <sup>a</sup>
Diethylenetriamine (dien)	3	8.40 <sup>b</sup>
Triethylenetetramine (trien)	4	10.63 <sup>a</sup>
Tetraethylenepentamine (tetren)	5	14.00 <sup>a</sup>
N,N,N',N'-tetra(2-aminoethyl) ethylenediamine (penten)	6	16.10 <sup>b</sup>
Iminodiacetic acid (IDA)	3	5.71 <sup>a</sup>
Nitrilotriacetic acid (NTA)	4	9.78 <sup>a</sup>
N-(2-Hydroxyethyl) ethylenediamine-N,N',N'-triacetic acid (HEDTA)	5	13.10 <sup>a</sup>
Ethylenediaminetetraacetic acid (EDTA)	6	16.36 <sup>a</sup>
Diethylenetrinitriolpentaacetic acid (DTPA)	8	19.00 <sup>a</sup>

<sup>a</sup>25°C, 0.1.<sup>b</sup>20°C, 0.1. Data from Martell and Smith (47).

At low concentrations of chelant, the chelate effect is an important contribution to stability. For the formation of a hexacoordinate Cd complex with a monodentate ligand Cd + 6A → CdA<sub>6</sub> the equilibrium constant is:

$$\beta_{(A)_6} = (CdA_6)/(Cd)(A)^6$$

At a decreasing concentration of ligand, the amount of complex will decrease with (A)<sup>6</sup>, i.e., extremely fast, and for biological ligands (except water) (A)<sup>6</sup> is a very small number. For the formation of the analog complex with a hexadentate ligand containing 6A groups:



the equilibrium constant is

$$\beta_{(LA)_6} = (CdLA_6)/(Cd)(LA_6)$$

and the amount of complex formed is directly proportional to the concentration of the ligand.

Experimentally, the gain in stability due to chelate ring formation has been found to be in the order of 10<sup>2</sup> per ring formed, i.e., β<sub>LA<sub>6</sub></sub> could be 10<sup>10</sup> β<sub>A<sub>6</sub></sub> or larger. The obvious consequence for chelation therapy during acute Cd intoxication is that at the nontoxic chelant concentrations the efficiency of chelation therapy can be improved most effectively by increasing the β value. This improvement can be obtained by increasing the number of chelate rings formed. According to Table 6, the introduction of one extra ring may increase β by a factor up to 10<sup>4</sup> (note that the complex MLA<sub>n</sub> is compared with the complex MLA<sub>n+1</sub>, thus only part of the increased stability is due to the chelate effect). The other way would be to increase the softness of the chelant's donor groups. According to Table 3, a change of "N" to "S" can increase β by approximately 4 orders of magnitude for complexes with soft metals. Table 7

Table 7. Stability of complexes between Cd<sup>2+</sup> and substituted IDA.

Compound:	Equilibrium	Log β
Aliphatic CH <sub>3</sub> -N-(CH <sub>2</sub> COO) <sub>2</sub>	ML/M·L ML <sup>2</sup> /M·L <sup>2</sup>	6.75 <sup>a</sup> 12.43 <sup>a</sup>
Alcohol HO-CH <sub>2</sub> -CH <sub>2</sub> -N(CH <sub>2</sub> COO) <sub>2</sub>	ML/M·L ML <sub>2</sub> M·L <sup>2</sup>	7.24 <sup>a</sup> 12.31 <sup>a</sup>
Ether CH-O-CH <sub>2</sub> -CH <sub>2</sub> -N(CH <sub>2</sub> COO) <sub>2</sub>	ML/M·L ML <sub>2</sub> /M·L <sup>2</sup>	7.53 <sup>b</sup> 13.18 <sup>b</sup>
Ketone CH <sub>3</sub> -CO-CH <sub>2</sub> -N(CH <sub>2</sub> COO) <sub>2</sub>	ML/M·L ML <sub>2</sub> /M·L <sup>2</sup>	7.37 <sup>a</sup> 12.34 <sup>a</sup>
Carboxyl OOC-CH <sub>2</sub> -N-(CH <sub>2</sub> COO) <sub>2</sub>	ML/M·L ML <sub>2</sub> /ML <sup>2</sup>	9.78 <sup>a</sup> 14.39 <sup>a</sup>
Amine H <sub>2</sub> N-CH <sub>2</sub> -CH <sub>2</sub> -N(CH <sub>2</sub> COO) <sub>2</sub>	ML/M·L ML <sub>2</sub> /M·L <sup>2</sup>	10.53 <sup>b</sup> 16.59 <sup>b</sup>
Mercapto HS-CH <sub>2</sub> -CH <sub>2</sub> -N(CH <sub>2</sub> COO) <sub>2</sub>	ML/M·L ML <sub>2</sub> /M·L <sup>2</sup>	16.72 <sup>b</sup> 22.33 <sup>b</sup>

<sup>a</sup>25°C, 0.1.<sup>b</sup>20°C, 0.1. Data from Martell and Smith (47).

demonstrates that the stability of analogous complexes of Cd<sup>2+</sup> with different compounds formed by substitution of the methyl group in N-methyliminodiacetic acid (MIDA) with donor groups of increasing softness, increases as expected: The change of -COOH to -CH<sub>2</sub>SH yields an increase in stability of 7 to 8 orders of magnitude. It therefore seems a logical step to test chelants with as many SH groups as possible. The highest number of S atoms which can at the same time coordinate with Cd<sup>2+</sup> seems to be 5. In hexacoordinate complexes, Cd<sup>2+</sup> can coordinate with S<sub>4</sub>N<sub>2</sub> and S<sub>4</sub>O<sub>2</sub> (Table 1).

Effective SH-containing chelants for Cd<sup>2+</sup> should have two vicinal SH groups (50). The tox-

Table 8. Rate constants for ligand exchange reactions.<sup>a</sup>

Metal ion	Ionic index	Exchange rate constant $k_M \text{H}_2\text{O}, \text{sec}^{-1}$		Formation rate constant $K_f, \text{M}^{-1}\text{sec}^{-1}$	
		$\text{H}_2\text{O}-\text{H}_2\text{O}$	$\text{L}-\text{H}_2\text{O}$	$\text{NTA}^{3-}$	$\text{HEDTA}^{3-}$
Be <sup>2+</sup>	9.09	$2.1 \times 10^3$	102b	—	—
Mg <sup>2+</sup>	4.87	$5.3 \times 10^5$	$3 \times 10^{5b}$	—	—
Ca <sup>2+</sup>	4.04	$4 \times 10^8$	$6-9 \times 10^{8c}$	—	$1-1.7 \times 10^7$
Ni <sup>2+</sup>	7.79	$3 \times 10^4$	$1-2 \times 10^{4b}$	$4.8 \times 10^5$	$1 \times 10^6$
Fe <sup>2+</sup>	5.40	$3.2 \times 10^6$	$1-2 \times 10^{6b}$	—	—
Co <sup>2+</sup>	5.55	$1.1-2.4 \times 10^6$	$2 \times 10^5-2 \times 10^{6b}$	—	$4 \times 10^6$
Mn <sup>2+</sup>	5.00	$3.1 \times 10^7$	$4 \times 10^6-5 \times 10^{7b}$	$5 \times 10^8$	—
Zn <sup>2+</sup>	5.40	$3 \times 10^7$	$3 \times 10^{7b}$	$2 \cdot 10^9-10^{10}$	$1-2 \times 10^9$
Cu <sup>2+</sup>	5.55	$5 \times 10^9$	$2-5 \times 10^{9b}$	$2 \times 10^8$	$8.4 \times 10^7$
Cd <sup>2+</sup>	4.12	$3 \times 10^8$	$2.5 \times 10^{8c}$	$10^9-2 \times 10^{10}$	$2.6 \cdot 10^8-6.4 \times 10^9$
Pb <sup>2+</sup>	3.33	—	$7.5 \times 10^{9c}$	$1.5 \times 10^{11}$	$23 \times 10^9-6 \times 10^{11}$
Hg <sup>2+</sup>	3.63	$2 \times 10^9$	$2 \times 10^{9d}$	—	—

<sup>a</sup>Data from Margerum et al. (52).

<sup>b</sup>SO<sub>4</sub><sup>2-</sup>.

<sup>c</sup>CH<sub>3</sub>COO<sup>-</sup>.

<sup>d</sup>Cl<sup>-</sup>.

icity of propanetrithiol is much larger than that of BAL and related compounds, most likely due to adduct formation with proteins (50). Consequently, the toxicity may be reduced by introducing two pairs of vicinal SH groups, favouring tetrahedral coordination as in the Cd-MT complex. The efficiency of charged chelants as, e.g., the polyaminopolycarboxylic acids against the acute toxicity of parenterally administered Cd<sup>2+</sup> decreases rapidly concomitantly with cellular uptake of Cd<sup>2+</sup>, and not with the kinetics of MT induction (51). High efficiency of SH-containing chelants against aged Cd<sup>2+</sup> deposits requires lipophilicity of the agent (50). Thus, the critical step in successful chelation of Cd<sup>2+</sup>, except immediately after administration, seems to be transport over the cell membrane, first of the chelant and subsequently of the chelate.

By combining chemical considerations, such as those reviewed above with biological experiments it will hopefully be possible to design new and more effective chelating agents for Cd<sup>2+</sup>, in order to antagonize the acute toxicity of Cd<sup>2+</sup> as well as to detoxify aged Cd<sup>2+</sup> deposits in the liver without inducing renal damage.

## Rate Effects

The formation of coordination complexes in aqueous solution (in complex biological systems it could be called "ligand selection") is dependent on rate constants and equilibrium constants. Table 8 demonstrates a correlation between the aptitude of different ions for hydration and the ionic index,  $Z^2/r$ . The rate of exchange of water molecules from the inner hydration layer is largest for the less stable hydration complexes, and inspection of

Table 8 demonstrates that also the rate of simple displacement reactions, as expected, is inversely related to the ionic index. The ligand exchange rates for cadmium are generally large for water exchange, simple ligand exchange and complex formation with polydentate ligands (Table 8).

Thus, equilibrium will be reached rather fast, and after some time the distribution of Cd<sup>2+</sup> between different ligands will reflect (1) the affinity of Cd<sup>2+</sup> for the ligands as discussed above, i.e., mainly softness, and (2) stabilizing effects, i.e., mainly the chelate effect. Thus, during the initial phase of Cd<sup>2+</sup> distribution in the body, equilibrium constants dominate, e.g., during complex formation with serum albumin in the blood stream, and subsequent ligand exchange from serum albumin to metallothionein (MT) in the liver. It has been demonstrated, however, that the process of *in vitro* chelation of Cd<sup>2+</sup> out of the Cd-MT complex with EDTA is determined by the very slow rate of exchange, and not by the equilibrium (53). It is therefore of importance to consider rate constants for ligand exchange during the discussion of redistribution of Cd<sup>2+</sup> after Cd<sup>2+</sup> has been bound to MT.

Ligand exchanges are multistep reactions, and different mechanisms exist. The attacking ligand needs an initial coordination site on the complex. This can be achieved by solvent displacement or by displacing one of the coordination groups of the residential ligand. In the first case, the increased stability of the mixed complex will contribute to facilitate chelate ring opening. In the second case, the ring opening and subsequent displacement normally occur at a much slower rate. During the next step the attacking ligand must become chelated to decrease the rate of dissociation from the

mixed complex. Steric hindrance often decreases the exchange rate during reaction steps where several bulky groups simultaneously approach the metal ion. The kinetics of ligand exchange reactions have been extensively reviewed by Margerum et al. (52).

## Effects of Competing Species on Chelate Stability

In biological systems, the conditional or effective stability constants  $\beta'$  for metal chelates may be several orders of magnitude lower than indicated by the standard stability constants,  $\beta$  shown in the tables, since competition by other ligands decreases the concentration of free metal ion, and competition by other cations, e.g., the  $H^+$  ion and other metal ions, decreases the concentration of noncomplexed ligand available for chelation. The most important interfering ions in biological systems are normally the  $OH^-$  and  $H^+$  ions ( $\approx 10^7 M$ ) and the  $Ca^{2+}$  ion ( $\approx 10^{-3} M$ ). The term  $\alpha$  is used for the ratio of the sum of concentrations of all species of the ion involved in complex formation with a competing ligand over the concentration of the free compound available for chelation, and analogous for competing ions:  $\alpha_L(L) = \Sigma(L_i)$  and  $\alpha_M(M) = \Sigma(M_i)$ . For formation of the complex  $L_aM_b$  with one competing chelant and one competing ion present, the conditional stability constant,  $\beta'$  is:

$$\beta' = \frac{(L_aM_b)}{[\Sigma(L_i)]^a[\Sigma(M_i)]^b} = \frac{(L_aM_b)}{\alpha_L^a(L)^a\alpha_M^b(M)^b}$$

and

$$\beta' = \log \beta - a \log \alpha_L - b \log \alpha_M$$

Calculation of  $\beta'$  for chelation of  $Cd^{2+}$  with EDTA at pH = 7.4 with 2.5 mM  $Ca^{2+}$  present as in human serum (54) requires that four different equilibria be considered for the hydrogenated species of EDTA, giving

$$\alpha_{L/H} = \frac{(EDTA) + (HEDTA) + (H_2EDTA) + (H_3EDTA) + (H_4EDTA)}{(EDTA)}$$

$$= 1 + \beta_1(H) + \beta_2(H)^2 + \beta_3(H)^3 + \beta_4(H)^4$$

where the stability constants for the different hydrogenated forms of EDTA are:  $\log \beta_1 = 10.17$ ,  $\log \beta_2 = 6.11$ ,  $\log \beta_3 = 2.68$ ,  $\log \beta_4 = 2.0$  (47), and  $\log \alpha_{L/H} = 2.77$ . Four different hydroxyl complexes of  $Cd^{2+}$  should be considered, having  $\beta_1 = 4.1$ ,  $\beta_2 = 7.7$ ,  $\beta_3 = 10.3$  and  $\beta_4 = 12.0$  (47). The  $\alpha$  value can be calculated analogously to the  $\alpha_{L/H}$ -value:

$$\alpha_{M/OH} = \frac{1 + \beta_1(OH) + \beta_2(OH)^2 + \beta_3(OH)^3 + \beta_4(OH)^4}{\text{with } \log \alpha_{M/OH} \approx 0.}$$

Thus, competition by  $OH^-$  ions does not interfere with chelation reactions involving Cd at physiological pH.

There is only one EDTA species of  $Ca^{2+}$  competing for Cd, having  $\beta = 10.61$  (47). We can now calculate  $\alpha_{L/Ca} = 1 + \beta_1[Ca^{2+}]$  and by approximating  $[Ca^{2+}] \approx 2.5 \times 10^{-3}$  we find  $\log \alpha_{L/Ca} = 7.01$ .

For the CdEDTA complex,  $\beta = 16.36$  (47). At the conditions specified the value of  $\beta'$  can be calculated to  $16.36 - 2.77 - 7.01 = 6.58$  without taking into account mutual interactions between competing species: However, due to mutual interactions between ions the standard stability constants are not valid, and calculation of  $\alpha$ -values in order to determine conditional stability constants becomes highly speculative. In serum also  $Mg^{2+}$  ( $\approx 0.86$  mM) (54),  $Cu^{2+} + Cu^+$  ( $\approx 19$   $\mu M$ ) (54), and  $Zn^{2+}$  ( $\approx 46$   $\mu M$ ) (54) compete for  $Cd^{2+}$ . These metal ions are partially protein-bound, and exact knowledge about how available they are for competition is not present. Furthermore, chelation of  $Cd^{2+}$  with serum albumin during the initial phase after  $Cd^{2+}$  exposure and with metallothionein later will reduce the conditional stability constants for complexes of  $Cd^{2+}$  with chelating agents even more.

## Binding of $Cd^{2+}$ to Biological Ligands

*In vivo*,  $Cd^{2+}$  can bind to nucleic acids, or to proteins (Table 4). Several investigations have demonstrated that  $Cd^{2+}$  is not found associated with low molecular weight compounds like free amino acids, etc., to any appreciable extent, except maybe during a very brief period immediately after ingress of  $Cd^{2+}$ . Proteins offer much more favorable binding sites for  $Cd^{2+}$  than nucleic acids due to the soft binding sites formed by cysteine residues.

In nucleic acids, the phosphodiester groups offer negatively charged coordination sites for hard ions like  $Mg^{2+}$  and  $Ca^{2+}$ , leading to bonding with mainly ionic character, while nitrogen-containing sites on the bases represent donor sites favored by  $Cu^{2+}$ ,  $Cd^{2+}$  and  $Hg^{2+}$ . Intermediate ions like  $Co^{2+}$ ,  $Mn^{2+}$  and  $Zn^{2+}$  can bind to either the phosphates or the bases. The deoxyribose moieties, however, are not suited for complex formation with metal ions.

A few investigations have yielded information about complex formation between  $Cd^{2+}$  and free



bases, nucleosides, and nucleotides. With cytidine monophosphate  $\text{Cd}^{2+}$  forms a complex,  $[\text{Cd}(5'\text{CMP})(\text{H}_2\text{O})]_n$ , in which  $\text{Cd}^{2+}$  is pentacoordinate, binding to phosphate groups from three different 5'-CMP molecules, a water molecule and to N-3 of a fourth 5'-CMP molecule (55). With 5' - AMP and  $\text{NO}_3^-$ ,  $\text{Cd}^{2+}$  forms a complex  $[\text{5'AMP}\text{Cd}(\text{NO}_3)_2\cdot\text{H}_2\text{O}]_2^{2+}$ , in which two  $\text{Cd}^{2+}$  ions are octacoordinated and bind to N-3 and N-9 from two protonated adenine residues, arranged point symmetrically. Furthermore, the  $\text{Cd}^{2+}$  ions coordinate with the same two water molecules, and each  $\text{Cd}^{2+}$  ion coordinates with two of four  $\text{NO}_3^-$  ions. (15,56).

Less is known about binding of  $\text{Cd}^{2+}$  to DNA. In intact chromosomes from eukaryotic organisms, the DNA is present as a highly organized complex with a large number of different chromosomal proteins, histones as well as nonhistone proteins, the latter containing SH groups, which offer binding sites for  $\text{Cd}^{2+}$ . A very limited amount of information is available concerning interaction of  $\text{Cd}^{2+}$  with chromosomal proteins. *In vitro* exposure of human lymphocytes to  $\text{Cd}^{2+}$  can, however, inactivate the mitotic spindle (57,58), most likely by binding of  $\text{Cd}^{2+}$  to SH groups in spindle proteins. This inhibition of spindle function, which was also observed with  $\text{Hg}^{2+}$ , was counteracted by metallothionein induction after prolonged *in vitro* exposure to  $\text{Cd}^{2+}$  or  $\text{Hg}^{2+}$  (57,58). Sephadex G-75 chromatography of lysates of human lymphocyte cultures after 72 hr exposure to  $^{109}\text{Cd}$  *in vitro* demonstrated binding of Cd to MT (Fig. 1).

The affinity of  $\text{Cd}^{2+}$  for proteins parallels to some degree the size of the stability constants for the complexes between the  $\text{Cd}^{2+}$  ion and the amino acids in the proteins. However, both the carboxyl groups and the amino groups are modified by peptide bond formation. Due to the uniformity of the peptide bond, functional side groups in the amino acids become major determinants of the affinity of metal ions for proteins. Table 9 shows the stability constants for the complexes of several amino acids with  $\text{Cd}^{2+}$ . Only histidine and cysteine offer good binding sites for  $\text{Cd}^{2+}$  in proteins.  $\text{Cd}^{2+}$  can bind strongly to proteins if two SH groups are adjacent, allowing chelation. In metallothionein,  $\text{Cd}^{2+}$  is bound very strongly ( $\log \beta = 25.5$ ) (59). Optical studies (60) and  $^{113}\text{Cd}$ -NMR studies (61) have indicated that in the metallothionein molecule  $\text{Cd}^{2+}$  is tetrahedrally coordinated to four deprotonated cysteinyl side groups by thiolate bonds, and that adjacent  $\text{Cd}^{2+}$  ions are linked through single thiolate bridges.

This strong binding of  $\text{Cd}^{2+}$  to MT is in sharp




FIGURE 1. Sephadex G-75 chromatograms of lymphocyte lysates after *in vitro* exposure to  $^{109}\text{Cd}$ : ( $\Delta$ )  $5 \times 10^{-5}$  M  $\text{Cd}^{2+}$  4 hr before harvest; ( $\circ$ )  $5 \times 10^{-5}$  M  $\text{Cd}^{2+}$  72 hr before harvest; (1) elution volume of serum albumin; (2) elution volume of metallothionein; (3) elution volume of low molecular weight compounds.

Table 9. Stability of complexes between free amino acids and  $\text{Cd}^{2+}$ .<sup>a</sup>

Amino acid	ML/M-L $\log \beta_1$	ML <sub>2</sub> /M·L <sup>2</sup> $\log \beta_2$
Alanine	3.80b	7.10b
Valine	3.46b	6.46b
Leucine	3.84b	6.54b
Isoleucine	7.00b	8.80b
Proline	—	(8.0)d
Phenylalanine	3.87c	6.73c
Tryptophane	4.47c	8.18c
Methionine	3.67c	7.03c
Glycine	4.22c	7.69c
Serine	(4.84)d	(7.4)d
Threonine	(4.43)d	(7.2)d
Cysteine	—	(9.9)d
Tyrosine	3.57c	6.08c
Asparagine	—	(7.1–8.6)d
Glutamine	—	(7.4)d
Aspartic acid	4.39c	7.55c
Glutamic acid	3.90c	(7.9)d
Lysine	—	(5.8)d
Arginine	—	(6.7)d
Histidine	5.39c	9.66c

<sup>a</sup>Values from Martell and Smith (47), except as noted.

<sup>b</sup>25°C, 0.5.

<sup>c</sup>25°C, 0.1.

<sup>d</sup>Data from Squibb et al. (62).

contrast with the kinetics of ligand exchange reactions involving  $Zn^{2+}$ . Removal of Zn from MT by EDTA during pseudo first-order kinetics for EDTA showed triphasic kinetics with  $k_1$  fast,  $k_2 = 10^{-3} \text{ sec}^{-1}$  and  $K_3 = 10^{-4} \text{ sec}^{-1}$ , while the rate constant for removal of  $Cd^{2+}$  was approximately  $3 \times 10^{-6} \text{ sec}^{-1}$  (53), and the rate constant for transfer of zinc to apo-carbonic anhydrase was found to be about  $10^3$ , i.e., about the same rate as the reaction with free zinc, suggesting that a physiological role for MT could be acting as a zinc donor to zinc-requiring macromolecules (53). Due to the high degree of resistance of the Cd-MT complex against ligand exchange reactions, both favored by the large stability constant (59) and the sluggish reaction of the Cd-MT complex with competing ligands (53),  $Cd^{2+}$  is not likely to be removed from MT *in vivo* to any appreciable degree, except after translocation of the Cd-MT complex to the kidneys, where the MT moiety is broken down, and  $Cd^{2+}$  is rapidly reassociated with renal MT (63-65).

The efficiency of chelating agents for increasing renal excretion of  $Cd^{2+}$  during acute  $Cd^{2+}$  intoxication is sharply reduced concomitantly with cellular uptake of  $Cd^{2+}$ , mainly in the liver (51). Already 12 hr after administration of  $Cd^{2+}$ , only a limited amount of the dose is chelatable by antidotes like DTPA, EDTA and dimercaptosuccinic acid (66). It is, however, possible to chelate  $Cd^{2+}$  out of metallothionein and reduce the whole body burden as well as the concentration of  $Cd^{2+}$  in liver and kidneys by daily chelation therapy with BAL, initiated several days after cessation of Cd exposure (67). The main excretion route during BAL-induced mobilization of aged Cd deposits in the liver is fecal, due to increased biliary excretion (68).

## Effects of Chelation on $Cd^{2+}$ Toxicity

The possibilities of chelating agents for modifying the acute toxicity of  $Cd^{2+}$  depend on the chelating agent used, the molar ratio between the chelant and  $Cd^{2+}$ , the route of exposure and the time lapse between exposure to  $Cd^{2+}$  and to the chelating agent. Reported effects of chelating agents on acute  $Cd^{2+}$  toxicity include antagonistic effects, no apparent effect, and even enhancement of toxicity.

## Parenteral Exposure

Nitrilotriacetic acid (NTA) increased the acute toxicity of  $Cd^{2+}$  in rats after subcutaneous expo-

sure (69). Several studies have demonstrated that subcutaneous exposure of mice to  $Cd^{2+}$  administered as a complex with NTA (70) or sodium tripolyphosphate (STPP) (70,71) increased the acute toxicity of  $Cd^{2+}$ , as indicated by mortality within 24 hr. The reason for the enhanced toxicity is probably a high hepatotoxicity of these complexes of Cd, since extensive hepatic necrosis was observed at doses of  $Cd^{2+}$  (3.2 or 3.35 mg/kg body weight), exerting mild or no hepatotoxicity when administered alone (70,71). The liver damage induced by the Cd-STPP complex was characterized by extensive centrilobular necrosis seen at 24 hr (71). During a time-course study of the induction of hepatic necrosis after subcutaneous administration of the STPP-Cd complex to mice, Andersen et al. (71) investigated livers of mice killed 2, 4, 6, 8, 12 and 24 hr after administration of 30  $\mu\text{mole/kg}$  Cd with 90  $\mu\text{mole/kg}$  STPP and observed pronounced centrilobular blood stasis and early centrilobular necroses with fragmentation and lysis of nuclei and swollen hepatocytes already 6 and 8 hr after administration. This picture progressed to a more advanced state at 12 hr, and already before 16 hr, 2/10 animals were dead. At 24 hr, 7/10 animals were dead. In animals given the same dose of  $Cd^{2+}$  without STPP the liver histology was normal during the first 12 hr. At 24 hr all animals had survived, and 4/8 animals had normal liver histology, while four livers showed necrosis of single or several hepatocytes with eosinophile, swollen cells and with admixture of neutrophiles. Focal necroses were randomly localized. All control animals had normal livers, and the kidneys of all animals appeared normal. The events inducing liver necrosis during Cd-STPP exposure must take place very rapidly after administration, since the minimum period required for induction of hepatic necrosis is 6-8 hr. The enhancement of toxicity exerted by STPP is most likely due to induction of increased rate of interorgan distribution of  $Cd^{2+}$  (71): Figure 2 demonstrates that both the transport away from the injection site and the accumulation of  $Cd^{2+}$  in liver and kidneys take place much faster when  $Cd^{2+}$  is administered as a complex with STPP than when  $Cd^{2+}$  is administered alone.

Investigation of the binding of  $Cd^{2+}$  to macromolecules by Sephadex G-75 chromatography demonstrated that STPP delayed and decreased the binding of  $Cd^{2+}$  to metallothionein in the liver (Fig. 3). Thus, soon after exposure, when metallothionein induction has not yet taken place, a high concentration of  $Cd^{2+}$  is present in the liver, able to exert a full toxic potential. The decrease in




FIGURE 2. Effect of STPP on Cd transport and distribution among organs: (○) Cd, 30  $\mu\text{mole/kg}$ ; ( $\Delta$ ) Cd, 30  $\mu\text{mole/kg}$  and STPP, 90  $\mu\text{mole/kg}$ . Each point represents the average of determinations on four animals. The SEM is shown when larger than the height of the symbol.

the liver content of  $\text{Cd}^{2+}$  observed at later times most likely is due to leakage of  $\text{Cd}^{2+}$  from necrotic liver cells.

Sephadex G-75 chromatography (Fig. 4) of blood plasma, liver homogenate and kidney homogenate from animals killed 20 min after subcutaneous (SC) injection of  $\text{Cd}^{2+}$  or  $\text{Cd}^{2+}$  + STPP demonstrated that Cd in blood plasma was distributed in the MW area from serum albumin down to low molecular weight compounds. In the




FIGURE 3. Binding of Cd to proteins: (○) Cd, 30  $\mu\text{mole/kg}$ ; ( $\Delta$ ) Cd, 30  $\mu\text{mole/kg}$  and STPP, 90  $\mu\text{mole/kg}$ . From elution profiles like those shown in Fig. 4 the amount of  $^{109}\text{Cd}$  bound to MT was calculated as % of total  $^{109}\text{Cd}$  eluted from the column. The values obtained are shown versus time after injection.

liver,  $\text{Cd}^{2+}$  was solely found associated with high molecular weight proteins (fractions 12–14), while in kidneys most of the  $\text{Cd}^{2+}$  was associated with high molecular weight compounds, but a small amount eluted in the MT region (fraction 23–25). In liver and kidneys  $\text{Cd}^{2+}$  was not found associated with low molecular weight compounds. There was no difference between the elution pattern between animals injected with Cd and Cd + STPP. One hour after injection, similar elution patterns were observed for liver and kidney homogenates, while all  $^{109}\text{Cd}$  in plasma eluted with high molecular weight compounds, irrespectively of whether Cd or Cd + STPP had been injected. After Sephadex G-75 chromatography,  $^{109}\text{Cd}$  and  $^{109}\text{Cd}$  + STPP eluted in the low molecular weight area (= fraction 40).  $^{109}\text{Cd}$  + STPP mixed with serum albumin eluted in the high molecular weight area. The unexpected elution pattern of Cd in plasma observed 20 min after injection has been verified in several trials. It may be due to splitting of high molecular weight complexes during chromatography, leading to extensive trailing.

In a study of the effect of cysteine on the renal uptake of Cd(72), SC coinjection of  $\text{Cd}^{2+}$  and cysteine decreased the liver concentration of  $\text{Cd}^{2+}$  and increased the kidney concentration of  $\text{Cd}^{2+}$  at 0.5–7 hr as well as 24 hr after injection, as compared with the same dose of  $\text{Cd}^{2+}$  alone. In contrast to the enhancing effect of NTA and STPP on acute  $\text{Cd}^{2+}$  toxicity, SC exposure of mice to  $\text{Cd}^{2+}$




FIGURE 4. Distribution of  $^{109}\text{Cd}$  among protein fraction after Sephadex G-75 gel chromatography. Same doses as in Figs. 2 and 3.

as a complex with EDTA reduced the acute toxicity, as well as the liver and kidney concentrations of  $\text{Cd}^{2+}$  after 21 days, both when the  $\text{CdEDTA}$  and when the  $\text{Cd}_2\text{EDTA}$  complex was dominating, as compared with the same dose of  $\text{Cd}^{2+}$  administered alone (73). Suzuki and Yamamura (74) injected rats intraperitoneally with  $\text{Cd}^{2+}$ , uncomplexed or complexed with chelating agents with different affinity for  $\text{Cd}^{2+}$ , and found after 4 days that the higher the stability constant of the complex with  $\text{Cd}^{2+}$ , the larger the ratio between the kidney and liver concentrations of  $\text{Cd}^{2+}$ , thus suggesting that the stability constant of the complex would determine the interorgan distribution.

### Gastrointestinal Exposure

Eighteen months of chronic exposure of mice to  $\text{Cd}^{2+}$  (50 ppm in drinking water) administered as a complex with NTA (500 ppm), STPP (500 ppm) or EDTA (50 ppm) did not change the toxicity of  $\text{Cd}^{2+}$ , measured as mortality and pathological proteinuria, and only slight changes in the concentrations of  $\text{Cd}^{2+}$  in liver and kidneys were found, as compared with administration of  $\text{Cd}^{2+}$  alone (70). Chelation of  $\text{Cd}^{2+}$  with EDTA or BAL reduced the absorption of  $\text{Cd}^{2+}$  from ligated intestine after injection into duodenum in rats, as measured by  $^{109}\text{Cd}$  uptake into intestinal tissue. In addition, the liver and blood concentrations of  $\text{Cd}^{2+}$  were decreased. BAL reduced the amount of

$\text{Cd}^{2+}$  in the kidneys, while EDTA increased the amount of  $\text{Cd}^{2+}$  in kidneys, yielding a maximum at around 1.5 hr, while other much weaker chelating agents (cystein and glutathion) only marginally changed  $\text{Cd}^{2+}$  absorption (75).

The mortality in mice during 21 days, after administration of  $\text{Cd}^{2+}$ , either by stomach tube or by direct injection into the duodenum was significantly decreased when  $\text{Cd}^{2+}$  was given as the  $\text{CdEDTA}$  complex ( $\text{Cd}:\text{EDTA} = 1:4$ , molar ratio). However, when the predominating complex was  $\text{Cd}_2\text{EDTA}$  ( $\text{Cd}:\text{EDTA} = 1:0.4$ , molar ratio), the toxicity was identical to that exerted by the same dose of  $\text{Cd}^{2+}$  administered alone (73). The reduced toxicity of the  $\text{CdEDTA}$  complex was accompanied by a reduced liver and kidney burden of Cd. The reason the toxicity of the  $\text{Cd}_2\text{EDTA}$  complex approaches that of the  $\text{Cd}^{2+}$  ion is most likely that the  $\text{Cd}_2\text{EDTA}$  complex can split off one  $\text{Cd}^{2+}$  ion (73). The mortality during 3 weeks after a single dose of 60 mg/kg  $\text{Cd}^{2+}$  given by stomach tube was not changed when  $\text{Cd}^{2+}$  was administered as a complex with NTA (600 mg/kg) but was totally abolished when  $\text{Cd}^{2+}$  was complexed with STPP (600 mg/kg) (76).

### Translocation of $\text{Cd}^{2+}$ from Liver to Kidneys

In experimental animals, the liver has been found to contain the largest amount of  $\text{Cd}^{2+}$  dur-

ing chronic Cd<sup>2+</sup> exposure although the kidneys after prolonged exposure eventually may contain the largest concentration of Cd<sup>2+</sup> (77,78). The Cd<sup>2+</sup> deposit in kidneys is stable and urinary Cd<sup>2+</sup> excretion is low before tubular damage has been induced (79,80). Thus, Cd<sup>2+</sup> concentrations of the order of 200–300 mg/kg in kidneys may be reached, and such concentrations have been suggested as the critical concentration for induction of kidney dysfunction (6,81–84). After induction of tubular damage, urinary Cd<sup>2+</sup> rises sharply (85), and Cd<sup>2+</sup> bound to MT is excreted into the urine (86,87). During the period before induction of tubular damage the interorgan distribution of Cd<sup>2+</sup> is mainly due to slow translocation of Cd–MT (and the small amount of non–MT bound Cd<sup>2+</sup>) from the liver to the kidneys (78). The retention half-life of a single dose of Cd<sup>2+</sup> in rats has been measured to be 73 days in the liver and at least 1240 days in the kidneys (88). Most likely this translocation involves transport of liver Cd–MT complex, because circulating Cd–MT complex has been demonstrated in the plasma of mice and rats after prolonged Cd<sup>2+</sup> exposure (86,87,89–94), and MT has been demonstrated in the plasma of rats already 1 day after a single dose of Cd<sup>2+</sup>, by the use of a radioimmunoassay (95). MT has also been demonstrated in plasma and urine of Cd exposed humans (94,96–100).

Parenterally administered Cd–MT complex is preferentially absorbed by renal tubules (101–104) and exert a much higher renal toxicity than Cd<sup>2+</sup> administered alone (79,103,105–107). After glomerular filtration and tubular resorption, the Cd–MT complex is rapidly degraded, and Cd<sup>2+</sup> is re-incorporated into renal MT (63–65,108). During the period of slow translocation of Cd<sup>2+</sup> from liver to kidneys, biliary excretion seems to contribute considerably to Cd<sup>2+</sup> detoxification (109).

Nordberg et al. (110) concluded that, after long-term Cd<sup>2+</sup> exposure, the largest amount of soluble Cd<sup>2+</sup> in mouse blood cell hemolysate was Cd–MT complex in erythrocytes. *In vitro* exposure of human lymphocytes to <sup>109</sup>Cd<sup>2+</sup> showed that induction of MT synthesis and Cd–MT complex formation occur in lymphocyte cytoplasm (57,111). Quantitation of Cd<sup>2+</sup> in plasma as well as in red and white blood cells, 60 hr after a single dose of Cd<sup>2+</sup> to rats, showed that, although the highest concentration of Cd<sup>2+</sup> was found in white blood cells, the most important contribution to the blood content of Cd<sup>2+</sup> was Cd<sup>2+</sup> in erythrocytes, mainly bound to cell membranes. Cd<sup>+</sup> in the erythrocyte cytosol was partly bound to high molecular weight proteins, partly to a protein with MW like MT (112).

Table 10. Effect of dietary PCB exposure on Cd transition from liver to kidneys.

Expt. No.	PCB in diet, ppm	Group	<sup>109</sup> Cd, % of initial dose ± SD (p) <sup>a</sup>	
			Liver	Kidneys
1 <sup>b</sup>	0	1 (n=4)	50.27 ± 1.72	8.7 ± 1.10
		2	47.42 ± 3.84	10.80 ± 0.92 (p<0.005)
	10	3 (n=5)	50.14 ± 3.59 (ns)	9.15 ± 0.65 (ns)
		4	49.18 ± 0.93 (ns)	11.30 ± 1.77 (p=0.01)
	25	5 (n=8)	48 ± 2.33 (p<0.025)	11.31 ± 1.49 (p<0.0025)
		6 (n=5)	46.36 ± 2.36 (p<0.01)	11.66 ± 1.21 (p<0.0025)
2 <sup>c</sup>	0	7 (n=6)	17.74 ± 4.03	29.76 ± 4.0
		8	15.43 ± 2.72 (ns)	26.61 ± 3.63 (ns)
	10	9 (n=5)	14.25 ± 4.53 (ns)	27.28 ± 1.22 (ns)
		10	10.53 ± 2.65 (p<0.0025)	26.82 ± 5.25 (ns)
	25	11 (n=5)	11.42 ± 4.52 (p<0.05)	30.54 ± 3.84 (ns)
		12 (n=7)	9.02 ± 2.46 (p<0.0005)	32.48 ± 3.49 (ns)

<sup>a</sup>Student's *t*-test, two-sided.

<sup>b</sup><sup>109</sup>Cd dose, SC; 10 μmole/kg; 10 μCi/animal; animals killed after 15 days.

<sup>c</sup><sup>109</sup>Cd dose, SC; 0.1 μmole/kg; 1 μCi/animal; animals killed after 15 days.

## Effect of Hepatic Damage on Cd<sup>2+</sup> Metabolism

Due to the large deposit of Cd<sup>2+</sup> present in the liver during Cd<sup>2+</sup> exposure, it is likely that hepatic disorders may alter the tissue distribution of Cd<sup>2+</sup>. This has been verified in a few investigations where acute liver damage resulted in rapid and substantial translocation of aged Cd<sup>2+</sup> deposits from liver to kidneys. Induction of hepatic necrosis by aflatoxin B<sub>1</sub> lead to translocation of Cd<sup>2+</sup> administered 1 month earlier from liver to kidneys. Approximately 40% of the decreased liver burden of Cd<sup>2+</sup> was recovered as increased kidney burden, most likely transferred as the MT–Cd complex (113). Intragastric administration of 1 mL/kg CCl<sub>4</sub> to rats at least 2 weeks after the end of exposure to 200 pm Cd<sup>2+</sup> in the drinking water for 1 or 4 months also induced a decrease in the liver content of Cd<sup>2+</sup>, paralleled with recovery of some of the Cd<sup>2+</sup> in the kidneys. In this study a strongly increased urinary excretion of Cd<sup>2+</sup> as well as proteinuria was also in-

duced by  $\text{CCl}_4$  (114). The proteinuria was largest in the group of rats exposed to  $\text{Cd}^{2+}$  for 4 months.

In a similar study, aged  $\text{Cd}^{2+}$  deposits in the liver of rats were also mobilized by hepatic necrosis induced by 2.5 mL/kg of  $\text{CCl}_4$  administered 1 day after cessation of 1–8 weeks exposure to repeated SC injections of  $\text{Cd}^{2+}$ . The hepatic concentration of  $\text{Cd}^{2+}$  was almost reduced by 50% 24 hr after  $\text{CCl}_4$  exposure. Highly increased plasma and urinary concentrations of  $\text{Cd}^{2+}$  were found, and Sephadex G-75 chromatography indicated that the Cd-MT complex accounted for most of the  $\text{Cd}^{2+}$  present in plasma and urine. The kidney concentration of  $\text{Cd}^{2+}$  was also increased. Only a small part of the mobilized liver  $\text{Cd}^{2+}$  content was, however, recovered in kidneys. The  $\text{Cd}^{2+}$  content in other organs (spleen, small intestine, pancreas) and in feces was not increased, compared with control animals (115,116). Since  $\text{CCl}_4$  is nephrotoxic and induces proximal tubular damage (117), evaluation of these studies in relation to the toxicity of Cd-MT to the proximal renal tubule is difficult.

Little is known about effects of chronic liver damage on  $\text{Cd}^{2+}$  translocation. In an unpublished study, Andersen et al. (118) found that the rate of translocation of  $\text{Cd}^{2+}$  from liver to kidneys was increased in mice with hepatic necrosis induced by chronic PCB exposure. Mice fed diets containing 0, 10, 25, 50, 100 or 200 ppm PCB were given a single SC dose of  $^{109}\text{Cd}$  (see Table 10). Dietary PCB exposure for 36 weeks induced dose-dependent pathological changes. In the group fed 200 ppm PCB centrilobular liver cell enlargement with increased diameter of both cytoplasm and nuclei was seen, as well as focal necrosis with Kupffer cell proliferation. Some confluent necroses, a few macroscopically visible, were also observed. In mice killed 2 weeks after a single SC dose of  $^{109}\text{Cd}^{2+}$ , slightly lower amounts of  $\text{Cd}^{2+}$  were present in the liver and slightly larger amounts were found in kidneys in animals exposed to 100 and 200 ppm PCB, as compared to control animals. Fifteen weeks after SC  $^{109}\text{Cd}^{2+}$  exposure, a dose-dependent, highly significant reduction in hepatic  $\text{Cd}^{2+}$  content was observed; this decrease was, however, not followed by a comparable increase in the kidney content of  $\text{Cd}^{2+}$  (Table 10). Thus, during the presumably very slightly increased rate of translocation of  $\text{Cd}^{2+}$  due to PCB-induced chronic liver damage, some of the mobilized  $\text{Cd}^{2+}$  seem to be excreted, probably in the urine. Whole body counting of the animals could not demonstrate PCB-induced changes in the rate of excretion of SC dose of  $^{109}\text{Cd}^{2+}$  during 15 weeks. The difference in liver +

kidney burden of  $\text{Cd}^{2+}$  between the control group and the group exposed to 200 ppm PCB at 15 weeks is, however, only about 6% of the initial dose (Table 10), thereby explaining why a possible slight increase in detoxification rate was not observed.

So far the effect of acute or chronic liver damage on the renal toxicity of chronic low level  $\text{Cd}^{2+}$  exposure is virtually unknown. The studies reviewed here indicate that hepatic  $\text{Cd}^{2+}$  mobilized due to acute or chronic liver damage is not quantitatively recovered in the kidneys. The possibility does, however, exist that humans with liver damage, e.g., due to excessive alcohol consumption have an increased risk for tubular kidney damage if they are exposed to  $\text{Cd}^{2+}$ . Furthermore, people with a high liver burden of  $\text{Cd}^{2+}$  due to former or present  $\text{Cd}^{2+}$  exposure may be at risk for kidney damage if acute liver damage occurs, e.g., due to hepatitis. Further experimental investigation of this aspect of  $\text{Cd}^{2+}$  toxicodynamics is needed.

## Chelation Therapy

According to recent reviews (51,119,120) only a limited knowledge is presently available concerning the possibility for chelate treatment of  $\text{Cd}^{2+}$  intoxication in humans. Although BAL and EDTA increase the renal excretion of  $\text{Cd}^{2+}$  in experimental animals, both antidotes have been reported also to increase the renal injury produced by  $\text{Cd}^{2+}$  (121,122). Intravenous administration of  $\text{CaNa}_2\text{-EDTA}$  (which is superior to the sodium salt of EDTA, which may induce hypocalcemic tetany as a complicating side effect) increases the renal excretion of zinc considerably, and the excretion of iron and copper slightly (119). Extensive destruction of proximal tubular epithelium has been observed in the kidneys of humans and experimental animals after EDTA treatment (123–125). Due to the mobilizing effect on aged  $\text{Cd}^{2+}$  deposits, chelation therapy of acute  $\text{Cd}^{2+}$  intoxication is contraindicated in humans with any significant previous  $\text{Cd}^{2+}$  exposure (6,119).

In order to maximize the efficiency of chelation therapy against a toxic metal ion and to minimize the unwanted excretion of essential metal ions (i.e., to obtain a high value of  $\beta'$ , the conditional stability constant for the complex between the toxic ion and the selected chelating antidote), knowledge about the stabilities of complexes with competing metal ions is necessary. In Table 11 the stability constants for complexes between the most frequently used antidotes and competing

**Table 11. Stabilities of complexes between important chelating agents and Cd<sup>2+</sup> or other biologically important ions: M + L → ML.<sup>a</sup>**

Compound	log β				
	Ca <sup>2+</sup>	Mg <sup>2+</sup>	Cu <sup>2+</sup>	Zn <sup>2+</sup>	Cd <sup>2+</sup>
Sodium tripolyphosphate (STPP)	5.20	5.76	8.3	7.5	6.58
2,3-Dimercaptopropanol (BAL)	—	—	—	13.5	—
Penicillamine (PEN) <sup>b</sup>	—	—	19.5	9.40	10.8
Nitrilotriacetic acid (NTA)	6.39	5.47	12.94	10.66	9.78
Ethylenediaminetetraacetic acid (EDTA)	10.61	8.83	18.70	16.44	16.36
Diethylenetriaminetetraacetic acid (DTPA)	10.57	9.34	21.38	18.29	19.0
Ethylene bis(oxyethylenetriolo)tetraacetic acid (EGTA)	10.86	5.28	17.57	12.6	16.5

<sup>a</sup>Data from Martell and Smith, 25°C, 0.1, except as noted.

<sup>b</sup>20°C, 0.1.

metal ions are listed. DTPA, which has proven superior as an antagonist toward the acute toxicity of Cd<sup>2+</sup> (126,127), forms a complex with Cd<sup>2+</sup> several orders of magnitude more stable than EDTA, while STPP and NTA, which in fact increase the acute toxicity of parenterally administered Cd<sup>2+</sup> (69–71) and also PEN, which is without beneficial effects on acute Cd<sup>2+</sup> toxicity (126,127), form much weaker complexes with Cd<sup>2+</sup>.

As indicated above, mobilization of aged Cd<sup>2+</sup> deposits in the liver by chelating agents may increase the risk for development of tubular damage. Cherian and co-workers (50,68) and Von Burg and Smith (128) have demonstrated that renal excretion of Cd<sup>2+</sup> can be circumvented by the use of BAL or structurally related di- and trimercaptans, which increase the biliary excretion of Cd<sup>2+</sup>. The requirement for induction of efficient detoxification of the Cd-MT complex in liver seems to be a relatively lipophilic chelating agent with two adjacent SH groups. It is presently not known whether these interesting results pertain to the human situation, since the high toxicity of this kind of compound in laboratory animals seems to contraindicate extensive human use.

Most published experiments evaluating antagonizing effects of chelating agents on acute Cd<sup>2+</sup> toxicity have been performed by administration of Cd<sup>2+</sup> to experimental animals by the intravenous or other parenteral routes. Accidental acute Cd<sup>2+</sup> intoxication in humans is, however, always due to either gastrointestinal or pulmonary exposure. Effects of chelating agents on acute Cd<sup>2+</sup> toxicity after administration via these routes are very poorly investigated, and a very limited knowledge is available.

Future experiments investigating effects of chelating agents on acute Cd<sup>2+</sup> toxicity, biliary and urinary excretion and organ distribution of Cd<sup>2+</sup> after administration via these routes are of great importance for developing chelation ther-

apy regimens applicable to treatment of acute human Cd<sup>2+</sup> poisoning.

The author wants to thank Professor Gunnar Nordberg for valuable help and discussions during the experiments reported as part of this review. Thanks are due to Ranja Bjerring for excellent technical assistance and to Inge Bøgelund and Yrsa Kildeberg for typing the manuscript.

#### REFERENCES

1. Friberg, L., Piscator, M., Nordberg, G. F., and Kjellström, T. Cadmium in the Environment, 2nd ed. CRC Press, Cleveland, 1974.
2. NIOSH. Criteria for a Recommended Standard. Occupational Exposure to Cadmium. HEW No 76-192. National Institute for Occupational Safety and Health, U.S. Department of Health, Education and Welfare, Rockville, MD, 1976.
3. CEC. Evaluation of Impact of Cadmium on the Health of Man. Commission of the European Communities, Luxembourg, 1977.
4. WHO. Environmental Health Criteria for Cadmium. Interim Report. World Health Organization, Geneva, 1979.
5. EPA. Health Assessment Document for Cadmium. EPA-600/8-79-003. U.S. Environmental Protection Agency, Research Triangle Park, N.C., 1979.
6. Friberg, L., Kjellström, T., Nordberg, G. F., and Piscator, M. Cadmium. In: Handbook on the Toxicology of Metals (L. Friberg, G. F. Nordberg and V. B. Vouk, Eds.), Elsevier-North Holland, Amsterdam, 1979, pp. 355–381.
7. Webb, M., Ed. The Chemistry, Biochemistry and Biology of Cadmium. Elsevier North Holland, Amsterdam, 1979.
8. WHO. Health-based Permissible Levels for Occupational Exposure to Cadmium. World Health Organization, Geneva, 1980.
9. Kägi, J. H. R., and Nordberg, M., Eds. Metallothionein. Birkhäuser Verlag, Basel, 1979.
10. Foulkes, E. C., Ed. Biological roles of metallothionein. In: Developments in Toxicology and Environmental Sciences, Vol. 9. Elsevier-North Holland, Amsterdam, 1982.
11. Martell, A. E. Chemistry of carcinogenic metals. Environ. Health Perspect. 40: 207–226 (1981).
12. Schubert, J. Chelating agents in biological systems. Environ. Health Perspect. 40: 227–232 (1981).
13. Nieboer, E., and Richardson, D. H. S. The replacement of the nondescript term "heavy metals" by a biologically and chemically significant classification of metal ions. Environ. Pollut. 1: 3–26 (1980).
14. Tuck, D. G. The stereochemistry of the coordination

- compounds of cadmium(II). *Rev. Inorg. Chem.* 1: 209–243 (1979).
- Jacobson, K. B., and Turner, J. E. The interaction of cadmium and certain other metal ions with proteins and nucleic acids. *Toxicology* 16: 1–37 (1980).
  - Jørgensen, C. K. Electric polarizability, innocent ligands and spectroscopic oxidation states. *Struct. Bond.* 1: 234–248 (1966).
  - Akishin, P. A., Spiridonov, V. P., Naumov, V. A., and Rambidi, N. G. Electronographic investigation of molecular structure III. Cadmium halides. *Zhur. Fiz. Khim.* 30: 155–160 (1956).
  - Davies, J. E. D., and Long, D. A. Vibrational spectra of the halides and complex halides of the group IIB elements. The vibrational spectra of  $\text{CdI}_3^-$ ,  $\text{CdBr}_2$ ,  $\text{CdBr}_3^-$ ,  $\text{CdCl}_2$ ,  $\text{CdCl}_3^-$ ,  $\text{CdCl}_4^{2-}$  and  $\text{MgCl}_4^{2-}$ . *J. Chem. Soc. A:* 2054–2058.
  - Calvo, C., and Stephens, J. S. Crystal structure of  $\text{CdZn}_2(\text{PO}_4)_2$  and  $\text{Cd}_2\text{Zn}(\text{PO}_4)_2$ . *Can. J. Chem.* 46: 903–915 (1968).
  - Iwamoto, T. Crystal structure of catena- $\mu$ -ethylenediamine cadmium(II), tetracyano cadmate(II) dibenzene chlateate  $\text{Cd}(\text{en})\text{Cd}(\text{CN})_4 \cdot 2\text{C}_6\text{H}_6$ . *Chem. Letters* 7: 723–726 (1973).
  - Pecs, L., Ozols, J., Ievins, A., and Sturis, A. X-ray diffraction study of cadmium 2-methyl-8-mercaptoquinolate (thiooxinate) ( $\text{Cd}(\text{C}_9\text{H}_8\text{CH}_3\text{N})_2$ ) crystals. *Latv. PSR Zinat. Akad. Vestis, Kim. Ser.* 6: 745–746 (1975).
  - Nardelli, M., Cavalca, L., and Braibanti, A. Structure of bis(thiourea) cadmium chloride. *Gazz. Chim. Ital.* 87: 137–145 (1957).
  - Iimura, Y., Ito, T., and Hagihara, H. Crystal structure of cadmium ethylxanthate. *Acta Cryst. B*28: 2271–2279 (1972).
  - Agre, V. M., Shugam, E. A., and Rukhadze, E. G. Structure of intercomplex complex compounds with the metal-sulfur bonds. I. Crystallographic data for some alkylthiocarbamates of bivalent metals. *Zhur. Strukt. Khim.* 7: 897–898 (1966).
  - Bürgi, H. B., Stereochemistry of polynuclear cadmium(II) thioglycolates. Crystal structure of cadmium(II) bithioglycolate. *Helv. Chim. Acta* 57: 513–519 (1974).
  - Ohtaki, M., and Maeda, M. X-ray diffraction studies on the structures of the hydrated cadmium and tetraiodo-cadmate ions in aqueous solution. *Proc. Int. Conf. Coord. Chem.* 16: 2.37a; *Chem. Abstr.* 85: 39601b (1974).
  - Maslen, E. N., Greaney, T. M., Raston, C. L., and White, A. H. Crystal structure of catena-di- $\mu$ -acetylacetonato-cadmium(II). *J. Chem. Soc. Dalton Trans.* 5: 400–402 (1975).
  - Antti, B.-M., Lundberg, B. K. S., and Ingri, N. J. X-ray crystal structure of hexakis(imidazole)cadmium(II) carbonate pentahydrate, a supposed hydroxynitrate. *J. Chem. Soc. Chem. Commun.* 1972: 712–713 (1972).
  - Cannas, M., Carta, G., Cristini, A., and Marongiu, G. J. Three-co-ordinate thiocyanate in cadmium dithiocyanate. *Chem. Soc. Dalton Trans* 4: 300–301 (1976).
  - Nardelli, M., Fava, G., and Boldrini, P. Complexes with sulfur bridges: structure of bis(thiourea)-cadmium formate. *Gazz. Chim. Ital.* 92: 1392–1400 (1962).
  - Flook, R. J., Freeman, H. C., Moore, C. J., and Scudder, M. L. Model compounds for metal-protein interaction. Crystal structures of seven cadmium(II) complexes of amino acid and peptides. *J. Chem. Soc. Chem. Commun.* 19: 753–754 (1973).
  - Freeman, H. C., Huq, F. and Stevens, G. N. Metal binding by D-penicillamine: crystal structure of D-penicillaminatocadmium(II) hydrate. *J. Chem. Soc. Chem. Commun.* 3: 90–91 (1976).
  - Harrison, W., and Trotter, J. J. Crystal and molecular structure of cadmium diacetate dihydrate. *Chem. Soc. Dalton Trans.* 8/9: 956–960 (1972).
  - Post, M. L., and Trotter, J. J. Crystal and molecular structure of cadmium(II)-maleate dihydrate. *Chem. Soc. Dalton Trans.* 7: 674–678 (1974).
  - Matkovic, B., Ribar, B., Zelenco, B., and Peterson, S. W. Refinement of the structure of  $\text{Cd}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$ . *Acta Cryst.* 21: 719–725 (1966).
  - Caira, M. R., Nassimbeni, L. R., and Orpen, G. The crystal structure of *cis*-catena- $\mu$ -sulphato-aquotris(imidazole)cadmium(II). *Acta Cryst. B*32: 140–144 (1976).
  - Schwarzenbach, G. Organic complex-forming compounds. *Experientia. (Suppl.)* 5: 162–192 (1956).
  - Ahrland, S., Chatt, J., and Davies, N. R. The relative affinities of ligand atoms for acceptor molecules and ions. *Quant. Rev. Chem. Soc.* 12: 265–276 (1958).
  - Pearson, R. G., Ed. *Hard and Soft Acid and Bases*. Dowden, Hutchinson and Ross, Stroudsburg, PA, 1973.
  - Craig, D. P., and Nyholm, R. S. Nature of the metal-ligand bond. In: *Chelating Agents and Metal Chelates* (F. P. Dwyer and D. P. Mellor, Eds.), Academic Press, New York, 1964, pp. 60–71.
  - Williams, R. J. P., and Hale, J. D. The classification of acceptors and donors in inorganic reactions. *Struct. Bond.* 1: 249–281 (1966).
  - Pearson, R. G., and Mawby, R. J. *Halogen Chemistry, Vol. 3* (V. Gutman, Ed.), Academic Press, New York 1965, p. 59.
  - Klopman, G. Chemical reactivity and the concept of charge- and frontier-controlled reactions. *J. Am. Chem. Soc.* 90: 223–234 (1968).
  - Ahrland, S. Scales of softness for acceptors and donors. *Chem. Phys. Letters* 2: 303–306 (1968).
  - Ahrland, S. Thermodynamics of complex formation between hard and soft acceptors and donors. *Struct. Bond.* 5: 118–149 (1968).
  - Weast, R. C., Ed. *CRC Handbook of Chemistry and Physics*. CRC Press, Cleveland, 1977–1978.
  - Martell, A. E., and Smith, R. E., Eds. *Critical Stability Constants Vols. 1–4*. Plenum Press, New York, 1974–1977.
  - Anderegg, G. Multidentate ligands. In: *Coordination Chemistry, Vol. 1*, A. E. Martell, Ed. ACS Monograph 168, Van Nostrand-Reinhold, New York 1971, pp. 427–490.
  - Schwarzenbach, G. The chelate effect. *Helv. Chim. Acta* 35: 2344–2359 (1952).
  - Cherian, M. G. Chelation of cadmium without increased renal deposition. *Environ. Health Perspect.* 54: 243–248 (1984).
  - Klaassen, C. D., Waalkes, M. P., and Cantilena, L. R., Jr. Alteration of tissue deposition of cadmium by chelating agents. *Environ. Health Perspect.* 54: 233–242 (1984).
  - Margerum, D. W., Cayley, G. R., Weatherburn, D. C., and Pagenkopf, G. K., Kinetics and mechanisms of complex formation and ligand exchange. In: *Coordination Chemistry, Vol. 2* (A. E. Martell, Ed.), ACS Monograph 174, American Chemical Society, Washington, 1978, pp. 1–220.
  - Li, T.-Y., Kraker, A. J., Shaw, C. F., III, and Petering, D. H. Ligand substitution reactions of metallothioneins with EDTA and apo-carbonic anhydrase. *Proc. Natl. Acad. Sci. (U.S.)* 77: 6334–6338 (1980).
  - Altman, P. L., and Dittmer, D. S., Eds. *Blood and Other Body Fluids*. Fed. Am. Soc. Exp. Biol., 1971, p. 22.



55. Goodgame, D. M. L., Jeeves, I., Reynolds, C. D., and Skapski, A. C. Heavy metal-pyrimidine nucleotide interaction. X-ray structure of a cadmium derivative of cytidine-5 monophosphate. *Biochem. J.* 151: 467-468 (1975).
56. Wei, C. H., and Jacobson, K. B. Paper presented at Am. Cryst. Assn. Meet., March 1980, Lakeport Convention Center, Ala.
57. Andersen, O., Rønne, M., and Nordberg, G. F. Effects of inorganic metal salts on chromosome length in human lymphocytes. *Hereditas* 98: 65-70 (1983).
58. Andersen, O., and Rønne, M. Quantitation of spindle inhibiting effects of metal compounds by chromosome length measurements. *Hereditas* 98: 215-218 (1983).
59. Kägi, J. H., and Vallee, B. L. Metallothionein: a cadmium and zinc-containing protein from equine renal cortex. II. Physico-chemical properties. *J. Biol. Chem.* 236: 2435-2442 (1961).
60. Vasak, M., Kägi, H. R., and Hill, H. A. O. Zinc (II), cadmium (II) and mercury (II) thiolate transitions in metallothionein. *Biochemistry* 20: 2852-2856 (1981).
61. Otvos, J. D., and Armitage, I. M. Structure of metal clusters in rabbit liver metallothionein. *Proc. Natl. Acad. Sci. (U.S.)* 77: 7094-7098 (1980).
62. Sillén, L. G., and Martell, A. E. Stability Constants of Metal Ion Complexes. Spec. Publ. No. 25, The Chemical Society, London, 1971.
63. Shaikh, Z. A. The low molecular weight cadmium-, mercury- and zinc-binding proteins (metallothioneins): biosynthesis, metabolism, and possible role in metal toxicity. In: *Metallothionein* (J. H. R. Kägi and M. Nordberg, Eds.), Birkhäuser Verlag, Basel, 1979, pp. 331-336.
64. Webb, M. The metallothioneins. In: *The Chemistry, Biochemistry and Biology of Cadmium* (M. Webb, Ed.), Elsevier-North Holland, Amsterdam, 1980.
65. Squibb, K. S., Pritchard, J. B., and Fowler, B. A. Renal metabolism and toxicity of metallothionein. In: *Biological Roles of Metallothionein* (E. C. Foulkes, Ed), Toxicology and Environmental Sciences, Vol. 9, Elsevier-North Holland, Amsterdam, 1982, pp. 181-192.
66. Cantilena, L. R., and Klaassen, C. D. Decreased effectiveness of chelation therapy with time after acute cadmium poisoning. *Toxicol. Appl. Pharmacol.* 63: 173-180 (1982).
67. Cherian, M. G. and Rodgers, K. Chelation of cadmium from metallothionein *in vivo* and its excretion in rats repeatedly injected with cadmium chloride. *J. Pharmacol. Exptl. Therap.* 222: 699-704 (1982).
68. Cherian, M. G., Onosaka, S., Carson, G. K., and Dean, P. A. W. Biliary excretion of cadmium in rats. V. Effects of structurally related mercaptols on chelation of cadmium from metallothionein. *J. Toxicol. Environ. Health* 9: 389-399 (1982).
69. Chernoff, N., and Courtney, K. D. Maternal and fetal effects of NTA, NTA and Cadmium, NTA and Mercury, NTA and nutritional imbalance in mice and rats. *Natl. Inst. Environ. Health Sci. Progr. Rept.*, Dec. 1, 1970 (Jan. 18, 1971).
70. Engström, B., and Nordberg, G. F. Effects of detergent formula chelating agents on the metabolism and toxicity of cadmium in mice. *Acta Pharmacol. Toxicol* 43: 387-397 (1978).
71. Andersen, O., Hägerstrand, I., and Nordberg, G. F. Effect of the chelating agent sodium tripolyphosphate on cadmium toxicity in mice. *Environ. Res.* 29: 54-61 (1982).
72. Mucakemi, M., and Webb, M. A morphological and biochemical study of the effects of L-cysteine on the renal uptake and nephrotoxicity of cadmium. *Brit. J. Exptl. Pathol.* 62: 115-130 (1981).
73. Engström, B., Norin, H., Jawaid, M., and Ingram, F. Influence of different Cd-EDTA complexes on distribution and toxicity of cadmium in mice on oral or parenteral administration. *Acta Pharmacol. Toxicol.* 46: 219-234 (1980).
74. Suzuki, K. T., and Yamamura, M. Distribution of cadmium in liver and kidneys by loading of various Cd-complexes and relative metal ratios in the induced metallothioneins. *Biomed. Pharmacol.* 28: 3643-3649 (1979).
75. McGivern, J., and Mason, I. The effect of chelation on the absorption of cadmium from rat intestine *in vivo*. *J. Comp. Pathol.* 89: 293-300 (1979).
76. Engström, B. Influence of chelating agents on toxicity and distribution of cadmium among proteins of mouse liver and kidney following oral or subcutaneous exposure. *Acta Pharmacol. Toxicol.* 48: 108-117 (1981).
77. Gunn, S. A., and Gould, T. C. Selective accumulation of <sup>115</sup>Cd by cortex of rat kidney. *Proc. Soc. Exptl. Biol. Med.* 96: 820-833 (1957).
78. Frazier, J. M. The role of metallothionein in the systemic distribution of cadmium. In: *Biological Roles of Metallothionein* (E. C. Foulkes, Ed.), Elsevier-North Holland, Amsterdam, 1982, pp. 141-153.
79. Nordberg, G. F. Cadmium metabolism and toxicity. *Environ. Physiol. Biochem.* 2: 7-36 (1972).
80. Bernard, A., Goret, A., Buchet, J. P., Roels, H., and Lauwerys, R. Significance of cadmium levels in blood and urine during longterm exposure of rats to cadmium. *J. Toxicol. Environ. Health.* 6: 157-166 (1980).
81. WHO. Task Group on Environmental Health. Criteria for Cadmium. Geneva, 1977.
82. Nordberg, G. F. Effects and dose-response relationships of Toxic metals. Elsevier, Amsterdam, 1976, pp. 64-72.
83. Roels, H., Bernard, A., Buchet, J. P., Goret, A., Lauwerys, R., Chettle, D. R., Harvey, T. C., and A. Haddad, I. Critical concentration of cadmium in renal cortex and urine. *Lancet* i: 221 (1979).
84. Lauwerys, R., Bernard, A., Buchet, J. P., and Roels, H. Dose-response relationships for the nephrotoxic action of cadmium in man. In: *International Conference on the Management and Control of Heavy Metals in the Environment*, London, 1979, pp. 19-23.
85. Friberg, L. Further investigations on chronic cadmium poisoning. A study on rabbits with radioactive cadmium. *Arch. Ind. Hyg. Occup. Med.* 5: 30-36 (1952).
86. Shaikh, Z. A., and Hirayama, K. Metallothionein in the extracellular fluids as an index of cadmium toxicity. *Environ. Health Perspect.* 28: 267-271 (1979).
87. Suzuki, Y. Metal-binding properties of metallothionein in extracellular fluids and its role in cadmium-exposed rats. In: *Biological Roles of Metallothionein* (E. C. Foulkes, Ed.), Elsevier-North Holland, Amsterdam, 1982, pp. 25-35.
88. Shaikh, Z. A. Metallothionein as a storage protein for cadmium: its toxicological implications. In: *Biological Roles of Metallothionein* (E. C. Foulkes, Ed.), Elsevier-North Holland, Amsterdam, 1982, pp. 69-76.
89. Nordberg, G. F., Piscator, M., and Nordberg, M. On the distribution of cadmium in blood. *Acta Pharmacol.* 30: 289-295 (1971).
90. Goyer, R. A., Cherian, M. G., and Delaquerriere-Richardson, L. Renal effects of cadmium. In: *Proceedings of the First International Cadmium Conference*, San Francisco. Metal Bulletin, Ltd., London, 1978, pp. 183-185.
91. Cherian, M. G., Goyer, R. A., and Delaquerriere-Richardson, L. Relationship between plasma cadmium-thionein and cadmium-induced nephropathy. *Toxicol. Appl. Pharmacol.* 41: 145-146 (1977).

92. Suzuki, Y. Cadmium, copper, and zinc distribution in blood of rats after long term cadmium administration. *J. Toxicol. Environ. Health* 7: 251-262 (1981).
93. Vander Mallie, R. J., and Garvey, J. S. Radioimmunoassay of metallothioneins. *J. Biol. Chem.* 254: 8416-8421 (1979).
94. Suzuki, Y., Toda, K., Koike, S., and Yoshikawa, H. Distribution of metals in urine of the welders using silver solder. *Japan J. Ind. Health* 19: 381 (1977).
95. Garvey, J. S., and Chang, C. H. Detection of circulating metallothionein in rats injected with zinc or cadmium. *Science* 214: 805-807 (1981).
96. Nordberg, G. F., Garvey, J. S., and Chang, C. C. Metallothionein in plasma and urine of cadmium workers. *Environ. Res.* 28: 179-182 (1982).
97. Chang, C. C., Vander Mallie, R. J., and Garvey, J. S. A radioimmunoassay for human metallothionein. *Toxicol. Appl. Pharmacol.* 55: 94-102 (1980).
98. Tohyama, C., Shaikh, Z. A., Nogawa, K., Kobayashi, E., and Honda, R. Elevated urinary excretion of metallothionein due to environmental cadmium exposure. *Toxicology* 20: 189-297 (1981).
99. Tohyama, C., Shaikh, Z. A., Ellis, K. J., and Cohn, S. H. Metallothionein excretion in urine upon cadmium exposure: its relationship with liver and kidney cadmium. *Toxicology* 22: 181-191 (1981).
100. Chang, C. C., Lauwerys, R., Bernard, A., Roels, H., Buchet, J. P., and Garvey, J. S. Metallothionein in cadmium-exposed workers. *Environ. Res.* 23: 422-428 (1980).
101. Tanaka, K., Sueda, K., Onosaka, S., and Okahara, K. Fate of  $^{109}\text{Cd}$  labelled metallothionein in rats. *Toxicol. Appl. Pharmacol.* 33: 258-266 (1975).
102. Cherian, M. G., and Shaikh, Z. A. Metabolism of intravenously injected cadmium-binding protein. *Biochem. Biophys. Res. Commun.* 65: 863-869 (1975).
103. Nordberg, G. F., Goyer, R. A., and Nordberg, M. Comparative toxicity of cadmium-metallothionein and cadmium chloride in mouse kidney. *Arch. Pathol.* 99: 192-197 (1975).
104. Johnson, D. R., and Foulkes, E. C. On the proposed role of metallothionein in the transport of cadmium. *Environ. Res.* 21: 360-365 (1980).
105. Fowler, B. R., and Nordberg, G. F. The renal toxicity of cadmium metallothionein: morphometric and x-ray microanalytical studies. *Toxicol. Appl. Pharmacol.* 46: 609-623 (1978).
106. Cherian, M. G., Goyer, R. A., and Delaquerriere-Richardson, L. Cadmium-metallothionein-induced nephropathy. *Toxicol. Appl. Pharmacol.* 38: 399-408 (1976).
107. Webb, M., and Etienne, A. T. Studies on the toxicity and metabolism of cadmium-thionein. *Biochem. Pharmacol.* 26: 25 (1977).
108. Suzuki, K. T., and Yamamura, M. Degradation and re-synthesis of injected liver cadmium-thioneins in rat kidney. *Biochem. Pharmacol.* 28: 1806-1807 (1979).
109. Elinder, C. G., and Pannone, M. Biliary excretion of cadmium. *Environ. Health Perspect.* 28: 123-126 (1979).
110. Nordberg, G. F., Piscator, M., and Nordberg, M. On the distribution of cadmium in blood. *Acta Pharmacol.* 30: 289-295 (1971).
111. Hildebrand, C. E., and Cram, L. S. Distribution of cadmium in human blood cultured in low levels of  $\text{CdCl}_2$ : accumulation of Cd in lymphocytes and preferential binding to metallothionein. *Proc. Soc. Exptl. Biol. Med.* 161: 438-443 (1979).
112. Garty, M., Wong, K.-L., and Klaassen, C. D. Redistribution of cadmium to blood of rats. *Toxicol. Appl. Pharmacol.* 59: 548-554 (1981).
113. Cain, K., and Griffiths, B. Transfer of liver cadmium to the kidney after aflatoxin induced liver damage. *Biochem. Pharmacol.* 29: 1852-1855 (1980).
114. Bernard, A. M., and Lauwerys, R. L. The effects of sodium chromate and carbon tetrachloride on the urinary excretion and tissue distribution of cadmium in cadmium-pretreated rats. *Toxicol. Appl. Pharmacol.* 57: 30-38 (1981).
115. Tanaka, K., Nomura, H., Onosaka, S., and Min, K.-S. Release of hepatic cadmium by carbon tetrachloride treatment. *Toxicol. Appl. Pharmacol.* 59: 535-539 (1981).
116. Tanaka, K. Effects of hepatic disorder on the fate of cadmium in rats. In: *Biological Roles of Metallothionein* (E. C. Foulkes, Ed.), Elsevier-North Holland, Amsterdam, 1982, pp. 237-249.
117. Hook, J. B. Toxic responses of the kidney. In: *Cassarett and Doull's Toxicology, The Basic Science of Poisons*. (J. Doull, C. D. Klaassen and M. O. Amdur, Eds.), Macmillan, New York, 1980, pp. 232-245.
118. Andersen, O., P. Lindegaard, M. Unger, and G. F. Nordberg. Effects of dietary PCB exposure on cadmium metabolism. Unpublished results.
119. Klaassen, C. D. Heavy metals and heavy-metal antagonists. In: *The Pharmacological Basis of Therapeutics*. 6th ed. (A. G. Gilman and L. S. Goodman, Eds.), New York, 1980, pp. 1615-1637.
120. Nordberg, G. F. Chelating agents and cadmium toxicity: Problems and prospects. *Environ. Health Perspect.* 54: 213-218 (1984).
121. Gilman, A., Philips, F. S., Allen, R. P., and Koelle, E. The treatment of acute cadmium intoxication in rabbits with 2,3-dimercaptopropanol (BAL) and other mercaptans. *J. Pharmacol. Exptl. Therap.* 87 (Suppl.): 85-101 (1946).
122. Friberg, L. Edathamil calcium-disodium in cadmium poisoning. *Arch. Ind. Health* 13: 13-23 (1956).
123. Dudley, H. R., Ritchie, A., Schilling, A., and Baker, W. H. Pathological changes associated with the use of sodium ethylenediaminetetraacetate in the treatment of hypercalcemia. *N. Eng. J. Med.* 252: 331-337 (1955).
124. Foreman, H., Finnegan, C., and Luskbaugh, C. C. Nephrotoxic hazards from uncontrolled edathamil calcium disodium therapy. *J. Am. Med. Assoc.* 160: 1042-1046 (1956).
125. Altman, J., Watkin, K. G., and Winkelmann, R. K. Effects of edathamil disodium on the kidney. *J. Invest. Dermatol.* 38: 215-218 (1962).
126. Cantilena, L. R., and Klaassen, C. D. Comparison of the effectiveness of several chelators after single administration on the toxicity, excretion and distribution of cadmium. *Toxicol. Appl. Pharmacol.* 58: 452-460 (1981).
127. Basinger, M. A., Jones, M. M., and Shinobu, L. A. Structural requirements for chelate antidotes for acute cadmium intoxication. *J. Inorg. Nucl. Chem.* 43: 3039-3042 (1981).
128. Von Burg, R., and Smith, J. C. Biliary mobilization of cadmium by 2,3-dimercaptopropanol and some related compounds. *J. Toxicol. Environ. Health* 6: 75-85 (1980).