

Presented at

# Great Rivers Reference Condition Workshop

January 10-11, Cincinnati, OH

Sponsored by

The U.S. Environmental Protection Agency and The Council of State Governments



**EMAP**  
Great River Ecosystems






U.S. EPA Office of Research and Development


Environmental Monitoring and Assessment Program


# Using Available Historical Data for Developing Reference Condition

Robert M. Hughes  
Department of Fisheries & Wildlife  
Oregon State University  
Corvallis, OR




(from Bryce et al. 1999)






(from Maret & Mebane 2005)



(from Hughes et al. 1993)




(from MacCoy & Blew 2005)





1939



2000








(from Jungwirth et al. 2003)

# Historical Changes in Large River Fish Assemblages of the Americas




John N. Rinne, Robert M. Hughes, and  
Bob Calamusso, editors

American Fisheries Society Symposium 45



(from Jungwirth et al. 2003)



(from Jungwirth et al. 2003)


(from Aadland  
et al. 2005; 184 kg)




(from Maret & Mebane  
2005; 286 kg)






(from Ebel et al. 1989)




(from Galat et al. 2005)




(from Freeman et al. 2005)











(from Karr et al. 1985)




(from Hughes et al. 2005)




(from Thomas et al. 2005)

(from Brown & Moyle 2005)

Mountains	Mountains	Foothills	Valley
Fishless	Rainbow trout assemblage	Pikeminnow-hardhead-sucker assemblage	Deep-bodied fishes assemblage
> 1,000 m	450 - 1,000 m	30 - 450 m	< 30 m



Mountains	Foothills	Valley
Rainbow trout assemblage	Pikeminnow-hardhead-sucker assemblage	Alien fishes dominant
> 450 m	30 - 450 m	< 30 m





(from Brown & Moyle 2005)

# Native Fishes of the Lower Colorado Main Stem


(from Mueller et al. 2005)

Species	Historical Abundance	Current Status
Bonytail	abundant	endangered
Roundtail chub	rare	extirpated
Humpback chub	rare	endangered
Colorado pikeminnow	common	endangered
Woundfin	common	endangered
Razorback sucker	abundant	endangered
Flannelmouth sucker	rare	uncommon
Desert pupfish	abundant	endangered
Sonoran topminnow	common	endangered
Gila topminnow	abundant	endangered




(from Kilgour & Stanfield 2006)






(from Contreras-Balderas et al. 2005)



(from VanSickle et al. 2004)

# % Reduction in Native Fish Fauna Diversity Within a 6-digit HUC




- 50% and greater reduction
- 25-49% reduction
- 10-24% reduction
- 1-9% reduction
- 0% reduction
- Fishless


NOTE: Current species numbers are based on a count of species that have at least one "current" record for that species within any 8-digit HUC within the 6-digit HUC


100 0 100 200 300 Miles

(from Reed & Czech 2005)




(from Freeman et al. 2005)






Number of fish species introduced into USGS 4-digit HUC subregions. Fuller et al. 1999




# EMAP-West Aquatic Vertebrate Proportionate Abundance by Proportion of Stream Length (from Lomnicky et al. In Review)






# Predominate Species in Washington (from Lomnicky et al In Review)



# Predominate Species in North Dakota

(from Lomnicky et al. In Review)



# Predominate Species in Arizona (from Lomnicky et al. In Review)

