

Presented at

Great Rivers Reference Condition Workshop

January 10-11, Cincinnati, OH

Sponsored by

The U.S. Environmental Protection Agency and The Council of State Governments

EMAP
Great River Ecosystems



U.S. EPA Office of Research and Development

Environmental Monitoring and Assessment Program



Predictive Models for Bioassessment of Large and Great Rivers

Lester L. Yuan

National Center for Environmental Assessment, Office of Research
and Development, US EPA

Outline

- Overview of predictive models
 - River Invertebrate Prediction and Classification System (RIVPACS)
- Examples from National Wadeable Streams Assessment (NWSA)
- Application of models to Great Rivers

Biological Assessment

The composition of stream organisms is used increasingly to assess stream condition.

A predictive bioassessment model is one method for analyzing biological data to assess stream condition.

RIVPACS*-type methods

compare the *observed* composition (**O**) to *expectations* (**E**) derived from reference sites.

Sites at which the assemblage composition differs significantly from expectation are declared impaired.

*Clarke, R.T., Wright, J.F, and Furse, M.T. (2002) RIVPACS models for predicting the expected macroinvertebrate fauna and assessing the ecological quality of rivers. *Ecological Modelling* 160:219-233.



Predictive Bioassessment Models

1. Identify and sample *reference* sites.
2. Group reference sites into clusters with similar biological assemblages.
3. Examine how natural factors vary with clusters of reference sites.
4. For each *test* site, use natural factors to predict the clusters in which the site would most likely be grouped.
5. The *expected* biological composition of the test site is predicted to be the same as that observed in the reference clusters. Expected composition is expressed as the capture probability, p_i , for each taxon, i .

Predictive Bioassessment Models

6. Compare *observed* taxa list to *expected* taxa list.

$$O/E_{TAXA} = \frac{\sum Y_i}{\sum p_i}$$


- $Y_i = 1$ if taxon i is present and $Y_i = 0$ if absent.
- p_i is the capture probability predicted by the model.

$O/E_{TAXA} \sim 1$ in reference sites, and decreases as taxa are lost in test sites.

National Wadeable Streams Assessment

- Statistically representative sample of small, wadeable streams across the U.S.
- Macroinvertebrates sampled on reach-wide transects.
- Reference sites sources:
 - Statistical sites attaining reference criteria.
 - Hand-picked sites.
 - USGS sites attaining reference criteria.


Reference Sites for National WSA



RESEARCH & DEVELOPMENT

Building a scientific foundation for sound environmental decisions


Super-Ecoregions



RESEARCH & DEVELOPMENT


Building a scientific foundation for sound environmental decisions

Clusters for Eastern Highlands



11 clusters with 9-25 reference sites per cluster.





Clusters in the Eastern Highlands



RESEARCH & DEVELOPMENT

Building a scientific foundation for sound environmental decisions


Predictor Variables within Clusters



RESEARCH & DEVELOPMENT

Building a scientific foundation for sound environmental decisions

Predicted clusters for a single test site



RESEARCH & DEVELOPMENT



Building a scientific foundation for sound environmental decisions

Computing Probability of Occurrence (Acroneuria)

Frequency of Occurrence in Cluster	Probability of Cluster Membership
0.58	0.09
1.00	0.31
0.18	0.46
0.50	0.13

$$p = 0.58 \times 0.09 + 1.00 \times 0.31 + 0.18 \times 0.46 + 0.50 \times 0.13 = 0.51$$

Model Results: Eastern Highlands




Useful Properties of O/E

- Index value has an intuitive ecological meaning.
 - *e.g., $O/E = 0.5$ suggests that only half of the taxa that are expected were observed.*
- Index value is expressed in scaled units and can be compared across different locations.
- One model can account for variability across a large region.
- Model is developed without any assumptions regarding stressor gradients.

Other properties

- Predictions can be sensitive to sampling method.



O/E values at state reference sites differ because of different sampling methods.

Other properties (2)

- Predictive models are most useful for controlling for many different natural gradients.
- In regions where natural gradients are weak, or dominated by a single factor, simpler models can be as effective as RIVPACS.

	WEST	PLAINS	EAST HIGH
% improvement over null model	17%	8%	27%

Conclusions

- RIVPACS-style predictive models can provide a very useful measure of biological condition.
- Further testing is required to establish whether and how predictive models can be applied in Great River.