


**Figure 1**  
**Whole Body Black Crappie**  
**Highest Concentrations by Location**  
**Collected by TVA between 04/09/2009 and 05/05/2009**



Row Labels	Total Samples	Detects	Min	Max	Average	Frequency of Detection
Arsenic	20	20	0.45	1.5	0.8895	100%
CRM8.0	5	5	0.86	1.4	1.072	100%
ERM0.9	5	5	0.81	1.5	0.96	100%
ERM2.0	5	5	0.59	0.82	0.714	100%
ERM8.0	5	5	0.45	1.2	0.812	100%
Mercury	20	9	0.07	0.24	0.126	45%
CRM8.0	5	3	0.087	0.24	0.149	60%
ERM0.9	5	1	0.13	0.13	0.13	20%
ERM2.0	5	1	0.07	0.07	0.07	20%
ERM8.0	5	4	0.096	0.15	0.1215	80%
Selenium	20	20	1.2	2.9	2.08	100%
CRM8.0	5	5	1.2	2.4	1.64	100%
ERM0.9	5	5	1.5	2.8	2.24	100%
ERM2.0	5	5	2.3	2.9	2.5	100%
ERM8.0	5	5	1.6	2.4	1.94	100%

Note: Samples where an analyte is not detected are reported at the lowest concentration that laboratory instruments can reliably measure.

- All results are verified and reported on a dry-weight basis; data are pending rigorous third-party data validation and may be updated based upon the findings from this process.

- Samples were split with the Oak Ridge National Laboratory and Tennessee State Laboratory.