

Chapter 8. Bibliography

ATSDR public health assessment guidance manual. 2005. Available at:

<http://www.atsdr.cdc.gov/HAC/PHAManual/toc.html>.

Anderson HA, Falk C, Hanrahan L, Olson J, Burse V, Needham L, et al. 1998. Profiles of Great Lakes critical pollutants: a sentinel analysis of human blood and urine—The Great Lakes Consortium. *Environ Health Perspect* 106(5):279–89.

Baibergenova A, Kudryakov R, Zdeb M, Carpenter D. 2003. Low birth weight and residential proximity to PCB-contaminated waste sites. *Environ Health Perspect* 111:135–57.

Bloom MS, Vena JE, Swanson MK, Moysich KB, Olson JR. 2005. Profiles of ortho-polychlorinated biphenyl congeners, dichlorodiphenyldichloroethylene, hexachlorobenzene, and Mirex among male Lake Ontario sportfish consumers: the New York State Angler Cohort Study. *Environ Res* 97:178–94.

Buck GM, Vena JE, Greizerstein HB, Weiner JM, McGuinness B, Mendola PP. et al. 2002. PCB congeners and pesticides and female fecundity: New York State Angler Prospective Pregnancy Study. *Environ Toxicol Pharmacol* 12: 83–92.

Bureau of the Census. 2000. Demographic data. Available at:

<http://factfinder.census.gov/>. Washington DC: US Department of Commerce.

Courval JM, DeHoog JV, Stein AD, Tay EM, He J, Humphrey HEB, and Paneth N. 1999. Sport-caught fish consumption and conception delay in licensed Michigan anglers. *Environ Res* 80 (Suppl 2):183–88.

Decaprio AP, Johnson GW, Tarbell AM, Carpenter DO, Chiarenzelli JR, Morse GS, Akwesasneet al. 2005. Task Force on the Environment. Polychlorinated Bipenyl (PCB) Exposure assessment by multivariate statistical analysis of congener profiles in an adult Native American population. *Environ Res* 98:284–302.

Dellinger JA. 2004. Exposure assessment and initial intervention regarding fish consumption of tribal members of the Upper Great Lakes Region in the United States. *Environ Res* 95(3):325–40.

Fein GG, Jacobson JL, Jacobson SW, Schwartz PM, Dowler JK. 1984. Prenatal exposure to polychlorinated biphenyls, effects on birth size and gestational age. *J Pediatr* 105:315320–20.

Fitzgerald EF, Hwanf SA, Lamvert G, Gomez M, Tarbell A. 2005. PCB Exposure and in vivo CYP1A2 activity among Native Americans. *Environ Health Perspect* 113(3):1–6.

Gardiner JC. 2001. Impairments memory and learning in older adults exposed to polychlorinated biphenyls via consumption of Great Lakes Fish. *Environ Health Perspect* 109:605–11.

HazDat. Agency for Toxic Substances and Disease Registry Hazardous Substance Release and Health Effects Database. Available at: <http://www.atsdr.cdc.gov/hazdat.html> [cited 2008 Dec 5].

- Lonky E, Reihman J, Darvill T, Mather JS, Daly H. 1996. Neonatal behavioral assessment scale performance in humans influenced by maternal consumption of environmentally contaminated Lake Ontario fish. *J Great Lakes Res* 22 (2):198–221.
- Mendola P, Buck GM, Sever LE, Zielezny M, Vena JE. 1997. Consumption of PCB-contaminated freshwater fish and shortened menstrual cycle length. *Am J Epidemiol*; 146: 55–60.
- Mendola P, Robinson LK, Buck GM, Druschel CM, Fitzgerald EF, Sever LE, et al. 2005. Birth defects associated with maternal sport fish consumption: potential effect modification by offspring. *Environ Res* 97(2):133–140.
- Schantz SL, Gasior DM, Polverejan E, McCaffrey RJ, Sweeney AM, Humphrey HEB, Gardiner et al. 2001. Impairments of memory and learning in older adults exposed to poly-chlorinated biphenyls via consumption of Great Lakes fish. *Environ Health Perspect* 109(6):605–11.
- Smith B.J. 1984. PCB levels in human fluids: Sheboygan Case Study. TR WIS-SG-83-240. Madison, Wisconsin: University of Wisconsin Sea Grant Institute.
- Stewart PW, Sargent DM, Reihman J, Gump BB, Lonkey E, Darvill T, et al. 2006. Reinforcement inhibition during differential reinforcement of low rates (DRL) schedules may be sensitive to low-level PCB, MeHg, and Pb exposure in children. *Environ Health Perspect* 114:1923–29.
- Stewart P, Fitzgerald S, Reihman J, Gump B, Lonky E, Darvill T, et al. 2003. Prenatal PCB exposure, the corpus callosum and response inhibition. *Environ Health Perspect* 111(13):1670–77.
- Stewart P, Reihman J, Lonky E, Darvill T, Pagano J. 2000. Prenatal PCB exposure and neonatal behavioral assessment scale (NBAS) performance. *Neurotoxicol Teratol* 22:21–9.
- Tilden J, Hanrahan LP, Anderson H, Palit C, Olson J, Kenzie WM. 1997. Health advisories for consumers of Great Lakes Sport fish: is the message being received? *Environ Health Perspect* 105(12):13601365–65.
- Turyk M, Anderson HA, Hanrahan LP, Falk C, Steenport DN, Needham LL Jr. 2006. Relationship serum levels of individual PCB, dioxin, and furan congeners and DDE with Great Lakes sport-caught fish consumption. *Environ Res* 100:173–83.
- US E.P.A. US Environmental Protection Agency. Great Lakes National Program Office . Available at: <http://www.epa.gov/glnpo/aoc/> (for AOC location information)).
- US E.P.A. US Environmental Protection Agency Toxics Release Inventory (TRI). Available at: <http://www.epa.gov/tri/>. 2003 Public Data Release data set (frozen March 3, 2003 and released to the public June 30, 2003).
- US E.P.A. 2003. NPL Fact Sheets. Available at: <http://www.epa.gov/epahome/whereyoulive.htm#regiontext>.
- Vasiliu O, Cameron L, Gardiner J, DeGuire P, and Karmaus W. 2006. Polybrominated biphenyls, polychlorinated biphenyls, body weight, and incidence of adult-onset diabetes mellitus. *Epidemiol* 17(4):352–59.

Xhu F, Holzman C, Rahbar MH, Trosko K, and Fischer L. 2006. Maternal fish consumption, mercury levels and risk of preterm delivery. *Environ Health Perspect*;115(1):42-7.