

Table II-9
ALL SITES (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	470.3	554.3	414.4	-0.7 ^d	-1.6 ^d	0.1
White	479.6	556.5	429.8	-0.5 ^d	-1.5 ^d	0.3
White Hispanic ^b	360.2	429.1	318.5	-0.8 ^d	-1.4 ^d	-0.3
White Non-Hispanic ^b	490.9	563.2	443.7	-0.4	-1.4 ^d	0.4
Black	515.8	689.2	400.1	-1.2 ^d	-1.9 ^d	-0.4
Asian/Pacific Islander	336.6	385.9	302.8	-0.7 ^d	-1.3 ^d	0.2
Amer Ind/Alaska Nat	237.7	263.2	222.5	-2.8 ^d	-3.9 ^d	-1.7 ^d
Hispanic ^b	351.3	419.8	309.9	-0.6 ^d	-1.2 ^d	-0.1

Mortality^c

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	199.8	251.1	166.7	-1.0 ^d	-1.5 ^d	-0.7 ^d
White	196.9	245.5	165.5	-0.9 ^d	-1.4 ^d	-0.7 ^d
White Hispanic ^b	139.7	177.8	114.4	-0.6 ^d	-0.8 ^d	-0.4 ^d
White Non-Hispanic ^b	200.1	249.7	167.9	-0.8 ^d	-1.3 ^d	-0.6 ^d
Black	252.5	347.3	196.5	-1.4 ^d	-1.9 ^d	-0.8 ^d
Asian/Pacific Islander	122.0	151.2	100.5	-1.6 ^d	-1.9 ^d	-1.1 ^d
Amer Ind/Alaska Nat	134.9	167.0	113.4	-0.5	-0.7	-0.4
Hispanic ^b	136.5	174.0	111.6	-0.5 ^d	-0.8 ^d	-0.3 ^d

The APC is the Annual Percent Change over the time interval.

- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).

^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.

^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).

^d The APC is significantly different from zero (p<.05).

Table III-9
BRAIN AND OTHER NERVOUS SYSTEM CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	6.4	7.6	5.3	-0.7	-0.7	-0.6
White	7.1	8.4	6.0	-0.4	-0.5	-0.3
White Hispanic ^b	5.3	6.0	4.7	0.2	0.4	-0.3
White Non-Hispanic ^b	7.5	9.0	6.2	-0.2	-0.3	-0.1
Black	4.0	4.6	3.4	-0.6	0.1	-0.9
Asian/Pacific Islander	3.3	4.0	2.7	-2.2 ^d	-0.9	-3.6 ^d
Amer Ind/Alaska Nat	2.5	3.0	2.3	-	-	-
Hispanic ^b	5.1	5.8	4.6	0.3	0.4	-0.1

Mortality^c

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	4.6	5.6	3.8	-0.8 ^d	-0.7 ^d	-1.1 ^d
White	4.9	6.0	4.0	-0.7 ^d	-0.6 ^d	-1.0 ^d
White Hispanic ^b	3.1	3.7	2.6	0.6	0.5	0.6
White Non-Hispanic ^b	5.1	6.3	4.2	-0.7 ^d	-0.5 ^d	-0.9 ^d
Black	2.7	3.3	2.3	-0.6	-0.8 ^d	-0.5
Asian/Pacific Islander	1.9	2.3	1.5	-0.6	1.8	-3.1
Amer Ind/Alaska Nat	2.2	2.7	1.8	2.0	3.0	-
Hispanic ^b	3.0	3.6	2.5	0.7	0.6	0.8

The APC is the Annual Percent Change over the time interval.

- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).

^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.

^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).

^d The APC is significantly different from zero (p<.05).

Table IV-12
FEMALE BREAST CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001 Rate per 100,000 persons		Trend 1992-2001 APC (%)	
	<u>Females</u>		<u>Females</u>	
All Races	135.2		0.6 ^d	
White	141.7		0.8 ^d	
White Hispanic ^b	92.1		0.5	
White Non-Hispanic ^b	149.9		1.0 ^d	
Black	119.9		-0.3	
Asian/Pacific Islander	96.8		1.7 ^d	
Amer Ind/Alaska Nat	54.2		-3.7 ^d	
Hispanic ^b	89.6		0.7	

Mortality^c

RACE/ETHNICITY	Rate 1997-2001 Rate per 100,000 persons		Trend 1992-2001 APC (%)	
	<u>Females</u>		<u>Females</u>	
All Races	27.0		-2.4 ^d	
White	26.4		-2.6 ^d	
White Hispanic ^b	17.7		-1.8 ^d	
White Non-Hispanic ^b	26.7		-2.5 ^d	
Black	35.4		-1.2 ^d	
Asian/Pacific Islander	12.6		-0.9	
Amer Ind/Alaska Nat	13.6		-1.8	
Hispanic ^b	17.3		-1.8 ^d	

- The APC is the Annual Percent Change over the time interval.
- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.
- ^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).
- ^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.
- ^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).
- ^d The APC is significantly different from zero (p<.05).

Table IV-13
 FEMALE BREAST CANCER (*In Situ*)

AGE-ADJUSTED SEER INCIDENCE

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001	Trend 1992-2001
	Rate per 100,000 persons	APC (%)
	<u>Females</u>	<u>Females</u>
All Races	30.0	5.9 ^c
White	31.0	5.8 ^c
White Hispanic ^b	17.0	5.9 ^c
White Non-Hispanic ^b	32.2	5.8 ^c
Black	27.4	6.5 ^c
Asian/Pacific Islander	24.5	7.8 ^c
Amer Ind/Alaska Nat	9.4	-
Hispanic ^b	16.7	6.3 ^c

- The APC is the Annual Percent Change over the time interval. Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).

^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.

^c The APC is significantly different from zero (p<.05).

Table V-7
CERVIX UTERI CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001	Trend 1992-2001
	Rate per 100,000 persons	APC (%)
	<u>Females</u>	<u>Females</u>
All Races	9.3	-2.6 ^d
White	8.9	-2.2 ^d
White Hispanic ^b	16.8	-3.4 ^d
White Non-Hispanic ^b	7.3	-2.5 ^d
Black	11.8	-3.1 ^d
Asian/Pacific Islander	9.5	-4.7 ^d
Amer Ind/Alaska Nat	6.0	-7.0
Hispanic ^b	16.2	-3.3 ^d

Mortality^c

RACE/ETHNICITY	Rate 1997-2001	Trend 1992-2001
	Rate per 100,000 persons	APC (%)
	<u>Females</u>	<u>Females</u>
All Races	2.9	-3.0 ^d
White	2.6	-2.5 ^d
White Hispanic ^b	3.7	-3.1 ^d
White Non-Hispanic ^b	2.5	-2.4 ^d
Black	5.6	-5.1 ^d
Asian/Pacific Islander	2.8	-2.7 ^d
Amer Ind/Alaska Nat	2.8	-5.9 ^d
Hispanic ^b	3.6	-2.9 ^d

- The APC is the Annual Percent Change over the time interval. Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.
- ^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).
- ^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.
- ^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).
- ^d The APC is significantly different from zero (p<.05).

Table VI-14
COLON AND RECTUM CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	53.7	63.4	46.4	-0.8 ^d	-1.1 ^d	-0.5
White	53.3	63.1	45.9	-0.8 ^d	-1.2 ^d	-0.6
White Hispanic ^b	40.7	51.0	33.1	0.1	0.3	-0.2
White Non-Hispanic ^b	54.4	63.8	47.2	-0.7 ^d	-1.1 ^d	-0.5
Black	63.0	72.9	56.5	-0.3	-0.6	0.0
Asian/Pacific Islander	46.3	56.3	38.6	-0.6 ^d	-0.7 ^d	-0.4
Amer Ind/Alaska Nat	35.2	38.3	32.7	-1.9	-2.8	-0.9
Hispanic ^b	39.7	49.6	32.5	0.3	0.5	-0.1

Mortality^c

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	20.8	25.3	17.7	-1.8 ^d	-2.1 ^d	-1.7 ^d
White	20.3	24.8	17.1	-1.9 ^d	-2.2 ^d	-1.8 ^d
White Hispanic ^b	14.6	18.4	11.8	-0.1	-0.2	0.2
White Non-Hispanic ^b	20.4	24.8	17.2	-1.8 ^d	-2.1 ^d	-1.7 ^d
Black	28.3	34.3	24.5	-0.7 ^d	-0.8 ^d	-0.7 ^d
Asian/Pacific Islander	13.0	15.8	10.8	-2.0 ^d	-1.8 ^d	-2.1 ^d
Amer Ind/Alaska Nat	13.9	17.1	11.7	0.6	2.5	-0.8
Hispanic ^b	14.3	18.0	11.6	0.0	-0.1	0.2

The APC is the Annual Percent Change over the time interval.

- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).

^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.

^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).

^d The APC is significantly different from zero (p<.05).

Table VI-15
COLON CANCER (Invasive) and RECTUM CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE

By Race/Ethnicity and Sex

Colon Incidence^a

RACE/ETHNICITY	Rate 1997-2001 Rate per 100,000 persons			Trend 1992-2001 APC (%)		
	Total	Males	Females	Total	Males	Females
	All Races	38.8	44.3	34.8	-0.9 ^d	-1.3 ^d
White	38.6	44.2	34.4	-0.9 ^d	-1.4 ^d	-0.6 ^d
White Hispanic ^b	28.3	34.4	23.8	0.4	0.9	-0.2
White Non-Hispanic ^b	39.6	44.9	35.6	-0.8 ^d	-1.3 ^d	-0.5
Black	48.7	55.2	44.6	-0.3	-0.5	-0.2
Asian/Pacific Islander	31.3	36.2	27.6	-0.6	-0.8	-0.2
Amer Ind/Alaska Nat	24.9	25.3	24.7	-2.9	-5.1	-0.8
Hispanic ^b	27.7	33.5	23.5	0.6	1.0	0.1

Rectum Incidence^a

RACE/ETHNICITY	Rate 1997-2001 Rate per 100,000 persons			Trend 1992-2001 APC (%)		
	Total	Males	Females	Total	Males	Females
	All Races	14.8	19.1	11.6	-0.5	-0.7
White	14.8	18.9	11.5	-0.5	-0.7	-0.6
White Hispanic ^b	12.4	16.6	9.3	-0.5	-0.8	-0.4
White Non-Hispanic ^b	14.8	18.8	11.6	-0.5	-0.6	-0.6
Black	14.3	17.7	11.9	-0.3	-0.8	0.4
Asian/Pacific Islander	15.0	20.1	11.0	-0.6	-0.4	-0.7
Amer Ind/Alaska Nat	10.2	13.0	8.0	0.5	1.7	-
Hispanic ^b	12.1	16.1	9.0	-0.4	-0.6	-0.4

- The APC is the Annual Percent Change over the time interval.
- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.
- ^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).
- ^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.
- ^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).
- ^d The APC is significantly different from zero (p<.05).

Table VII-7
CORPUS AND UTERUS, NOS CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001	Trend 1992-2001
	Rate per 100,000 persons	APC (%)
	Females	Females
All Races	24.5	-0.1
White	26.1	-0.1
White Hispanic ^b	17.3	0.3
White Non-Hispanic ^b	27.1	-0.1
Black	18.2	0.9
Asian/Pacific Islander	17.2	1.4 ^d
Amer Ind/Alaska Nat	10.2	-
Hispanic ^b	16.8	0.5

Mortality^c

RACE/ETHNICITY	Rate 1997-2001	Trend 1992-2001
	Rate per 100,000 persons	APC (%)
	Females	Females
All Races	4.1	-0.2
White	3.9	-0.2
White Hispanic ^b	3.3	0.5
White Non-Hispanic ^b	3.8	-0.1
Black	6.9	-0.1
Asian/Pacific Islander	2.1	0.4
Amer Ind/Alaska Nat	2.3	-1.8
Hispanic ^b	3.2	0.4

- The APC is the Annual Percent Change over the time interval.
 - Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.
^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).
^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.
^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).
^d The APC is significantly different from zero (p<.05).

Table VIII-9
ESOPHAGUS CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	4.5	7.7	2.0	0.3	0.5	-0.9
White	4.5	7.7	1.9	1.5 ^d	1.9 ^d	-0.2
White Hispanic ^b	3.3	6.2	1.2	-0.2	-0.6	-
White Non-Hispanic ^b	4.7	7.9	2.0	2.0 ^d	2.3 ^d	0.1
Black	6.7	10.6	3.8	-5.2 ^d	-5.8 ^d	-4.1 ^d
Asian/Pacific Islander	2.9	5.1	1.1	-0.7	-1.3	-
Amer Ind/Alaska Nat	3.2	4.9	-	-	-	-
Hispanic ^b	3.2	6.0	1.2	-0.1	-0.4	-

Mortality^c

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	4.4	7.7	1.8	0.6 ^d	0.7 ^d	0.0
White	4.2	7.4	1.7	1.6 ^d	1.7 ^d	0.8 ^d
White Hispanic ^b	2.5	4.5	1.0	-0.3	-1.2 ^d	2.7
White Non-Hispanic ^b	4.3	7.6	1.7	1.9 ^d	1.9 ^d	1.2 ^d
Black	6.7	11.7	3.2	-4.0 ^d	-4.3 ^d	-3.3 ^d
Asian/Pacific Islander	2.1	3.5	1.0	-3.0	-3.1	-1.9
Amer Ind/Alaska Nat	2.9	4.8	1.4	2.8	0.9	-
Hispanic ^b	2.4	4.4	1.0	-0.4	-1.2 ^d	2.2

The APC is the Annual Percent Change over the time interval.

- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).

^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.

^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).

^d The APC is significantly different from zero (p<.05).

Table IX-9
HODGKIN LYMPHOMA (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	2.7	3.0	2.4	-0.5	-0.3	-0.8
White	2.9	3.3	2.6	-0.4	-0.3	-0.5
White Hispanic ^b	2.6	3.2	2.1	0.8	1.0	0.6
White Non-Hispanic ^b	3.1	3.4	2.9	-0.3	-0.2	-0.4
Black	2.4	2.9	2.0	-0.8	1.2	-3.1
Asian/Pacific Islander	1.1	1.2	1.0	2.8	-0.6	6.6 ^d
Amer Ind/Alaska Nat	-	-	-	-	-	-
Hispanic ^b	2.5	3.1	2.0	1.1	1.4	0.6

Mortality^c

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	0.5	0.6	0.4	-3.5 ^d	-4.0 ^d	-2.9 ^d
White	0.5	0.6	0.4	-3.4 ^d	-3.7 ^d	-3.0 ^d
White Hispanic ^b	0.6	0.8	0.4	-2.8	-2.2	-4.1
White Non-Hispanic ^b	0.5	0.6	0.4	-3.0 ^d	-3.6 ^d	-2.3 ^d
Black	0.4	0.6	0.3	-4.5 ^d	-5.5 ^d	-3.3 ^d
Asian/Pacific Islander	0.2	0.2	0.2	-	-	-
Amer Ind/Alaska Nat	-	-	-	-	-	-
Hispanic ^b	0.5	0.7	0.4	-2.8	-2.2	-3.9

- The APC is the Annual Percent Change over the time interval.
 - Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.
 a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).
 b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.
 c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).
 d The APC is significantly different from zero (p<.05).

Table XI-9
KIDNEY AND RENAL PELVIS CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	11.3	15.7	7.8	1.4 ^d	1.3 ^d	1.5 ^d
White	11.7	16.2	8.1	1.6 ^d	1.4 ^d	1.4 ^d
White Hispanic ^b	11.5	15.5	8.5	1.9	1.1	2.9 ^d
White Non-Hispanic ^b	11.5	16.0	7.8	1.7 ^d	1.7 ^d	1.4 ^d
Black	13.3	18.8	9.4	2.2 ^d	1.9	2.8 ^d
Asian/Pacific Islander	6.4	8.9	4.3	1.4	0.4	3.1 ^d
Amer Ind/Alaska Nat	10.0	13.9	7.0	-5.6 ^d	-5.9 ^d	-
Hispanic ^b	11.2	15.1	8.2	2.2 ^d	1.4	3.0 ^d

Mortality^c

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	4.2	6.1	2.8	-0.2	-0.1	-0.4
White	4.3	6.2	2.8	-0.2	-0.1	-0.4
White Hispanic ^b	3.9	5.7	2.5	0.3	0.8	-0.8
White Non-Hispanic ^b	4.3	6.3	2.9	-0.1	-0.1	-0.4
Black	4.2	6.3	2.7	0.2	0.5	-0.1
Asian/Pacific Islander	2.0	2.8	1.3	1.0	0.8	1.6
Amer Ind/Alaska Nat	4.7	6.8	3.2	-0.3	-0.4	-1.7
Hispanic ^b	3.8	5.5	2.4	0.4	0.8	-0.6

The APC is the Annual Percent Change over the time interval.

- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).

^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.

^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).

^d The APC is significantly different from zero (p<.05).

Table XII-9
LARYNX CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	3.8	6.7	1.4	-2.9 ^d	-3.4 ^d	-1.9
White	3.8	6.6	1.5	-2.8 ^d	-3.3 ^d	-1.5
White Hispanic ^b	2.9	5.7	0.9	-0.9	-1.6 ^d	-
White Non-Hispanic ^b	3.7	6.5	1.5	-2.9 ^d	-3.4 ^d	-1.7
Black	6.4	12.1	2.4	-3.1 ^d	-2.9 ^d	-3.4
Asian/Pacific Islander	1.7	3.4	0.5	-2.5	-2.8	-
Amer Ind/Alaska Nat	-	-	-	-	-	-
Hispanic ^b	2.8	5.4	0.9	-0.7	-1.5 ^d	-

Mortality^c

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	1.4	2.6	0.5	-2.0 ^d	-2.3 ^d	-1.7 ^d
White	1.3	2.3	0.5	-1.9 ^d	-2.2 ^d	-1.5 ^d
White Hispanic ^b	1.1	2.3	0.2	-1.7	-1.9	-
White Non-Hispanic ^b	1.3	2.3	0.5	-1.6 ^d	-2.0 ^d	-1.1
Black	2.7	5.4	0.9	-2.7 ^d	-2.8 ^d	-2.4 ^d
Asian/Pacific Islander	0.5	1.0	0.1	-0.6	1.1	-
Amer Ind/Alaska Nat	1.3	2.4	-	-	-	-
Hispanic ^b	1.0	2.2	0.2	-1.8	-2.0	-

The APC is the Annual Percent Change over the time interval.

- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).

^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.

^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).

^d The APC is significantly different from zero (p<.05).

Table XIII-15
LEUKEMIA (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	12.1	15.9	9.4	-1.0 ^d	-1.3 ^d	-0.8 ^d
White	12.9	16.9	9.9	-0.9 ^d	-1.3 ^d	-0.5
White Hispanic ^b	9.6	11.8	7.9	-1.2	-1.0	-1.3
White Non-Hispanic ^b	12.8	17.0	9.7	-0.9 ^d	-1.2 ^d	-0.7
Black	9.9	12.8	7.9	-0.9	-1.2	-0.5
Asian/Pacific Islander	7.5	9.6	5.9	-1.4	-0.1	-2.9 ^d
Amer Ind/Alaska Nat	5.0	5.6	4.5	3.3	-	-
Hispanic ^b	9.2	11.4	7.6	-1.2	-0.9	-1.3

Mortality^c

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	7.6	10.2	5.9	-0.5 ^d	-0.7 ^d	-0.5 ^d
White	7.8	10.5	6.0	-0.4 ^d	-0.6 ^d	-0.4 ^d
White Hispanic ^b	5.5	7.0	4.4	0.2	0.7	-0.5
White Non-Hispanic ^b	7.9	10.6	6.1	-0.4 ^d	-0.7 ^d	-0.3 ^d
Black	6.8	9.0	5.4	-1.0 ^d	-1.1 ^d	-0.8 ^d
Asian/Pacific Islander	4.2	5.3	3.3	-1.4	-1.2	-1.4
Amer Ind/Alaska Nat	3.9	4.8	3.1	-2.1	-3.9	-1.2
Hispanic ^b	5.4	6.8	4.3	0.2	0.7	-0.4

- The APC is the Annual Percent Change over the time interval.
- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.
- ^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).
- ^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.
- ^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).
- ^d The APC is significantly different from zero (p<.05).

Table XIV-9
LIVER AND INTRAHEPATIC BILE DUCT CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	6.0	9.1	3.5	3.4 ^d	3.2 ^d	3.3 ^d
White	4.8	7.2	2.9	3.7 ^d	3.2 ^d	3.9 ^d
White Hispanic ^b	9.5	13.9	5.9	2.8 ^d	1.7	4.3 ^d
White Non-Hispanic ^b	4.1	6.2	2.4	3.1 ^d	2.6 ^d	3.3 ^d
Black	7.2	11.8	3.9	3.6 ^d	4.6 ^d	1.9
Asian/Pacific Islander	13.7	21.1	7.7	0.2	0.8	-0.9
Amer Ind/Alaska Nat	6.4	8.3	4.8	-2.8	-	-
Hispanic ^b	9.2	13.5	5.8	2.9 ^d	1.9	4.3 ^d

Mortality^c

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	4.6	6.7	2.9	1.9 ^d	2.0 ^d	1.2 ^d
White	4.2	6.1	2.7	2.0 ^d	2.1 ^d	1.1 ^d
White Hispanic ^b	7.7	10.8	5.2	2.2 ^d	1.6 ^d	2.5 ^d
White Non-Hispanic ^b	4.0	5.8	2.6	1.8 ^d	2.0 ^d	0.8
Black	6.1	9.3	3.8	1.3 ^d	1.1	1.1
Asian/Pacific Islander	10.7	15.6	6.6	-0.8	-0.8	-0.4
Amer Ind/Alaska Nat	6.1	8.3	4.3	3.0 ^d	2.5	2.5
Hispanic ^b	7.5	10.6	5.1	2.2 ^d	1.7 ^d	2.4 ^d

The APC is the Annual Percent Change over the time interval.

- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).

^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.

^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).

^d The APC is significantly different from zero (p<.05).

Table XV-14
LUNG AND BRONCHUS (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	61.7	79.1	49.1	-1.3 ^d	-2.3 ^d	-0.2
White	62.4	77.9	51.3	-1.2 ^d	-2.3 ^d	-0.2
White Hispanic ^b	33.3	46.0	24.5	-2.3 ^d	-2.9 ^d	-1.6
White Non-Hispanic ^b	64.1	79.1	53.4	-1.1 ^d	-2.1 ^d	-0.1
Black	79.8	117.2	54.5	-1.5 ^d	-2.6 ^d	0.5
Asian/Pacific Islander	42.4	60.5	28.5	-1.1 ^d	-1.3 ^d	-0.1
Amer Ind/Alaska Nat	33.1	46.0	23.4	-3.9 ^d	-4.3 ^d	-3.1 ^d
Hispanic ^b	32.6	45.2	23.9	-2.1 ^d	-2.7 ^d	-1.5

Mortality^c

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	56.2	77.9	40.8	-0.8 ^d	-1.9 ^d	0.6 ^d
White	56.2	76.6	41.6	-0.7 ^d	-1.7 ^d	0.7 ^d
White Hispanic ^b	25.7	40.4	15.3	-0.9 ^d	-1.5 ^d	0.1
White Non-Hispanic ^b	58.4	79.5	43.3	-0.5 ^d	-1.6 ^d	0.8 ^d
Black	65.2	104.1	39.9	-1.5 ^d	-2.4 ^d	0.4
Asian/Pacific Islander	28.2	40.2	19.2	-1.2 ^d	-1.7 ^d	0.1
Amer Ind/Alaska Nat	36.3	49.8	26.6	-0.6	-1.8	1.0
Hispanic ^b	25.2	39.6	14.9	-0.9 ^d	-1.4 ^d	0.1

The APC is the Annual Percent Change over the time interval.

- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).

^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.

^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).

^d The APC is significantly different from zero (p<.05).

Table XVI-9
MELANOMA OF THE SKIN (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	17.0	21.4	13.8	2.5 ^d	2.7 ^d	2.5 ^d
White	20.3	25.4	16.8	3.1 ^d	3.2 ^d	3.2 ^d
White Hispanic ^b	4.5	4.6	4.6	3.1 ^d	3.5 ^d	2.8
White Non-Hispanic ^b	23.2	28.4	19.5	3.2 ^d	3.1 ^d	3.4 ^d
Black	1.0	1.5	0.7	-0.7	-	-
Asian/Pacific Islander	1.6	1.7	1.5	3.0	1.7	4.5
Amer Ind/Alaska Nat	1.9	-	-	-	-	-
Hispanic ^b	4.3	4.4	4.5	3.3 ^d	3.4 ^d	3.3 ^d

Mortality^c

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	2.7	3.9	1.8	-0.2	0.0	-0.6 ^d
White	3.0	4.4	2.0	0.0	0.2	-0.4
White Hispanic ^b	0.9	1.2	0.7	0.0	0.6	-1.5
White Non-Hispanic ^b	3.2	4.7	2.1	0.1	0.3	-0.2
Black	0.5	0.5	0.5	-0.8	-0.7	-0.7
Asian/Pacific Islander	0.4	0.4	0.3	-4.0	-	-
Amer Ind/Alaska Nat	0.5	0.8	-	-	-	-
Hispanic ^b	0.8	1.1	0.6	0.0	0.5	-1.4

The APC is the Annual Percent Change over the time interval.

- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).

^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.

^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).

^d The APC is significantly different from zero (p<.05).

Table XVII-9
MYELOMA (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	5.6	6.9	4.6	-0.7	-1.0	-0.6
White	5.2	6.6	4.2	-0.5	-0.7	-0.7
White Hispanic ^b	5.1	6.1	4.4	-1.6 ^d	-2.2	-0.9
White Non-Hispanic ^b	5.2	6.7	4.1	-0.3	-0.5	-0.5
Black	11.0	12.9	9.8	-1.7 ^d	-1.7	-1.6
Asian/Pacific Islander	3.5	4.1	3.0	0.0	-2.1	2.7
Amer Ind/Alaska Nat	3.2	-	3.4	-	-	-
Hispanic ^b	5.1	6.0	4.4	-1.3	-1.8	-0.7

Mortality^c

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	3.8	4.7	3.2	-0.4	-0.6	-0.3
White	3.5	4.4	2.9	-0.3	-0.4	-0.3
White Hispanic ^b	3.3	3.8	2.8	1.0	0.2	1.5
White Non-Hispanic ^b	3.6	4.5	2.9	-0.3	-0.4	-0.3
Black	7.5	9.0	6.6	-0.5	-1.0 ^d	-0.1
Asian/Pacific Islander	1.8	2.2	1.6	-0.5	-1.8	0.9
Amer Ind/Alaska Nat	3.0	3.4	2.7	1.0	-	-
Hispanic ^b	3.2	3.8	2.8	1.2	0.5	1.6

The APC is the Annual Percent Change over the time interval.

- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).

^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.

^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).

^d The APC is significantly different from zero (p<.05).

Table XVIII-13
NON-HODGKIN LYMPHOMA (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	19.1	23.3	15.7	0.0	-0.5	0.8 ^d
White	20.1	24.4	16.6	0.0	-0.5	0.8 ^d
White Hispanic ^b	16.3	19.1	13.9	-0.7	-1.6	0.5
White Non-Hispanic ^b	20.6	25.1	16.9	0.0	-0.5	0.8 ^d
Black	14.6	18.3	11.7	0.2	-1.5	2.9 ^d
Asian/Pacific Islander	13.7	16.7	11.4	0.8	0.3	1.3
Amer Ind/Alaska Nat	8.7	9.7	7.7	2.6	-	-
Hispanic ^b	15.8	18.5	13.5	-0.6	-1.4	0.6

Mortality^c

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	8.4	10.5	6.9	-0.4	-0.3	-0.5
White	8.7	10.8	7.2	-0.4	-0.3	-0.5
White Hispanic ^b	6.7	8.2	5.5	-0.3	-0.5	0.0
White Non-Hispanic ^b	8.8	11.0	7.2	-0.2	0.0	-0.4
Black	5.7	7.4	4.6	-0.3	-0.3	-0.2
Asian/Pacific Islander	5.2	6.7	4.1	-1.1 ^d	-1.4	-0.6
Amer Ind/Alaska Nat	5.0	5.9	4.4	5.1 ^d	5.0 ^d	5.1 ^d
Hispanic ^b	6.5	8.1	5.3	-0.2	-0.4	0.1

- The APC is the Annual Percent Change over the time interval.
- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.
- ^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).
- ^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.
- ^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).
- ^d The APC is significantly different from zero (p<.05).

Table XIX-9
ORAL CAVITY AND PHARYNX CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	10.7	15.7	6.4	-2.0 ^d	-2.4 ^d	-1.4 ^d
White	10.7	15.7	6.4	-1.8 ^d	-2.2 ^d	-1.5 ^d
White Hispanic ^b	6.6	9.9	3.9	-2.7 ^d	-3.4 ^d	-1.8
White Non-Hispanic ^b	11.1	16.3	6.7	-1.8 ^d	-2.1 ^d	-1.8 ^d
Black	11.9	19.5	6.2	-3.2 ^d	-3.4 ^d	-2.1
Asian/Pacific Islander	8.7	12.2	5.8	-1.7	-2.3	-0.2
Amer Ind/Alaska Nat	6.8	10.7	3.8	-5.8 ^d	-7.8 ^d	-
Hispanic ^b	6.4	9.6	3.9	-2.6 ^d	-3.4 ^d	-1.5

Mortality^c

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	2.8	4.3	1.6	-2.6 ^d	-2.8 ^d	-2.5 ^d
White	2.6	3.9	1.6	-2.3 ^d	-2.5 ^d	-2.3 ^d
White Hispanic ^b	1.8	3.0	0.9	-3.0 ^d	-4.3 ^d	0.0
White Non-Hispanic ^b	2.7	4.0	1.7	-2.1 ^d	-2.3 ^d	-2.1 ^d
Black	4.3	7.5	2.0	-4.2 ^d	-4.4 ^d	-3.3 ^d
Asian/Pacific Islander	2.4	3.6	1.3	-2.6 ^d	-2.5 ^d	-2.7
Amer Ind/Alaska Nat	2.3	3.8	1.1	-1.4	-1.0	-
Hispanic ^b	1.8	2.9	0.9	-3.0 ^d	-4.2 ^d	0.0

The APC is the Annual Percent Change over the time interval.

- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).

^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.

^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).

^d The APC is significantly different from zero (p<.05).

Table XX-7
OVARY CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001	Trend 1992-2001
	Rate per 100,000 persons	APC (%)
	<u>Females</u>	<u>Females</u>
All Races	13.9	-1.0 ^d
White	14.8	-0.8 ^d
White Hispanic ^b	11.7	-1.2
White Non-Hispanic ^b	15.3	-0.7 ^d
Black	10.1	-1.6
Asian/Pacific Islander	10.3	-0.6
Amer Ind/Alaska Nat	9.0	-
Hispanic ^b	11.4	-1.0

Mortality^c

RACE/ETHNICITY	Rate 1997-2001	Trend 1992-2001
	Rate per 100,000 persons	APC (%)
	<u>Females</u>	<u>Females</u>
All Races	8.9	-0.6 ^d
White	9.2	-0.5 ^d
White Hispanic ^b	6.3	-0.7
White Non-Hispanic ^b	9.3	-0.4
Black	7.5	-0.9 ^d
Asian/Pacific Islander	4.6	-0.5
Amer Ind/Alaska Nat	4.9	-0.1
Hispanic ^b	6.1	-0.6

- The APC is the Annual Percent Change over the time interval.
- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.
- ^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).
- ^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.
- ^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).
- ^d The APC is significantly different from zero (p<.05).

Table XXI-9
PANCREAS CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	11.0	12.6	9.7	-0.5	-0.4	-0.6
White	10.8	12.5	9.4	-0.3	0.0	-0.6
White Hispanic ^b	10.1	10.7	9.6	-0.8	-1.3	-0.5
White Non-Hispanic ^b	10.7	12.5	9.2	-0.2	0.3	-0.8
Black	15.4	17.1	13.9	-2.2 ^d	-2.0 ^d	-2.5 ^d
Asian/Pacific Islander	9.3	10.2	8.6	-0.4	-2.9 ^d	2.3 ^d
Amer Ind/Alaska Nat	8.3	8.3	8.4	-	-	-
Hispanic ^b	9.8	10.4	9.4	-0.8	-1.3	-0.4

Mortality^c

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	10.5	12.2	9.2	-0.1	-0.3 ^d	-0.1
White	10.3	12.0	8.9	0.0	-0.1	0.0
White Hispanic ^b	8.5	9.6	7.7	-0.2	-0.7	0.1
White Non-Hispanic ^b	10.3	12.0	8.9	0.1	0.0	0.1
Black	14.2	16.0	12.8	-1.1 ^d	-1.6 ^d	-0.7 ^d
Asian/Pacific Islander	7.3	8.1	6.6	-0.8	-2.2 ^d	0.6
Amer Ind/Alaska Nat	6.2	6.5	5.9	0.0	1.8	-0.9
Hispanic ^b	8.4	9.4	7.5	-0.1	-0.5	0.1

The APC is the Annual Percent Change over the time interval.

- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).

^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.

^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).

^d The APC is significantly different from zero (p<.05).

Table XXII-7
PROSTATE CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001		Trend 1992-2001	
	Rate per 100,000 persons		APC (%)	
	<u>Males</u>		<u>Males</u>	
All Races	172.3		-2.5 ^d	
White	167.4		-2.6 ^d	
White Hispanic ^b	141.6		-1.3 ^d	
White Non-Hispanic ^b	166.7		-2.6 ^d	
Black	271.3		-2.3 ^d	
Asian/Pacific Islander	100.7		-1.9	
Amer Ind/Alaska Nat	51.2		-7.9 ^d	
Hispanic ^b	140.0		-1.1 ^d	

Mortality^c

RACE/ETHNICITY	Rate 1997-2001		Trend 1992-2001	
	Rate per 100,000 persons		APC (%)	
	<u>Males</u>		<u>Males</u>	
All Races	31.5		-3.6 ^d	
White	28.8		-3.7 ^d	
White Hispanic ^b	23.8		-2.3 ^d	
White Non-Hispanic ^b	29.1		-3.7 ^d	
Black	70.4		-2.3 ^d	
Asian/Pacific Islander	13.0		-4.8 ^d	
Amer Ind/Alaska Nat	20.2		-3.9 ^d	
Hispanic ^b	23.5		-2.3 ^d	

The APC is the Annual Percent Change over the time interval.

- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).

^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.

^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).

^d The APC is significantly different from zero (p<.05).

Table XXIII-9
STOMACH CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	8.9	12.6	6.1	-1.6 ^d	-2.3 ^d	-0.9
White	7.5	10.8	5.0	-1.8 ^d	-2.3 ^d	-1.3
White Hispanic ^b	13.6	18.4	10.3	-2.2 ^d	-2.9 ^d	-1.4
White Non-Hispanic ^b	6.5	9.7	4.1	-2.2 ^d	-2.5 ^d	-2.1 ^d
Black	13.6	18.8	9.9	-1.4	-2.6 ^d	0.0
Asian/Pacific Islander	16.5	21.9	12.4	-3.3 ^d	-3.5 ^d	-2.7 ^d
Amer Ind/Alaska Nat	11.8	15.7	8.9	2.1	-	-
Hispanic ^b	13.2	17.8	10.0	-2.1 ^d	-2.8 ^d	-1.2

Mortality^c

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	4.7	6.6	3.3	-2.9 ^d	-3.4 ^d	-2.6 ^d
White	4.1	5.8	2.8	-3.2 ^d	-3.6 ^d	-2.8 ^d
White Hispanic ^b	7.3	9.9	5.4	-1.9 ^d	-2.1 ^d	-1.6
White Non-Hispanic ^b	3.8	5.4	2.6	-3.3 ^d	-3.7 ^d	-3.1 ^d
Black	9.1	13.3	6.3	-2.6 ^d	-2.9 ^d	-2.3 ^d
Asian/Pacific Islander	9.1	11.9	7.0	-3.6 ^d	-3.5 ^d	-3.6 ^d
Amer Ind/Alaska Nat	5.4	7.3	4.1	-1.5	-1.2	-1.4
Hispanic ^b	7.1	9.7	5.3	-1.8 ^d	-2.0 ^d	-1.6

The APC is the Annual Percent Change over the time interval.

- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).

^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.

^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).

^d The APC is significantly different from zero (p<.05).

Table XXIV-7
TESTIS CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001	Trend 1992-2001
	Rate per 100,000 persons	APC (%)
	<u>Males</u>	<u>Males</u>
All Races	5.2	1.2 ^d
White	6.2	1.3 ^d
White Hispanic ^b	3.7	1.0
White Non-Hispanic ^b	7.0	1.8 ^d
Black	1.5	6.4 ^d
Asian/Pacific Islander	2.1	2.3
Amer Ind/Alaska Nat	2.3	-
Hispanic ^b	3.6	1.1

Mortality^c

RACE/ETHNICITY	Rate 1997-2001	Trend 1992-2001
	Rate per 100,000 persons	APC (%)
	<u>Males</u>	<u>Males</u>
All Races	0.3	-1.3
White	0.3	-1.4
White Hispanic ^b	0.2	-4.0
White Non-Hispanic ^b	0.3	-0.8
Black	0.2	2.3
Asian/Pacific Islander	-	-
Amer Ind/Alaska Nat	-	-
Hispanic ^b	0.2	-3.9

The APC is the Annual Percent Change over the time interval.

- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).

^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.

^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).

^d The APC is significantly different from zero (p<.05).

Table XXV-9
THYROID CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	7.2	3.8	10.4	3.8 ^d	2.4 ^d	4.3 ^d
White	7.5	4.0	10.9	4.2 ^d	2.8 ^d	4.8 ^d
White Hispanic ^b	6.7	3.0	10.3	2.0	1.9	2.2 ^d
White Non-Hispanic ^b	7.7	4.2	11.2	4.7 ^d	3.3 ^d	5.3 ^d
Black	4.1	2.2	5.7	3.2 ^d	0.8	3.8 ^d
Asian/Pacific Islander	8.1	3.9	11.8	1.3	0.7	1.3
Amer Ind/Alaska Nat	4.0	-	6.3	0.9	-	2.6
Hispanic ^b	6.5	2.9	9.9	2.2	1.8	2.5 ^d

Mortality^c

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	0.5	0.4	0.5	0.6	2.3 ^d	-0.3
White	0.5	0.5	0.4	0.5	2.3 ^d	-0.5
White Hispanic ^b	0.6	0.5	0.7	-0.9	2.8	-2.4
White Non-Hispanic ^b	0.4	0.5	0.4	0.8	2.7 ^d	-0.2
Black	0.5	0.4	0.5	2.5 ^d	4.0 ^d	2.0
Asian/Pacific Islander	0.7	0.5	0.8	-3.0	-	-3.1
Amer Ind/Alaska Nat	-	-	-	-	-	-
Hispanic ^b	0.6	0.5	0.7	-0.9	2.8	-2.3

The APC is the Annual Percent Change over the time interval.

- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).

^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.

^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).

^d The APC is significantly different from zero (p<.05).

Table XXVI-9
URINARY BLADDER CANCER

AGE-ADJUSTED SEER INCIDENCE (Invasive and *In Situ*) AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	20.4	36.1	9.1	-0.1	-0.1	-0.5 ^d
White	22.5	39.9	9.8	0.1	0.0	-0.4 ^d
White Hispanic ^b	10.8	19.1	5.1	-0.9	-0.8	-1.1
White Non-Hispanic ^b	23.4	41.3	10.2	0.3	0.1	-0.1
Black	12.6	20.4	7.7	-0.2	-0.3	0.3
Asian/Pacific Islander	9.6	16.3	4.5	-0.2	0.1	0.2
Amer Ind/Alaska Nat	5.6	9.5	-	-	-	-
Hispanic ^b	10.5	18.6	5.0	-0.7	-0.6	-1.1

Mortality^c

RACE/ETHNICITY	Rate 1997-2001			Trend 1992-2001		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	4.4	7.6	2.3	-0.4 ^d	-0.6 ^d	-0.6 ^d
White	4.5	7.9	2.3	-0.2 ^d	-0.5 ^d	-0.5
White Hispanic ^b	2.5	4.2	1.3	-0.3	-0.7	0.3
White Non-Hispanic ^b	4.5	8.0	2.3	-0.1	-0.4 ^d	-0.2
Black	3.9	5.6	2.9	-1.6 ^d	-2.2 ^d	-1.0
Asian/Pacific Islander	1.8	2.9	1.0	-1.1	0.3	-3.0
Amer Ind/Alaska Nat	1.5	2.4	1.0	-0.2	-	-
Hispanic ^b	2.4	4.1	1.3	-0.4	-0.8	0.4

- The APC is the Annual Percent Change over the time interval.
- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.
- ^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).
- ^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.
- ^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).
- ^d The APC is significantly different from zero (p<.05).