

SEER Incidence* and US Death Rates,# 1997-2001 5-Year Relative Survival Rates,^ 1995-2000 All Cancer Combined, by Race and Sex




Figure 1-9


* Incidence rates are from the SEER 12 areas and are age-adjusted to the 2000 US standard population by 5-year age groups.

Death rates are from the NCHS public use data file for the total US and are age-adjusted to the 2000 US standard population by 5-year age groups.

^ Survival rates are from the SEER 9 areas. Relative survival rates are expressed as percents.

Figure I-11

5-Year Relative Survival Rates SEER Program, 1995-2000 Both Sexes, by Race




Source: SEER 9 areas.


SEER Cancer Incidence and US Death Rates, 1997-2001 By Cancer Site and Race

Incidence Cancer Site Mortality


Lung and Bronchus




Colon and Rectum



Breast (Female)



Prostate



Source: SEER 12 areas and NCHS public use data file for the total US.

^^ Hispanic is not mutually exclusive from Whites, Blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics excludes Detroit and Hawaii. Mortality data for Hispanics excludes cases from Connecticut, Maine, Maryland, Oklahoma, New York, North Dakota, New Hampshire, and Vermont.

Rates are per 100,000 and age-adjusted to the 2000 U.S. standard population by 5-year age groups.