

Table I-18
US PREVALENCE COUNTS, INVASIVE CANCERS ONLY, JANUARY 1, 2001^a
USING DIFFERENT TUMOR INCLUSION CRITERIA^b

Site/Sex	5-Year Limited Duration			26-Year Limited Duration	
	1st Invasive Tumor Ever ^c	1st Per Site in Previous 26 Years ^d	1st Per Site in Previous 5 Years ^e	1st Invasive Tumor Ever ^c	1st Per Site in Previous 26 Years ^d
All Sites	3,765,929	3,839,441	4,120,783	9,048,822	9,220,361
Male	1,898,532	1,924,231	2,057,870	4,163,637	4,212,021
Female	1,867,397	1,915,210	2,062,913	4,885,185	5,008,340
Oral Cavity & Pharynx	82,674	93,488	95,948	206,494	224,986
Male	55,409	62,109	63,607	133,057	143,709
Female	27,265	31,379	32,341	73,437	81,277
Esophagus	15,737	18,638	18,650	22,081	25,810
Male	11,952	13,945	13,945	16,515	19,027
Female	3,785	4,693	4,705	5,566	6,783
Stomach	29,431	34,023	34,105	55,827	62,559
Male	17,724	20,593	20,627	32,088	36,017
Female	11,707	13,430	13,478	23,739	26,542
Colon & Rectum	403,032	460,916	468,964	968,947	1,070,487
Male	201,690	231,008	234,709	471,439	517,261
Female	201,342	229,908	234,255	497,508	553,226
Liver & Intrahep	10,956	12,582	12,593	14,108	15,944
Male	7,118	8,193	8,204	8,706	9,905
Female	3,838	4,389	4,389	5,402	6,039
Pancreas	18,095	21,156	21,156	23,967	27,543
Male	8,680	10,372	10,372	11,163	13,073
Female	9,415	10,784	10,784	12,804	14,470
Larynx	34,076	39,252	39,432	92,464	101,321
Male	26,759	30,913	31,046	74,006	80,900
Female	7,317	8,339	8,386	18,458	20,421
Lung & Bronchus	201,687	243,497	247,320	330,991	388,538
Male	101,771	123,706	125,522	165,394	193,505
Female	99,916	119,791	121,798	165,597	195,033
Melanoma of the Skin	194,560	214,066	219,641	544,920	578,885
Male	103,566	115,336	118,977	269,150	287,927
Female	90,994	98,730	100,664	275,770	290,958
Breast					
Female	791,033	846,865	880,990	2,058,593	2,184,125
Cervix					
Female	46,131	48,226	48,288	174,429	179,437
Corpus & Uterus					
Female	140,432	157,422	157,475	474,773	510,266

^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER 9 registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census.

^{bcd} (b) Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion: (c) First invasive tumor ever; (d) First invasive tumor for each cancer site diagnosed during the previous 26 years (1975-2000); (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (1996-2000). For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 1997; Melanoma in 1998.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 26-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred 20 years ago.

In method (d) the 1981 melanoma is counted for the melanoma and all sites 26-year limited duration prevalence. The 1997 breast cancer is counted for the breast 5-year and 26-year limited duration prevalence.

In method (e) the 1997 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 1998 melanoma is counted for 5-year limited duration prevalence for melanoma.

Table I-18 - continued
US PREVALENCE COUNTS, INVASIVE CANCERS ONLY, JANUARY 1, 2001^a
USING DIFFERENT TUMOR INCLUSION CRITERIA^b

Site/Sex	5-Year Limited Duration			26-Year Limited Duration	
	1st Invasive Tumor Ever ^c	1st Per Site in Previous 26 Years ^d	1st Per Site in Previous 5 Years ^e	1st Invasive Tumor Ever ^c	1st Per Site in Previous 26 Years ^d
Ovary					
Female	53,367	61,207	61,212	137,976	153,863
Prostate					
Male	854,021	920,213	920,281	1,721,366	1,838,653
Testis					
Male	36,654	37,256	37,808	136,928	138,606
Urinary Bladder					
Male	182,170	217,670	218,962	460,530	521,945
Female	136,806	163,071	164,170	341,794	385,549
Female	45,364	54,599	54,792	118,736	136,396
Kidney & Renal Pelvis					
Male	85,215	102,069	102,742	196,981	225,128
Female	51,234	62,053	62,545	116,559	133,919
Female	33,981	40,016	40,197	80,422	91,209
Brain & Nervous System					
Male	34,921	36,642	36,751	87,853	90,304
Female	19,339	20,344	20,376	47,814	49,141
Female	15,582	16,298	16,375	40,039	41,163
Thyroid					
Male	82,780	88,485	88,667	261,607	273,642
Female	19,086	20,927	20,976	59,817	63,033
Female	63,694	67,558	67,691	201,790	210,609
Hodgkin Lymphoma					
Male	33,445	34,715	34,726	120,684	123,181
Female	17,766	18,435	18,435	62,988	64,279
Female	15,679	16,280	16,291	57,696	58,902
Non-Hodgkin Lymphoma					
Male	149,775	169,517	169,968	316,297	346,749
Female	78,674	89,136	89,398	163,797	178,734
Female	71,101	80,381	80,570	152,500	168,015
Myeloma					
Male	34,489	39,418	39,430	48,987	54,930
Female	18,725	21,516	21,528	26,710	30,013
Female	15,764	17,902	17,902	22,277	24,917
Leukemia					
Male	76,511	85,522	85,546	173,859	187,285
Female	43,969	49,297	49,309	97,991	105,642
Female	32,542	36,225	36,237	75,868	81,643
Acute Lymphocytic Leuk					
Male	13,506	13,685	13,685	42,330	42,610
Female	7,536	7,633	7,633	23,174	23,304
Female	5,970	6,052	6,052	19,156	19,306
Childhood (0-19)					
Male	54,418	54,527	54,800	204,131	204,569
Female	28,817	28,871	29,025	104,636	104,829
Female	25,601	25,656	25,775	99,495	99,740

^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER 9 registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census.

^{bcde} (b) Prevalence estimates are ambiguous for those with multiple cancers, unless the tumor inclusion criteria are understood. Depending on the application, different inclusion criteria may be appropriate. This table provides three different methods of tumor inclusion: (c) First invasive tumor ever; (d) First invasive tumor for each cancer site diagnosed during the previous 26 years (1975-2000); (e) First invasive tumor for each cancer site diagnosed during the previous 5 years (1996-2000). For definitions (d) and (e) all sites is treated as a separate cancer "site".

Consider a woman who had three invasive cancers: Melanoma in 1981; Breast cancer in 1997; Melanoma in 1998.

In method (c) the melanoma is the woman's first cancer, and is counted for the melanoma and all sites 26-year limited duration prevalence. For 5-year limited duration prevalence, the woman is not counted at all since her first cancer occurred 20 years ago.

In method (d) the 1981 melanoma is counted for the melanoma and all sites 26-year limited duration prevalence. The 1997 breast cancer is counted for the breast 5-year and 26-year limited duration prevalence.

In method (e) the 1997 breast cancer is counted for the breast cancer and all sites 5-year limited duration prevalence. The 1998 melanoma is counted for 5-year limited duration prevalence for melanoma.

Table II-12
ALL SITES (Invasive)Estimated United States Cancer Prevalence Counts^a on January 1, 2001
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^{e,h}
<u>Race/Sex</u>									
All Races ^b	3,765,929	2,409,640	1,410,983	832,269	542,570	6,522,090	9,048,822	760,218	9,809,040
Males	1,898,532	1,213,034	572,801	293,261	163,916	3,264,449	4,163,637	157,484	4,321,121
Females	1,867,397	1,196,606	838,182	539,008	378,654	3,257,641	4,885,185	602,734	5,487,919
White ^b	3,280,511	2,140,626	1,280,741	758,890	496,384	5,734,878	8,037,307	674,680	8,711,987
Males	1,641,392	1,071,727	522,570	268,582	151,104	2,852,303	3,675,672	145,211	3,820,883
Females	1,639,119	1,068,899	758,171	490,308	345,280	2,882,575	4,361,635	529,469	4,891,104
Black ^b	342,611	191,200	89,855	50,405	31,011	557,030	709,908	41,802	751,710
Males	185,889	104,653	35,472	16,917	8,403	300,563	352,415	5,205	357,620
Females	156,722	86,547	54,383	33,488	22,608	256,467	357,493	36,597	394,090
Asian/Pacific Islander ^c	79,924	42,826	+	+	+	128,513	+	+	+
Males	35,024	17,614	+	+	+	54,697	+	+	+
Females	44,900	25,212	+	+	+	73,816	+	+	+
Hispanic ^d	175,950	100,089	+	+	+	290,276	+	+	+
Males	84,529	46,015	+	+	+	136,143	+	+	+
Females	91,421	54,074	+	+	+	154,133	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	1.9251%	0.0759%	0.0994%	0.2014%	0.5219%	1.2325%	2.9832%	6.5481%	10.5728%	10.4983%	2.1054%
Males	1.9471%	0.0810%	0.1033%	0.1875%	0.4121%	0.8204%	2.4413%	7.5631%	14.6265%	15.6310%	2.4508%
Females	1.9037%	0.0706%	0.0954%	0.2160%	0.6347%	1.6377%	3.4978%	5.6497%	7.5352%	7.7750%	1.9058%
White ^c	2.0992%	0.0822%	0.1079%	0.2228%	0.5644%	1.2894%	3.0870%	6.8604%	10.9466%	10.6539%	2.1861%
Males	2.0946%	0.0878%	0.1109%	0.2096%	0.4534%	0.8637%	2.4918%	7.7550%	14.8675%	15.6128%	2.5002%
Females	2.1038%	0.0764%	0.1046%	0.2371%	0.6826%	1.7215%	3.6673%	6.0484%	7.9756%	8.0679%	2.0136%
Black ^c	1.3984%	0.0493%	0.0739%	0.1346%	0.3801%	1.0500%	2.9712%	6.8665%	10.6346%	9.1821%	2.0285%
Males	1.5986%	0.0509%	0.0797%	0.1163%	0.2588%	0.7471%	3.0449%	9.6854%	17.4957%	16.3750%	2.8247%
Females	1.2169%	0.0477%	0.0679%	0.1513%	0.4879%	1.3147%	2.9094%	4.6359%	6.1102%	5.8783%	1.5363%
Asian/Pacific Islander ^c	1.1940%	0.0640%	0.0730%	0.1198%	0.3279%	0.9177%	2.0300%	3.7514%	6.6994%	8.0337%	1.3926%
Males	1.1052%	0.0689%	0.0769%	0.1014%	0.2179%	0.4745%	1.2609%	3.7775%	9.0317%	11.6043%	1.4820%
Females	1.2769%	0.0588%	0.0688%	0.1376%	0.4312%	1.3140%	2.7052%	3.7301%	4.9490%	5.5513%	1.3560%
Hispanic ^d	0.7392%	0.0720%	0.0899%	0.1335%	0.3598%	0.8659%	1.9391%	4.2548%	7.6086%	7.8589%	1.4766%
Males	0.6686%	0.0757%	0.0931%	0.1250%	0.2624%	0.4589%	1.4142%	4.8558%	11.0489%	11.9357%	1.7297%
Females	0.8133%	0.0681%	0.0866%	0.1436%	0.4698%	1.2760%	2.4194%	3.7616%	5.1579%	5.6525%	1.3525%

^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas (d) SEER 11 Areas excluding Hawaii and Detroit.

^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{g h} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table III-12
BRAIN AND OTHER NERVOUS SYSTEM (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2001
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
<u>Race/Sex</u>									
All Races ^b	34,921	21,211	15,147	8,980	6,552	59,591	87,853	14,153	102,006
Males	19,339	11,972	7,962	4,703	3,317	33,247	47,814	7,258 ⁱ	55,072
Females	15,582	9,239	7,185	4,277	3,235	26,344	40,039	6,895 ⁱ	46,934
White ^b	31,194	18,671	13,350	7,959	5,928	52,770	78,053	10,986	89,039
Males	17,386	10,555	6,871	4,179	3,022	29,576	42,491	5,843 ⁱ	48,334
Females	13,808	8,116	6,479	3,780	2,906	23,194	35,562	5,143 ⁱ	40,705
Black ^b	2,430	1,644	1,337	795	453	4,522	6,746	+	+
Males	1,289	890	793	416	206	2,424	3,639	+	+
Females	1,141	754	544	379	247	2,098	3,107	+	+
Asian/Pacific Islander ^c	824	457	+	+	+	1,360	+	+	+
Males	441	262	+	+	+	754	+	+	+
Females	383	195	+	+	+	606	+	+	+
Hispanic ^d	2,791	1,559	+	+	+	4,620	+	+	+
Males	1,483	800	+	+	+	2,438	+	+	+
Females	1,308	759	+	+	+	2,182	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0181%	0.0121%	0.0189%	0.0142%	0.0196%	0.0224%	0.0229%	0.0205%	0.0177%	0.0104%	0.0182%
Males	0.0204%	0.0133%	0.0202%	0.0156%	0.0217%	0.0251%	0.0269%	0.0253%	0.0209%	0.0121%	0.0205%
Females	0.0159%	0.0109%	0.0175%	0.0128%	0.0174%	0.0198%	0.0191%	0.0162%	0.0153%	0.0095%	0.0160%
White ^c	0.0202%	0.0136%	0.0205%	0.0155%	0.0224%	0.0255%	0.0257%	0.0229%	0.0190%	0.0096%	0.0203%
Males	0.0226%	0.0149%	0.0219%	0.0168%	0.0244%	0.0283%	0.0297%	0.0287%	0.0225%	0.0112%	0.0227%
Females	0.0178%	0.0122%	0.0191%	0.0141%	0.0203%	0.0227%	0.0218%	0.0176%	0.0163%	0.0087%	0.0179%
Black ^c	0.0112%	0.0081%	0.0142%	0.0103%	0.0100%	0.0121%	0.0143%	0.0102%	0.0096%	0.0133%	0.0113%
Males	0.0125%	0.0092%	0.0145%	0.0115%	0.0116%	0.0149%	0.0190%	0.0092%	0.0087%	-	0.0125%
Females	0.0099%	0.0070%	0.0140%	0.0092%	0.0086%	0.0096%	0.0103%	0.0110%	0.0103%	0.0156%	0.0101%
Asian/Pacific Islander ^c	0.0106%	0.0070%	0.0126%	0.0100%	0.0100%	0.0112%	0.0109%	0.0119%	0.0133%	0.0157%	0.0108%
Males	0.0121%	0.0079%	0.0132%	0.0116%	0.0121%	0.0106%	0.0137%	0.0140%	0.0180%	0.0231%	0.0124%
Females	0.0093%	0.0061%	0.0120%	0.0084%	0.0081%	0.0117%	0.0085%	0.0101%	0.0097%	0.0106%	0.0094%
Hispanic ^d	0.0120%	0.0101%	0.0122%	0.0071%	0.0137%	0.0164%	0.0180%	0.0154%	0.0127%	0.0112%	0.0130%
Males	0.0123%	0.0105%	0.0127%	0.0075%	0.0135%	0.0165%	0.0194%	0.0175%	0.0103%	-	0.0134%
Females	0.0118%	0.0097%	0.0117%	0.0067%	0.0139%	0.0162%	0.0168%	0.0137%	0.0144%	0.0108%	0.0126%

^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.
^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.
^{g h i} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.
 - Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
 + Not available.

Table IV-17
BREAST CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2001
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
Race/Sex									
All Races ^b	795,912	543,613	373,231	206,928	127,643	1,427,054	2,069,330	141,227	2,210,557
Males	4,879	2,640	1,676	926	574	7,932	10,737	426	11,163
Females	791,033	540,973	371,555	206,002	127,069	1,419,122	2,058,593	140,801	2,199,394
White ^b	700,869	487,453	340,297	188,493	117,015	1,267,718	1,854,404	125,610	1,980,014
Males	4,255	2,313	1,475	848	530	6,917	9,465	145 ⁱ	9,610
Females	696,614	485,140	338,822	187,645	116,485	1,260,801	1,844,939	125,465	1,970,404
Black ^b	66,979	40,377	23,971	13,656	7,621	113,305	153,842	+	+
Males	508	237	184	67	37	805	1,031	+	+
Females	66,471	40,140	23,787	13,589	7,584	112,500	152,811	9,628	162,439
Asian/Pacific Islander ^c	19,379	11,081	+	+	+	32,099	+	+	+
Males	54	60	+	+	+	120	+	+	+
Females	19,325	11,021	+	+	+	31,979	+	+	+
Hispanic ^d	34,959	21,292	+	+	+	59,416	+	+	+
Males	155	59	+	+	+	224	+	+	+
Females	34,804	21,233	+	+	+	59,192	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race/Sex											
All Races ^c	0.4211%	-	0.0001%	0.0050%	0.0750%	0.3804%	0.9446%	1.4084%	1.8604%	2.0605%	0.4571%
Males	0.0048%	-	-	-	0.0003%	0.0023%	0.0086%	0.0210%	0.0298%	0.0426%	0.0061%
Females	0.8274%	-	0.0003%	0.0102%	0.1517%	0.7523%	1.8335%	2.6367%	3.2321%	3.1312%	0.8318%
White ^c	0.4635%	-	0.0001%	0.0047%	0.0734%	0.3880%	0.9836%	1.5017%	1.9871%	2.1824%	0.4804%
Males	0.0052%	-	-	-	0.0004%	0.0022%	0.0082%	0.0216%	0.0306%	0.0443%	0.0062%
Females	0.9185%	-	0.0002%	0.0097%	0.1511%	0.7795%	1.9346%	2.8452%	3.4697%	3.2974%	0.8814%
Black ^c	0.2859%	-	-	0.0077%	0.0948%	0.3623%	0.8425%	1.1809%	1.5083%	1.5897%	0.3947%
Males	0.0046%	-	-	-	-	0.0024%	0.0146%	0.0269%	0.0415%	0.0424%	0.0078%
Females	0.5410%	-	-	0.0146%	0.1787%	0.6766%	1.5363%	2.0940%	2.4755%	2.3004%	0.6822%
Asian/Pacific Islander ^c	0.2851%	-	-	0.0047%	0.0659%	0.3452%	0.7472%	0.9535%	1.1215%	1.0811%	0.3188%
Males	0.0024%	-	-	-	-	0.0019%	0.0036%	0.0084%	0.0167%	0.0280%	0.0032%
Females	0.5490%	-	-	0.0093%	0.1279%	0.6522%	1.3999%	1.7253%	1.9507%	1.8133%	0.5783%
Hispanic ^d	0.1527%	-	-	0.0036%	0.0485%	0.2682%	0.6085%	0.9032%	1.2305%	1.3962%	0.3041%
Males	0.0011%	-	-	-	-	0.0015%	0.0038%	0.0089%	0.0130%	0.0238%	0.0030%
Females	0.3120%	-	-	0.0079%	0.1032%	0.5369%	1.1618%	1.6370%	2.0978%	2.1390%	0.5464%

- ^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
- ^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.
- ^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
- ^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.
- ^{g h i} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.
- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
- + Not available.

Table V-10
CERVIX UTERI CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2001
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
<u>Race</u>									
All Races ^b	46,131	38,200	34,144	25,961	25,140	91,890	174,429	47,572 ⁱ	222,001
White ^b	35,988	31,218	28,217	21,334	20,503	73,391	141,320	35,052 ⁱ	176,372
Black ^b	7,012	4,559	4,090	3,573	3,621	12,582	23,533	+	+
Asian/Pacific Islander ^c	1,979	1,538	+	+	+	3,763	+	+	+
Hispanic ^d	7,684	6,073	+	+	+	14,831	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted ^f All Ages
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	
<u>Race</u>											
All Races ^c	0.0629%	-	-	0.0124%	0.0787%	0.1268%	0.1236%	0.1163%	0.0953%	0.0629%	0.0631%
White ^c	0.0630%	-	-	0.0138%	0.0873%	0.1312%	0.1174%	0.1052%	0.0799%	0.0501%	0.0625%
Black ^c	0.0591%	-	-	0.0110%	0.0656%	0.1169%	0.1269%	0.1483%	0.1431%	0.1489%	0.0680%
Asian/Pacific Islander ^c	0.0617%	-	-	0.0028%	0.0334%	0.0982%	0.1543%	0.1565%	0.1700%	0.1261%	0.0634%
Hispanic ^d	0.0779%	-	-	0.0115%	0.0982%	0.2142%	0.2340%	0.2352%	0.2266%	0.1451%	0.1116%

- ^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
- ^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.
- ^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
- ^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.
- ^{g h i} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.
- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
- + Not available.

Table VI-18
COLON AND RECTUM CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2001
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
Race/Sex									
All Races ^b	403,032	232,979	164,231	102,328	58,789	674,705	968,947	67,744	1,036,691
Males	201,690	114,621	81,118	46,565	24,428	335,541	471,439	25,149	496,588
Females	201,342	118,358	83,113	55,763	34,361	339,164	497,508	42,595	540,103
White ^b	349,697	205,229	146,168	92,231	52,753	589,287	853,052	60,764	913,816
Males	176,447	101,593	72,947	42,127	22,009	295,196	417,930	22,763	440,693
Females	173,250	103,636	73,221	50,104	30,744	294,091	435,122	38,001	473,123
Black ^b	37,684	19,814	13,069	7,334	4,247	60,609	82,571	4,634	87,205
Males	16,812	8,878	5,536	2,967	1,549	27,105	35,883	1,595	37,478
Females	20,872	10,936	7,533	4,367	2,698	33,504	46,688	3,039	49,727
Asian/Pacific Islander ^c	10,944	5,603	+	+	+	17,347	+	+	+
Males	5,817	2,889	+	+	+	9,134	+	+	+
Females	5,127	2,714	+	+	+	8,213	+	+	+
Hispanic ^d	17,254	8,553	+	+	+	27,040	+	+	+
Males	9,468	4,468	+	+	+	14,498	+	+	+
Females	7,786	4,085	+	+	+	12,542	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race/Sex											
All Races ^c	0.1983%	-	0.0003%	0.0028%	0.0182%	0.0694%	0.2390%	0.6733%	1.2444%	1.7269%	0.2207%
Males	0.2017%	-	0.0002%	0.0025%	0.0189%	0.0736%	0.2703%	0.8184%	1.5065%	1.9713%	0.2600%
Females	0.1949%	-	0.0004%	0.0030%	0.0175%	0.0652%	0.2093%	0.5448%	1.0479%	1.5972%	0.1907%
White ^c	0.2108%	-	0.0003%	0.0026%	0.0177%	0.0658%	0.2310%	0.6712%	1.2573%	1.7548%	0.2205%
Males	0.2133%	-	0.0003%	0.0025%	0.0184%	0.0689%	0.2664%	0.8171%	1.5281%	1.9968%	0.2607%
Females	0.2082%	-	0.0003%	0.0028%	0.0168%	0.0626%	0.1964%	0.5387%	1.0521%	1.6286%	0.1893%
Black ^c	0.1516%	-	-	0.0031%	0.0187%	0.0846%	0.3031%	0.7724%	1.2697%	1.4860%	0.2318%
Males	0.1455%	-	-	0.0024%	0.0174%	0.0919%	0.3275%	0.8823%	1.3820%	1.6996%	0.2583%
Females	0.1572%	-	-	0.0037%	0.0197%	0.0783%	0.2826%	0.6854%	1.1957%	1.3878%	0.2145%
Asian/Pacific Islander ^c	0.1681%	-	-	0.0032%	0.0203%	0.0771%	0.2327%	0.6063%	1.1037%	1.5983%	0.2051%
Males	0.1832%	-	-	0.0025%	0.0218%	0.0871%	0.2440%	0.7607%	1.3966%	1.8898%	0.2466%
Females	0.1540%	-	-	0.0039%	0.0190%	0.0682%	0.2228%	0.4802%	0.8839%	1.3957%	0.1734%
Hispanic ^d	0.0681%	-	0.0004%	0.0024%	0.0138%	0.0491%	0.1772%	0.4749%	0.9084%	1.1848%	0.1586%
Males	0.0708%	-	-	0.0029%	0.0145%	0.0500%	0.2020%	0.6171%	1.2114%	1.3476%	0.1964%
Females	0.0653%	-	-	0.0019%	0.0129%	0.0482%	0.1545%	0.3582%	0.6925%	1.0968%	0.1311%

^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{g h} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table VII-10
CORPUS AND UTERUS, NOS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2001
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
<u>Race</u>									
All Races ^b	140,432	105,849	82,158	67,121	64,672	264,462	474,773	96,080	570,853
White ^b	127,016	97,961	77,056	63,723	61,587	242,104	441,307	86,284	527,591
Black ^b	7,805	4,561	3,045	1,892	1,630	12,930	19,237	1,871	21,108
Asian/Pacific Islander ^c	3,225	1,968	+	+	+	5,467	+	+	+
Hispanic ^d	5,965	3,852	+	+	+	10,546	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	All Ages	0-9	10-19	20-29	Age Specific (Crude)						Age-Adjusted ^f
					30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races ^c	0.1492%	-	-	0.0025%	0.0180%	0.0802%	0.2827%	0.5864%	0.6946%	0.5967%	0.1508%
White ^c	0.1717%	-	-	0.0022%	0.0175%	0.0806%	0.3069%	0.6594%	0.7710%	0.6471%	0.1649%
Black ^c	0.0611%	-	-	0.0029%	0.0149%	0.0440%	0.1267%	0.3214%	0.3958%	0.2936%	0.0806%
Asian/Pacific Islander ^c	0.0937%	-	-	0.0031%	0.0209%	0.1054%	0.2505%	0.3098%	0.3185%	0.2788%	0.0985%
Hispanic ^d	0.0546%	-	-	0.0034%	0.0225%	0.0650%	0.1991%	0.3601%	0.3846%	0.3114%	0.0957%

^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{g h} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table VIII-12
ESOPHAGUS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2001
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
<u>Race/Sex</u>									
All Races ^b	15,737	3,909	1,497	658	288	20,116	22,081	285	22,366
Males	11,952	2,822	1,103	438	209	15,097	16,515	140	16,655
Females	3,785	1,087	394	220	79	5,019	5,566	145	5,711
White ^b	13,553	3,396	1,185	497	245	17,303	18,880	238	19,118
Males	10,435	2,435	914	320	197	13,119	14,305	132	14,437
Females	3,118	961	271	177	48	4,184	4,575	106	4,681
Black ^b	1,773	432	271	155	40	2,307	2,660	77	2,737
Males	1,178	317	168	111	12	1,561	1,775	25	1,800
Females	595	115	103	44	28	746	885	52	937
Asian/Pacific Islander ^c	276	65	+	+	+	350	+	+	+
Males	217	51	+	+	+	275	+	+	+
Females	59	14	+	+	+	75	+	+	+
Hispanic ^d	621	168	+	+	+	808	+	+	+
Males	474	139	+	+	+	632	+	+	+
Females	147	29	+	+	+	176	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0057%	-	-	0.0001%	0.0005%	0.0026%	0.0107%	0.0249%	0.0330%	0.0245%	0.0063%
Males	0.0086%	-	-	-	0.0008%	0.0044%	0.0184%	0.0415%	0.0554%	0.0399%	0.0105%
Females	0.0028%	-	-	-	0.0002%	0.0008%	0.0034%	0.0103%	0.0161%	0.0164%	0.0028%
White ^c	0.0061%	-	-	-	0.0005%	0.0026%	0.0111%	0.0253%	0.0339%	0.0256%	0.0064%
Males	0.0093%	-	-	-	0.0008%	0.0045%	0.0195%	0.0420%	0.0572%	0.0422%	0.0108%
Females	0.0029%	-	-	-	-	0.0008%	0.0029%	0.0102%	0.0161%	0.0170%	0.0028%
Black ^c	0.0052%	-	-	-	-	0.0038%	0.0145%	0.0323%	0.0380%	0.0160%	0.0074%
Males	0.0072%	-	-	-	-	0.0058%	0.0211%	0.0489%	0.0640%	0.0212%	0.0114%
Females	0.0034%	-	-	-	-	0.0020%	0.0090%	0.0191%	0.0209%	0.0136%	0.0045%
Asian/Pacific Islander ^c	0.0034%	-	-	-	-	0.0012%	0.0044%	0.0166%	0.0240%	0.0179%	0.0040%
Males	0.0055%	-	-	-	-	0.0025%	0.0076%	0.0324%	0.0380%	0.0280%	0.0070%
Females	0.0015%	-	-	-	-	-	-	0.0038%	0.0135%	0.0108%	0.0017%
Hispanic ^d	0.0021%	-	-	-	0.0005%	0.0018%	0.0078%	0.0152%	0.0228%	0.0224%	0.0045%
Males	0.0031%	-	-	-	0.0008%	0.0032%	0.0139%	0.0281%	0.0360%	0.0360%	0.0076%
Females	0.0010%	-	-	-	-	-	0.0023%	0.0046%	0.0134%	0.0150%	0.0020%

^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.
^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.
^{g h} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26.
 - Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
 + Not available.

Table IX-12
HODGKIN LYMPHOMA (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2001
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
<u>Race/Sex</u>									
All Races ^b	33,445	27,810	23,968	19,234	13,836	66,620	120,684	21,058	141,742
Males	17,766	14,305	12,290	10,238	7,171	34,994	62,988	11,295	74,283
Females	15,679	13,505	11,678	8,996	6,665	31,626	57,696	9,763	67,459
White ^b	29,104	24,800	21,925	17,920	12,868	58,725	108,868	19,251	128,119
Males	15,425	12,788	11,270	9,438	6,509	30,918	56,555	10,012	66,567
Females	13,679	12,012	10,655	8,482	6,359	27,807	52,313	9,239	61,552
Black ^b	3,421	2,463	1,603	1,100	820	6,352	9,510	1,292	10,802
Males	1,793	1,226	774	699	579	3,207	5,139	888	6,027
Females	1,628	1,237	829	401	241	3,145	4,371	404	4,775
Asian/Pacific Islander ^c	461	297	+	+	+	799	+	+	+
Males	254	141	+	+	+	406	+	+	+
Females	207	156	+	+	+	393	+	+	+
Hispanic ^d	2,546	1,579	+	+	+	4,464	+	+	+
Males	1,449	903	+	+	+	2,526	+	+	+
Females	1,097	676	+	+	+	1,938	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0197%	0.0008%	0.0082%	0.0304%	0.0363%	0.0250%	0.0187%	0.0182%	0.0166%	0.0118%	0.0194%
Males	0.0212%	0.0013%	0.0082%	0.0288%	0.0370%	0.0281%	0.0232%	0.0235%	0.0199%	0.0150%	0.0210%
Females	0.0183%	0.0003%	0.0082%	0.0320%	0.0355%	0.0220%	0.0145%	0.0135%	0.0141%	0.0101%	0.0180%
White ^c	0.0223%	0.0010%	0.0096%	0.0346%	0.0411%	0.0283%	0.0205%	0.0203%	0.0181%	0.0132%	0.0219%
Males	0.0238%	0.0016%	0.0093%	0.0325%	0.0415%	0.0312%	0.0249%	0.0262%	0.0214%	0.0164%	0.0234%
Females	0.0208%	0.0003%	0.0099%	0.0368%	0.0407%	0.0253%	0.0162%	0.0149%	0.0156%	0.0116%	0.0206%
Black ^c	0.0163%	-	0.0054%	0.0280%	0.0315%	0.0209%	0.0190%	0.0160%	0.0117%	-	0.0165%
Males	0.0176%	-	0.0064%	0.0260%	0.0336%	0.0256%	0.0255%	0.0183%	0.0121%	-	0.0183%
Females	0.0151%	-	0.0044%	0.0298%	0.0297%	0.0168%	0.0135%	0.0142%	0.0114%	-	0.0149%
Asian/Pacific Islander ^c	0.0063%	-	0.0030%	0.0087%	0.0105%	0.0069%	0.0055%	0.0052%	0.0088%	0.0073%	0.0060%
Males	0.0068%	-	0.0039%	0.0081%	0.0099%	0.0069%	0.0072%	0.0077%	0.0141%	0.0127%	0.0069%
Females	0.0057%	-	0.0020%	0.0093%	0.0111%	0.0069%	0.0040%	-	0.0048%	-	0.0053%
Hispanic ^d	0.0114%	0.0013%	0.0086%	0.0151%	0.0175%	0.0123%	0.0152%	0.0207%	0.0180%	0.0114%	0.0126%
Males	0.0126%	0.0022%	0.0092%	0.0148%	0.0180%	0.0152%	0.0177%	0.0224%	0.0332%	0.0199%	0.0148%
Females	0.0102%	-	0.0078%	0.0155%	0.0169%	0.0093%	0.0130%	0.0192%	0.0072%	-	0.0107%

^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.
^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.
^{g h} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26.
 - Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
 + Not available.

Table XI-12
KIDNEY AND RENAL PELVIS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2001
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
Race/Sex									
All Races ^b	85,215	52,025	30,386	17,501	10,439	144,822	196,981	14,013	210,994
Males	51,234	30,962	17,228	10,606	5,653	86,613	116,559	7,794	124,353
Females	33,981	21,063	13,158	6,895	4,786	58,209	80,422	6,219	86,641
White ^b	73,002	45,559	27,226	15,629	9,345	125,325	172,073	13,851	185,924
Males	44,504	27,342	15,592	9,638	5,149	75,831	103,061	6,876	109,937
Females	28,498	18,217	11,634	5,991	4,196	49,494	69,012	6,975	75,987
Black ^b	9,493	5,096	2,615	1,568	901	15,254	19,745	2,092	21,837
Males	5,051	2,789	1,335	778	398	8,190	10,384	722	11,106
Females	4,442	2,307	1,280	790	503	7,064	9,361	1,370	10,731
Asian/Pacific Islander ^c	1,491	695	+	+	+	2,272	+	+	+
Males	913	435	+	+	+	1,410	+	+	+
Females	578	260	+	+	+	862	+	+	+
Hispanic ^d	5,300	2,955	+	+	+	8,729	+	+	+
Males	3,045	1,726	+	+	+	4,962	+	+	+
Females	2,255	1,229	+	+	+	3,767	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race/Sex											
All Races ^c	0.0419%	0.0075%	0.0040%	0.0014%	0.0070%	0.0276%	0.0809%	0.1563%	0.2051%	0.1787%	0.0456%
Males	0.0512%	0.0067%	0.0038%	0.0013%	0.0077%	0.0351%	0.1083%	0.2114%	0.2840%	0.2692%	0.0615%
Females	0.0328%	0.0085%	0.0042%	0.0014%	0.0063%	0.0203%	0.0548%	0.1076%	0.1461%	0.1306%	0.0330%
White ^c	0.0458%	0.0080%	0.0042%	0.0013%	0.0070%	0.0288%	0.0833%	0.1656%	0.2178%	0.1842%	0.0478%
Males	0.0559%	0.0076%	0.0038%	0.0011%	0.0078%	0.0371%	0.1114%	0.2207%	0.3014%	0.2805%	0.0643%
Females	0.0358%	0.0085%	0.0045%	0.0015%	0.0062%	0.0205%	0.0559%	0.1157%	0.1545%	0.1340%	0.0344%
Black ^c	0.0365%	0.0079%	0.0056%	0.0020%	0.0092%	0.0330%	0.1028%	0.1667%	0.1944%	0.1734%	0.0498%
Males	0.0431%	0.0048%	0.0064%	0.0022%	0.0090%	0.0415%	0.1388%	0.2285%	0.2522%	0.2629%	0.0657%
Females	0.0306%	0.0112%	0.0047%	0.0018%	0.0094%	0.0256%	0.0727%	0.1178%	0.1563%	0.1324%	0.0383%
Asian/Pacific Islander ^c	0.0210%	0.0036%	-	0.0013%	0.0049%	0.0138%	0.0408%	0.0774%	0.1115%	0.1151%	0.0243%
Males	0.0268%	0.0020%	-	0.0020%	0.0062%	0.0166%	0.0553%	0.1157%	0.1599%	0.1585%	0.0335%
Females	0.0156%	0.0052%	-	-	0.0038%	0.0114%	0.0281%	0.0460%	0.0752%	0.0849%	0.0170%
Hispanic ^d	0.0220%	0.0061%	0.0024%	0.0012%	0.0051%	0.0244%	0.0691%	0.1452%	0.2163%	0.2026%	0.0441%
Males	0.0245%	0.0045%	0.0016%	0.0013%	0.0057%	0.0268%	0.0940%	0.2024%	0.2791%	0.2963%	0.0583%
Females	0.0194%	0.0078%	0.0032%	0.0012%	0.0044%	0.0220%	0.0464%	0.0984%	0.1715%	0.1519%	0.0335%

^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{g h} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XII-12
LARYNX CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2001
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
<u>Race/Sex</u>									
All Races ^b	34,076	23,259	16,532	10,943	6,730	61,341	92,464	7,032	99,496
Males	26,759	18,794	13,348	8,792	5,554	48,693	74,006	5,737	79,743
Females	7,317	4,465	3,184	2,151	1,176	12,648	18,458	1,295	19,753
White ^b	28,697	20,104	14,370	9,910	6,071	52,321	79,936	5,767	85,703
Males	22,728	16,237	11,727	7,992	5,007	41,733	64,346	4,696	69,042
Females	5,969	3,867	2,643	1,918	1,064	10,588	15,590	1,071	16,661
Black ^b	4,613	2,709	1,804	927	553	7,727	10,725	506	11,231
Males	3,413	2,199	1,375	710	464	5,912	8,250	417	8,667
Females	1,200	510	429	217	89	1,815	2,475	89	2,564
Asian/Pacific Islander ^c	450	254	+	+	+	734	+	+	+
Males	373	217	+	+	+	616	+	+	+
Females	77	37	+	+	+	118	+	+	+
Hispanic ^d	1,388	792	+	+	+	2,365	+	+	+
Males	1,135	668	+	+	+	1,965	+	+	+
Females	253	124	+	+	+	400	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0170%	-	-	0.0002%	0.0010%	0.0069%	0.0303%	0.0812%	0.1021%	0.0675%	0.0188%
Males	0.0274%	-	-	-	0.0011%	0.0102%	0.0512%	0.1392%	0.1910%	0.1534%	0.0340%
Females	0.0068%	-	-	0.0002%	0.0008%	0.0037%	0.0105%	0.0300%	0.0355%	0.0220%	0.0069%
White ^c	0.0183%	-	-	0.0002%	0.0011%	0.0072%	0.0302%	0.0852%	0.1038%	0.0687%	0.0193%
Males	0.0294%	-	-	-	0.0011%	0.0103%	0.0511%	0.1454%	0.1920%	0.1578%	0.0347%
Females	0.0073%	-	-	-	0.0010%	0.0040%	0.0099%	0.0305%	0.0369%	0.0222%	0.0071%
Black ^c	0.0190%	-	-	-	0.0008%	0.0110%	0.0536%	0.1187%	0.1430%	0.0787%	0.0276%
Males	0.0305%	-	-	-	-	0.0172%	0.0910%	0.2018%	0.2878%	0.1781%	0.0507%
Females	0.0086%	-	-	-	-	0.0056%	0.0222%	0.0529%	0.0475%	0.0331%	0.0113%
Asian/Pacific Islander ^c	0.0072%	-	-	-	-	0.0018%	0.0110%	0.0267%	0.0624%	0.0457%	0.0087%
Males	0.0126%	-	-	-	-	0.0034%	0.0194%	0.0473%	0.1245%	0.0963%	0.0169%
Females	0.0022%	-	-	-	-	-	0.0036%	0.0100%	0.0158%	0.0106%	0.0025%
Hispanic ^d	0.0057%	-	-	-	0.0010%	0.0035%	0.0155%	0.0555%	0.0738%	0.0483%	0.0125%
Males	0.0092%	-	-	-	0.0009%	0.0055%	0.0289%	0.1053%	0.1441%	0.1151%	0.0243%
Females	0.0021%	-	-	-	0.0011%	0.0016%	0.0033%	0.0146%	0.0237%	0.0121%	0.0037%

^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.
^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.
^{g h} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26.
 - Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
 + Not available.

Table XIII-18
LEUKEMIA (Invasive)Estimated United States Cancer Prevalence Counts^a on January 1, 2001
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
Race/Sex									
All Races ^b	76,511	45,977	25,608	15,529	9,150	128,514	173,859	6,611	180,470
Males	43,969	26,162	14,350	8,429	4,548	73,260	97,991	2,778	100,769
Females	32,542	19,815	11,258	7,100	4,602	55,254	75,868	3,833	79,701
White ^b	67,895	41,550	23,305	14,180	8,479	114,916	156,446	5,736	162,182
Males	39,112	23,826	13,242	7,718	4,291	65,845	88,701	2,403	91,104
Females	28,783	17,724	10,063	6,462	4,188	49,071	67,745	3,333	71,078
Black ^b	5,353	2,741	1,266	827	417	8,394	10,644	642	11,286
Males	2,809	1,371	556	464	158	4,276	5,380	298	5,678
Females	2,544	1,370	710	363	259	4,118	5,264	344	5,608
Asian/Pacific Islander ^c	1,607	884	+	+	+	2,661	+	+	+
Males	931	459	+	+	+	1,474	+	+	+
Females	676	425	+	+	+	1,187	+	+	+
Hispanic ^d	5,832	3,423	+	+	+	9,719	+	+	+
Males	3,289	1,985	+	+	+	5,584	+	+	+
Females	2,543	1,438	+	+	+	4,135	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race/Sex											
All Races ^c	0.0393%	0.0289%	0.0278%	0.0124%	0.0138%	0.0217%	0.0490%	0.0959%	0.1487%	0.1571%	0.0417%
Males	0.0457%	0.0311%	0.0290%	0.0141%	0.0164%	0.0270%	0.0624%	0.1286%	0.2011%	0.2084%	0.0523%
Females	0.0330%	0.0267%	0.0265%	0.0106%	0.0111%	0.0166%	0.0364%	0.0670%	0.1095%	0.1298%	0.0330%
White ^c	0.0441%	0.0323%	0.0309%	0.0138%	0.0139%	0.0229%	0.0525%	0.1057%	0.1643%	0.1702%	0.0454%
Males	0.0512%	0.0341%	0.0321%	0.0160%	0.0162%	0.0288%	0.0673%	0.1413%	0.2216%	0.2288%	0.0570%
Females	0.0370%	0.0303%	0.0296%	0.0115%	0.0115%	0.0169%	0.0382%	0.0735%	0.1208%	0.1396%	0.0358%
Black ^c	0.0217%	0.0120%	0.0175%	0.0067%	0.0114%	0.0167%	0.0357%	0.0662%	0.0910%	0.0886%	0.0266%
Males	0.0243%	0.0147%	0.0180%	0.0061%	0.0145%	0.0182%	0.0444%	0.0885%	0.1126%	0.0949%	0.0318%
Females	0.0193%	0.0092%	0.0170%	0.0073%	0.0086%	0.0154%	0.0284%	0.0486%	0.0768%	0.0857%	0.0225%
Asian/Pacific Islander ^c	0.0213%	0.0286%	0.0212%	0.0090%	0.0137%	0.0154%	0.0268%	0.0327%	0.0480%	0.0485%	0.0226%
Males	0.0247%	0.0324%	0.0231%	0.0093%	0.0174%	0.0178%	0.0281%	0.0375%	0.0715%	0.0662%	0.0269%
Females	0.0181%	0.0247%	0.0192%	0.0087%	0.0101%	0.0133%	0.0255%	0.0288%	0.0303%	0.0362%	0.0190%
Hispanic ^d	0.0256%	0.0323%	0.0329%	0.0127%	0.0144%	0.0168%	0.0310%	0.0476%	0.0763%	0.0823%	0.0301%
Males	0.0283%	0.0339%	0.0347%	0.0152%	0.0169%	0.0195%	0.0378%	0.0618%	0.1050%	0.0953%	0.0357%
Females	0.0228%	0.0307%	0.0310%	0.0097%	0.0116%	0.0141%	0.0248%	0.0360%	0.0559%	0.0752%	0.0253%

^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{g h} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XIII-19
ACUTE LYMPHOCYTIC LEUKEMIA (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2001
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
<u>Race/Sex</u>									
All Races ^b	13,506	9,779	8,068	6,298	4,270	24,934	42,330	+	+
Males	7,536	5,514	4,541	3,203	2,171	13,915	23,174	+	+
Females	5,970	4,265	3,527	3,095	2,099	11,019	19,156	+	+
White ^b	11,699	8,487	7,098	5,703	3,919	21,673	37,299	+	+
Males	6,473	4,871	4,034	2,932	2,033	12,135	20,539	+	+
Females	5,226	3,616	3,064	2,771	1,886	9,538	16,760	+	+
Black ^b	1,015	755	503	315	177	1,828	2,780	+	+
Males	605	357	247	159	75	988	1,459	+	+
Females	410	398	256	156	102	840	1,321	+	+
Asian/Pacific Islander ^c	522	375	+	+	+	984	+	+	+
Males	335	180	+	+	+	557	+	+	+
Females	187	195	+	+	+	427	+	+	+
Hispanic ^d	2,991	1,882	+	+	+	5,121	+	+	+
Males	1,666	1,100	+	+	+	2,926	+	+	+
Females	1,325	782	+	+	+	2,195	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0090%	0.0252%	0.0234%	0.0059%	0.0027%	0.0016%	0.0020%	0.0017%	0.0014%	0.0007%	0.0088%
Males	0.0102%	0.0272%	0.0246%	0.0076%	0.0034%	0.0017%	0.0023%	0.0020%	0.0018%	-	0.0097%
Females	0.0078%	0.0231%	0.0221%	0.0041%	0.0021%	0.0015%	0.0016%	0.0014%	0.0012%	0.0007%	0.0079%
White ^c	0.0099%	0.0284%	0.0266%	0.0068%	0.0030%	0.0018%	0.0020%	0.0017%	0.0015%	0.0007%	0.0100%
Males	0.0111%	0.0302%	0.0281%	0.0087%	0.0037%	0.0019%	0.0023%	0.0022%	0.0017%	-	0.0109%
Females	0.0087%	0.0265%	0.0250%	0.0048%	0.0023%	0.0017%	0.0017%	0.0012%	0.0013%	0.0008%	0.0090%
Black ^c	0.0047%	0.0096%	0.0127%	0.0026%	0.0014%	-	0.0012%	-	-	-	0.0041%
Males	0.0056%	0.0118%	0.0126%	0.0029%	0.0021%	-	-	-	-	-	0.0046%
Females	0.0040%	0.0073%	0.0129%	0.0023%	-	-	-	-	-	-	0.0036%
Asian/Pacific Islander ^c	0.0071%	0.0238%	0.0160%	0.0036%	0.0024%	0.0021%	0.0021%	0.0017%	-	-	0.0073%
Males	0.0084%	0.0275%	0.0173%	0.0047%	0.0026%	0.0024%	0.0027%	-	-	-	0.0083%
Females	0.0058%	0.0199%	0.0145%	0.0025%	0.0023%	0.0018%	-	-	-	-	0.0063%
Hispanic ^d	0.0137%	0.0294%	0.0284%	0.0065%	0.0041%	0.0023%	0.0033%	0.0023%	-	-	0.0109%
Males	0.0151%	0.0312%	0.0305%	0.0082%	0.0053%	0.0027%	0.0038%	-	-	-	0.0118%
Females	0.0122%	0.0274%	0.0263%	0.0045%	0.0027%	0.0020%	0.0030%	-	-	-	0.0097%

- ^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
- ^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.
- ^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
- ^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.
- ^{g h} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26.
- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
- + Not available.

Table XIV-12
LIVER AND INTRAHEPATIC BILE DUCT CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2001
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
<u>Race/Sex</u>									
All Races ^b	10,956	1,492	891	538	222	12,660	14,108	373	14,481
Males	7,118	868	367	237	94	8,114	8,706	69	8,775
Females	3,838	624	524	301	128	4,546	5,402	304	5,706
White ^b	8,097	1,145	713	443	156	9,376	10,540	179	10,719
Males	5,080	672	254	179	68	5,817	6,249	54 ⁱ	6,303
Females	3,017	473	459	264	88	3,559	4,291	125 ⁱ	4,416
Black ^b	1,381	129	120	49	45	1,549	1,727	+	+
Males	991	59	71	26	22	1,075	1,171	+	+
Females	390	70	49	23	23	474	556	+	+
Asian/Pacific Islander ^c	1,247	205	+	+	+	1,472	+	+	+
Males	873	130	+	+	+	1,016	+	+	+
Females	374	75	+	+	+	456	+	+	+
Hispanic ^d	1,218	232	+	+	+	1,477	+	+	+
Males	772	146	+	+	+	934	+	+	+
Females	446	86	+	+	+	543	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0053%	0.0015%	0.0006%	0.0005%	0.0008%	0.0038%	0.0099%	0.0196%	0.0254%	0.0180%	0.0057%
Males	0.0069%	0.0016%	0.0006%	0.0004%	0.0009%	0.0052%	0.0160%	0.0279%	0.0377%	0.0233%	0.0081%
Females	0.0036%	0.0015%	0.0005%	0.0005%	0.0007%	0.0024%	0.0042%	0.0122%	0.0163%	0.0152%	0.0037%
White ^c	0.0042%	0.0016%	0.0006%	0.0004%	0.0006%	0.0030%	0.0074%	0.0131%	0.0195%	0.0143%	0.0043%
Males	0.0053%	0.0016%	0.0007%	0.0004%	0.0007%	0.0038%	0.0118%	0.0185%	0.0276%	0.0167%	0.0059%
Females	0.0031%	0.0015%	0.0004%	0.0005%	0.0006%	0.0022%	0.0032%	0.0083%	0.0134%	0.0131%	0.0030%
Black ^c	0.0037%	-	0.0008%	-	-	0.0030%	0.0110%	0.0202%	0.0222%	0.0080%	0.0051%
Males	0.0054%	-	-	-	-	0.0054%	0.0196%	0.0329%	0.0319%	-	0.0079%
Females	0.0022%	-	0.0014%	-	-	-	0.0037%	0.0102%	0.0157%	-	0.0029%
Asian/Pacific Islander ^c	0.0139%	0.0024%	-	0.0008%	0.0017%	0.0100%	0.0267%	0.0610%	0.0728%	0.0589%	0.0159%
Males	0.0199%	0.0021%	-	-	0.0027%	0.0151%	0.0438%	0.0893%	0.1197%	0.0834%	0.0245%
Females	0.0084%	0.0027%	-	-	-	0.0055%	0.0116%	0.0379%	0.0375%	0.0419%	0.0090%
Hispanic ^d	0.0039%	0.0015%	0.0007%	-	0.0006%	0.0036%	0.0116%	0.0249%	0.0422%	0.0293%	0.0076%
Males	0.0047%	0.0018%	0.0011%	-	-	0.0046%	0.0195%	0.0331%	0.0591%	0.0300%	0.0103%
Females	0.0030%	0.0013%	-	-	-	0.0026%	0.0043%	0.0182%	0.0301%	0.0289%	0.0053%

^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.
^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.
^{g h i} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.
 - Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
 + Not available.

Table XV-17
LUNG AND BRONCHUS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2001
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
Race/Sex									
All Races ^b	201,687	65,597	33,792	18,636	9,864	275,731	330,991	16,540	347,531
Males	101,771	31,625	16,925	9,065	5,175	137,354	165,394	9,659	175,053
Females	99,916	33,972	16,867	9,571	4,689	138,377	165,597	6,881	172,478
White ^b	174,418	58,006	29,982	16,579	8,691	239,871	288,892	8,771	297,663
Males	86,667	27,574	14,762	7,854	4,532	117,662	142,117	4,546	146,663
Females	87,751	30,432	15,220	8,725	4,159	122,209	146,775	4,225	151,000
Black ^b	21,217	6,066	3,097	1,678	870	28,107	33,088	879	33,967
Males	11,525	3,183	1,749	967	499	15,160	18,014	510	18,524
Females	9,692	2,883	1,348	711	371	12,947	15,074	369	15,443
Asian/Pacific Islander ^c	4,269	1,016	+	+	+	5,426	+	+	+
Males	2,532	597	+	+	+	3,208	+	+	+
Females	1,737	419	+	+	+	2,218	+	+	+
Hispanic ^d	5,186	1,554	+	+	+	7,017	+	+	+
Males	2,805	765	+	+	+	3,696	+	+	+
Females	2,381	789	+	+	+	3,321	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race/Sex											
All Races ^c	0.0790%	-	0.0001%	0.0011%	0.0043%	0.0254%	0.1157%	0.3512%	0.5468%	0.3766%	0.0881%
Males	0.0806%	-	-	0.0008%	0.0039%	0.0256%	0.1206%	0.3908%	0.6464%	0.4991%	0.1012%
Females	0.0775%	-	-	0.0014%	0.0047%	0.0252%	0.1111%	0.3161%	0.4721%	0.3117%	0.0787%
White ^c	0.0851%	-	0.0002%	0.0012%	0.0042%	0.0250%	0.1134%	0.3663%	0.5642%	0.3789%	0.0900%
Males	0.0837%	-	-	0.0009%	0.0038%	0.0240%	0.1121%	0.3934%	0.6463%	0.4946%	0.1000%
Females	0.0864%	-	-	0.0015%	0.0047%	0.0259%	0.1147%	0.3417%	0.5019%	0.3185%	0.0831%
Black ^c	0.0689%	-	-	-	0.0061%	0.0410%	0.1787%	0.4027%	0.5670%	0.3226%	0.1012%
Males	0.0784%	-	-	-	0.0065%	0.0477%	0.2239%	0.5082%	0.7307%	0.5098%	0.1309%
Females	0.0603%	-	-	-	0.0058%	0.0351%	0.1409%	0.3191%	0.4590%	0.2365%	0.0807%
Asian/Pacific Islander ^c	0.0537%	-	-	0.0009%	0.0032%	0.0149%	0.0796%	0.2164%	0.4061%	0.3978%	0.0648%
Males	0.0661%	-	-	-	0.0025%	0.0181%	0.0973%	0.2855%	0.5999%	0.5385%	0.0882%
Females	0.0422%	-	-	0.0014%	0.0039%	0.0119%	0.0640%	0.1600%	0.2607%	0.2999%	0.0469%
Hispanic ^d	0.0177%	-	-	0.0008%	0.0026%	0.0109%	0.0387%	0.1499%	0.2648%	0.2278%	0.0409%
Males	0.0179%	-	-	-	0.0027%	0.0107%	0.0456%	0.1677%	0.3334%	0.3316%	0.0505%
Females	0.0174%	-	-	0.0010%	0.0024%	0.0110%	0.0323%	0.1353%	0.2160%	0.1717%	0.0343%

^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.
^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.
^{g h} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26.
 - Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
 + Not available.

Table XVI-12
MELANOMA OF THE SKIN (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2001
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
<u>Race/Sex</u>									
All Races ^b	194,560	133,626	99,181	67,971	43,512	350,172	544,920	52,891	597,811
Males	103,566	68,108	46,919	30,234	18,008	182,537	269,150	18,561	287,711
Females	90,994	65,518	52,262	37,737	25,504	167,635	275,770	34,330	310,100
White ^b	186,420	127,426	94,117	64,754	41,900	334,906	520,314	49,881	570,195
Males	99,434	65,089	44,825	28,807	17,292	175,034	257,637	17,662	275,299
Females	86,986	62,337	49,292	35,947	24,608	159,872	262,677	32,219	294,896
Black ^b	817	534	384	226	76	1,422	2,084	160	2,244
Males	370	281	131	50	18	678	856	60	916
Females	447	253	253	176	58	744	1,228	100	1,328
Asian/Pacific Islander ^c	562	258	+	+	+	860	+	+	+
Males	256	117	+	+	+	386	+	+	+
Females	306	141	+	+	+	474	+	+	+
Hispanic ^d	3,050	1,857	+	+	+	5,234	+	+	+
Males	1,220	704	+	+	+	2,044	+	+	+
Females	1,830	1,153	+	+	+	3,190	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0981%	0.0003%	0.0021%	0.0228%	0.0686%	0.1313%	0.1983%	0.2595%	0.3142%	0.3002%	0.1035%
Males	0.1046%	0.0003%	0.0017%	0.0159%	0.0522%	0.1236%	0.2235%	0.3394%	0.4458%	0.4678%	0.1209%
Females	0.0917%	0.0003%	0.0025%	0.0301%	0.0855%	0.1389%	0.1743%	0.1888%	0.2156%	0.2113%	0.0916%
White ^c	0.1201%	0.0003%	0.0025%	0.0278%	0.0835%	0.1595%	0.2348%	0.3087%	0.3649%	0.3354%	0.1226%
Males	0.1274%	0.0004%	0.0019%	0.0187%	0.0621%	0.1480%	0.2615%	0.3999%	0.5165%	0.5326%	0.1415%
Females	0.1128%	-	0.0032%	0.0376%	0.1064%	0.1711%	0.2087%	0.2259%	0.2501%	0.2325%	0.1101%
Black ^c	0.0037%	-	-	0.0011%	0.0027%	0.0035%	0.0079%	0.0172%	0.0151%	0.0213%	0.0049%
Males	0.0038%	-	-	-	0.0021%	0.0040%	0.0080%	0.0229%	0.0225%	-	0.0057%
Females	0.0035%	-	-	-	0.0033%	0.0030%	0.0077%	0.0126%	0.0103%	0.0222%	0.0042%
Asian/Pacific Islander ^c	0.0075%	-	-	0.0024%	0.0053%	0.0075%	0.0152%	0.0208%	0.0252%	0.0348%	0.0083%
Males	0.0072%	-	-	0.0029%	0.0024%	0.0054%	0.0163%	0.0249%	0.0311%	0.0417%	0.0087%
Females	0.0079%	-	-	0.0020%	0.0081%	0.0093%	0.0143%	0.0175%	0.0208%	0.0301%	0.0082%
Hispanic ^d	0.0134%	-	-	0.0034%	0.0131%	0.0275%	0.0444%	0.0494%	0.0759%	0.0887%	0.0229%
Males	0.0100%	-	-	0.0019%	0.0070%	0.0203%	0.0360%	0.0484%	0.0950%	0.0995%	0.0213%
Females	0.0169%	-	-	0.0051%	0.0199%	0.0347%	0.0520%	0.0502%	0.0623%	0.0829%	0.0251%

^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{g h} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XVII-12
MYELOMA (Invasive)Estimated United States Cancer Prevalence Counts^a on January 1, 2001
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
<u>Race/Sex</u>									
All Races ^b	34,489	9,759	3,114	1,185	369	45,237	48,987	605	49,592
Males	18,725	5,474	1,775	577	145	24,706	26,710	164	26,874
Females	15,764	4,285	1,339	608	224	20,531	22,277	441	22,718
White ^b	27,453	7,483	2,390	768	318	35,709	38,446	285	38,731
Males	15,470	4,248	1,374	326	142	20,112	21,566	155 ⁱ	21,721
Females	11,983	3,235	1,016	442	176	15,597	16,880	130 ⁱ	17,010
Black ^b	5,960	1,921	629	376	39	8,074	8,964	+	+
Males	2,711	1,015	364	238	0	3,828	4,340	+	+
Females	3,249	906	265	138	39	4,246	4,624	+	+
Asian/Pacific Islander ^c	639	177	+	+	+	831	+	+	+
Males	331	105	+	+	+	438	+	+	+
Females	308	72	+	+	+	393	+	+	+
Hispanic ^d	1,947	504	+	+	+	2,515	+	+	+
Males	1,092	311	+	+	+	1,434	+	+	+
Females	855	193	+	+	+	1,081	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0138%	-	-	0.0002%	0.0014%	0.0066%	0.0255%	0.0519%	0.0799%	0.0796%	0.0152%
Males	0.0153%	-	-	0.0002%	0.0018%	0.0084%	0.0307%	0.0644%	0.0987%	0.0982%	0.0188%
Females	0.0124%	-	-	-	0.0010%	0.0047%	0.0205%	0.0408%	0.0657%	0.0698%	0.0124%
White ^c	0.0136%	-	-	0.0002%	0.0015%	0.0058%	0.0230%	0.0489%	0.0768%	0.0739%	0.0143%
Males	0.0154%	-	-	-	0.0020%	0.0079%	0.0289%	0.0616%	0.0967%	0.0926%	0.0180%
Females	0.0119%	-	-	-	0.0010%	0.0038%	0.0173%	0.0373%	0.0617%	0.0642%	0.0112%
Black ^c	0.0203%	-	-	-	0.0014%	0.0144%	0.0564%	0.1013%	0.1465%	0.1510%	0.0300%
Males	0.0206%	-	-	-	-	0.0169%	0.0626%	0.1272%	0.1614%	0.1653%	0.0342%
Females	0.0200%	-	-	-	0.0018%	0.0122%	0.0513%	0.0807%	0.1367%	0.1443%	0.0268%
Asian/Pacific Islander ^c	0.0081%	-	-	-	0.0009%	0.0043%	0.0161%	0.0270%	0.0510%	0.0634%	0.0097%
Males	0.0091%	-	-	-	0.0016%	0.0045%	0.0177%	0.0302%	0.0668%	0.0840%	0.0120%
Females	0.0073%	-	-	-	-	0.0041%	0.0147%	0.0244%	0.0391%	0.0491%	0.0080%
Hispanic ^d	0.0065%	-	-	-	0.0020%	0.0068%	0.0207%	0.0453%	0.0759%	0.0713%	0.0139%
Males	0.0072%	-	-	-	0.0031%	0.0080%	0.0263%	0.0514%	0.0940%	0.1191%	0.0180%
Females	0.0058%	-	-	-	0.0008%	0.0056%	0.0156%	0.0403%	0.0630%	0.0454%	0.0110%

- ^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
- ^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.
- ^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
- ^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.
- ^{g h i} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26. (i) Age-specific completeness index was approximated using empirical data from historical Connecticut tumor registry.
- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
- + Not available.

Table XVIII-16
NON-HODGKIN LYMPHOMA (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2001
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
Race/Sex									
All Races ^b	149,775	80,690	45,179	25,601	13,353	241,752	316,297	15,268	331,565
Males	78,674	41,667	23,243	12,930	6,440	126,162	163,797	8,107	171,904
Females	71,101	39,023	21,936	12,671	6,913	115,590	152,500	7,161	159,661
White ^b	134,163	72,482	41,063	23,565	12,328	216,877	285,203	15,295	300,498
Males	70,304	37,249	21,234	11,885	5,994	112,824	147,500	8,528	156,028
Females	63,859	35,233	19,829	11,680	6,334	104,053	137,703	6,767	144,470
Black ^b	9,940	5,151	2,788	1,269	775	15,861	19,986	1,054	21,040
Males	5,389	2,787	1,327	653	328	8,570	10,493	488	10,981
Females	4,551	2,364	1,461	616	447	7,291	9,493	566	10,059
Asian/Pacific Islander ^c	3,138	1,670	+	+	+	5,009	+	+	+
Males	1,691	824	+	+	+	2,616	+	+	+
Females	1,447	846	+	+	+	2,393	+	+	+
Hispanic ^d	7,840	3,742	+	+	+	12,174	+	+	+
Males	4,256	1,917	+	+	+	6,463	+	+	+
Females	3,584	1,825	+	+	+	5,711	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race/Sex											
All Races ^c	0.0724%	0.0022%	0.0081%	0.0129%	0.0289%	0.0609%	0.1237%	0.2243%	0.3315%	0.3448%	0.0782%
Males	0.0769%	0.0032%	0.0112%	0.0156%	0.0340%	0.0723%	0.1444%	0.2548%	0.3720%	0.3883%	0.0900%
Females	0.0681%	0.0011%	0.0048%	0.0100%	0.0238%	0.0497%	0.1040%	0.1974%	0.3012%	0.3217%	0.0679%
White ^c	0.0804%	0.0022%	0.0085%	0.0135%	0.0301%	0.0640%	0.1313%	0.2449%	0.3555%	0.3604%	0.0832%
Males	0.0846%	0.0030%	0.0117%	0.0162%	0.0354%	0.0756%	0.1527%	0.2765%	0.3971%	0.4063%	0.0954%
Females	0.0762%	0.0013%	0.0051%	0.0105%	0.0245%	0.0523%	0.1105%	0.2163%	0.3240%	0.3364%	0.0724%
Black ^c	0.0406%	0.0017%	0.0057%	0.0120%	0.0278%	0.0582%	0.1032%	0.1352%	0.1572%	0.1583%	0.0519%
Males	0.0456%	0.0028%	0.0080%	0.0159%	0.0326%	0.0730%	0.1275%	0.1578%	0.1704%	0.1755%	0.0615%
Females	0.0359%	-	0.0033%	0.0085%	0.0235%	0.0454%	0.0828%	0.1174%	0.1485%	0.1503%	0.0439%
Asian/Pacific Islander ^c	0.0457%	0.0022%	0.0080%	0.0098%	0.0198%	0.0371%	0.0722%	0.1374%	0.2319%	0.2674%	0.0522%
Males	0.0490%	0.0036%	0.0107%	0.0119%	0.0227%	0.0418%	0.0811%	0.1528%	0.2683%	0.3059%	0.0598%
Females	0.0426%	-	0.0051%	0.0076%	0.0171%	0.0329%	0.0643%	0.1248%	0.2046%	0.2407%	0.0459%
Hispanic ^d	0.0312%	0.0020%	0.0067%	0.0085%	0.0204%	0.0427%	0.0859%	0.1639%	0.2558%	0.2488%	0.0572%
Males	0.0321%	0.0030%	0.0091%	0.0097%	0.0241%	0.0460%	0.0996%	0.1792%	0.2905%	0.2220%	0.0626%
Females	0.0302%	0.0010%	0.0041%	0.0071%	0.0161%	0.0393%	0.0733%	0.1513%	0.2311%	0.2634%	0.0521%

^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{g h} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XIX-12
ORAL CAVITY AND PHARYNX CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2001
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
<u>Race/Sex</u>									
All Races ^b	82,674	51,214	33,360	22,078	14,999	142,122	206,494	24,203	230,697
Males	55,409	32,984	20,511	13,562	9,130	93,734	133,057	14,767	147,824
Females	27,265	18,230	12,849	8,516	5,869	48,388	73,437	9,436	82,873
White ^b	71,166	45,290	29,753	19,920	13,467	123,767	181,536	20,660	202,196
Males	47,891	29,439	18,484	12,453	8,412	82,143	118,028	12,723	130,751
Females	23,275	15,851	11,269	7,467	5,055	41,624	63,508	7,937	71,445
Black ^b	7,237	3,752	2,307	1,341	876	11,576	15,673	1,120	16,793
Males	4,861	2,213	1,305	673	351	7,410	9,470	498	9,968
Females	2,376	1,539	1,002	668	525	4,166	6,203	622	6,825
Asian/Pacific Islander ^c	2,682	1,401	+	+	+	4,287	+	+	+
Males	1,712	869	+	+	+	2,709	+	+	+
Females	970	532	+	+	+	1,578	+	+	+
Hispanic ^d	3,244	1,861	+	+	+	5,383	+	+	+
Males	2,116	1,128	+	+	+	3,398	+	+	+
Females	1,128	733	+	+	+	1,985	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0426%	0.0003%	0.0016%	0.0048%	0.0127%	0.0374%	0.0913%	0.1537%	0.1822%	0.1590%	0.0460%
Males	0.0568%	0.0003%	0.0014%	0.0048%	0.0147%	0.0520%	0.1357%	0.2290%	0.2648%	0.2435%	0.0668%
Females	0.0287%	0.0004%	0.0018%	0.0047%	0.0105%	0.0231%	0.0491%	0.0871%	0.1204%	0.1141%	0.0288%
White ^c	0.0451%	0.0003%	0.0017%	0.0044%	0.0115%	0.0348%	0.0917%	0.1625%	0.1905%	0.1652%	0.0468%
Males	0.0601%	0.0002%	0.0013%	0.0043%	0.0135%	0.0490%	0.1364%	0.2415%	0.2757%	0.2555%	0.0680%
Females	0.0302%	0.0004%	0.0021%	0.0045%	0.0094%	0.0203%	0.0481%	0.0908%	0.1259%	0.1181%	0.0288%
Black ^c	0.0297%	0.0007%	0.0015%	0.0047%	0.0115%	0.0411%	0.0908%	0.1300%	0.1332%	0.0681%	0.0388%
Males	0.0401%	-	0.0021%	0.0054%	0.0136%	0.0549%	0.1415%	0.1967%	0.2108%	0.0849%	0.0573%
Females	0.0201%	-	-	0.0041%	0.0096%	0.0291%	0.0482%	0.0771%	0.0820%	0.0604%	0.0247%
Asian/Pacific Islander ^c	0.0376%	-	0.0010%	0.0067%	0.0186%	0.0487%	0.0842%	0.1066%	0.1395%	0.1396%	0.0414%
Males	0.0492%	-	-	0.0076%	0.0207%	0.0669%	0.1214%	0.1554%	0.1954%	0.2008%	0.0578%
Females	0.0269%	-	-	0.0059%	0.0165%	0.0325%	0.0515%	0.0668%	0.0975%	0.0970%	0.0280%
Hispanic ^d	0.0136%	-	0.0010%	0.0027%	0.0066%	0.0179%	0.0448%	0.0920%	0.1107%	0.1057%	0.0262%
Males	0.0167%	-	-	0.0025%	0.0071%	0.0234%	0.0640%	0.1393%	0.1630%	0.1429%	0.0371%
Females	0.0103%	-	0.0015%	0.0028%	0.0061%	0.0123%	0.0273%	0.0531%	0.0735%	0.0855%	0.0177%

^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{g h} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XX-10
OVARY CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2001
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
<u>Race</u>									
All Races ^b	53,367	28,304	21,790	18,522	13,537	86,619	137,976	29,026	167,002
White ^b	47,519	25,140	20,105	16,667	12,202	77,384	123,803	25,354	149,157
Black ^b	3,578	1,936	1,105	1,254	974	5,647	9,037	2,465	11,502
Asian/Pacific Islander ^c	1,586	858	+	+	+	2,557	+	+	+
Hispanic ^d	3,061	1,647	+	+	+	4,971	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	All Ages	Age Specific (Crude)									Age-Adjusted ^f	
		0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages	
<u>Race</u>												
All Races ^c	0.0521%	0.0004%	0.0046%	0.0118%	0.0203%	0.0544%	0.1142%	0.1653%	0.1744%	0.1153%	0.0527%	
White ^c	0.0574%	0.0004%	0.0043%	0.0118%	0.0206%	0.0575%	0.1226%	0.1809%	0.1864%	0.1210%	0.0562%	
Black ^c	0.0274%	-	0.0046%	0.0098%	0.0159%	0.0325%	0.0618%	0.0929%	0.1248%	0.0837%	0.0332%	
Asian/Pacific Islander ^c	0.0425%	-	0.0068%	0.0133%	0.0223%	0.0536%	0.0984%	0.1195%	0.1181%	0.0728%	0.0435%	
Hispanic ^d	0.0263%	0.0006%	0.0050%	0.0111%	0.0163%	0.0412%	0.0821%	0.1268%	0.1403%	0.0885%	0.0410%	

^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{g h} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXI-12
PANCREAS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2001
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
<u>Race/Sex</u>									
All Races ^b	18,095	3,104	1,515	690	460	21,576	23,967	549	24,516
Males	8,680	1,346	575	287	221	10,177	11,163	216	11,379
Females	9,415	1,758	940	403	239	11,399	12,804	333	13,137
White ^b	15,117	2,577	1,124	536	381	18,030	19,814	441	20,255
Males	7,306	1,109	438	237	197	8,554	9,323	186	9,509
Females	7,811	1,468	686	299	184	9,476	10,491	255	10,746
Black ^b	2,191	344	331	107	78	2,569	3,065	253	3,318
Males	1,012	137	122	39	23	1,162	1,343	104	1,447
Females	1,179	207	209	68	55	1,407	1,722	149	1,871
Asian/Pacific Islander ^c	547	82	+	+	+	643	+	+	+
Males	247	40	+	+	+	291	+	+	+
Females	300	42	+	+	+	352	+	+	+
Hispanic ^d	877	261	+	+	+	1,155	+	+	+
Males	336	117	+	+	+	464	+	+	+
Females	541	144	+	+	+	691	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0068%	-	0.0002%	0.0003%	0.0013%	0.0038%	0.0113%	0.0264%	0.0372%	0.0361%	0.0075%
Males	0.0065%	-	-	-	0.0013%	0.0039%	0.0127%	0.0292%	0.0379%	0.0390%	0.0079%
Females	0.0071%	-	0.0002%	0.0005%	0.0014%	0.0038%	0.0100%	0.0240%	0.0367%	0.0345%	0.0071%
White ^c	0.0070%	-	0.0001%	0.0004%	0.0013%	0.0037%	0.0111%	0.0267%	0.0359%	0.0341%	0.0073%
Males	0.0067%	-	-	-	0.0012%	0.0041%	0.0123%	0.0297%	0.0360%	0.0382%	0.0078%
Females	0.0072%	-	-	0.0006%	0.0014%	0.0034%	0.0099%	0.0239%	0.0358%	0.0320%	0.0069%
Black ^c	0.0062%	-	-	-	0.0013%	0.0049%	0.0134%	0.0305%	0.0452%	0.0470%	0.0090%
Males	0.0058%	-	-	-	0.0018%	0.0044%	0.0168%	0.0359%	0.0467%	0.0254%	0.0092%
Females	0.0064%	-	-	-	-	0.0053%	0.0105%	0.0261%	0.0443%	0.0569%	0.0086%
Asian/Pacific Islander ^c	0.0063%	-	-	-	0.0013%	0.0031%	0.0109%	0.0210%	0.0401%	0.0477%	0.0075%
Males	0.0059%	-	-	-	-	0.0022%	0.0123%	0.0201%	0.0433%	0.0563%	0.0078%
Females	0.0066%	-	-	-	-	0.0040%	0.0096%	0.0218%	0.0377%	0.0418%	0.0073%
Hispanic ^d	0.0030%	-	-	0.0005%	0.0010%	0.0030%	0.0075%	0.0164%	0.0410%	0.0426%	0.0066%
Males	0.0023%	-	-	-	0.0010%	0.0017%	0.0072%	0.0158%	0.0376%	0.0476%	0.0062%
Females	0.0038%	-	-	0.0010%	0.0010%	0.0043%	0.0078%	0.0168%	0.0435%	0.0399%	0.0070%

^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.
^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.
^{g h} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26.
 - Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
 + Not available.

Table XXII-10
PROSTATE CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2001
 By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
<u>Race</u>									
All Races ^b	854,021	616,832	182,507	49,579	16,895	1,531,453	1,721,366	6,481	1,727,847
White ^b	714,334	533,805	165,767	44,467	15,196	1,303,231	1,474,940	6,764	1,481,704
Black ^b	109,319	67,353	14,006	4,447	1,550	181,277	196,813	561	197,374
Asian/Pacific Islander ^c	12,286	6,931	+	+	+	19,761	+	+	+
Hispanic ^d	35,737	20,470	+	+	+	57,990	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
 By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races ^c	0.9028%	0.0002%	0.0002%	0.0002%	0.0008%	0.0477%	0.7545%	3.9101%	8.7396%	9.0628%	1.1958%
White ^c	0.9465%	-	-	0.0003%	0.0008%	0.0419%	0.7351%	3.9010%	8.7085%	8.7840%	1.1798%
Black ^c	0.9585%	-	-	-	0.0018%	0.1226%	1.4371%	6.5504%	12.9982%	12.0044%	1.8322%
Asian/Pacific Islander ^c	0.4350%	-	-	-	-	0.0134%	0.2221%	1.4682%	4.7206%	6.5065%	0.6327%
Hispanic ^d	0.2823%	-	-	-	-	0.0201%	0.4200%	2.5868%	7.0666%	7.4846%	0.9163%

^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{g h} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXIII-12
STOMACH CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2001
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
Race/Sex									
All Races ^b	29,431	11,709	7,320	4,576	2,574	42,701	55,827	3,372	59,199
Males	17,724	6,478	3,873	2,448	1,451	24,999	32,088	1,953	34,041
Females	11,707	5,231	3,447	2,128	1,123	17,702	23,739	1,419	25,158
White ^b	22,534	8,958	5,560	3,529	2,134	32,585	42,890	2,734	45,624
Males	13,816	4,975	3,003	1,923	1,211	19,361	25,022	1,510	26,532
Females	8,718	3,983	2,557	1,606	923	13,224	17,868	1,224	19,092
Black ^b	4,254	1,639	1,027	723	235	6,159	7,907	445	8,352
Males	2,346	928	499	378	144	3,409	4,305	262	4,567
Females	1,908	711	528	345	91	2,750	3,602	183	3,785
Asian/Pacific Islander ^c	2,291	974	+	+	+	3,410	+	+	+
Males	1,350	547	+	+	+	1,979	+	+	+
Females	941	427	+	+	+	1,431	+	+	+
Hispanic ^d	3,056	1,146	+	+	+	4,378	+	+	+
Males	1,694	587	+	+	+	2,367	+	+	+
Females	1,362	559	+	+	+	2,011	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race/Sex											
All Races ^c	0.0159%	-	0.0001%	0.0005%	0.0028%	0.0084%	0.0211%	0.0524%	0.0973%	0.1170%	0.0176%
Males	0.0188%	-	-	0.0004%	0.0029%	0.0098%	0.0281%	0.0725%	0.1348%	0.1601%	0.0238%
Females	0.0131%	-	-	0.0005%	0.0026%	0.0071%	0.0144%	0.0345%	0.0692%	0.0942%	0.0129%
White ^c	0.0134%	-	-	0.0004%	0.0023%	0.0068%	0.0170%	0.0436%	0.0741%	0.0934%	0.0140%
Males	0.0159%	-	-	0.0003%	0.0025%	0.0077%	0.0227%	0.0614%	0.1038%	0.1283%	0.0190%
Females	0.0110%	-	-	0.0004%	0.0022%	0.0058%	0.0114%	0.0276%	0.0516%	0.0752%	0.0101%
Black ^c	0.0156%	-	-	0.0012%	0.0034%	0.0146%	0.0281%	0.0629%	0.1333%	0.1413%	0.0234%
Males	0.0181%	-	-	-	0.0036%	0.0182%	0.0389%	0.0886%	0.1898%	0.1461%	0.0308%
Females	0.0133%	-	-	-	0.0033%	0.0115%	0.0191%	0.0426%	0.0961%	0.1391%	0.0179%
Asian/Pacific Islander ^c	0.0335%	-	-	-	0.0046%	0.0142%	0.0437%	0.1030%	0.2449%	0.3319%	0.0410%
Males	0.0401%	-	-	-	0.0045%	0.0170%	0.0587%	0.1403%	0.3354%	0.4245%	0.0543%
Females	0.0272%	-	-	-	0.0047%	0.0117%	0.0306%	0.0726%	0.1769%	0.2675%	0.0308%
Hispanic ^d	0.0112%	-	-	0.0004%	0.0047%	0.0121%	0.0276%	0.0667%	0.1259%	0.2042%	0.0250%
Males	0.0117%	-	-	-	0.0044%	0.0125%	0.0307%	0.0886%	0.1674%	0.2675%	0.0316%
Females	0.0107%	-	-	-	0.0050%	0.0116%	0.0249%	0.0486%	0.0963%	0.1699%	0.0204%

- ^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
- ^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.
- ^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
- ^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.
- ^{g h} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26.
- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
- + Not available.

Table XXIV-10
TESTIS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2001
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
<u>Race</u>									
All Races ^b	36,654	31,755	29,060	21,694	15,575	74,666	136,928	20,421	157,349
White ^b	34,466	30,101	28,019	20,927	15,006	70,615	130,559	19,622	150,181
Black ^b	1,021	622	417	367	256	1,723	2,747	201	2,948
Asian/Pacific Islander ^c	562	494	+	+	+	1,122	+	+	+
Hispanic ^d	3,236	2,389	+	+	+	5,990	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races ^c	0.0467%	0.0021%	0.0040%	0.0494%	0.1083%	0.0930%	0.0438%	0.0175%	0.0096%	0.0062%	0.0450%
White ^c	0.0560%	0.0021%	0.0045%	0.0602%	0.1307%	0.1099%	0.0513%	0.0202%	0.0113%	0.0071%	0.0537%
Black ^c	0.0099%	-	0.0032%	0.0090%	0.0202%	0.0238%	0.0108%	0.0060%	-	-	0.0101%
Asian/Pacific Islander ^c	0.0177%	0.0033%	0.0019%	0.0166%	0.0376%	0.0373%	0.0159%	0.0054%	-	-	0.0167%
Hispanic ^d	0.0306%	0.0024%	0.0037%	0.0420%	0.0708%	0.0527%	0.0241%	0.0105%	-	-	0.0294%

^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{g h} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXV-12
 THYROID CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2001
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
Race/Sex									
All Races ^b	82,780	60,518	45,668	35,741	31,427	153,434	261,607	48,887	310,494
Males	19,086	14,644	9,924	7,887	6,965	35,894	59,817	11,488	71,305
Females	63,694	45,874	35,744	27,854	24,462	117,540	201,790	37,399	239,189
White ^b	72,726	53,622	40,943	32,272	28,291	135,451	232,837	44,245	277,082
Males	17,284	13,350	9,116	7,116	6,449	32,602	54,530	9,807	64,337
Females	55,442	40,272	31,827	25,156	21,842	102,849	178,307	34,438	212,745
Black ^b	4,901	3,354	2,129	1,661	1,544	8,763	13,847	1,999	15,846
Males	758	680	378	382	164	1,565	2,399	204	2,603
Females	4,143	2,674	1,751	1,279	1,380	7,198	11,448	1,795	13,243
Asian/Pacific Islander ^c	3,505	2,518	+	+	+	6,440	+	+	+
Males	684	439	+	+	+	1,196	+	+	+
Females	2,821	2,079	+	+	+	5,244	+	+	+
Hispanic ^d	6,625	4,587	+	+	+	11,974	+	+	+
Males	1,202	840	+	+	+	2,198	+	+	+
Females	5,423	3,747	+	+	+	9,776	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
 By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race/Sex											
All Races ^c	0.0487%	0.0001%	0.0036%	0.0266%	0.0632%	0.0840%	0.0934%	0.0877%	0.0770%	0.0521%	0.0496%
Males	0.0226%	-	0.0012%	0.0079%	0.0219%	0.0355%	0.0508%	0.0583%	0.0533%	0.0378%	0.0243%
Females	0.0742%	-	0.0062%	0.0461%	0.1056%	0.1318%	0.1338%	0.1137%	0.0949%	0.0596%	0.0740%
White ^c	0.0520%	0.0001%	0.0039%	0.0295%	0.0690%	0.0899%	0.0959%	0.0892%	0.0757%	0.0493%	0.0520%
Males	0.0248%	-	0.0012%	0.0089%	0.0241%	0.0387%	0.0541%	0.0615%	0.0548%	0.0359%	0.0259%
Females	0.0789%	-	0.0067%	0.0517%	0.1168%	0.1418%	0.1367%	0.1144%	0.0916%	0.0562%	0.0782%
Black ^c	0.0219%	-	0.0015%	0.0092%	0.0265%	0.0377%	0.0574%	0.0533%	0.0538%	0.0320%	0.0258%
Males	0.0088%	-	-	0.0023%	0.0085%	0.0135%	0.0275%	0.0259%	0.0397%	0.0212%	0.0118%
Females	0.0337%	-	0.0022%	0.0155%	0.0425%	0.0588%	0.0825%	0.0749%	0.0631%	0.0370%	0.0375%
Asian/Pacific Islander ^c	0.0524%	-	0.0043%	0.0240%	0.0591%	0.0860%	0.1037%	0.1005%	0.0991%	0.0935%	0.0537%
Males	0.0206%	-	-	0.0055%	0.0194%	0.0316%	0.0454%	0.0567%	0.0488%	0.0617%	0.0228%
Females	0.0821%	-	0.0078%	0.0418%	0.0964%	0.1346%	0.1548%	0.1362%	0.1368%	0.1156%	0.0809%
Hispanic ^d	0.0310%	-	0.0031%	0.0190%	0.0452%	0.0715%	0.0819%	0.0748%	0.0864%	0.0621%	0.0428%
Males	0.0108%	-	0.0007%	0.0059%	0.0116%	0.0218%	0.0378%	0.0404%	0.0549%	0.0416%	0.0178%
Females	0.0523%	-	0.0057%	0.0346%	0.0831%	0.1215%	0.1223%	0.1030%	0.1088%	0.0732%	0.0670%

^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{g h} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

Table XXVI-12
URINARY BLADDER CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2001
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
Race/Sex									
All Races ^b	182,170	120,521	78,521	47,739	28,041	320,828	460,530	29,928	490,458
Males	136,806	89,589	58,708	34,535	19,731	239,991	341,794	19,677	361,471
Females	45,364	30,932	19,813	13,204	8,310	80,837	118,736	10,251	128,987
White ^b	170,466	113,783	74,937	45,510	26,687	301,386	434,668	23,918	458,586
Males	128,819	84,688	56,105	32,947	18,815	226,458	323,642	15,447	339,089
Females	41,647	29,095	18,832	12,563	7,872	74,928	111,026	8,471	119,497
Black ^b	7,038	4,273	2,482	1,390	794	11,991	16,070	493	16,563
Males	4,427	3,016	1,789	962	583	7,864	10,846	393	11,239
Females	2,611	1,257	693	428	211	4,127	5,224	100	5,324
Asian/Pacific Islander ^c	2,089	1,291	+	+	+	3,554	+	+	+
Males	1,538	987	+	+	+	2,655	+	+	+
Females	551	304	+	+	+	899	+	+	+
Hispanic ^d	4,733	2,916	+	+	+	8,125	+	+	+
Males	3,546	2,216	+	+	+	6,116	+	+	+
Females	1,187	700	+	+	+	2,009	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^f
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race/Sex											
All Races ^c	0.0897%	0.0001%	0.0002%	0.0016%	0.0062%	0.0277%	0.1028%	0.3124%	0.6191%	0.7157%	0.1001%
Males	0.1367%	-	0.0003%	0.0021%	0.0087%	0.0423%	0.1640%	0.5220%	1.1063%	1.4331%	0.1781%
Females	0.0438%	-	-	0.0011%	0.0036%	0.0132%	0.0447%	0.1269%	0.3351%	0.0431%	
White ^c	0.1067%	0.0001%	0.0002%	0.0021%	0.0070%	0.0316%	0.1174%	0.3598%	0.6921%	0.7707%	0.1121%
Males	0.1619%	-	0.0004%	0.0026%	0.0096%	0.0476%	0.1848%	0.5949%	1.2360%	1.5710%	0.1989%
Females	0.0519%	-	-	0.0015%	0.0043%	0.0154%	0.0517%	0.1465%	0.2800%	0.3533%	0.0479%
Black ^c	0.0300%	-	-	-	0.0034%	0.0143%	0.0551%	0.1399%	0.2633%	0.3552%	0.0468%
Males	0.0428%	-	-	-	0.0058%	0.0228%	0.0911%	0.2340%	0.4375%	0.6136%	0.0785%
Females	0.0184%	-	-	-	-	0.0068%	0.0249%	0.0655%	0.1484%	0.2365%	0.0257%
Asian/Pacific Islander ^c	0.0355%	-	-	-	0.0027%	0.0111%	0.0356%	0.1101%	0.2894%	0.3913%	0.0445%
Males	0.0548%	-	-	-	0.0042%	0.0192%	0.0588%	0.1867%	0.5001%	0.6741%	0.0763%
Females	0.0176%	-	-	-	0.0013%	0.0039%	0.0152%	0.0474%	0.1312%	0.1947%	0.0203%
Hispanic ^d	0.0201%	-	-	0.0007%	0.0033%	0.0110%	0.0444%	0.1394%	0.2869%	0.4102%	0.0485%
Males	0.0292%	-	-	0.0006%	0.0042%	0.0143%	0.0746%	0.2463%	0.5322%	0.7658%	0.0870%
Females	0.0104%	-	-	0.0009%	0.0024%	0.0077%	0.0167%	0.0517%	0.1122%	0.2177%	0.0213%

^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.
^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).
^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.
^{g h} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26.
 - Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.
 + Not available.

Table XXVII-12
 CHILDHOOD CANCER (Invasive, <20 Years old at diagnosis)

Estimated United States Cancer Prevalence Counts^a on January 1, 2001
 By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^f	Complete ^g
<u>Race/Sex</u>									
All Races ^b	54,418	45,273	38,363	32,455	28,901	107,593	204,131	+	+
Males	28,817	23,755	20,104	16,244	13,509	57,026	104,636	+	+
Females	25,601	21,518	18,259	16,211	15,392	50,567	99,495	+	+
White ^b	45,606	37,927	32,874	28,432	25,519	90,272	174,485	+	+
Males	24,091	20,001	17,353	14,394	12,004	47,985	89,833	+	+
Females	21,515	17,926	15,521	14,038	13,515	42,287	84,652	+	+
Black ^b	5,708	4,948	3,684	2,804	2,392	11,380	19,946	+	+
Males	3,026	2,485	1,762	1,294	1,046	5,848	9,760	+	+
Females	2,682	2,463	1,922	1,510	1,346	5,532	10,186	+	+
Asian/Pacific Islander ^c	1,811	1,410	+	+	+	3,497	+	+	+
Males	988	689	+	+	+	1,828	+	+	+
Females	823	721	+	+	+	1,669	+	+	+
Hispanic ^d	8,073	5,728	+	+	+	14,784	+	+	+
Males	4,270	3,137	+	+	+	8,000	+	+	+
Females	3,803	2,591	+	+	+	6,784	+	+	+

^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^{f g} (f) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <26 and >=26.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.