
## Percentages of HIV/AIDS Cases and Population among Adolescents 13 to 19 Years of Age, by Race/Ethnicity Diagnosed in 2007—34 States





Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 34 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays.

\*Includes 10 persons of unknown race or multiple races.


†Includes Asian and Pacific Islander legacy cases.

§Hispanics/Latinos can be of any race.





### Percentages of HIV/AIDS Cases and Population among Young Adults 20 to 24 Years of Age, by Race/Ethnicity Diagnosed in 2007—34 States





Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 34 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays.

\*Includes 41 persons of unknown race or multiple races.


†Includes Asian and Pacific Islander legacy cases.

§Hispanics/Latinos can be of any race.





#### Percentages of HIV/AIDS Cases among Adults and Adolescents, by Sex and Age Group Diagnosed in 2007—34 States





Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis.

Data from 34 states with confidential name-based HIV infection reporting since at least 2003.

Data based on person's age at diagnosis.

Data have been adjusted for reporting delays.






#### Estimated Numbers of HIV/AIDS Cases among Male Adolescents and Young Adults, by Transmission Category 2004-2007-34 States

	13–19 years		20-24 years	
Transmission category	N	%	N	%
Male-to-male sexual contact	3,171	87	10,226	83
Injection drug use	122	3	558	5
Male-to-male sexual contact and injection drug use	130	4	551	4
High-risk heterosexual contact*	221	6	910	7
Other/not identified <sup>†</sup>	7	<1	19	<1
Total	3,651		12,264	



Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 34 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays and missing risk-factor information.







## Estimated Numbers of HIV/AIDS Cases among Female Adolescents and Young Adults, by Transmission Category 2004–2007—34 States

	13–19	years	20-24 years	
Transmission category	N	%	N	%
Injection drug use	219	11	555	13
High-risk heterosexual contact*	1,694	88	3,846	87
Other/not identified <sup>†</sup>	7	<1	16	<1
Total	1,920		4,417	




Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis. Data from 34 states with confidential name-based HIV infection reporting since at least 2003. Data have been adjusted for reporting delays and missing risk-factor information.

\*Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

†Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



#### Estimated Numbers of HIV/AIDS Cases among Adolescents 13 to 19 Years of Age, 2007—34 States





Note. Data include persons with a diagnosis of HIV infection regardless of their AIDS status at diagnosis.

Data from 34 states with confidential name-based HIV infection reporting since at least 2003.


Data have been adjusted for reporting delays.


Data based on person's age at diagnosis.






### Estimated Numbers of Adolescents 13 to 19 Years of Age Living with HIV/AIDS, 2007—34 States









### Reported AIDS Cases among Adolescents 13 to 19 Years of Age, by Sex,1985–2007—United States and Dependent Areas N=7,200



### Reported AIDS Cases among Young Adults 20 to 24 Years of Age, by Sex,1985–2007—United States and Dependent Areas N=38,483



# Percentages of AIDS Cases and Population among Adolescents 13 to 19 Years of Age, by Race/Ethnicity Reported in 2007—50 States and DC




Population, 50 States and DC N = 29,775,943**15%** 18% 60% Hispanic/Latino 1 Native Hawaiian/Other Pacific Islander




Note. Data based on person's age at AIDS diagnosis. \*Includes 6 persons of unknown race or multiple races. †Includes Asian and Pacific Islander legacy cases. §Hispanics/Latinos can be of any race.



# Percentages of AIDS Cases and Population among Young Adults 20 to 24 Years of Age, by Race/Ethnicity Reported in 2007—50 States and DC




Note. Data based on person's age at AIDS diagnosis.
\*Includes17 persons of unknown race or multiple races.
†Includes Asian and Pacific Islander legacy cases.
§Hispanics/Latinos can be of any race.







# Percentages of AIDS Cases among Adults and Adolescents, by Sex and Age Group, Reported in 2007—United States and Dependent Areas







# Estimated Numbers of AIDS Cases among Male Adolescents and Young Adults, by Transmission Category 2003–2007—United States and Dependent Areas

	13–19 years		20–24 years	
Transmission category	N	%	N	%
Male-to-male sexual contact	860	64	4,711	77
Injection drug use	56	4	374	6
Male-to-male sexual contact and injection drug use	37	3	365	6
High-risk heterosexual contact*	100	7	583	10
Other/not identified <sup>†</sup>	291	22	67	1
Total	1,343		6,101	



Note. Data have been adjusted for reporting delays and missing risk-factor information. \*Heterosexual contact with a person known to have, or to be at high risk for, HIV infection. †Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.





# Estimated Numbers of AIDS Cases among Female Adolescents and Young Adults, by Transmission Category 2003–2007—United States and Dependent Areas

Transmission category	13–19 years		20-24 years		
	N	%	N	%	
Injection drug use	84	9	359	15	
High-risk heterosexual contact*	527	54	2,033	82	
Other/not identified <sup>†</sup>	364	37	76	3	
Total	975		2,468		




Note. Data have been adjusted for reporting delays and missing risk-factor information.

\*Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

†Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.



#### Reported AIDS Cases among Adolescents 13 to 19 Years of Age, 2007—United States and Dependent Areas





#### Adolescents 13 to 19 Years of Age Reported to be Living with AIDS, 2007—United States and Dependent Areas

