

Revised Oil-Spill Risk Analysis: Beaufort Sea Outer Continental Shelf Lease Sale 170



Revised Oil-Spill Risk Analysis: Beaufort Sea Outer Continental Shelf Lease Sale 170

By: Cheryl M. Anderson
Walter R. Johnson
Charles F. Marshall
Eileen M. Lear (Editor)

Contents

	Page
Introduction	1
Summary of the Proposed Action	2
Summary of the Alternatives	2
Framework of the Analysis	2
Environmental Resource Areas	2
Estimated Volume of Oil Resources	4
Oil-Spill Risk Analysis	4
Probability of Oil Spills Occurring	5
Oil-Spill Trajectory Simulations	7
Conditional Probabilities	10
Combined Analysis of Oil-Spill Occurrence and Oil-Spill Trajectory Simulations	11
Discussion	12
References Cited	16
Appendix A: Environmental Resource Areas	41
Appendix B: Seasonal Conditional Probabilities of Contact to Environmental Resource Areas	51
Appendix C: Seasonal Conditional Probabilities of Contact to Land Segments	61

Figures

1. Study area and proposed lease sale area for proposed Beaufort Sea OCS Lease Sale 170	18
2. Enlarged representation of launch areas (L1-L8) for proposed Beaufort Sea OCS Lease Sale 170	19
3. Enlarged representation of pipeline route segments (P1-P7) for proposed Beaufort Sea OCS Lease Sale 170	20
4. Enlarged representation of Kaktovik Deferral Alternative for proposed Beaufort Sea OCS Lease Sale 170	21
5. Enlarged representation of Area Offshore of the ANWR for proposed Beaufort Sea OCS Lease Sale 170	22
6. Enlarged representation of Cross Island Area for proposed Beaufort Sea OCS Lease Sale 170	23
7. Study area coastline divided into 61 land segments for proposed Beaufort Sea OCS Lease Sale 170	24

Tables

1. Oil-spill occurrence probability estimates for spills greater than or equal to 1,000 barrels resulting over the assumed production life of the proposed action, Beaufort Sea OCS Lease Sale 170	25
2.-9. Annual conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location will contact:	
2. A certain environmental resource area within 3 days, Beaufort Sea OCS Lease Sale 170	26
3. A certain environmental resource area within 10 days, Beaufort Sea OCS Lease Sale 170	27
4. A certain environmental resource area within 30 days, Beaufort Sea OCS Lease Sale 170	28
5. A certain environmental resource area within 180 days, Beaufort Sea OCS Lease Sale 170	29

Tables

6.	A certain land segment within 3 days, Beaufort Sea OCS Lease Sale 170	30
7.	A certain land segment within 10 days, Beaufort Sea OCS Lease Sale 170	30
8.	A certain land segment within 30 days, Beaufort Sea OCS Lease Sale 170	31
9.	A certain land segment within 180 days, Beaufort Sea OCS Lease Sale 170	31
10-11.	Annual combined probabilities (expressed as percent chance) of one or more spills greater than or equal to 1,000 barrels, and the estimated number of spills (mean), occurring and contacting:	
10.	A certain environmental resource area over the assumed production life of Beaufort Sea OCS Lease Sale 170	32
11.	A certain land segment over the assumed production life of Beaufort Sea OCS Lease Sale 170	33
12.	A certain environmental resource area over the assumed production life of Beaufort Sea OCS Lease Sale 170, Kaktovik Deferral	34
13.	A certain land segment over the assumed production life of Beaufort Sea OCS Lease Sale 170, Kaktovik Deferral	35
14.	A certain environmental resource area over the assumed production life of Beaufort Sea OCS Lease Sale 170, Area Offshore of the ANWR	36
15.	A certain land segment over the assumed production life of Beaufort Sea OCS Lease Sale 170, Area Offshore of the ANWR	37
16.	A certain environmental resource area over the assumed production life of Beaufort Sea OCS Lease Sale 170, Cross Island Area	38
17.	A certain land segment over the assumed production life of Beaufort Sea OCS Lease Sale 170, Cross Island Area	39

Appendix A

Figures

A-1.	Locations of Ice/Sea Segments 1-17 for proposed Beaufort Sea OCS Lease Sale 170	43
A-2.	Locations of Peard Bay, Elson Lagoon, Simpson Lagoon, Gwydyr Bay, Jago Lagoon, and Beaufort Lagoon for proposed Beaufort Sea OCS Lease Sale 170	44
A-3.	Locations of Subsistence Resource Area A, Fall Feeding Area, Subsistence Resource Area C, and Summer Feeding Area 1 for proposed Beaufort Sea OCS Lease Sale 170	45
A-4.	Locations of Subsistence Resource Area B, Subsistence Resource Area D, and Summer Feeding Area 2 for proposed Beaufort Sea OCS Lease Sale 170	46
A-5.	Location of Northern Spring Lead System (SLS) Area for proposed Beaufort Sea OCS Lease Sale 170	47
A-6.	Location of Southern Spring Lead System (SLS) Area for proposed Beaufort Sea OCS Lease Sale 170	48
A-7.	Locations of Boundary Segments (B1-B38) for proposed Beaufort Sea OCS Lease Sale 170	49

Appendix B

Tables

B-1.-B-8. Seasonal conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location in the:

B-1.	Winter season will contact a certain environmental resource area within 3 days, Beaufort Sea OCS Lease Sale 170	53
B-2.	Winter season will contact a certain environmental resource area within 10 days, Beaufort Sea OCS Lease Sale 170	54
B-3.	Winter season will contact a certain environmental resource area within 30 days, Beaufort Sea OCS Lease Sale 170	55
B-4.	Winter season will contact a certain environmental resource area within 180 days, Beaufort Sea OCS Lease Sale 170	56
B-5.	Summer season will contact a certain environmental resource area within 3 days, Beaufort Sea OCS Lease Sale 170	57

B-6.	Summer season will contact a certain environmental resource area within 10 days, Beaufort Sea OCS Lease Sale 170	58
B-7.	Summer season will contact a certain environmental resource area within 30 days, Beaufort Sea OCS Lease Sale 170	59
B-8.	Summer season will contact a certain environmental resource area within 180 days, Beaufort Sea OCS Lease Sale 170	60

Appendix C

C-1.-C-8. Seasonal conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location in the:

C-1.	Winter season will contact a certain land segment within 3 days, Beaufort Sea OCS Lease Sale 170	63
C-2.	Winter season will contact a certain land segment within 10 days, Beaufort Sea OCS Lease Sale 170	63
C-3.	Winter season will contact a certain land segment within 30 days, Beaufort Sea OCS Lease Sale 170	64
C-4.	Winter season will contact a certain land segment within 180 days, Beaufort Sea OCS Lease Sale 170	64
C-5.	Summer season will contact a certain land segment within 3 days, Beaufort Sea OCS Lease Sale 170	65
C-6.	Summer season will contact a certain land segment within 10 days, Beaufort Sea OCS Lease Sale 170	65
C-7.	Summer season will contact a certain land segment within 30 days, Beaufort Sea OCS Lease Sale 170	66
C-8.	Summer season will contact a certain land segment within 180 days, Beaufort Sea OCS Lease Sale 170	66

Introduction

The Federal Government proposes to offer Outer Continental Shelf (OCS) lands in the Beaufort Sea for oil and gas leasing. The estimated amounts of oil resources likely to be leased, discovered, and produced as a result of Beaufort Sea OCS Lease Sale 170 (Proposed Action) are assumed to range between 0.35 billion barrels (Bbbl) and 0.67 Bbbl. This report examines the probabilities of hypothetical oil-spill occurrence and contact if leases are issued and commercial quantities of oil are found and produced from OCS Lease Sale 170.

Because oil spills may occur from activities associated with offshore oil production and transportation, the Minerals Management Service (MMS) formally assesses the risk of hypothetical oil-spills. When evaluating the significance of accidental oil spills, it is important to remember that the occurrence of such spills is fundamentally a matter of probability. No one can certify the volume of oil that would be produced, or the size or likelihood of a spill that would occur, during the assumed 21-year production life (2006-2026) of Beaufort Sea OCS Lease Sale 170. Neither can factors that affect spills such as the wind, ocean current, or ice conditions be known for certain. Although some of this uncertainty is due to incomplete and imperfect data, a considerable amount exists simply because future events are difficult to estimate. For example, a probabilistic event such as an oil-spill occurrence cannot be predicted— only an estimate of its likelihood or probability can be quantified. However, the possible effects of oil spills that could occur from oil and gas production must be considered in the leasing decision. To maintain perspective, each potential effect must be associated with a quantitative estimate of its probability of occurrence.

This report summarizes the results of the oil-spill risk analysis (OSRA) conducted for the proposed Beaufort Sea OCS Lease Sale 170. The objective of this analysis was to estimate the relative oil-spill risks associated with oil and gas production from the leasing alternatives proposed for the lease sale. To address proposed changes to the alternative actions that have been made in response to public concerns raised in the draft EIS review process, the MMS is issuing this report, which revises its original OSRA report for this lease sale (Anderson, et al., 1996). The MMS will analyze the oil-spill risks in the final environmental impact statement (EIS) prepared for the lease sale. A description of the OSRA model used in this analysis can be found in previous papers (Smith et al., 1982; LaBelle and Anderson, 1985).

The analysis for proposed OCS Lease Sale 170 was conducted in three parts, corresponding to different aspects of the overall situation.

- The first part addressed the probability of oil-spill occurrence.
- The second part addressed trajectories of oil spills from hypothetical spill sites to various environmental resource areas or land segments (for probability of contact).
- The third part used the results of the first two to estimate the combined risk of oil-spill occurrence and contact given that there is oil production as a result of the lease sale.

Summary of the Proposed Action

The Proposed Action (Alternative 1) offers for lease a portion of OCS lands in the Beaufort Sea. The OSRA study area extends from lat. 68° N. to 74° N. and from long. 134° W. to 176° W. (fig. 1). The proposed leasing area consists of 363 blocks (688,000 hectares) located in the Beaufort Sea (fig. 1). This area is located 4.8-40.3 km offshore and lies in waters that are 7.6-36.6 m deep. Trajectories were simulated from 344 hypothetical spill sites (spill sites) in the proposed sale area. These spill sites were grouped into 8 larger hypothetical launch areas (L1 through L8), which were used to represent areas for hypothetical platform spill occurrences (fig. 2). The scenario (that which is likely to be leased, discovered, and produced) for the Proposed Action assumes production and transportation ranging from 0.35 Bbbl to 0.67 Bbbl of oil.

The transportation scenario assumes that the crude oil produced in L1 through L8 will be transported by pipeline to shore with potential landfalls at Oliktok Point, Point McIntyre, and Bullen Point (fig. 3). These hypothetical offshore pipeline route segments (P1-P7) are used to represent spill risks from oil transportation. It is assumed that the oil is then transported by onshore pipeline to Pump Station #1 of the Trans-Alaska Pipeline System.

Summary of the Alternatives

Only alternatives that were judged to be "economic" were analyzed in the OSRA. The alternatives analyzed are the Kaktovik Deferral Alternative (fig. 4), the Area Offshore of the Alaska National Wildlife Refuge (ANWR) Alternative (fig. 5), and the Cross Island Area Alternative (fig. 6). The estimated volumes of oil resources (leased, discovered, and produced) for these alternatives were assumed to be 0.24-0.48 Bbbl (Kaktovik Deferral), 0.21-0.45 Bbbl (Area Offshore of the ANWR), and 0.28-0.55 Bbbl (Cross Island Area). The alternatives were analyzed by using the combined probabilities (representing oil-spill occurrence and contact), which excluded oil resources assumed (over the low end of the resource range) to be located in those portions of the hypothetical launch areas (as shown in fig. 2) that overlay the respective areas. Oil resources were excluded from hypothetical L7 and L8 in the analysis of the Kaktovik Deferral, from L5 through L8 for the Area Offshore of the ANWR, and from L3 and L4 for the Cross Island Area.

Framework of the Analysis

The oil-spill risk analysis depends not only on the meteorologic, oceanographic, geographic, and sea-ice conditions of the study area, but also on the environmental resource areas at risk from oil spills and the estimated volume of oil resources assumed to be leased, discovered, produced, and transported.

Environmental Resource Areas

Environmental resource areas were selected by the MMS analysts in the Alaska OCS Region who prepare the EIS for the lease sale. Appendix A (figs. A-1 through A-7) contains maps showing the digitized environmental resource area locations. The locations of 32 environmental resource areas were digitized in the same coordinate system, or base map, used for the trajectory simulations. These environmental resource areas are composed of Ice/Sea Segments 1-17, 6 bays and lagoons (Peard Bay, Elson Lagoon, Simpson Lagoon, Gwydry Bay, Jago Lagoon, and Beaufort Lagoon), Subsistence Resource Areas A-D, Fall Feeding Area, Summer Feeding Areas 1-2, and Northern and Southern Spring Lead Areas. Each environmental resource area that is typically present year-round was treated as being vulnerable to oil all year. This method assumes that the environmental resource area is sensitive to oil-spill effects throughout the entire year rather than seasonally. Each environmental resource area not usually present year-round was treated as vulnerable to contact from oil spills only during the months it is likely to be present. Several environmental resource areas were analyzed as being vulnerable on both an annual and a seasonal basis. The digitized environmental resource areas, their months of vulnerability, and the illustration depicting their locations are listed below.

Because the trajectory model simulates an oil spill as a point, each environmental resource area was digitized with an areal extent slightly greater than it actually occupies. For example, the shoreline environmental resource areas extend a short distance offshore, thus allowing the model to simulate a spill that approaches and partially contacts the environmental resource area (or boundary segment), then withdraws and continues along its path. For this analysis, the trajectory simulations were calculated for two seasons—winter (October-June) and summer (July-September).

Environmental Resource Area	Months Assumed Vulnerable	Figure
Ice/Sea Segments 1-17	January-December	A-1
Ice/Sea Segments 1-4*	May-June	A-1
Peard Bay	June-October	A-2
Elson Lagoon, Simpson Lagoon, Gwydry Bay, Jago Lagoon, and Beaufort Lagoon	July-September	A-2
Subsistence Resource Areas A-D	January-December	A-3, A-4
Fall Feeding Area	September-October	A-3
Summer Feeding Areas 1 and 2	June-September	A-3, A-4
Southern and Northern Spring Lead System Areas	January-December	A-5, A-6
Northern Spring Lead System Area*	May-June	A-5

* These resources were analyzed as vulnerable year-round and separately as vulnerable only during specific months.

In order to quantify spill trajectories that may travel out of the study area, the locations of 38 boundary segments (B1-B38) located along the edges of the study area were also

digitized in the same coordinate system used for the trajectory simulations (fig. A-7). Each boundary segment was treated as being vulnerable all year.

Also included in the analysis was an additional environmental resource area, land, which comprised the entire study area coastline. Upon contacting land, the simulated trajectory ends. Land was further analyzed by dividing the study area coastline into 61 land segments of approximately equal lengths (24 km), as shown in figure 7.

Estimated Volume of Oil Resources

For this analysis, both benefits and risks are functions of the volume of oil and are mutually dependent. For example, greater volumes of oil are associated with greater economic benefits as well as greater risks. If the benefits are evaluated by assuming production of a specific amount of oil, then the corresponding risks should be stated conditionally, such as "the risks are . . . , given that the volume is" Any statements about the likelihood of a particular volume of oil being present apply equally to the likelihood of the corresponding benefits and risks.

The estimated volumes of oil resources used for oil-spill risk calculations in this report correspond to those that will be used in the final EIS for the lease sale (Alternative Options revised by MMS in August 1997). Revisions to the original resource estimates found in Anderson et al. (1996) represent an increased volume of oil estimated to be deferred by the Kaktovik Deferral Alternative due to the incorporation of information on recent development. The revisions also introduced two new alternatives, for the Area Offshore of the ANWR and Cross Island Area.

The hypothetical development and transportation scenario for the Proposed Action is contingent on an assumed range of 0.35-0.67 Bbbl of oil. The development and transportation scenarios for the deferral alternatives assume reduced ranges of 0.24-0.48 Bbbl (Kaktovik Deferral), 0.21-0.45 Bbbl (Area Offshore of the ANWR), and 0.28-0.55 Bbbl (Cross Island Area).

It should be emphasized that these estimates and the following analyses assume that geologically recoverable quantities of oil are present. If no geologically recoverable quantities of oil are present, no crude oil-spill risks exist from OCS production.

Oil-Spill Risk Analysis (OSRA)

The OSRA consists of three parts: (1) the probability of oil spills occurring, (2) the trajectory simulation of oil spills, and (3) the combination of oil-spill occurrence probabilities and trajectory simulations. Contingent upon actual discovery of geologically recoverable quantities of oil, exploration and development are assumed to take place over 12 years (1999-2010), and production is assumed to span a period of 21 years (2006-2026).

Probability of Oil Spills Occurring

The probability of oil spills occurring (given that geologically recoverable quantities of oil are present) assumes that spills occur independently of each other as a Poisson process. The Poisson process is a statistical distribution commonly used to model random events. The probability of oil spills occurring is based on a spill rate derived from past OCS platform and pipeline experience in U.S. waters, and depends on the volume of oil produced and transported. All types of accidental spills greater than or equal to 1,000 barrels (bbl) were considered in this analysis. These spills include not only well blowouts but also other accidents that occur on platforms and during transportation of oil to shore. These accidents were classified as either platform or pipeline spills. This classification allows the analyst to compare the risks from each spill source between the Proposed Action and any alternatives.

Anderson and LaBelle (1994) examined oil-spill occurrence rates applicable to the U.S. Outer Continental Shelf. Their results, adjusted for recent experience and based upon more complete databases than were available for earlier analyses (Anderson and LaBelle, 1990; Lanfear and Amstutz, 1983), indicated some significant changes in the spill rates for platforms and pipelines. This report uses these updated spill occurrence rates.

Spill rates are expressed as number of spills per billion barrels (spills/Bbbl) of oil produced or transported. Only spills greater than or equal to 1,000 bbl are addressed because smaller spills may not persist long enough to be simulated by trajectory modeling. Another consideration is that a large spill is likely to be identified and reported; therefore, these records are more comprehensive than those of smaller spills. (Small, chronic spillage is addressed in MMS's environmental analyses without the use of trajectory modeling.)

Two basic criteria were used in selecting the risk exposure variable: (1) the exposure variable should be simple to define and (2) it should be a quantity that can be estimated. The volume of oil produced or transported was the chosen exposure variable primarily for the following reasons: historic volumes of oil produced and transported are well documented, using these volumes makes the calculation of the estimated oil-spill occurrence rate simple—the ratio of the number of historic spills to volume of oil produced or transported, and future volumes of oil production and transportation are routinely estimated. Oil volume estimates are prepared by MMS Resource Evaluation program analysts; their function and expertise involve the assessment of oil resources by using comprehensive geological and geophysical databases and related models. In addition, the MMS analysts estimate most other exposure variables (such as number of platforms and tanker trips) as a function of the volume of oil produced or transported.

Anderson and LaBelle (1994) analyzed platform and pipeline spills that occurred on the U.S. Outer Continental Shelf from 1964 through 1992. In these analyses, every spill record was examined and verified to the furthest extent possible. Each spill was classified according to its applicability to the analysis for size, product spilled, and spill source.

In an earlier analysis (Anderson and LaBelle, 1990), nonparametric tests were applied to determine whether OCS platform and pipeline spills from 1964 through 1987 were random and independent. For these observations, the volume of oil produced and transported between spills appeared to be nonrandom, increasing over time. Extending the data through 1992, Anderson and LaBelle (1994) showed that the OCS platform spill rate continued to decline. However, with the occurrence of four pipeline spills between 1988 and 1992, there was no longer any evidence of a corresponding decrease in the estimated OCS pipeline spill occurrence rate.

For spills greater than or equal to 1,000 bbl, the spill rates are 0.45 spills/Bbbl of produced oil for U.S. Outer Continental Shelf platforms and 1.32 spills/Bbbl of transported oil for U.S. Outer Continental Shelf pipelines. The platform spill rate reflects the observed decline in spill occurrence mentioned above. This rate is based on the most recent 73 percent of the production record, which includes only 27 percent of the historic platform spills (3 spills/6.6 Bbbl). The pipeline spill rate of 1.32 spills/Bbbl is based on the entire OCS production (transportation) and pipeline spill records (12 spills/9.1 Bbbl).

In summary, the spill rates (expressed as number of spills \geq 1,000 bbl/Bbbl of produced or transported oil) used in this report are as follows:

Platforms—0.45 spills/Bbbl
Pipelines—1.32 spills/Bbbl

Oil-spill occurrences (spills \geq 1,000 bbl) are considered to be governed by a Poisson process (Smith et al., 1982; Lanfear and Amstutz, 1983). The probability of a specific number of spills $p(n)$ occurring is described by the Poisson distribution:

$$p(n) = \frac{e^{-\lambda} * \lambda^n}{n!}$$

where n is the specific number of spills (0, 1, 2, ..., n), e is the base of the natural logarithm, and λ is the parameter of the Poisson distribution. For oil spills, the Poisson parameter (λ) is equal to the spill rate multiplied by the volume of oil to be produced or transported. The spill rate has dimensions of number of spills/Bbbl, and the volume is expressed in Bbbl. Therefore, λ denotes the mean number of spills estimated to occur as a result of production or transportation of a specific volume of oil.

Oil-spill occurrence estimates for spills greater than or equal to 1,000 bbl were calculated for production and transportation of oil during the assumed production life of the Proposed Action (2006-2026). Table 1 shows the mean number of spills estimated to occur and the probability of one or more spills occurring in the study area over the assumed production life of the proposed lease sale.

This analysis addresses only the risk of platform and pipeline spills from the production and transportation (via offshore pipeline) of Beaufort OCS Lease Sale 170 resources to the Trans-Alaska Pipeline System. Historically, North Slope crude transported via the Trans-Alaska Pipeline System has been loaded onto tankers at Valdez, Alaska (the pipeline terminus) and shipped to the U.S. west coast and other destinations. These crude oil movements have a historic spill rate of 1.10 spills ($\geq 1,000$ bbl) per billion barrels handled (Anderson and LaBelle, 1994). Using this rate and assuming Beaufort OCS Lease Sale 170 resources are transported by tanker from Valdez, the mean number of spills and the probability of one or more spills associated with the movement of the North Slope crude from Valdez occurring outside the study area are estimated as follows:

<u>Scenario</u>	<u>Mean Number of Spills</u>	<u>Probability of One or More Spills</u>
Proposed Action	0.38-0.74	32-52%
Kaktovik Deferral	0.26-0.53	23-41%
Area Offshore of the ANWR	0.23-0.50	21-39%
Cross Island Area	0.31-0.61	27-46%

Oil-Spill Trajectory Simulations

The trajectory simulation portion of the model consists of many hypothetical oil-spill trajectories that collectively represent the mean surface transport and the variability of the surface transport as a function of time and space. The trajectories represent the Lagrangian motion that a particle on the surface might take under given wind, ice, and ocean current conditions. Multiple trajectories are simulated to give a statistical representation, over time and space, of possible transport under the range of wind, ice, and ocean current conditions that exist in the area.

Trajectories are constructed from simulations of wind-driven and density-induced ocean flow fields, and the ice motion field. The basic approach is to simulate these time and spatially dependent currents separately, then combine them through linear superposition to produce an oil-transport vector. This vector is then used to create a trajectory. Simulations are performed for two seasons, winter (October-June) and summer (July-September). The choice of this seasonal division was based on meteorological, climatological, and biological cycles, as well as consultation with Alaska Region EIS analysts. The modeling of each ice motion field and ocean current component is detailed in Hedström, Haidvogel and Signorini (1995) and Hedström (1994). Brief summaries of the method and assumptions follow.

For cases where the ice concentration is below 80 percent, each trajectory is constructed using vector addition of the ocean current field and 3.5 percent of the instantaneous wind field—a method based on work done by Huang and Monastero (1982), Smith et al. (1982), and Stolzenbach et al. (1977). For cases where the ice concentration is 80 percent or greater, the model ice velocity is used to transport the oil. Equations 1 and 2 show the components of motion that are simulated and used to describe the oil transport:

$$U_{oil} = U_{current} + 0.035 U_{wind} \quad (1)$$

or

$$U_{oil} = U_{ice} \quad (2)$$

where: U_{oil} = oil drift vector

$U_{current}$ = current vector (when ice concentration < 80%)

U_{wind} = wind speed at 10 m above the sea surface

U_{ice} = ice vector (when ice concentration \geq 80%)

The wind drift factor was estimated to be 0.035, with a variable drift angle ranging from 0° to 25° clockwise. The drift angle was computed as a function of wind speed according to the formula in Samuels et al. (1982). (The drift angle is inversely related to wind speed.)

For each trajectory simulation, the start time for the first trajectory was the first day of the season (winter or summer) of the first year of wind data (1978) at 6 a.m. Greenwich Mean Time (GMT). Each subsequent trajectory was started every 1.5 days on average, at 6 a.m. GMT. A total of 2,000 trajectories (1,500 in winter, 500 in summer) was launched from each of the 344 hypothetical spill sites distributed over the proposed lease sale area over the 9 years of wind data (1978-1986). Results of these trajectory simulations were combined into eight hypothetical launch areas (L1-L8) to represent platform risk (fig. 2). Transportation risks were represented by 2,000 trajectories launched uniformly along each hypothetical pipeline route (P1-P7—fig.3).

For the Beaufort/Chukchi Sea, the $U_{current}$ and U_{ice} are simulated using a three-dimensional coupled ice-ocean hydrodynamic model (Hedström et al., 1995; Hedström, 1994). The model is based on the ocean model of Haidvogel et al. (1991) and the ice model of Hibler (1979). The location of each trajectory at each time interval is used to select the appropriate ice concentration. Depending on the ice concentration, either the ice or water velocity with wind drift from the stored results of the Haidvogel et al. (1991) coupled ice-ocean model is used (see equations (1) and (2) above). Surface transport of the oil slick for each spill was simulated as a series of straight-line displacements in 3-hour increments of a point governed by the U_{oil} vectors.

The trajectories age while they are in the water/on the ice. For each day that the hypothetical spill is in the water, the spill ages—up to a total of 30 days. While the spill is in the ice (\geq 80% concentration), the aging process is suspended. The maximum time allowed for the transport of oil in the ice is 180 days; after this, the trajectory is terminated. The 30-day limit is maintained for spill trajectories in open water.

Summer trajectories are those that start between the beginning of July and the end of September. Therefore, if any contact to an environmental resource area or land segment is made by a trajectory that began at the end of September, it is considered a summer contact and is counted along with the rest of the contacts from spills launched in the summer.

The wind data set used came from the National Weather Service Limited Fine Mesh (LFM) model (Gerrity, 1977), and the 9-year simulation covered both the low frequency variability and interannual variability. A major assumption used in this analysis is that the ice motion velocities and the ocean daily flows calculated by the coupled ice-ocean model adequately represent the flow components. Sensitivity tests and comparisons with data illustrate that the ice-ocean model captures the first-order transport and the dominant flow (Hedström et al., 1995).

After quality assurance checks (performed by MMS) were passed, the trajectories were used in the OSRA model structure. The OSRA model was run, given the land/sea segments and environmental resource areas specified for this analysis. Plots of trajectories and overlays of land/sea segments and environmental resource areas were examined to ensure that contacts were properly established and tabulated. The quality assurance checks provide an important means of gaining information and insight into the behavior of the simulated trajectories. The conditional probabilities of contact (probabilities that an oil spill will contact an environmental resource area within a given travel time) offer the analyst much oceanographic information. Some summary information is provided below to help identify the particular trajectory behavior.

When examining any one trajectory, it is important to be aware of the relative scales or sizes of the flow components that have been combined to produce the final U_{oil} vector. The wind-induced surface drift is the dominant component of U_{oil} except where the density-induced circulation is very strong or where the ice concentration is greater than or equal to 80 percent. In addition, the wind-driven flow component contains seasonal trends as well as a large degree of variability. Because many other factors are related to wind forcing (such as mixing in the surface layer and biological, chemical, and physical processes), it is useful to compare the envelopes of the trajectories to seasonal mixing regimes and processes. Some information can be gained by comparing trajectories from winter versus summer (when wind speeds and ice concentration are high versus low).

The second dominant transport factor is the density-induced flow. This forcing results in well-organized, coherent envelopes that follow the general trend of the coastline. The degree to which wind forcing plays a role in the nearshore areas has been studied as part of the Beaufort Mesoscale Circulation Study (Aagaard et al., 1990), and there is often a reinforcement of the wind-driven component by the density-driven component.

An important test is to compare observed drifter tracks with trajectories. Some historic data exist (Colony and Thorndike, 1984; Aagaard et al., 1990), but comparing drifter tracks with trajectories must be done carefully, especially considering the ice station trajectories. Ice stations on large flows that are at the ice edge move differently than ice stations in the pack ice. In general, the results of the coupled ice-ocean hydrodynamic model show these features at the correct location and magnitude, meeting the zero-th order test of the basic model. As expected, the simulated trajectories appear more variable than these drifter tracks. To a large degree, the Uoil transport vector has components of wind-driven and surface-driven motion. Thus, the observations and simulations contain similar trends, but an exact match should not be expected.

In conclusion, the simulated spill trajectories for Beaufort Sea OCS Lease Sale 170 show distinct variations in response to the seasonal wind patterns and the strength of density-driven currents. Hypothetical spills on the shelf demonstrate the wind-induced variability and the relatively important density-driven current along the Beaufort Sea coast. Landfall or contact of the simulated trajectories is generally highest year-round for points located west of the hypothetical areas, with contacts east of these areas occurring in the summer.

As the simulated oil spills moved, any contacts with environmental resource area were recorded. Spill movement continued until the spill contacted land, moved out of the study area, or aged more than 30 days in open water or 180 days in ice conditions.

The trajectories simulated by the model represent hypothetical pathways of oil slicks; they do not directly consider cleanup, dispersion, or weathering processes that could determine the quantity or properties of the oil that might eventually contact environmental resource areas or land segments. An implicit analysis of weathering and decay can be considered by noting the age of a simulated trajectory when it contacts an environmental resource area. For this analysis, the periods selected were 3, 10, and 30 days (180 days in ice conditions) to represent implicit measures of oil weathering as well as matters relating to containment and cleanup.

Conditional Probabilities

The probability that an oil spill will contact a specific environmental resource area within a given time of travel from a certain location or spill site is termed a conditional probability, the condition being that a spill is assumed to have occurred. Conditional probabilities of contact for 3, 10, and 30 days (180 days in ice conditions) were calculated for the eight hypothetical launch areas shown in figure 2 (referred to as L1-L8) and for the pipeline route segments shown in figure 3 (referred to as P1-P7). These conditional probabilities of contact with environmental resource areas and land

segments are presented in tables 2 through 9 (on an annual basis) and in appendices B and C (on a seasonal basis).

Combined Analysis of Oil-Spill Occurrence and Oil-Spill Trajectory Simulations

A critical difference exists between the conditional probabilities and the combined probabilities calculated in this part of the OSRA. Conditional probabilities assume that a spill has occurred and, thus, depend only on the winds, ice, and ocean currents in the study area. Combined probabilities, on the other hand, depend not only on the physical conditions, but also on the chance of spill occurrence, the estimated volume of oil to be produced or transported, and the oil transportation scenario. These additional variables make the combined probabilities sensitive to the course of action chosen by the decisionmaker, that is, choosing to lease or not to lease certain tracts or implementing specific transportation stipulations. The annual combined probabilities for the various leasing alternatives (the Proposed Action, the Kaktovik Deferral, the Area Offshore of the ANWR, and the Cross Island Area) are presented in tables 10-17.

In calculating the combined probabilities, those that represent probabilities of both oil-spill occurrence and contact, the following steps are performed.

1. For a set of n_l environmental resource areas and n_l spill sites, the conditional probabilities can be represented in a matrix form. Let [C] be an $n_l \times n_l$ matrix, where each element $c_{i,j}$ is the probability that an oil spill will contact environmental resource area i , given that a spill occurs at hypothetical spill site j . Note that spill sites can represent potential spill starting points at production areas or transportation routes.
2. Spill occurrence can be represented by another matrix [S]. With n_l spill sites and n_s production sites, the dimensions of [S] are $n_l \times n_s$. Let each element $s_{j,k}$ be the estimated mean number of spills occurring at spill site j owing to production of a unit volume (1 Bbbl) of oil at site k . These spills can result from either production or transportation. The $s_{j,k}$ can be determined as a function of the volume of oil (spills/Bbbl). Each column of [S] corresponds to one production site and one transportation route. If alternative and mutually exclusive transportation routes are considered for the same production site, they can be represented by additional columns of [S], thus increasing n_s .
3. Matrix [U] is defined as:

$$[U] = [C] \times [S]$$

Matrix [U]—which has dimensions $n_t \times n_s$ —is termed the unit risk matrix. Each element $u_{i,k}$ corresponds to the estimated mean number of spills occurring and contacting environmental resource area i , owing to the production of a unit volume (1 Bbbl) of oil at site k .

4. With [U], the mean number of contacts to each environmental resource area are estimated, given a set of oil volumes at each site. Let [V] be a vector of dimension n_s where each element v_k corresponds to the volume of oil expected to be found at production site k . Then, if [L] is a vector of dimension n_t , where each element λ_i corresponds to the mean number of contacts to environmental resource area i , the formula is:

$$[L] = [U] \times [V]$$

Thus, estimates of the mean number of oil spills ($\geq 1,000$ bbl) that will both occur and contact environmental resource areas (or land segments) can be calculated. (Note that as a statistical parameter, the mean number can assume a fractional value, even though a fraction of an oil spill has no physical meaning.)

Using Bayesian techniques, Devanney and Stewart (1974) showed that the probability of n oil-spill contacts can be described by a negative binomial distribution. However, Smith et al. (1982) noted that when actual exposure is much less than historic exposure, as is the case here, the negative binomial distribution can be approximated by a Poisson distribution. The Poisson distribution has a significant advantage in calculations because it is defined by only one parameter, the assumed number of spills. Thus, the matrix [L] contains all the information needed to use the Poisson distribution—if $P(n,i)$ is the probability of exactly n contacts to environmental resource area i , then:

$$P(n,i) = \frac{\lambda_i^n * e^{-\lambda_i}}{n!}$$

Discussion

The Proposed Action, which assumes production and transportation of 0.35-0.67 Bbbl of oil, has the possibility of an estimated mean number of 0.62-1.19 spills greater than or equal to 1,000 bbl occurring (0.16-0.30 from platforms and 0.46-0.89 from pipelines), as shown in table 1. The alternatives reduce the estimated mean number of spills to 0.43-0.85 spills (Kaktovik Deferral), 0.37-0.79 spills (Area Offshore of the ANWR), and 0.50-0.97 spills (Cross Island Area), as also shown in table 1.

There is an estimated 46- to 70-percent probability (expressed as percent chance) that one or more spills ($\geq 1,000$ bbl) may occur as a result of the Proposed Action (see table 1). The deferral alternatives reduce this estimated probability to 35-57 percent (Kaktovik Deferral), 31-55 percent (Area Offshore of the ANWR), and 39-62 percent (Cross Island Area).

There are estimated conditional probabilities (expressed as percent chance) of 7-19 percent that a spill originating from hypothetical coastal launch areas (L2, L4, L6, L8) will contact land within 3 days; however, hypothetical launch areas farther offshore (L1, L3, L5, L7) have probabilities of 2-3 percent (table 2). Within a 10-day period, these probabilities increase to 12-26 percent from the hypothetical coastal launch areas and 7-8 percent from the hypothetical launch areas further offshore (table 3). The estimated probabilities of a spill originating from any of these hypothetical launch areas contacting land within 30 days are 12-35 percent (table 4). The estimated probabilities of a spill originating from any of the eight hypothetical launch areas contacting land within 180 days (in ice conditions) are 23-58 percent (table 5).

The highest estimated conditional probabilities of contact to land from transportation are from hypothetical pipeline route segments P5 through P7 (all three are pipeline route segments that make landfall—see fig. 3). The estimated conditional probabilities that a spill originating from these three pipeline route segments will contact land within 3 days are 8-15 percent (table 2). These probabilities increase to 15-20 percent within 10 days, 22-26 percent within 30 days (the longest period that spills were assumed to persist in open water), and 50-51 percent within 180 days (in ice conditions), as shown in tables 3, 4, and 5 respectively. The remaining pipeline route segments have estimated probabilities of contact similar to those of the hypothetical launch areas in which they are located.

In general, only environmental resource areas located over areas of the lease offering have high ($\geq 12\%$) probabilities of being contacted by a spill from the proposed lease area or its related transportation segments within 3 days (table 2). For example, Subsistence Resource Area C has a 99.5-percent-or-greater estimated conditional probability of being contacted within 3 days from spills originating at L2, L4, P1, P2 and P5-P7, and Subsistence Area D has a 99.5-percent-or-greater estimated conditional probability of being contacted within 3 days from spills originating at L6-L8, P3, P4, and P7.

Tables 6 through 9 list the annual conditional probabilities of a spill contacting a specific land segment (fig. 4) within a 3-, 10-, and 30-day period (the longest period that spills were assumed to persist in open water) and a 180-day period (in ice conditions). Generally, each land segment appears to be at greatest risk of contact by spills originating at sources immediately to its north. The conditional probability of any one land segment being contacted within 30 days does not exceed 11 percent from any hypothetical launch area (from L8 to land segment 40) or 16 percent from any pipeline segment (from P6 to land segment 34). Within 180 days, the probabilities of an oil spill contacting any one land segment increase to a maximum of 16 percent from any hypothetical launch area (from L8 to land segment 38) and 27 percent from any pipeline segment (from P6 to land segment 34).

Table 10 presents the range of estimated combined probabilities of one or more spills ($\geq 1,000$ bbl) occurring and contacting environmental resource areas for the Proposed Action as a result of the production and transportation scenarios. (A few of the Table 10

values have changed slightly compared to the original report (Anderson et al., 1996) because of revisions MMS made to oil resource allocations.) The combined probabilities that one or more spills ($\geq 1,000$ bbl) will occur and contact land within 3 days are 5-9 percent. These probabilities of occurrence and contact are 8-15 percent within 10 days, and 12-22 percent within 30 days (the longest period that spills were assumed to persist in open water). Within 180 days (in ice conditions), these probabilities are 23-40 percent. The chance of one or more spills occurring and contacting an environmental resource area within 3 days appears to be greatest for Subsistence Resource Areas C (39%-62%) and D (30%-47%). These probabilities increase by no more than 2 percentage points within 10 and 30 days, and they increase by no more than 6 percentage points within 180 days (42-65% for Area C and 34-53% for Area D).

Table 11 shows the range of estimated combined probabilities of one or more spills ($\geq 1,000$ bbl) occurring and contacting land segments as a result of the production and transportation scenarios. (A few of the Table 11 values have changed slightly compared to the original report (Anderson et al., 1996) because of revisions MMS made to oil resource allocations.) These probabilities do not exceed 3-5 percent within a 30-day period (maximum time in open water). Within 180 days (in ice conditions), the probabilities of one or more spills occurring and contacting land segments do not exceed 5-10 percent. The highest probability of one or more spills occurring and contacting a land segment within 30 days is for Land Segment 34 (3-5%), which is located between Oliktok Point and Point McIntyre (fig. 3).

Table 12 presents the range of estimated combined probabilities of one or more spills ($\geq 1,000$ bbl) occurring and contacting environmental resource areas for the Kaktovik Deferral Alternative as a result of the production and transportation scenarios. (A few of the Table 12 values have changed slightly compared to the original report (Anderson et al., 1996) as a result of revisions MMS made to oil resource allocations.) As compared to the Proposed Action, the Kaktovik Deferral Alternative results in an estimated reduction of 1-2 percentage points in the chance of one or more spills ($\geq 1,000$ bbl) occurring and contacting land within 3 days, and up to a 6-9 percentage point reduction in the chance of one or more spills occurring and contacting land within 180 days. This deferral is estimated to result in a 15-21 percentage point reduction in the chance of one or more spills ($\geq 1,000$ bbl) occurring and contacting Subsistence Area D (a portion of which overlays the deferral area) within 3 days. In general, the OSRA shows that environmental resource areas adjacent to the deferral area, particularly those superimposed southward or westward, would have reduced likelihoods of such spill occurrence and contact.

Table 13 shows the range of estimated combined probabilities of one or more spills ($\geq 1,000$ bbl) occurring and contacting land segments for the Kaktovik Deferral Alternative as a result of the production and transportation scenarios. (A few of the Table 13 values have changed slightly compared to the original report (Anderson et al., 1996) as a result of revisions MMS made to oil resource allocations.) As compared to the Proposed Action, the Kaktovik Deferral Alternative results in an estimated reduction of up to 1-2 percentage points in the chance of one or more spills ($\geq 1,000$ bbl)

occurring and contacting Land Segments 33-41 (the land segments adjacent to the deferral area, specifically those southward or westward) within the 3-, 10-, 30- and 180-day time periods.

Table 14 presents the range of estimated combined probabilities of one or more spills ($\geq 1,000$ bbl) occurring and contacting environmental resource areas for the Area Offshore of the ANWR Alternative as a result of the production and transportation scenarios. As compared to the Proposed Action, the Area Offshore of the ANWR Alternative results in an estimated reduction of 2 percentage points in the chance of one or more spills ($\geq 1,000$ bbl) occurring and contacting land within 3 days, and up to a 8-11 percentage point reduction within 180 days. This alternative is estimated to result in a 19- to 25-percent reduction in the chance of one or more spills ($\geq 1,000$ bbl) occurring and contacting Subsistence Area D (a portion of which overlays the deferral area) within 3 days. In general, it appears that environmental resource areas adjacent to the Area Offshore of the ANWR, particularly those superimposed southward or westward, would have reduced likelihoods of such spill occurrence and contact.

Table 15 shows the range of estimated combined probabilities of one or more spills ($\geq 1,000$ bbl) occurring and contacting land segments for the Area Offshore of the ANWR Alternative as a result of the production and transportation scenarios. As compared to the Proposed Action, the Area Offshore of the ANWR Alternative results in an estimated reduction of up to 1-2 percentage points in the chance one or more spills ($\geq 1,000$ bbl) occurring and contacting Land Segments 33-41 (the land segments adjacent to the alternative area, specifically those southward or westward) within the 3-, 10-, 30- and 180-day time periods. As compared to the Kaktovik Deferral Alternative (a subset of the Area Offshore of the ANWR Alternative), there is up to an additional 1 percentage point in the chance of one or more spills occurring and contacting Land Segments 34, 36, 38, and 39 within 10 or more days.

Table 16 presents the range of estimated combined probabilities of one or more spills ($\geq 1,000$ bbl) occurring and contacting environmental resource areas for the Cross Island Area Alternative as a result of the production and transportation scenarios. As compared to the Proposed Action, it appears that the Cross Island Area Alternative results in an estimated reduction of 1-2 percentage points in the chance of one or more spills ($\geq 1,000$ bbl) occurring and contacting land within 3 days, and up to a 4-6 percentage point reduction within 180 days. This alternative is estimated to result in 8-9 percentage point reduction in the chance of one or more spills ($\geq 1,000$ bbl) occurring and contacting Subsistence Area C (a portion of which overlays the alternative area) within 3 days. In general, the OSRA shows that environmental resource areas adjacent to the Cross Island Area, particularly those superimposed southward or westward, appear to have reduced likelihoods of such spill occurrence and contact.

Table 17 shows the range of estimated combined probabilities of one or more spills ($\geq 1,000$ bbl) occurring and contacting land segments for the Cross Island Area Alternative as a result of the production and transportation scenarios. As compared to the Proposed Action, the Cross Island Area Alternative results in an estimated reduction of 3-5 percentage points in the chance of one or more spills ($\geq 1,000$ bbl)

occurring and contacting Land Segment 34 (the land segment beneath the deferral area) within 180 days, and a reduction of up to 1-2 percentage points in the chance of one or more spills ($\geq 1,000$ bbl) occurring and contacting Land Segments 28 and 32-37 (the land segments adjacent to the Cross Island Area, specifically those southward or westward) within the 3-, 10-, 30- and 180-day time periods.

References Cited

- Aagaard, K., C.H. Pease, A.T. Roach, and S.A. Salo, 1990, Beaufort Sea mesoscale circulation study, final report, OCS Study MMS 90-0002, 136 pp. plus appendices.
- Anderson, C.M., W.R. Johnson, C.F. Marshall, and E.M. Lear, 1996, Oil-Spill Risk Analysis: Beaufort Sea Outer Continental Shelf Lease Sale 170, OCS Report MMS 96-0068, 58 pp.
- Anderson, C.M., and R.P. LaBelle, 1990, Estimated occurrence rates for accidental oil spills on the U.S. Outer Continental Shelf: *Oil and Chemical Pollution* 6:21-35.
- Anderson, C.M., and R.P. LaBelle, 1994, Comparative occurrence rates for offshore oil spills: *Spill Science and Technology Bulletin* 1(2):131-141.
- Colony, R., and A.S. Thorndike, 1984, An estimation of the mean field of ice motion: *Journal of Geophysical Research* 89:10623-10626.
- Devanney, M.W., III, and R.J. Stewart, 1974, Analysis of oilspill statistics: Massachusetts Institute of Technology (Cambridge), prepared for the Council on Environmental Quality, Report No. MITSG-74-20.
- Gerrity, J.P., Jr., 1977, The LFM model—1976: A documentation: NOAA Technical Memorandum NWS NMC 60, NOAA National Weather Service.
- Haidvogel, D.B., J.L. Wilkin, and R. Young, 1991, A semi-spectral primitive equation ocean circulation model using vertical sigma and orthogonal curvilinear horizontal coordinates: *Journal of Computational Physics* 94:151-185.
- Hedström, K.S., 1994, Technical manual for a coupled sea-ice/ocean circulation model (version 1), Technical Report, Institute of Marine and Coastal Sciences, Rutgers University, prepared for the Minerals Management Service, OCS Study MMS 94-0020, 117 p.
- Hedström, K.S., D.B. Haidvogel, and S. Signorini, 1995, Model simulations of ocean/sea-ice interaction in the western arctic in 1983, final report to the Minerals Management Service, OCS Study MMS 95-0001, 78 p.

- Hibler, W.D., III, 1979, A dynamic thermodynamic sea ice model: *Journal of Physical Oceanography* 17:815-849.
- Huang, J.C., and F.C. Monastero, 1982, Review of the state-of-the-art of oil spill simulation models: Final report submitted to American Petroleum Institute.
- LaBelle, R.P., and C.M. Anderson, 1985, The application of oceanography to oil-spill modeling for the Outer Continental Shelf oil and gas leasing program: *Marine Technology Society* 19(2):19-26.
- Lanfear, K.J., and D.E. Amstutz, 1983, A reexamination of occurrence rates for accidental oil spills on the U.S. Outer Continental Shelf: Proceedings of the Eighth Conference on the Prevention, Behavior, Control, and Cleanup of Oil Spills, San Antonio, Tex., February 28-March 3, 1983.
- Samuels, W.B., N.E. Huang, and D.E. Amstutz, 1982, An oil spill trajectory analysis model with a variable wind deflection angle: *Ocean Engineering* 9:347-360.
- Smith, R.A., J.R. Slack, T. Wyant, and K.J. Lanfear, 1982, The oil spill risk analysis model of the U.S. Geological Survey: U.S. Geological Survey Professional Paper 1227.
- Stolzenbach, K.D., O.S. Madsen, E.E. Adams, A.M. Pollack, and C.K. Cooper, 1977, A review and evaluation of basic techniques for predicting the behavior of surface oil slicks: Ralph M. Parsons Laboratories, Report No. 222.




Figure 1. OSRA study area and proposed lease sale area (shaded) for proposed Beaufort Sea OCS Lease Sale 170.




Figure 2. Enlarged representation of launch areas (L1-L8) for proposed Beaufort Sea OCS Lease Sale 170.




Figure 3. Enlarged representation of pipeline route segments (P1-P7) for proposed Beaufort Sea OCS Lease Sale 170.




Figure 4. Enlarged representation of the Kaktovik Deferral Alternative for proposed Beaufort Sea OCS Lease Sale 170.




Figure 5. Enlarged representation of the Alaska National Wildlife Refuge (ANWR) Deferral Alternative for proposed Beaufort Sea OCS Lease Sale 170




Figure 6. Enlarged representation of the Cross Island Deferral Alternative for proposed Beaufort Sea OCS Lease Sale 170.




Figure 7. Study area coastline divided into 61 land segments for proposed Beaufort Sea OCS Lease Sale 170.

Table 1. Oil-spill occurrence probability estimates for spills greater than or equal to 1,000 barrels resulting over the assumed production life or the proposed action and alternative actions, Beaufort Sea OCS Lease Sale 170

<u>Source</u>	<u>Probability of One or More Spills</u>	<u>Mean Number of Spills</u>
<u>Proposed Action</u>		
Platforms	15-26%	0.16-0.30
Pipelines	37-59%	0.46-0.89
All Sources	46-70%	0.62-1.19
<u>Kaktovik Deferral</u>		
Platforms	10-20%	0.11-0.22
Pipelines	27-47%	0.32-0.63
All Sources	35-57%	0.43-0.85
<u>Area Offshore of the ANWR Alternative</u>		
Platforms	9-18%	0.09-0.20
Pipelines	24-45%	0.28-0.59
All Sources	31-55%	0.37-0.79
<u>Cross Island Area Alternative</u>		
Platforms	12-22%	0.13-0.25
Pipelines	31-51%	0.37-0.72
All Sources	39-62%	0.50-0.97
ANWR = Alaska National Wildlife Refuge		
Volume of Oil: 0.35-0.67 (Proposed Action)		0.24-0.48 (Kaktovik)
0.21-0.45 (Area Offshore ANWR)		0.28-0.55 (Cross Island Area)

Table 2. Annual conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location will contact a certain environmental resource area within 3 days, Beaufort Sea OCS Lease Sale 170

Environmental Resource Area	Hypothetical Spill Location															
	L1	L2	L3	L4	L5	L6	L7	L8	P1	P2	P3	P4	P5	P6	P7	
Land	3	7	2	8	2	7	3	19	3	3	3	5	9	15	8	
Ice/Sea Segment 1	
Ice/Sea Segment 2	
Ice/Sea Segment 3	
Ice/Sea Segment 4	
Ice/Sea Segment 5	
Ice/Sea Segment 6	
Ice/Sea Segment 7	9	3	2	5	1	.	.	5	.	.	
Ice/Sea Segment 8	6	3	51	5	7	1	.	.	4	12	3	.	1	6	1	
Ice/Sea Segment 9	.	.	1	1	39	16	60	4	.	1	16	12	.	.	7	
Ice/Sea Segment 10	2	
Ice/Sea Segment 11	
Ice/Sea Segment 12	
Ice/Sea Segment 13	
Ice/Sea Segment 14	
Ice/Sea Segment 15	
Ice/Sea Segment 16	
Ice/Sea Segment 17	
Ice/Sea Seg. 1 SLS	
Ice/Sea Seg. 2 SLS	
Ice/Sea Seg. 3 SLS	
Ice/Sea Seg. 4 SLS	
Peard Bay	
Elson Lagoon	
Simpson Lagoon	1	4	.	1	1	.	.	.	11	5	.	
Gwydyr Bay	.	.	.	1	.	3	1	.	.	.	12	
Jago Lagoon	1	8	.	.	.	2	.	.	.	
Beaufort Lagoon	
Subsis. Res. Area A	
Subsis. Res. Area B	
Subsis. Res. Area C	80	**	50	**	70	83	6	2	**	**	87	5	**	**	**	
Subsis. Res. Area D	1	2	18	48	78	**	**	**	2	38	**	**	.	6	**	
Fall Feeding Area	
Summer Feed. Area 1	
Summer Feed. Area 2	
Southern SLS Area	
Northern SLS Area	
Northern SLS	

Notes: ** = Greater than 99.5%; . = Less than 0.5%; SLS = spring lead system.
Boundary Segments with all values less than 0.5% are not shown.

Table 3. Annual conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location will contact a certain environmental resource area within 10 days, Beaufort Sea OCS Lease Sale 170

Environmental Resource Area	Hypothetical Spill Location														
	L1	L2	L3	L4	L5	L6	L7	L8	P1	P2	P3	P4	P5	P6	P7
Land	8	12	7	14	8	13	8	26	8	9	9	11	15	20	15
Ice/Sea Segment 1
Ice/Sea Segment 2
Ice/Sea Segment 3
Ice/Sea Segment 4
Ice/Sea Segment 5
Ice/Sea Segment 6	2	1	1	1	1	.	.	2	.	.
Ice/Sea Segment 7	15	7	5	2	1	1	.	.	10	3	1	.	9	3	1
Ice/Sea Segment 8	9	6	55	9	11	4	1	1	8	18	6	1	4	10	4
Ice/Sea Segment 9	1	1	3	4	42	19	64	8	1	3	20	18	1	2	11
Ice/Sea Segment 10	.	.	1	1	3	2	6	3	.	1	3	4	.	1	2
Ice/Sea Segment 11	1	2	2	.	.	.	2	.	.	1
Ice/Sea Segment 12
Ice/Sea Segment 13
Ice/Sea Segment 14
Ice/Sea Segment 15
Ice/Sea Segment 16
Ice/Sea Segment 17
Ice/Sea Seg. 1 SLS
Ice/Sea Seg. 2 SLS
Ice/Sea Seg. 3 SLS
Ice/Sea Seg. 4 SLS
Peard Bay
Elson Lagoon
Simpson Lagoon	2	5	1	1	3	1	.	.	12	5	.
Gwydyr Bay	.	.	1	2	1	3	.	.	.	1	1	1	.	1	13
Jago Lagoon	1	2	3	9	.	.	1	4	.	.	1
Beaufort Lagoon
Subsis. Res. Area A
Subsis. Res. Area B
Subsis. Res. Area C	84	**	58	**	75	85	14	4	**	**	89	9	**	**	**
Subsis. Res. Area D	3	5	21	50	81	**	**	**	5	40	**	**	2	9	**
Fall Feeding Area
Summer Feed. Area 1
Summer Feed. Area 2
Southern SLS Area
Northern SLS Area
Northern SLS

Notes: ** = Greater than 99.5%; . = Less than 0.5%; SLS = spring lead system.
Boundary Segments with all values less than 0.5% are not shown.

Table 4. Annual conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location will contact a certain environmental resource area within 30 days, Beaufort Sea OCS Lease Sale 170

Environmental Resource Area	Hypothetical Spill Location														
	L1	L2	L3	L4	L5	L6	L7	L8	P1	P2	P3	P4	P5	P6	P7
Land	13	20	12	21	14	21	15	35	14	14	15	18	22	26	23
Ice/Sea Segment 1
Ice/Sea Segment 2
Ice/Sea Segment 3
Ice/Sea Segment 4
Ice/Sea Segment 5	1	1	.	.	.	1	.	.
Ice/Sea Segment 6	6	4	2	1	1	.	.	.	3	1	1	.	5	1	.
Ice/Sea Segment 7	22	11	10	5	2	1	1	1	16	7	2	1	13	6	1
Ice/Sea Segment 8	12	9	61	14	17	7	3	1	12	26	11	2	6	13	7
Ice/Sea Segment 9	2	2	4	5	44	22	70	11	3	4	27	25	1	3	13
Ice/Sea Segment 10	1	1	2	3	5	5	10	5	1	2	5	8	1	2	3
Ice/Sea Segment 11	.	.	1	1	2	2	4	3	.	1	2	4	.	.	1
Ice/Sea Segment 12	1	1	.	.	.	1	.	.	.
Ice/Sea Segment 13	1	1	1	1	.	.	1	1	.	.	1
Ice/Sea Segment 14
Ice/Sea Segment 15
Ice/Sea Segment 16
Ice/Sea Segment 17
Ice/Sea Seg. 1 SLS
Ice/Sea Seg. 2 SLS
Ice/Sea Seg. 3 SLS
Ice/Sea Seg. 4 SLS
Peard Bay
Elson Lagoon
Simpson Lagoon	3	7	2	2	1	.	.	.	4	3	.	.	15	7	.
Gwydyr Bay	.	1	1	3	1	4	1	1	1	2	1	1	.	1	15
Jago Lagoon	.	.	1	1	2	3	4	12	.	1	2	5	.	1	2
Beaufort Lagoon	1	1	.	.	.	1	.	.	.
Subsis. Res. Area A
Subsis. Res. Area B
Subsis. Res. Area C	87	**	68	**	80	88	26	8	**	**	92	20	**	**	**
Subsis. Res. Area D	5	7	24	53	84	**	**	**	8	44	**	**	4	13	**
Fall Feeding Area
Summer Feed. Area 1
Summer Feed. Area 2
Southern SLS Area
Northern SLS Area
Northern SLS
Boundary Segment 2	1	1	1	.	.	1	1	.	.	.

Notes: ** = Greater than 99.5%; . = Less than 0.5%; SLS = spring lead system.
Boundary Segments with all values less than 0.5% are not shown.

Table 5. Annual conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location will contact a certain environmental resource area within 180 days, Beaufort Sea OCS Lease Sale 170

Environmental Resource Area	Hypothetical Spill Location															
	L1	L2	L3	L4	L5	L6	L7	L8	P1	P2	P3	P4	P5	P6	P7	
Land	26	44	23	44	29	46	29	58	28	29	34	37	50	51	51	
Ice/Sea Segment 1	
Ice/Sea Segment 2	
Ice/Sea Segment 3	.	.	1	.	1	
Ice/Sea Segment 4	1	.	1	.	1	1	1	.	1	1	1	1	.	.	.	
Ice/Sea Segment 5	2	1	1	1	1	1	1	.	1	1	1	.	1	1	.	
Ice/Sea Segment 6	18	11	5	2	2	1	1	1	9	3	2	1	14	1	1	
Ice/Sea Segment 7	48	28	25	15	3	2	2	1	36	20	3	2	28	16	3	
Ice/Sea Segment 8	25	25	80	37	37	19	8	4	34	59	32	6	18	32	16	
Ice/Sea Segment 9	4	4	6	10	54	42	87	37	6	7	48	61	3	4	31	
Ice/Sea Segment 10	5	3	9	8	16	13	32	17	6	8	15	22	2	5	8	
Ice/Sea Segment 11	.	.	4	1	8	7	13	9	2	5	6	10	.	.	6	
Ice/Sea Segment 12	2	3	6	5	.	.	3	5	.	.	3	
Ice/Sea Segment 13	1	1	2	2	.	.	1	3	.	.	1	
Ice/Sea Segment 14	
Ice/Sea Segment 15	
Ice/Sea Segment 16	
Ice/Sea Segment 17	
Ice/Sea Seg. 1 SLS	
Ice/Sea Seg. 2 SLS	
Ice/Sea Seg. 3 SLS	
Ice/Sea Seg. 4 SLS	
Peard Bay	
Elson Lagoon	
Simpson Lagoon	6	15	7	7	3	1	.	.	9	10	2	.	20	14	1	
Gwydyr Bay	.	1	2	8	3	9	3	2	2	5	3	4	.	5	21	
Jago Lagoon	.	.	1	1	5	6	9	21	.	1	6	13	.	1	2	
Beaufort Lagoon	1	1	.	.	.	1	.	.	.	
Subsis. Res. Area A	
Subsis. Res. Area B	2	1	1	1	1	.	.	.	1	1	.	.	1	1	.	
Subsis. Res. Area C	90	**	78	**	88	92	47	26	**	**	96	49	**	**	**	
Subsis. Res. Area D	10	20	32	67	88	**	**	**	22	56	**	**	5	33	**	
Fall Feeding Area	
Summer Feed. Area 1	3	4	.	.	.	4	.	.	.	
Summer Feed. Area 2	
Southern SLS Area	
Northern SLS Area	1	1	1	1	1	1	1	.	1	1	1	1	1	1	1	
Northern SLS	
Boundary Segment 2	1	1	1	.	.	1	1	.	.	.	

Notes: ** = Greater than 99.5%; . = Less than 0.5%; SLS = spring lead system.
Boundary Segments with all values less than 0.5% are not shown.

Table 6. Annual conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location will contact a certain land segment within 3 days, Beaufort Sea OCS Lease Sale 170

Land Segment	Hypothetical Spill Location															
	L1	L2	L3	L4	L5	L6	L7	L8	P1	P2	P3	P4	P5	P6	P7	
32	1	1	2	.	.	
33	1	2	1	.	.	.	5	.	.	
34	1	3	1	2	1	2	.	.	1	12	.	
35	.	.	.	3	.	1	.	.	.	1	.	.	.	1	1	
36	.	.	.	3	1	1	1	2	
37	.	.	.	1	1	2	1	1	.	.	5	
38	2	1	4	.	.	1	2	.	.	1	
39	4	.	.	.	1	.	.	.	
40	1	8	.	.	.	1	.	.	.	
41	1	2	.	.	.	1	.	.	.	

Notes: ** = Greater than 99.5%; . = Less than 0.5%; SLS = spring lead system.
Boundary Segments with all values less than 0.5% are not shown.

Table 7. Annual conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location will contact a certain land segment within 10 days, Beaufort Sea OCS Lease Sale 170

Land Segment	Hypothetical Spill Location															
	L1	L2	L3	L4	L5	L6	L7	L8	P1	P2	P3	P4	P5	P6	P7	
29	1	1	1	.	.	
30	1	1	1	.	.	.	1	.	.	
31	1	1	.	.	
32	1	2	1	.	.	.	3	.	.	
33	1	3	1	2	1	.	.	6	.	.	
34	1	3	2	3	1	.	.	.	2	3	1	.	2	14	.	
35	1	1	1	4	1	1	.	.	1	2	1	.	.	2	1	
36	.	.	1	4	1	2	1	.	1	1	1	.	.	2	3	
37	.	.	1	1	1	3	1	1	.	1	2	2	.	1	6	
38	1	3	2	6	.	.	2	3	.	.	2	
39	1	1	5	.	.	1	1	.	.	.	
40	1	1	1	9	.	.	1	2	.	.	1	
41	1	1	2	4	.	.	.	2	.	.	.	

Notes: ** = Greater than 99.5%; . = Less than 0.5%; SLS = spring lead system.
Boundary Segments with all values less than 0.5% are not shown.

Table 8. Annual conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location will contact a certain land segment within 30 days, Beaufort Sea OCS Lease Sale 170

Land Segment	Hypothetical Spill Location														
	L1	L2	L3	L4	L5	L6	L7	L8	P1	P2	P3	P4	P5	P6	P7
28	1	1	1	.	.
29	2	1	1	.	.	.	1	.	.
30	1	1	1	.	.	.	2	.	.
31	1	1	1	.	.
32	1	2	1	1	.	.	.	3	.	.
33	2	4	2	1	2	1	.	.	7	.	.
34	2	5	3	4	1	1	.	.	2	5	1	.	2	16	1
35	1	2	1	6	2	2	1	.	1	2	3	.	2	2	2
36	1	1	1	4	2	3	1	.	2	1	2	1	1	2	4
37	.	.	1	2	1	4	2	1	.	2	2	3	.	2	8
38	1	1	1	1	2	5	2	9	1	1	3	4	.	.	3
39	1	1	1	6	.	.	1	1	.	.	1
40	1	2	2	11	.	.	2	3	.	.	1
41	1	1	3	5	.	1	1	3	.	.	1
42	1	1

Notes: ** = Greater than 99.5%; . = Less than 0.5%; SLS = spring lead system.
Boundary Segments with all values less than 0.5% are not shown.

Table 9. Annual conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location will contact a certain land segment within 180 days, Beaufort Sea OCS Lease Sale 170

Land Segment	Hypothetical Spill Location														
	L1	L2	L3	L4	L5	L6	L7	L8	P1	P2	P3	P4	P5	P6	P7
23	1
27	1	1	1	1	.
28	3	3	1	1	1	.	.	.	2	1	.	.	2	2	.
29	8	6	3	1	1	.	.	.	9	2	.	.	8	1	.
30	2	3	1	2	1	.	.	7	.	.
31	1	1	2	.	.
32	1	4	2	1	2	1	.	.	5	2	.
33	2	8	4	3	2	1	.	.	3	4	1	.	10	3	1
34	2	8	4	11	5	4	2	.	3	7	4	1	4	27	3
35	2	3	1	10	5	7	3	1	1	2	7	3	7	3	7
36	1	4	1	5	2	8	4	3	3	1	5	4	1	3	11
37	.	.	2	6	2	6	3	4	.	4	3	7	.	6	14
38	1	1	1	2	2	9	3	16	1	2	3	4	.	.	5
39	2	4	1	6	.	.	2	1	.	.	7
40	4	4	3	15	.	.	5	6	.	.	1
41	1	1	6	8	.	1	1	7	.	.	1
42	2	1	.	.	.	1	.	.	.
43	1	1

Notes: ** = Greater than 99.5%; . = Less than 0.5%; SLS = spring lead system.
Boundary Segments with all values less than 0.5% are not shown.

Table 10. Annual combined probabilities (expressed as percent chance) of one or more spills greater than or equal to 1,000 barrels, and the estimated number of spills (mean), occurring and contacting a certain environmental resource area over the assumed production life of Beaufort Sea OCS Lease Sale 170

Environmental Resource Area	Within 3 days		Within 10 days		Within 30 days		Within 180 days	
	Prob	Mean	Prob	Mean	Prob	Mean	Prob	Mean
Land	5- 9	0.1- 0.1	8-15	0.1- 0.2	12-22	0.1- 0.2	23-40	0.3- 0.5
Ice/Sea Segment 1	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0
Ice/Sea Segment 2	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0
Ice/Sea Segment 3	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0
Ice/Sea Segment 4	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	. 1	0.0- 0.0
Ice/Sea Segment 5	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	. 1	0.0- 0.0
Ice/Sea Segment 6	. . .	0.0- 0.0	. 1	0.0- 0.0	1- 2	0.0- 0.0	2- 4	0.0- 0.0
Ice/Sea Segment 7	1- 1	0.0- 0.0	2- 4	0.0- 0.0	3- 6	0.0- 0.1	7-14	0.1- 0.2
Ice/Sea Segment 8	3- 5	0.0- 0.1	4- 9	0.0- 0.1	6-12	0.1- 0.1	14-26	0.2- 0.3
Ice/Sea Segment 9	5- 9	0.1- 0.1	7-12	0.1- 0.1	8-15	0.1- 0.2	15-25	0.2- 0.3
Ice/Sea Segment 10	. . .	0.0- 0.0	1- 2	0.0- 0.0	2- 4	0.0- 0.0	6-11	0.1- 0.1
Ice/Sea Segment 11	. . .	0.0- 0.0	. 1	0.0- 0.0	1- 1	0.0- 0.0	3- 5	0.0- 0.0
Ice/Sea Segment 12	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	1- 2	0.0- 0.0
Ice/Sea Segment 13	. . .	0.0- 0.0	. . .	0.0- 0.0	. 1	0.0- 0.0	. 1	0.0- 0.0
Ice/Sea Segment 14	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0
Ice/Sea Segment 15	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0
Ice/Sea Segment 16	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0
Ice/Sea Segment 17	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0
Ice/Sea Seg. 1 SLS*	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0
Ice/Sea Seg. 2 SLS*	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0
Ice/Sea Seg. 3 SLS*	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0
Ice/Sea Seg. 4 SLS*	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0
Peard Bay#	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0
Elson Lagoon***	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0
Simpson Lagoon***	1- 3	0.0- 0.0	2- 3	0.0- 0.0	2- 4	0.0- 0.0	4- 8	0.0- 0.1
Gwydyr Bay***	2- 3	0.0- 0.0	2- 3	0.0- 0.0	2- 4	0.0- 0.0	4- 8	0.0- 0.1
Jago Lagoon***	. 1	0.0- 0.0	1- 2	0.0- 0.0	1- 2	0.0- 0.0	2- 4	0.0- 0.0
Beaufort Lagoon	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0
Subsis. Res. Area A	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0
Subsis. Res. Area B	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	. 1	0.0- 0.0
Subsis. Res. Area C	39-62	0.5- 1.0	39-62	0.5- 1.0	40-63	0.5- 1.0	42-65	0.5- 1.1
Subsis. Res. Area D	30-47	0.4- 0.6	31-48	0.4- 0.7	31-49	0.4- 0.7	34-53	0.4- 0.8
Fall Feeding Area+	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0
Summer Feed. Area 1++	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	. 1	0.0- 0.0
Summer Feed. Area 2++	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0
Southern SLS Area	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0
Northern SLS Area	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	. 1	0.0- 0.0
Northern SLS*	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0	. . .	0.0- 0.0

Note:

. = than 0.5%; SLS = Spring lead system.

Boundary Segments with values less than 0.5% probability of contacts within 180 days are not shown.

Environmental Resources are vulnerable year-round unless noted as shown below:

* May-June # June-October *** July-September + September-October ++ June-September

Table 11. Annual combined probabilities (expressed as percent chance) of one or more spills greater than or equal to 1,000 barrels, and the estimated number of spills (mean), occurring and contacting a certain land segment over the assumed production life of Beaufort Sea OCS Lease Sale 170

Land Segment	Within 3 days		Within 10 days		Within 30 days		Within 180 days	
	Prob	Mean	Prob	Mean	Prob	Mean	Prob	Mean
28	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	1- 1	0.0- 0.0
29	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	1- 2	0.0- 0.0
30	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	1- 1	0.0- 0.0
31	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0
32	.- .	0.0- 0.0	.- 1	0.0- 0.0	.- 1	0.0- 0.0	1- 2	0.0- 0.0
33	.- 1	0.0- 0.0	1- 1	0.0- 0.0	1- 1	0.0- 0.0	2- 3	0.0- 0.0
34	2- 3	0.0- 0.0	2- 4	0.0- 0.0	3- 5	0.0- 0.1	5-10	0.1- 0.1
35	.- 1	0.0- 0.0	1- 1	0.0- 0.0	1- 2	0.0- 0.0	3- 6	0.0- 0.1
36	1- 1	0.0- 0.0	1- 2	0.0- 0.0	1- 2	0.0- 0.0	3- 5	0.0- 0.1
37	1- 1	0.0- 0.0	1- 2	0.0- 0.0	2- 3	0.0- 0.0	4- 6	0.0- 0.1
38	.- 1	0.0- 0.0	1- 1	0.0- 0.0	1- 2	0.0- 0.0	2- 3	0.0- 0.0
39	.- .	0.0- 0.0	.- 1	0.0- 0.0	.- 1	0.0- 0.0	1- 2	0.0- 0.0
40	.- 1	0.0- 0.0	1- 1	0.0- 0.0	1- 1	0.0- 0.0	1- 3	0.0- 0.0
41	.- .	0.0- 0.0	.- 1	0.0- 0.0	1- 1	0.0- 0.0	1- 2	0.0- 0.0

Note: . = Less than 0.5%.
Segments with all values less than 0.5% are not shown.

Table 12. Annual combined probabilities (expressed as percent chance) of one or more spills greater than or equal to 1,000 barrels, and the estimated number of spills (mean), occurring and contacting a certain environmental resource area over the assumed production life of Beaufort Sea OCS Lease Sale 170, Kaktovik Deferral

Environmental Resource Area	Within 3 days		Within 10 days		Within 30 days		Within 180 days	
	Prob	Mean	Prob	Mean	Prob	Mean	Prob	Mean
US Land	4- 7	0.0- 0.1	6-11	0.1- 0.1	8-16	0.1- 0.2	17-31	0.2- 0.4
Ice/Sea Segment 1	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Segment 2	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Segment 3	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Segment 4	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Segment 5	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0
Ice/Sea Segment 6	.- .	0.0- 0.0	.- 1	0.0- 0.0	1- 2	0.0- 0.0	2- 4	0.0- 0.0
Ice/Sea Segment 7	1- 1	0.0- 0.0	2- 3	0.0- 0.0	3- 6	0.0- 0.1	7-14	0.1- 0.1
Ice/Sea Segment 8	2- 5	0.0- 0.0	4- 8	0.0- 0.1	5-10	0.1- 0.1	12-22	0.1- 0.3
Ice/Sea Segment 9	2- 4	0.0- 0.0	3- 5	0.0- 0.1	4- 7	0.0- 0.1	7-12	0.1- 0.1
Ice/Sea Segment 10	.- .	0.0- 0.0	1- 1	0.0- 0.0	1- 2	0.0- 0.0	3- 6	0.0- 0.1
Ice/Sea Segment 11	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0	1- 2	0.0- 0.0
Ice/Sea Segment 12	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0
Ice/Sea Segment 13	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Segment 14	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Segment 15	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Segment 16	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Segment 17	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Seg. 1 SLS*	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Seg. 2 SLS*	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Seg. 3 SLS*	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Seg. 4 SLS*	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Peard Bay#	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Elson Lagoon***	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Simpson Lagoon***	1- 3	0.0- 0.0	2- 3	0.0- 0.0	2- 4	0.0- 0.0	4- 8	0.0- 0.1
Gwydyr Bay***	1- 2	0.0- 0.0	1- 2	0.0- 0.0	2- 3	0.0- 0.0	3- 5	0.0- 0.1
Jago Lagoon***	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0	1- 1	0.0- 0.0
Beaufort Lagoon	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Subsis. Res. Area A	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Subsis. Res. Area B	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0
Subsis. Res. Area C	33-55	0.4- 0.8	33-55	0.4- 0.8	33-56	0.4- 0.8	34-56	0.4- 0.8
Subsis. Res. Area D	15-26	0.2- 0.3	16-27	0.2- 0.3	17-29	0.2- 0.3	20-34	0.2- 0.4
Fall Feeding Area+	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Summer Feed. Area 1++	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Summer Feed. Area 2++	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Southern SLS Area	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Northern SLS Area*	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0
Northern SLS	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0

Note:

. = than 0.5%; SLS = Spring lead system.

Boundary Segments with values less than 0.5% probability of contacts within 180 days are not shown.

Environmental Resources are vulnerable year-round unless noted as shown below:

* May-June # June-October *** July-September + September-October ++ June-September

Table 13. Annual combined probabilities (expressed as percent chance) of one or more spills greater than or equal to 1,000 barrels, and the estimated number of spills (mean), occurring and contacting a certain land segment over the assumed production life of Beaufort Sea OCS Lease Sale 170, Kaktovik Deferral

Land Segment	Within 3 days		Within 10 days		Within 30 days		Within 180 days	
	Prob	Mean	Prob	Mean	Prob	Mean	Prob	Mean
28	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	1- 1	0.0- 0.0
29	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	1- 2	0.0- 0.0
30	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	1- 1	0.0- 0.0
31	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0
32	.- .	0.0- 0.0	.- 1	0.0- 0.0	.- 1	0.0- 0.0	1- 2	0.0- 0.0
33	.- 1	0.0- 0.0	1- 1	0.0- 0.0	1- 1	0.0- 0.0	1- 3	0.0- 0.0
34	2- 3	0.0- 0.0	2- 4	0.0- 0.0	3- 5	0.0- 0.1	5- 9	0.0- 0.1
35	.- 1	0.0- 0.0	1- 1	0.0- 0.0	1- 2	0.0- 0.0	2- 4	0.0- 0.0
36	.- 1	0.0- 0.0	1- 1	0.0- 0.0	1- 2	0.0- 0.0	2- 3	0.0- 0.0
37	.- 1	0.0- 0.0	1- 1	0.0- 0.0	1- 2	0.0- 0.0	2- 4	0.0- 0.0
38	.- .	0.0- 0.0	.- 1	0.0- 0.0	1- 1	0.0- 0.0	1- 2	0.0- 0.0
39	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	1- 1	0.0- 0.0
40	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0

Note: . = Less than 0.5%.
 Segments with all values less than 0.5% are not shown.

Table 14. Annual combined probabilities (expressed as percent chance) of one or more spills greater than or equal to 1,000 barrels, and the estimated number of spills (mean), occurring and contacting a certain environmental resource area over the assumed production life of Beaufort Sea OCS Lease Sale 170, Area Offshore of the ANWR

Environmental Resource Area	Within 3 days		Within 10 days		Within 30 days		Within 180 days	
	Prob	Mean	Prob	Mean	Prob	Mean	Prob	Mean
US Land	3- 7	0.0- 0.1	5-11	0.1- 0.1	7-15	0.1- 0.2	15-29	0.2- 0.3
Ice/Sea Segment 1	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Segment 2	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Segment 3	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Segment 4	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Segment 5	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0
Ice/Sea Segment 6	.- .	0.0- 0.0	.- 1	0.0- 0.0	1- 1	0.0- 0.0	2- 4	0.0- 0.0
Ice/Sea Segment 7	1- 1	0.0- 0.0	2- 3	0.0- 0.0	3- 6	0.0- 0.1	7-14	0.1- 0.1
Ice/Sea Segment 8	2- 5	0.0- 0.0	4- 7	0.0- 0.1	5-10	0.1- 0.1	11-22	0.1- 0.2
Ice/Sea Segment 9	1- 3	0.0- 0.0	2- 4	0.0- 0.0	3- 6	0.0- 0.1	4-10	0.0- 0.1
Ice/Sea Segment 10	.- .	0.0- 0.0	.- 1	0.0- 0.0	1-2	0.0- 0.0	2- 5	0.0- 0.1
Ice/Sea Segment 11	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0	1- 2	0.0- 0.0
Ice/Sea Segment 12	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0
Ice/Sea Segment 13	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Segment 14	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Segment 15	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Segment 16	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Segment 17	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Seg. 1 SLS*	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Seg. 2 SLS*	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Seg. 3 SLS*	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Seg. 4 SLS*	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Peard Bay#	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Elson Lagoon***	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Simpson Lagoon***	1- 3	0.0- 0.0	2- 3	0.0- 0.0	2- 4	0.0- 0.0	4- 8	0.0- 0.1
Gwydyr Bay***	1- 1	0.0- 0.0	1- 2	0.0- 0.0	1- 2	0.0- 0.0	2- 5	0.0- 0.0
Jago Lagoon***	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0	.- 1	0.0- 0.0
Beaufort Lagoon***	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Subsis. Res. Area A	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Subsis. Res. Area B	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0
Subsis. Res. Area C	30-53	0.4- 0.8	30-53	0.4- 0.8	30-54	0.4- 0.8	31-54	0.4- 0.8
Subsis. Res. Area D	11-22	0.1- 0.3	11-23	0.1- 0.3	12-25	0.1- 0.3	16-30	0.2- 0.4
Fall Feeding Area+	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Summer Feed. Area 1++	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Summer Feed. Area 2++	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Southern SLS Area	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Northern SLS Area	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0
Northern SLS*	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0

Note:

ANWR = Alaska National Wildlife Refuge.

. = than 0.5%; SLS = Spring lead system.

Boundary Segments with values less than 0.5% probability of contacts within 180 days are not shown.

Environmental Resources are vulnerable year-round unless noted as shown below:

* May-June # June-October *** July-September + September-October ++ June-September

Table 15. Annual combined probabilities (expressed as percent chance) of one or more spills greater than or equal to 1,000 barrels, and the estimated number of spills (mean), occurring and contacting a certain land segment over the assumed production life of Beaufort Sea OCS Lease Sale 170, Area Offshore of the ANWR

Land Segment	Within 3 days		Within 10 days		Within 30 days		Within 180 days	
	Prob	Mean	Prob	Mean	Prob	Mean	Prob	Mean
28	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	1- 1	0.0- 0.0
29	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	1- 2	0.0- 0.0
30	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	1- 1	0.0- 0.0
31	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0
32	.- .	0.0- 0.0	.- 1	0.0- 0.0	.- 1	0.0- 0.0	1- 2	0.0- 0.0
33	.- 1	0.0- 0.0	1- 1	0.0- 0.0	1- 1	0.0- 0.0	1- 3	0.0- 0.0
34	2- 3	0.0- 0.0	2- 4	0.0- 0.0	3- 5	0.0- 0.1	4- 9	0.0- 0.1
35	.- 1	0.0- 0.0	1- 1	0.0- 0.0	1- 2	0.0- 0.0	2- 4	0.0- 0.0
36	.- 1	0.0- 0.0	1- 1	0.0- 0.0	1- 2	0.0- 0.0	1- 3	0.0- 0.0
37	.- 1	0.0- 0.0	1- 1	0.0- 0.0	1- 2	0.0- 0.0	2- 4	0.0- 0.0
38	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0	1- 1	0.0- 0.0
39	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0
40	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0

Note: ANWR = Alaska National Wildlife Refuge.

. = Less than 0.5%.

Segments with all values less than 0.5% are not shown.

Table 16. Annual combined probabilities (expressed as percent chance) of one or more spills greater than or equal to 1,000 barrels, and the estimated number of spills (mean), occurring and contacting a certain environmental resource area over the assumed production life of Beaufort Sea OCS Lease Sale 170, Cross Island Area

Environmental Resource Area	Within 3 days		Within 10 days		Within 30 days		Within 180 days	
	Prob	Mean	Prob	Mean	Prob	Mean	Prob	Mean
US Land	4- 7	0.0- 0.1	6-12	0.1- 0.1	10-18	0.1- 0.2	19-34	0.2- 0.4
Ice/Sea Segment 1	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Segment 2	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Segment 3	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Segment 4	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Segment 5	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0
Ice/Sea Segment 6	.- .	0.0- 0.0	.- 1	0.0- 0.0	1- 1	0.0- 0.0	2- 4	0.0- 0.0
Ice/Sea Segment 7	1- 1	0.0- 0.0	1- 3	0.0- 0.0	2- 5	0.0- 0.1	5-11	0.1- 0.1
Ice/Sea Segment 8	2- 3	0.0- 0.0	3- 6	0.0- 0.1	4- 9	0.0- 0.1	10-20	0.1- 0.2
Ice/Sea Segment 9	5- 9	0.1- 0.1	7-11	0.1- 0.1	8-14	0.1- 0.1	15-25	0.2- 0.3
Ice/Sea Segment 10	.- .	0.0- 0.0	1- 2	0.0- 0.0	2- 4	0.0- 0.0	6-10	0.1- 0.1
Ice/Sea Segment 11	.- .	0.0- 0.0	.- 1	0.0- 0.0	1- 1	0.0- 0.0	3- 5	0.0- 0.0
Ice/Sea Segment 12	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	1- 2	0.0- 0.0
Ice/Sea Segment 13	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0	.- 1	0.0- 0.0
Ice/Sea Segment 14	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Segment 15	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Segment 16	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Segment 17	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Seg. 1 SLS*	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Seg. 2 SLS*	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Seg. 3 SLS*	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Ice/Sea Seg. 4 SLS*	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Peard Bay#	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Elson Lagoon***	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Simpson Lagoon***	1- 2	0.0- 0.0	1- 3	0.0- 0.0	1- 3	0.0- 0.0	3- 6	0.0- 0.1
Gwydyr Bay***	2- 3	0.0- 0.0	2- 3	0.0- 0.0	2- 4	0.0- 0.0	4- 6	0.0- 0.1
Jago Lagoon***	.- 1	0.0- 0.0	1- 1	0.0- 0.0	1- 2	0.0- 0.0	2- 4	0.0- 0.0
Beaufort Lagoon***	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Subsis. Res. Area A	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Subsis. Res. Area B	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0
Subsis. Res. Area C	31-53	0.4- 0.8	31-54	0.4- 0.8	32-55	0.4- 0.8	35-57	0.4- 0.9
Subsis. Res. Area D	29-45	0.3- 0.6	29-46	0.3- 0.6	29-46	0.3- 0.6	31-49	0.4- 0.7
Fall Feeding Area+	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Summer Feed. Area 1++	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0
Summer Feed. Area 2++	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Southern SLS Area	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0
Northern SLS Area	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0
Northern SLS*	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0

Note:

. = than 0.5%; SLS = Spring lead system.

Boundary Segments with values less than 0.5% probability of contacts within 180 days are not shown.

Environmental Resources are vulnerable year-round unless noted as shown below:

* May-June # June-October *** July-September + September-October ++ June-September

Table 17. Annual combined probabilities (expressed as percent chance) of one or more spills greater than or equal to 1,000 barrels, and the estimated number of spills (mean), occurring and contacting a certain land segment over the assumed production life of Beaufort Sea OCS Lease Sale 170, Cross Island Area

Land Segment	Within 3 days		Within 10 days		Within 30 days		Within 180 days	
	Prob	Mean	Prob	Mean	Prob	Mean	Prob	Mean
28	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	.- 1	0.0- 0.0
29	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	1- 2	0.0- 0.0
30	.- .	0.0- 0.0	.- .	0.0- 0.0	.- .	0.0- 0.0	1- 1	0.0- 0.0
32	.- .	0.0- 0.0	.- 1	0.0- 0.0	.- 1	0.0- 0.0	1- 1	0.0- 0.0
33	.- 1	0.0- 0.0	.- 1	0.0- 0.0	1- 1	0.0- 0.0	1- 3	0.0- 0.0
34	1- 1	0.0- 0.0	1- 2	0.0- 0.0	1- 3	0.0- 0.0	2- 5	0.0- 0.1
35	.- .	0.0- 0.0	.- 1	0.0- 0.0	1- 2	0.0- 0.0	2- 5	0.0- 0.0
36	.- 1	0.0- 0.0	1- 1	0.0- 0.0	1- 2	0.0- 0.0	2- 5	0.0- 0.0
37	1- 1	0.0- 0.0	1- 2	0.0- 0.0	2- 3	0.0- 0.0	3- 5	0.0- 0.1
38	.- 1	0.0- 0.0	1- 1	0.0- 0.0	1- 2	0.0- 0.0	2- 3	0.0- 0.0
39	.- .	0.0- 0.0	.- 1	0.0- 0.0	.- 1	0.0- 0.0	1- 2	0.0- 0.0
40	.- 1	0.0- 0.0	1- 1	0.0- 0.0	1- 1	0.0- 0.0	1- 2	0.0- 0.0
41	.- .	0.0- 0.0	.- 1	0.0- 0.0	1- 1	0.0- 0.0	1- 2	0.0- 0.0

Note: ANWR = Alaska National Wildlife Refuge.
 . = Less than 0.5%.
 Segments with all values less than 0.5% are not shown.

Appendix A

Environmental Resource Areas




Figure A-1. Locations of Ice/Sea Segments 1-17 for proposed Beaufort Sea OCS Lease Sale 170.




Figure A-2. Locations of Peard Bay, Elson Lagoon, Simpson Lagoon, Gwydyr Bay, Jago Lagoon, and Beaufort Lagoon for proposed Beaufort Sea OCS Lease Sale 170.




Figure A-3. Locations of Subsistence Resource Area A, Fall Feeding Area, Subsistence Resource Area C, and Summer Feeding Area 1 for proposed Beaufort Sea OCS Lease Sale 170




Figure A-4. Locations of Subsistence Resource Area B, Subsistence Resource Area D, and Summer Feeding Area 2 for proposed Beaufort Sea OCS Lease Sale 170.




Figure A-5. Location of Northern Spring Lead System (SLS) Area for proposed Beaufort Sea OCS Lease Sale 170.




Figure A-6. Location of Southern Spring Lead System (SLS) Area for proposed Beaufort Sea OCS Lease Sale 170.




Figure A-7. Locations of Boundary Segments (B1-B38) for proposed Beaufort Sea OCS Lease Sale 170.

Appendix B

Seasonal Conditional Probabilities of Contact to Environmental Resource Areas

Table B-1. Seasonal conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location in the winter season will contact a certain environmental resource area within 3 days, Beaufort Sea OCS Lease Sale 170

Environmental Resource Area	Hypothetical Spill Location														
	L1	L2	L3	L4	L5	L6	L7	L8	P1	P2	P3	P4	P5	P6	P7
Land	2	5	2	7	2	6	2	17	3	3	2	4	9	14	6
Ice/Sea Segment 1
Ice/Sea Segment 2
Ice/Sea Segment 3
Ice/Sea Segment 4
Ice/Sea Segment 5
Ice/Sea Segment 6
Ice/Sea Segment 7	4	1	1	2	1	.	.	2	.	.
Ice/Sea Segment 8	4	1	47	2	5	1	.	.	1	6	1	.	.	2	1
Ice/Sea Segment 9	37	12	57	2	.	.	10	5	.	.	2
Ice/Sea Segment 10	1
Ice/Sea Segment 11
Ice/Sea Segment 12
Ice/Sea Segment 13
Ice/Sea Segment 14
Ice/Sea Segment 15
Ice/Sea Segment 16
Ice/Sea Segment 17
Ice/Sea Seg. 1 SLS
Ice/Sea Seg. 2 SLS
Ice/Sea Seg. 3 SLS
Ice/Sea Seg. 4 SLS
Peard Bay
Elson Lagoon
Simpson Lagoon
Gwydyr Bay
Jago Lagoon
Beaufort Lagoon
Subsis. Res. Area A
Subsis. Res. Area B
Subsis. Res. Area C	80	**	50	**	70	83	6	2	**	**	86	5	**	**	**
Subsis. Res. Area D	.	.	16	44	77	**	**	**	.	34	**	**	.	2	**
Fall Feeding Area
Summer Feed. Area 1
Summer Feed. Area 2
Southern SLS Area
Northern SLS Area
Northern SLS

Notes: ** = Greater than 99.5%; . = Less than 0.5%; SLS = spring lead system.
Boundary Segments with all values less than 0.5% are not shown

Table B-2. Seasonal conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location in the winter season will contact a certain environmental resource area within 10 days, Beaufort Sea OCS Lease Sale 170

Environmental Resource Area	Hypothetical Spill Location														
	L1	L2	L3	L4	L5	L6	L7	L8	P1	P2	P3	P4	P5	P6	P7
Land	6	9	5	11	5	9	6	22	6	6	6	7	12	16	10
Ice/Sea Segment 1
Ice/Sea Segment 2
Ice/Sea Segment 3
Ice/Sea Segment 4
Ice/Sea Segment 5
Ice/Sea Segment 6	1	1	1	1	1	.	.	1	.	.
Ice/Sea Segment 7	8	3	3	1	1	.	.	.	5	2	1	.	3	1	1
Ice/Sea Segment 8	5	2	51	4	8	2	1	1	2	11	3	1	1	4	2
Ice/Sea Segment 9	.	.	1	1	38	13	61	4	.	1	13	11	.	.	4
Ice/Sea Segment 10	1	.	1	1	.	.	.
Ice/Sea Segment 11	1
Ice/Sea Segment 12
Ice/Sea Segment 13
Ice/Sea Segment 14
Ice/Sea Segment 15
Ice/Sea Segment 16
Ice/Sea Segment 17
Ice/Sea Seg. 1 SLS
Ice/Sea Seg. 2 SLS
Ice/Sea Seg. 3 SLS
Ice/Sea Seg. 4 SLS
Peard Bay
Elson Lagoon
Simpson Lagoon	1	.	.
Gwydyr Bay	1
Jago Lagoon
Beaufort Lagoon
Subsis. Res. Area A
Subsis. Res. Area B
Subsis. Res. Area C	83	**	58	**	75	85	16	4	**	**	89	9	**	**	**
Subsis. Res. Area D	1	1	17	45	81	**	**	**	1	35	**	**	.	3	**
Fall Feeding Area
Summer Feed. Area 1
Summer Feed. Area 2
Southern SLS Area
Northern SLS Area
Northern SLS

Notes: ** = Greater than 99.5%; . = Less than 0.5%; SLS = spring lead system.
Boundary Segments with all values less than 0.5% are not shown

Table B-3. Seasonal conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location in the winter season will contact a certain environmental resource area within 30 days, Beaufort Sea OCS Lease Sale 170

Environmental Resource Area	Hypothetical Spill Location														
	L1	L2	L3	L4	L5	L6	L7	L8	P1	P2	P3	P4	P5	P6	P7
Land	8	13	6	14	7	13	7	27	7	7	7	10	14	17	13
Ice/Sea Segment 1
Ice/Sea Segment 2
Ice/Sea Segment 3
Ice/Sea Segment 4
Ice/Sea Segment 5	1	1
Ice/Sea Segment 6	3	1	1	1	1	.	.	2	1	.
Ice/Sea Segment 7	12	4	6	2	2	1	1	.	8	4	1	.	4	2	1
Ice/Sea Segment 8	5	2	54	5	11	2	3	1	3	16	5	1	1	5	2
Ice/Sea Segment 9	.	.	1	1	39	13	66	5	.	1	15	16	.	.	4
Ice/Sea Segment 10	1	1	2	1	.	.	1	1	.	.	.
Ice/Sea Segment 11	1	1	.	.	.
Ice/Sea Segment 12
Ice/Sea Segment 13
Ice/Sea Segment 14
Ice/Sea Segment 15
Ice/Sea Segment 16
Ice/Sea Segment 17
Ice/Sea Seg. 1 SLS
Ice/Sea Seg. 2 SLS
Ice/Sea Seg. 3 SLS
Ice/Sea Seg. 4 SLS
Peard Bay
Elson Lagoon
Simpson Lagoon	3	.	.
Gwydyr Bay	4
Jago Lagoon	1
Beaufort Lagoon
Subsis. Res. Area A
Subsis. Res. Area B
Subsis. Res. Area C	87	**	68	**	81	88	29	7	**	**	92	20	**	**	**
Subsis. Res. Area D	1	1	18	46	83	**	**	**	1	36	**	**	1	3	**
Fall Feeding Area
Summer Feed. Area 1
Summer Feed. Area 2
Southern SLS Area
Northern SLS Area
Northern SLS

Notes: ** = Greater than 99.5%; . = Less than 0.5%; SLS = spring lead system.
Boundary Segments with all values less than 0.5% are not shown

Table B-4. Seasonal conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location in the winter season will contact a certain environmental resource area within 180 days, Beaufort Sea OCS Lease Sale 170

Environmental Resource Area	Hypothetical Spill Location														
	L1	L2	L3	L4	L5	L6	L7	L8	P1	P2	P3	P4	P5	P6	P7
Land	24	44	20	43	25	45	25	56	25	26	31	33	50	50	49
Ice/Sea Segment 1
Ice/Sea Segment 2
Ice/Sea Segment 3	1
Ice/Sea Segment 4	1	.	1	.	.	.	1	1	.	.	.
Ice/Sea Segment 5	2	1	1	.	1	1	1	1	1	1	1	1	1	.	.
Ice/Sea Segment 6	17	10	5	1	2	1	1	1	8	2	1	1	13	1	1
Ice/Sea Segment 7	45	24	25	14	3	1	1	1	33	19	2	1	23	15	2
Ice/Sea Segment 8	21	22	78	33	36	16	9	4	29	57	31	7	16	28	14
Ice/Sea Segment 9	2	2	3	6	50	37	88	39	4	5	42	63	1	1	26
Ice/Sea Segment 10	4	3	8	5	12	9	27	14	6	7	11	18	1	3	6
Ice/Sea Segment 11	.	.	3	1	6	6	10	6	3	4	5	8	.	.	4
Ice/Sea Segment 12	2	3	5	4	.	.	2	5	.	.	2
Ice/Sea Segment 13	1	.	.	.	1	.	.	.
Ice/Sea Segment 14
Ice/Sea Segment 15
Ice/Sea Segment 16
Ice/Sea Segment 17
Ice/Sea Seg. 1 SLS
Ice/Sea Seg. 2 SLS
Ice/Sea Seg. 3 SLS
Ice/Sea Seg. 4 SLS
Peard Bay
Elson Lagoon
Simpson Lagoon	4	10	6	7	3	1	.	.	6	9	1	.	9	10	1
Gwydyr Bay	.	.	1	6	2	5	2	2	1	3	3	4	.	5	11
Jago Lagoon	3	4	6	11	.	.	5	9	.	.	.
Beaufort Lagoon	1
Subsis. Res. Area A
Subsis. Res. Area B	2	1	1	.	1	.	.	.	1	1	1	.	1	.	.
Subsis. Res. Area C	91	**	81	**	92	94	56	32	**	**	98	59	**	**	**
Subsis. Res. Area D	6	16	28	63	88	**	**	**	17	50	**	**	2	28	**
Fall Feeding Area
Summer Feed. Area 1	3	3	.	.	.	3	.	.	.
Summer Feed. Area 2
Southern SLS Area
Northern SLS Area	1	.	1	.	1	1	1	.	1	.	1	1	.	.	.
Northern SLS

Notes: ** = Greater than 99.5%; . = Less than 0.5%; SLS = spring lead system.
Boundary Segments with all values less than 0.5% are not shown

Table B-5. Seasonal conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location in the summer season will contact a certain environmental resource area within 3 days, Beaufort Sea OCS Lease Sale 170

Environmental Resource Area	Hypothetical Spill Location														
	L1	L2	L3	L4	L5	L6	L7	L8	P1	P2	P3	P4	P5	P6	P7
Land	4	10	3	12	4	11	6	25	5	5	6	9	12	19	13
Ice/Sea Segment 1
Ice/Sea Segment 2
Ice/Sea Segment 3
Ice/Sea Segment 4
Ice/Sea Segment 5
Ice/Sea Segment 6
Ice/Sea Segment 7	23	8	4	1	13	2	.	.	13	1	.
Ice/Sea Segment 8	12	10	63	15	13	3	.	.	14	29	8	.	5	16	2
Ice/Sea Segment 9	.	.	2	4	47	29	68	11	.	2	32	31	.	1	22
Ice/Sea Segment 10	1	.	5	1	.	.	1	2	.	.	1
Ice/Sea Segment 11	1	1
Ice/Sea Segment 12
Ice/Sea Segment 13
Ice/Sea Segment 14
Ice/Sea Segment 15
Ice/Sea Segment 16
Ice/Sea Segment 17
Ice/Sea Seg. 1 SLS
Ice/Sea Seg. 2 SLS
Ice/Sea Seg. 3 SLS
Ice/Sea Seg. 4 SLS
Peard Bay
Elson Lagoon
Simpson Lagoon	4	16	1	2	5	2	.	.	44	19	.
Gwydyr Bay	.	.	.	3	2	10	1	1	.	1	3	1	.	.	46
Jago Lagoon	1	2	6	30	.	.	1	7	.	.	.
Beaufort Lagoon
Subsis. Res. Area A
Subsis. Res. Area B
Subsis. Res. Area C	81	**	53	**	70	83	5	2	**	**	90	6	**	**	**
Subsis. Res. Area D	2	6	24	59	78	**	**	**	5	49	**	**	1	20	**
Fall Feeding Area
Summer Feed. Area 1
Summer Feed. Area 2
Southern SLS Area
Northern SLS Area
Northern SLS

Notes: ** = Greater than 99.5%; . = Less than 0.5%; SLS = spring lead system.
Boundary Segments with all values less than 0.5% are not shown

Table B-6. Seasonal conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location in the summer season will contact a certain environmental resource area within 10 days, Beaufort Sea OCS Lease Sale 170

Environmental Resource Area	Hypothetical Spill Location														
	L1	L2	L3	L4	L5	L6	L7	L8	P1	P2	P3	P4	P5	P6	P7
Land	13	21	13	25	15	26	17	38	15	17	19	22	23	33	30
Ice/Sea Segment 1
Ice/Sea Segment 2
Ice/Sea Segment 3
Ice/Sea Segment 4
Ice/Sea Segment 5	1
Ice/Sea Segment 6	6	3	1	3	1	.	.	4	.	.
Ice/Sea Segment 7	34	19	10	5	1	1	.	.	25	6	1	.	26	7	.
Ice/Sea Segment 8	21	20	70	26	21	11	3	1	25	42	15	2	15	28	11
Ice/Sea Segment 9	3	4	9	12	54	38	73	20	5	10	44	41	3	8	32
Ice/Sea Segment 10	1	2	3	5	10	9	18	9	1	3	12	13	1	4	7
Ice/Sea Segment 11	1	2	7	7	.	.	1	7	.	.	2
Ice/Sea Segment 12	1
Ice/Sea Segment 13
Ice/Sea Segment 14
Ice/Sea Segment 15
Ice/Sea Segment 16
Ice/Sea Segment 17
Ice/Sea Seg. 1 SLS
Ice/Sea Seg. 2 SLS
Ice/Sea Seg. 3 SLS
Ice/Sea Seg. 4 SLS
Peard Bay
Elson Lagoon
Simpson Lagoon	9	21	5	4	1	.	.	.	10	5	1	.	46	21	.
Gwydyr Bay	1	1	3	7	3	13	2	1	2	3	5	4	1	2	48
Jago Lagoon	.	.	1	2	5	7	11	36	.	1	5	15	.	1	5
Beaufort Lagoon	1	1	.	.	1	.	.	.
Subsis. Res. Area A
Subsis. Res. Area B
Subsis. Res. Area C	84	**	59	**	73	84	10	6	**	**	90	11	**	**	**
Subsis. Res. Area D	10	15	33	65	81	**	**	**	15	56	**	**	8	30	**
Fall Feeding Area
Summer Feed. Area 1
Summer Feed. Area 2
Southern SLS Area
Northern SLS Area
Northern SLS

Notes: ** = Greater than 99.5%; . = Less than 0.5%; SLS = spring lead system.
Boundary Segments with all values less than 0.5% are not shown

Table B-7. Seasonal conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location in the summer season will contact a certain environmental resource area within 30 days, Beaufort Sea OCS Lease Sale 170

Environmental Resource Area	Hypothetical Spill Location														
	L1	L2	L3	L4	L5	L6	L7	L8	P1	P2	P3	P4	P5	P6	P7
Land	31	40	30	42	36	46	38	59	32	34	40	45	44	50	53
Ice/Sea Segment 1
Ice/Sea Segment 2
Ice/Sea Segment 3
Ice/Sea Segment 4
Ice/Sea Segment 5	2	1	1	1	.	.	.	1	.	.
Ice/Sea Segment 6	15	11	4	2	2	1	1	.	7	3	1	.	14	1	1
Ice/Sea Segment 7	51	33	21	15	4	3	2	1	39	16	4	2	39	18	2
Ice/Sea Segment 8	33	31	82	41	33	21	5	4	42	57	30	4	23	39	20
Ice/Sea Segment 9	8	7	14	18	62	50	81	29	9	13	62	53	5	12	41
Ice/Sea Segment 10	3	4	9	10	18	17	35	19	4	8	19	28	3	8	12
Ice/Sea Segment 11	.	1	2	2	6	6	13	12	.	3	6	13	1	2	6
Ice/Sea Segment 12	1	2	3	3	.	.	1	3	.	.	2
Ice/Sea Segment 13	.	.	.	2	3	4	6	5	.	1	3	6	.	1	4
Ice/Sea Segment 14
Ice/Sea Segment 15
Ice/Sea Segment 16
Ice/Sea Segment 17
Ice/Sea Seg. 1 SLS
Ice/Sea Seg. 2 SLS
Ice/Sea Seg. 3 SLS
Ice/Sea Seg. 4 SLS
Peard Bay
Elson Lagoon
Simpson Lagoon	12	28	8	7	2	1	.	.	16	11	1	.	51	25	1
Gwydyr Bay	2	3	5	11	5	17	4	2	3	6	6	6	1	6	50
Jago Lagoon	.	1	2	3	9	12	16	46	1	3	8	22	.	3	7
Beaufort Lagoon	1	1	3	2	.	1	1	2	.	.	1
Subsis. Res. Area A
Subsis. Res. Area B	1
Subsis. Res. Area C	87	**	66	**	78	87	17	10	**	**	92	18	**	**	**
Subsis. Res. Area D	17	25	41	74	86	**	**	**	28	67	**	**	12	43	**
Fall Feeding Area
Summer Feed. Area 1	1	1	.	.	.	1	.	.	.
Summer Feed. Area 2
Southern SLS Area
Northern SLS Area
Northern SLS
Boundary Segment 1	1
Boundary Segment 2	2	2	4	3	.	.	2	4	.	.	2
Boundary Segment 3	2	1	1	.	.	.	2	.	.	.	1

Notes: ** = Greater than 99.5%; . = Less than 0.5%; SLS = spring lead system.
Boundary Segments with all values less than 0.5% are not shown

Table B-8. Seasonal conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location in the summer season will contact a certain environmental resource area within 180 days, Beaufort Sea OCS Lease Sale 170

Environmental Resource Area	Hypothetical Spill Location														
	L1	L2	L3	L4	L5	L6	L7	L8	P1	P2	P3	P4	P5	P6	P7
Land	34	44	32	46	40	50	42	63	36	37	43	49	49	53	57
Ice/Sea Segment 1
Ice/Sea Segment 2
Ice/Sea Segment 3	1	1	1	1	2	1	.	.	1	1	.
Ice/Sea Segment 4	1	1	1	1	1	1	.	.	2	1	1	.	1	1	1
Ice/Sea Segment 5	3	2	2	2	1	1	.	.	3	2	1	.	2	1	1
Ice/Sea Segment 6	19	17	5	3	3	2	1	1	11	5	3	1	19	3	2
Ice/Sea Segment 7	58	40	26	21	6	4	3	2	46	21	5	3	45	22	4
Ice/Sea Segment 8	37	35	86	49	39	26	6	5	49	66	36	4	24	43	23
Ice/Sea Segment 9	9	10	15	19	64	55	84	33	9	13	67	57	7	12	44
Ice/Sea Segment 10	7	6	14	14	26	23	48	27	8	12	27	37	5	12	16
Ice/Sea Segment 11	1	1	6	3	11	11	20	16	1	8	10	17	1	2	10
Ice/Sea Segment 12	3	4	8	7	.	.	6	7	.	.	6
Ice/Sea Segment 13	.	.	.	2	3	4	6	6	.	1	3	6	.	1	4
Ice/Sea Segment 14
Ice/Sea Segment 15
Ice/Sea Segment 16
Ice/Sea Segment 17
Ice/Sea Seg. 1 SLS
Ice/Sea Seg. 2 SLS
Ice/Sea Seg. 3 SLS
Ice/Sea Seg. 4 SLS
Peard Bay
Elson Lagoon
Simpson Lagoon	13	31	10	8	3	1	.	.	18	15	2	.	52	25	1
Gwydyr Bay	2	3	5	13	5	19	5	3	3	8	6	6	1	6	51
Jago Lagoon	.	1	2	3	10	13	18	48	1	3	8	24	.	3	7
Beaufort Lagoon	1	1	3	2	.	1	1	2	.	.	1
Subsis. Res. Area A
Subsis. Res. Area B	2	1	1	1	1	1	.	.	2	1	1
Subsis. Res. Area C	87	**	68	**	79	88	20	11	**	**	92	20	**	**	**
Subsis. Res. Area D	20	31	45	78	87	**	**	**	36	71	**	**	14	48	**
Fall Feeding Area
Summer Feed. Area 1	1	4	5	.	.	.	5	.	.	.
Summer Feed. Area 2
Southern SLS Area
Northern SLS Area	2	1	2	1	1	1	1	.	2	2	1	1	1	1	1
Northern SLS
Boundary Segment 1	1
Boundary Segment 2	2	2	4	3	.	.	2	4	.	.	2
Boundary Segment 3	2	1	1	.	.	.	2	.	.	.	1

Notes: ** = Greater than 99.5%; . = Less than 0.5%; SLS = spring lead system.
Boundary Segments with all values less than 0.5% are not shown

Appendix C

Seasonal Conditional Probabilities of Contact to Land Segments

Table C-1. Seasonal conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location in the winter season will contact a certain land segment area within 3 days, Beaufort Sea OCS Lease Sale 170

Land Segments	<u>Hypothetical Spill Location</u>														
	L1	L2	L3	L4	L5	L6	L7	L8	P1	P2	P3	P4	P5	P6	P7
32	1	1	3	.	.
33	1	2	1	1	.	.	4	.	.
34	.	2	1	2	1	2	.	.	1	11	.
35	.	.	.	2	.	1	.	.	.	1	.	.	.	1	1
36	.	.	.	3	.	1	1	2
37	2	1	1	.	.	3
38	2	1	5	.	.	1	1	.	.	.
39	5
40	6	.	.	.	1	.	.	.
41	1

Notes: . = Less than 0.5%.
 Rows with all values less than 0.5% are not shown

Table C-2. Seasonal conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location in the winter season will contact a certain land segment area within 10 days, Beaufort Sea OCS Lease Sale 170

Land Segments	<u>Hypothetical Spill Location</u>														
	L1	L2	L3	L4	L5	L6	L7	L8	P1	P2	P3	P4	P5	P6	P7
29	1	1	1	.	.
30	1	1	1	.	.	.	1	.	.
31	1	.	.
32	1	2	1	.	.	.	3	.	.
33	1	2	1	1	1	.	.	5	.	.
34	1	2	2	3	1	1	.	.	1	2	1	.	1	12	.
35	.	.	1	3	1	1	.	.	1	1	1	.	.	1	1
36	.	.	.	3	1	2	1	.	.	1	1	1	.	1	2
37	.	.	.	1	1	2	1	1	.	1	1	2	.	1	5
38	1	2	1	7	.	.	1	2	.	.	1
39	5	.	.	.	1	.	.	.
40	1	7	.	.	.	1	.	.	.
41	1	2	.	.	.	1	.	.	.

Notes: . = Less than 0.5%.
 Rows with all values less than 0.5% are not shown

Table C-3. Seasonal conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location in the winter season will contact a certain land segment area within 30 days, Beaufort Sea OCS Lease Sale 170

Land Segments	Hypothetical Spill Location														
	L1	L2	L3	L4	L5	L6	L7	L8	P1	P2	P3	P4	P5	P6	P7
28	.	1	1	.	.
29	1	1	1	.	.
30	1	1	1	.	.	.	1	.	.
31	1	.	.
32	1	2	1	1	.	.	3	.	.
33	1	3	1	1	1	.	.	5	.	.
34	1	3	2	3	1	1	.	.	2	2	1	.	1	13	.
35	.	1	1	5	1	2	1	.	1	1	2	.	.	1	2
36	.	.	.	3	1	3	1	.	.	1	1	1	.	2	3
37	.	.	1	1	1	3	1	1	.	1	1	2	.	1	6
38	1	3	1	9	.	.	1	3	.	.	1
39	6	.	.	1	1	.	.	.
40	1	8	.	.	.	1	.	.	.
41	1	2	.	.	.	1	.	.	.

Notes: . = Less than 0.5%.
 Rows with all values less than 0.5% are not shown

Table C-4. Seasonal conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location in the winter season will contact a certain land segment area within 180 days, Beaufort Sea OCS Lease Sale 170

Land Segments	Hypothetical Spill Location														
	L1	L2	L3	L4	L5	L6	L7	L8	P1	P2	P3	P4	P5	P6	P7
23	1
27	1	1	1	.	.
28	4	4	2	1	1	.	.	.	2	1	.	.	3	2	.
29	9	7	4	1	1	.	.	.	11	3	1	.	10	1	.
30	2	4	1	2	1	.	.	7	1	.
31	1	1	2	1	.
32	1	4	2	1	2	2	.	.	6	3	.
33	1	8	4	4	2	1	.	.	2	5	1	.	9	4	1
34	1	8	3	12	5	5	2	.	2	6	5	1	3	27	4
35	2	3	1	10	5	9	3	1	1	1	8	3	8	2	8
36	.	4	1	5	2	8	4	4	2	1	5	5	.	2	12
37	.	.	2	6	1	6	3	5	.	3	3	7	.	6	13
38	.	.	1	2	1	8	2	19	.	2	2	3	.	.	3
39	2	3	.	6	.	.	1	1	.	.	8
40	3	3	2	13	.	.	5	4	.	.	.
41	1	.	5	6	.	.	.	6	.	.	.
42	2	1	.	.	.	1	.	.	.

Notes: . = Less than 0.5%.
 Rows with all values less than 0.5% are not shown

Table C-5. Seasonal conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location in the summer season will contact a certain land segment area within 3 days, Beaufort Sea OCS Lease Sale 170

Land Segments	Hypothetical Spill Location														
	L1	L2	L3	L4	L5	L6	L7	L8	P1	P2	P3	P4	P5	P6	P7
32	1	1	2	.	.
33	1	3	2	.	.	.	7	.	.
34	1	5	1	2	2	2	.	.	2	15	.
35	.	1	1	4	2	.	.	.	3	.
36	.	.	.	3	1	1	1	.	.	1	2
37	.	.	.	1	1	4	1	.	.	1	1	.	.	.	9
38	1	4	2	3	.	.	2	3	.	.	2
39	1	1	4	.	.	.	1	.	.	.
40	1	1	12	.	.	1	2	.	.	.
41	2	5	.	.	.	3	.	.	.

Notes: . = Less than 0.5%.
 Rows with all values less than 0.5% are not shown

Table C-6. Seasonal conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location in the summer season will contact a certain land segment area within 10 days, Beaufort Sea OCS Lease Sale 170

Land Segments	Hypothetical Spill Location														
	L1	L2	L3	L4	L5	L6	L7	L8	P1	P2	P3	P4	P5	P6	P7
29	1	1	1	.	.	.	1	.	.
30	1	1	1	.	.
31	1	1	1	.	.
32	2	2	1	.	.	.	3	.	.
33	2	5	1	1	4	.	.	.	9	.	.
34	3	7	4	5	4	6	.	.	5	21	.
35	1	2	2	6	2	2	.	.	2	4	2	.	1	4	1
36	.	1	1	5	3	3	.	.	2	2	3	.	1	3	5
37	.	1	1	4	2	5	1	1	1	3	2	2	.	2	11
38	.	.	1	1	2	7	4	6	.	1	5	5	.	1	5
39	1	2	2	5	.	.	1	2	.	.	1
40	.	.	.	1	2	4	3	15	.	.	3	5	.	1	3
41	.	.	.	1	2	2	5	9	.	1	2	5	.	.	1
42	1	1	.	.	.	1	.	.	.

Notes: . = Less than 0.5%.
 Rows with all values less than 0.5% are not shown

Table C-7. Seasonal conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location in the summer season will contact a certain land segment area within 30 days, Beaufort Sea OCS Lease Sale 170

Land Segments	Hypothetical Spill Location														
	L1	L2	L3	L4	L5	L6	L7	L8	P1	P2	P3	P4	P5	P6	P7
24	1	1	.	.	.	1	.	.
26	.	1	1
27	1	1	1	1	1	1	.	1	1	.
28	2	1	.	1	1	.	.	.	2	.	.
29	4	2	1	1	4	.	.	.	2	.	.
30	1	2	4	.	.
31	2	1	1	.	.	.	3	.	.
32	2	2	1	2	.	.	.	3	.	.
33	3	6	4	1	1	.	.	.	5	1	.	.	12	1	.
34	5	10	6	8	3	1	.	.	5	11	.	.	7	27	1
35	4	5	3	8	4	3	1	.	2	5	6	1	5	4	2
36	2	4	2	5	4	5	1	.	5	2	4	.	2	5	8
37	1	1	3	7	3	6	2	1	1	5	3	5	1	5	16
38	2	2	3	3	4	9	5	8	2	3	8	7	2	2	8
39	.	1	1	1	2	4	2	5	1	.	1	2	.	.	4
40	.	.	1	1	4	6	5	19	.	1	6	9	.	1	3
41	.	1	2	2	3	3	8	13	1	2	2	9	.	2	3
42	.	.	.	1	1	1	2	2	.	1	1	1	.	1	1
43	.	.	.	1	1	1	1	1	.	1	2	1	.	.	1
44	1	2	1	.	.	1	1	.	.	1
45	1	1	.	.	.	1	.	.	.

Notes: . = Less than 0.5%.
 Rows with all values less than 0.5% are not shown

Table C-8. Seasonal conditional probabilities (expressed as percent chance) that an oil spill starting at a particular location in the summer season will contact a certain land segment area within 180 days, Beaufort Sea OCS Lease Sale 170

Land Segments	Hypothetical Spill Location														
	L1	L2	L3	L4	L5	L6	L7	L8	P1	P2	P3	P4	P5	P6	P7
24	1	1	.	.	.	1	.	.
26	.	1	1
27	1	1	1	1	1	1	.	1	1	.
28	2	1	.	1	1	.	.	.	2	.	.
29	5	2	1	1	6	.	.	.	2	.	.
30	2	2	1	4	.	.
31	2	1	1	.	.	.	4	.	.
32	2	2	1	2	.	.	.	3	.	.
33	3	7	5	1	1	.	.	.	5	2	.	.	13	1	.
34	5	11	6	9	3	1	.	.	5	11	2	.	8	27	1
35	4	5	3	9	5	3	1	.	2	5	6	1	6	4	2
36	2	5	2	6	4	6	2	.	7	2	4	.	2	5	8
37	1	1	4	9	3	7	3	2	1	7	3	5	1	7	18
38	2	2	3	4	4	10	5	9	2	3	8	8	2	2	10
39	.	1	1	1	3	4	2	5	1	.	3	2	.	.	4
40	.	.	1	1	6	7	5	20	.	1	6	10	.	1	3
41	.	1	2	2	3	3	9	14	1	2	2	11	.	2	3
42	.	.	.	1	1	1	2	2	.	1	1	1	.	1	1
43	.	.	.	1	1	1	2	1	.	1	2	1	.	.	1
44	1	2	1	.	.	1	1	.	.	1
45	1	1	.	.	.	1	.	.	.

Notes: . = Less than 0.5%.
 Rows with all values less than 0.5% are not shown



The Department of the Interior Mission

As the Nation's principal conservation agency, the Department of the Interior has responsibility for most of our nationally owned public lands and natural resources. This includes fostering sound use of our land and water resources; protecting our fish, wildlife, and biological diversity; preserving the environmental and cultural values of our national parks and historical places; and providing for the enjoyment of life through outdoor recreation. The Department assesses our energy and mineral resources and works to ensure that their development is in the best interests of all our people by encouraging stewardship and citizen participation in their care. The Department also has a major responsibility for American Indian reservation communities and for people who live in island territories under U.S. administration.



The Minerals Management Service Mission

As a bureau of the Department of the Interior, the Minerals Management Service's (MMS) primary responsibilities are to manage the mineral resources located on the Nation's Outer Continental Shelf (OCS), collect revenue from the Federal OCS and onshore Federal and Indian lands, and distribute those revenues.

Moreover, in working to meet its responsibilities, the **Offshore Minerals Management Program** administers the OCS competitive leasing program and oversees the safe and environmentally sound exploration and production of our Nation's offshore natural gas, oil and other mineral resources. The MMS **Minerals Revenue Management** meets its responsibilities by ensuring the efficient, timely and accurate collection and disbursement of revenue from mineral leasing and production due to Indian tribes and allottees, States and the U.S. Treasury.

The MMS strives to fulfill its responsibilities through the general guiding principles of: (1) being responsive to the public's concerns and interests by maintaining a dialogue with all potentially affected parties and (2) carrying out its programs with an emphasis on working to enhance the quality of life for all Americans by lending MMS assistance and expertise to economic development and environmental protection.