

**Table 5.1C Standard Errors of Estimated Numbers (in Thousands) of Persons Reporting Past Year Substance Dependence, by Age Group: 2000**

Past Year Dependence	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Any Illicit Drug <sup>1</sup>	135	28	44	128
Marijuana and Hashish	90	26	39	79
Cocaine	77	8	17	74
Heroin	43	3	9	42
Hallucinogens	19	7	13	11
Inhalants	29	7	4	27
Nonmedical Use of Any Psychotherapeutic <sup>2</sup>	91	12	19	89
Pain Relievers	60	11	15	58
Tranquilizers	44	5	7	45
Stimulants	63	6	11	62
Sedatives	32	4	4	31
Alcohol	222	25	52	214
Alcohol or Any Illicit Drug <sup>1</sup>	249	34	61	239
Alcohol and Any Illicit Drug <sup>1</sup>	85	16	22	81

\*Low precision; no estimate reported.

NOTE: Dependence is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

<sup>1</sup> Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>2</sup> Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.1D Standard Errors of Percentages Reporting Past Year Substance Dependence, by Age Group: 2000**

Past Year Dependence	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Any Illicit Drug <sup>1</sup>	0.06	0.12	0.15	0.08
Marijuana and Hashish	0.04	0.11	0.13	0.05
Cocaine	0.03	0.04	0.06	0.04
Heroin	0.02	0.01	0.03	0.02
Hallucinogens	0.01	0.03	0.05	0.01
Inhalants	0.01	0.03	0.01	0.02
Nonmedical Use of Any Psychotherapeutic <sup>2</sup>	0.04	0.05	0.07	0.05
Pain Relievers	0.03	0.05	0.05	0.03
Tranquilizers	0.02	0.02	0.02	0.03
Stimulants	0.03	0.03	0.04	0.04
Sedatives	0.01	0.02	0.01	0.02
Alcohol	0.10	0.11	0.18	0.13
Alcohol or Any Illicit Drug <sup>1</sup>	0.11	0.15	0.21	0.14
Alcohol and Any Illicit Drug <sup>1</sup>	0.04	0.07	0.08	0.05

\*Low precision; no estimate reported.

NOTE: Dependence is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

<sup>1</sup> Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>2</sup> Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.2C Standard Errors of Estimated Numbers (in Thousands) of Persons Reporting Past Year Any Illicit Drug or Alcohol Dependence, by Detailed Age Categories: 2000**

Age Category	TYPE OF PAST YEAR DEPENDENCE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
Total	135	222	249
12	4	3	5
13	7	5	8
14	8	6	10
15	14	12	16
16	15	13	18
17	16	14	19
18	20	18	24
19	19	21	26
20	20	19	25
21	16	18	23
22	16	21	25
23	13	20	22
24	11	14	17
25	11	12	16
26-29	32	48	57
30-34	32	51	58
35-39	54	91	103
40-44	52	87	104
45-49	65	88	101
50-54	59	93	93
55-59	18	64	66
60-64	*	33	33
65 or Older	24	67	71

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.2D Standard Errors of Percentages Reporting Past Year Any Illicit Drug or Alcohol Dependence, by Detailed Age Categories: 2000**

Age Category	TYPE OF PAST YEAR DEPENDENCE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
Total	0.06	0.10	0.11
12	0.11	0.09	0.13
13	0.18	0.11	0.20
14	0.20	0.16	0.24
15	0.34	0.30	0.40
16	0.38	0.33	0.44
17	0.43	0.39	0.54
18	0.46	0.42	0.57
19	0.50	0.53	0.67
20	0.52	0.48	0.64
21	0.47	0.52	0.65
22	0.48	0.61	0.72
23	0.38	0.58	0.64
24	0.33	0.43	0.52
25	0.34	0.36	0.47
26-29	0.23	0.34	0.40
30-34	0.17	0.27	0.31
35-39	0.25	0.42	0.48
40-44	0.23	0.39	0.47
45-49	0.33	0.44	0.50
50-54	0.31	0.49	0.49
55-59	0.13	0.47	0.49
60-64	*	0.32	0.32
65 or Older	0.08	0.21	0.23

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.3C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 12 or Older Reporting Past Year Any Illicit Drug or Alcohol Dependence, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR DEPENDENCE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	135	222	249
<b>AGE</b>			
12-17	28	25	34
18-25	44	52	61
26 or Older	128	214	239
<b>GENDER</b>			
Male	97	191	208
Female	101	116	139
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	116	192	216
Black Only	56	87	98
American Indian or Alaska Native Only	4	11	12
Native Hawaiian or Other Pacific Islander	4	4	5
Asian Only	11	48	50
More Than One Race	14	15	17
Hispanic	38	62	67

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.3D Standard Errors of Percentages of Persons Aged 12 or Older Reporting Past Year Any Illicit Drug or Alcohol Dependence, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR DEPENDENCE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.06	0.10	0.11
<b>AGE</b>			
12-17	0.12	0.11	0.15
18-25	0.15	0.18	0.21
26 or Older	0.08	0.13	0.14
<b>GENDER</b>			
Male	0.09	0.18	0.19
Female	0.09	0.10	0.12
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	0.07	0.12	0.13
Black Only	0.22	0.34	0.39
American Indian or Alaska Native Only	0.38	1.02	1.13
Native Hawaiian or Other Pacific Islander	0.76	0.71	0.95
Asian Only	0.14	0.64	0.65
More Than One Race	0.77	0.81	0.93
Hispanic	0.16	0.26	0.28

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.4C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 12 to 17 Reporting Past Year Any Illicit Drug or Alcohol Dependence, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR DEPENDENCE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	28	25	34
<b>GENDER</b>			
Male	23	18	26
Female	17	17	22
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	24	22	29
Black Only	8	5	9
American Indian or Alaska Native Only	3	2	3
Native Hawaiian or Other Pacific Islander	*	*	*
Asian Only	5	5	7
More Than One Race	2	4	5
Hispanic	11	9	13
<b>GENDER/RACE/HISPANIC ORIGIN</b>			
Male - White	20	16	23
Female - White	14	14	18
Male - Black	7	4	7
Female - Black	5	3	5
Male - Hispanic	8	7	9
Female - Hispanic	7	6	8

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.4D Standard Errors of Percentages of Persons Aged 12 to 17 Reporting Past Year Any Illicit Drug or Alcohol Dependence, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR DEPENDENCE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.12	0.11	0.15
<b>GENDER</b>			
Male	0.19	0.15	0.22
Female	0.15	0.15	0.19
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	0.16	0.14	0.19
Black Only	0.24	0.14	0.27
American Indian or Alaska Native Only	2.10	1.50	2.32
Native Hawaiian or Other Pacific Islander	*	*	*
Asian Only	0.65	0.58	0.80
More Than One Race	0.64	1.16	1.32
Hispanic	0.33	0.28	0.39
<b>GENDER/RACE/HISPANIC ORIGIN</b>			
Male - White	0.25	0.20	0.29
Female - White	0.19	0.19	0.24
Male - Black	0.39	0.23	0.44
Female - Black	0.28	0.17	0.32
Male - Hispanic	0.46	0.42	0.54
Female - Hispanic	0.47	0.35	0.53

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.



**Table 5.5C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 18 to 25 Reporting Past Year Any Illicit Drug or Alcohol Dependence, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR DEPENDENCE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	44	52	61
<b>GENDER</b>			
Male	35	41	51
Female	28	30	37
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	38	46	53
Black Only	16	16	22
American Indian or Alaska Native Only	2	2	3
Native Hawaiian or Other Pacific Islander	*	*	*
Asian Only	5	8	8
More Than One Race	6	7	8
Hispanic	16	18	23
<b>ADULT EDUCATION</b>			
< High School	23	26	32
High School Graduate	30	28	37
Some College	23	28	34
College Graduate	11	16	17
<b>CURRENT EMPLOYMENT</b>			
Full-Time	34	37	47
Part-Time	23	24	32
Unemployed	16	16	19
Other <sup>1</sup>	15	19	24

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

<sup>1</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.5D Standard Errors of Percentages of Persons Aged 18 to 25 Reporting Past Year Any Illicit Drug or Alcohol Dependence, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR DEPENDENCE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.15	0.18	0.21
<b>GENDER</b>			
Male	0.25	0.29	0.36
Female	0.19	0.21	0.26
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	0.20	0.24	0.27
Black Only	0.41	0.41	0.56
American Indian or Alaska Native Only	1.26	1.09	1.59
Native Hawaiian or Other Pacific Islander	*	*	*
Asian Only	0.47	0.77	0.75
More Than One Race	1.90	2.21	2.75
Hispanic	0.37	0.41	0.53
<b>ADULT EDUCATION</b>			
< High School	0.38	0.42	0.53
High School Graduate	0.29	0.27	0.35
Some College	0.25	0.31	0.37
College Graduate	0.31	0.46	0.51
<b>CURRENT EMPLOYMENT</b>			
Full-Time	0.22	0.24	0.30
Part-Time	0.34	0.35	0.47
Unemployed	1.00	0.96	1.19
Other <sup>1</sup>	0.29	0.36	0.44

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

<sup>1</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.6C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 26 or Older Reporting Past Year Any Illicit Drug or Alcohol Dependence, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR DEPENDENCE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	128	214	239
<b>GENDER</b>			
Male	87	185	198
Female	95	114	134
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	110	183	206
Black Only	52	85	96
American Indian or Alaska Native Only	2	11	12
Native Hawaiian or Other Pacific Islander	*	*	*
Asian Only	8	49	49
More Than One Race	13	13	14
Hispanic	33	60	61
<b>ADULT EDUCATION</b>			
< High School	69	123	128
High School Graduate	55	124	132
Some College	53	109	119
College Graduate	72	73	101
<b>CURRENT EMPLOYMENT</b>			
Full-Time	81	173	188
Part-Time	63	72	81
Unemployed	29	36	44
Other <sup>1</sup>	65	112	122

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

<sup>1</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.6D Standard Errors of Percentages of Persons Aged 26 or Older Reporting Past Year Any Illicit Drug or Alcohol Dependence, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR DEPENDENCE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.08	0.13	0.14
<b>GENDER</b>			
Male	0.11	0.23	0.24
Female	0.11	0.13	0.15
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	0.09	0.14	0.16
Black Only	0.29	0.47	0.53
American Indian or Alaska Native Only	0.29	1.41	1.58
Native Hawaiian or Other Pacific Islander	*	*	*
Asian Only	0.13	0.84	0.85
More Than One Race	1.07	1.07	1.17
Hispanic	0.20	0.37	0.38
<b>ADULT EDUCATION</b>			
< High School	0.24	0.42	0.44
High School Graduate	0.10	0.22	0.23
Some College	0.14	0.28	0.30
College Graduate	0.16	0.16	0.22
<b>CURRENT EMPLOYMENT</b>			
Full-Time	0.08	0.17	0.19
Part-Time	0.38	0.43	0.48
Unemployed	0.96	1.18	1.45
Other <sup>1</sup>	0.13	0.22	0.24

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

<sup>1</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.7C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 18 or Older Reporting Past Year Any Illicit Drug or Alcohol Dependence, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR DEPENDENCE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	136	222	249
<b>GENDER</b>			
Male	94	191	207
Female	101	114	138
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	115	192	216
Black Only	56	87	99
American Indian or Alaska Native Only	3	11	12
Native Hawaiian or Other Pacific Islander	4	4	5
Asian Only	9	48	49
More Than One Race	14	14	17
Hispanic	37	61	65
<b>ADULT EDUCATION</b>			
< High School	74	126	133
High School Graduate	65	127	139
Some College	57	113	125
College Graduate	73	74	103
<b>CURRENT EMPLOYMENT</b>			
Full-Time	89	178	196
Part-Time	66	74	86
Unemployed	34	40	49
Other <sup>1</sup>	67	114	125

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

<sup>1</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.7D Standard Errors of Percentages of Persons Aged 18 or Older Reporting Past Year Any Illicit Drug or Alcohol Dependence, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR DEPENDENCE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.07	0.11	0.12
<b>GENDER</b>			
Male	0.10	0.20	0.22
Female	0.10	0.11	0.13
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	0.08	0.13	0.15
Black Only	0.25	0.40	0.45
American Indian or Alaska Native Only	0.34	1.13	1.31
Native Hawaiian or Other Pacific Islander	0.79	0.74	1.01
Asian Only	0.13	0.71	0.72
More Than One Race	0.94	0.97	1.12
Hispanic	0.18	0.30	0.32
<b>ADULT EDUCATION</b>			
< High School	0.21	0.36	0.38
High School Graduate	0.10	0.19	0.21
Some College	0.12	0.23	0.26
College Graduate	0.15	0.15	0.21
<b>CURRENT EMPLOYMENT</b>			
Full-Time	0.08	0.15	0.17
Part-Time	0.28	0.32	0.37
Unemployed	0.73	0.85	1.06
Other <sup>1</sup>	0.12	0.20	0.22

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

<sup>1</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.8C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 12 or Older Reporting Past Year Any Illicit Drug or Alcohol Dependence, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR DEPENDENCE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	135	222	249
<b>GEOGRAPHIC DIVISION</b>			
Northeast	62	94	110
New England	44	43	53
Middle Atlantic	44	83	96
Midwest	41	76	86
East North Central	30	68	73
West North Central	28	32	44
South	63	143	154
South Atlantic	54	105	117
East South Central	18	64	65
West South Central	28	72	75
West	92	119	139
Mountain	31	44	42
Pacific	87	111	132
<b>COUNTY TYPE</b>			
Large Metro	110	173	197
Small Metro	60	101	113
250K - 1 Mil. Pop.	49	92	101
<250K Pop.	35	43	52
Nonmetro	47	95	105
Urbanized	19	37	39
Less Urbanized	42	75	84
Completely Rural	10	45	46

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.8D Standard Errors of Percentages of Persons Aged 12 or Older Reporting Past Year Any Illicit Drug or Alcohol Dependence, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR DEPENDENCE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.06	0.10	0.11
<b>GEOGRAPHIC DIVISION</b>			
Northeast	0.15	0.22	0.26
New England	0.39	0.39	0.47
Middle Atlantic	0.14	0.26	0.31
Midwest	0.08	0.15	0.17
East North Central	0.08	0.19	0.20
West North Central	0.18	0.21	0.28
South	0.08	0.18	0.19
South Atlantic	0.13	0.26	0.28
East South Central	0.13	0.46	0.47
West South Central	0.12	0.29	0.31
West	0.19	0.24	0.28
Mountain	0.22	0.31	0.30
Pacific	0.25	0.32	0.38
<b>COUNTY TYPE</b>			
Large Metro	0.10	0.16	0.18
Small Metro	0.09	0.15	0.16
250K - 1 Mil. Pop.	0.10	0.18	0.20
<250K Pop.	0.20	0.24	0.29
Nonmetro	0.10	0.21	0.23
Urbanized	0.14	0.27	0.29
Less Urbanized	0.15	0.28	0.31
Completely Rural	0.18	0.85	0.87

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.



**Table 5.9C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 12 to 17 Reporting Past Year Any Illicit Drug or Alcohol Dependence, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR DEPENDENCE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	28	25	34
<b>GEOGRAPHIC DIVISION</b>			
Northeast	11	9	13
New England	8	8	10
Middle Atlantic	7	6	8
Midwest	12	10	15
East North Central	10	8	11
West North Central	6	7	9
South	17	16	21
South Atlantic	11	8	13
East South Central	7	7	10
West South Central	11	12	13
West	16	14	18
Mountain	8	5	8
Pacific	14	13	16
<b>COUNTY TYPE</b>			
Large Metro	20	15	23
Small Metro	18	17	21
250K - 1 Mil. Pop.	13	12	16
<250K Pop.	11	11	13
Nonmetro	11	10	14
Urbanized	6	5	7
Less Urbanized	8	8	10
Completely Rural	3	4	4

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.9D Standard Errors of Percentages of Persons Aged 12 to 17 Reporting Past Year Any Illicit Drug or Alcohol Dependence, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR DEPENDENCE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.12	0.11	0.15
<b>GEOGRAPHIC DIVISION</b>			
Northeast	0.25	0.23	0.31
New England	0.73	0.68	0.93
Middle Atlantic	0.23	0.18	0.27
Midwest	0.22	0.19	0.27
East North Central	0.27	0.21	0.31
West North Central	0.36	0.39	0.53
South	0.21	0.19	0.25
South Atlantic	0.27	0.21	0.33
East South Central	0.51	0.52	0.69
West South Central	0.39	0.41	0.46
West	0.30	0.26	0.34
Mountain	0.47	0.29	0.51
Pacific	0.37	0.35	0.43
<b>COUNTY TYPE</b>			
Large Metro	0.18	0.13	0.20
Small Metro	0.25	0.23	0.29
250K - 1 Mil. Pop.	0.25	0.22	0.31
<250K Pop.	0.62	0.62	0.71
Nonmetro	0.22	0.21	0.28
Urbanized	0.45	0.38	0.56
Less Urbanized	0.28	0.28	0.36
Completely Rural	0.53	0.65	0.73

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.10C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 18 to 25 Reporting Past Year Any Illicit Drug or Alcohol Dependence, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR DEPENDENCE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	44	52	61
<b>GEOGRAPHIC DIVISION</b>			
Northeast	21	21	25
New England	11	12	14
Middle Atlantic	18	16	22
Midwest	16	23	27
East North Central	15	19	23
West North Central	7	13	14
South	26	32	37
South Atlantic	16	22	24
East South Central	11	12	14
West South Central	18	20	25
West	23	27	32
Mountain	11	11	15
Pacific	20	24	28
<b>COUNTY TYPE</b>			
Large Metro	33	39	45
Small Metro	24	29	34
250K - 1 Mil. Pop.	20	24	28
<250K Pop.	13	16	19
Nonmetro	18	21	26
Urbanized	9	12	13
Less Urbanized	15	17	22
Completely Rural	2	4	4

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.10D Standard Errors of Percentages of Persons Aged 18 to 25 Reporting Past Year Any Illicit Drug or Alcohol Dependence, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR DEPENDENCE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.15	0.18	0.21
<b>GEOGRAPHIC DIVISION</b>			
Northeast	0.41	0.40	0.50
New England	0.87	0.95	1.04
Middle Atlantic	0.47	0.43	0.57
Midwest	0.23	0.33	0.39
East North Central	0.30	0.40	0.48
West North Central	0.34	0.62	0.69
South	0.25	0.31	0.36
South Atlantic	0.32	0.44	0.49
East South Central	0.63	0.66	0.74
West South Central	0.50	0.57	0.70
West	0.34	0.40	0.47
Mountain	0.53	0.56	0.72
Pacific	0.44	0.52	0.60
<b>COUNTY TYPE</b>			
Large Metro	0.24	0.28	0.32
Small Metro	0.26	0.30	0.36
250K - 1 Mil. Pop.	0.31	0.36	0.43
<250K Pop.	0.47	0.55	0.66
Nonmetro	0.32	0.38	0.47
Urbanized	0.48	0.61	0.72
Less Urbanized	0.46	0.53	0.68
Completely Rural	0.46	0.77	0.91

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.11C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 26 or Older Reporting Past Year Any Illicit Drug or Alcohol Dependence, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR DEPENDENCE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	128	214	239
<b>GEOGRAPHIC DIVISION</b>			
Northeast	58	92	103
New England	42	41	49
Middle Atlantic	39	82	90
Midwest	34	73	78
East North Central	23	66	68
West North Central	25	30	39
South	53	138	147
South Atlantic	49	103	112
East South Central	11	58	59
West South Central	18	70	73
West	95	115	137
Mountain	32	41	43
Pacific	89	108	131
<b>COUNTY TYPE</b>			
Large Metro	109	170	192
Small Metro	50	93	102
250K - 1 Mil. Pop.	42	86	92
<250K Pop.	27	37	45
Nonmetro	41	90	97
Urbanized	17	32	34
Less Urbanized	37	70	77
Completely Rural	9	45	46

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.11D Standard Errors of Percentages of Persons Aged 26 or Older Reporting Past Year Any Illicit Drug or Alcohol Dependence, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR DEPENDENCE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.08	0.13	0.14
<b>GEOGRAPHIC DIVISION</b>			
Northeast	0.17	0.27	0.31
New England	0.48	0.47	0.56
Middle Atlantic	0.16	0.33	0.36
Midwest	0.09	0.18	0.20
East North Central	0.08	0.24	0.25
West North Central	0.21	0.25	0.33
South	0.09	0.23	0.24
South Atlantic	0.15	0.32	0.35
East South Central	0.11	0.54	0.56
West South Central	0.10	0.39	0.40
West	0.26	0.31	0.37
Mountain	0.30	0.38	0.41
Pacific	0.34	0.41	0.50
<b>COUNTY TYPE</b>			
Large Metro	0.13	0.20	0.23
Small Metro	0.10	0.18	0.20
250K - 1 Mil. Pop.	0.11	0.22	0.24
<250K Pop.	0.21	0.28	0.35
Nonmetro	0.12	0.26	0.28
Urbanized	0.17	0.32	0.33
Less Urbanized	0.18	0.33	0.37
Completely Rural	0.21	1.08	1.09

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.12C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 18 or Older Reporting Past Year Any Illicit Drug or Alcohol Dependence, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR DEPENDENCE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	136	222	249
<b>GEOGRAPHIC DIVISION</b>			
Northeast	61	94	110
New England	43	42	51
Middle Atlantic	43	84	97
Midwest	39	77	86
East North Central	28	70	74
West North Central	27	33	44
South	63	143	153
South Atlantic	54	105	114
East South Central	17	63	66
West South Central	28	74	77
West	96	118	139
Mountain	33	43	43
Pacific	90	110	132
<b>COUNTY TYPE</b>			
Large Metro	113	172	196
Small Metro	58	100	111
250K - 1 Mil. Pop.	49	91	100
<250K Pop.	32	41	50
Nonmetro	46	95	104
Urbanized	18	36	38
Less Urbanized	41	74	83
Completely Rural	9	45	45

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.12D Standard Errors of Percentages of Persons Aged 18 or Older Reporting Past Year Any Illicit Drug or Alcohol Dependence, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR DEPENDENCE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.07	0.11	0.12
<b>GEOGRAPHIC DIVISION</b>			
Northeast	0.16	0.24	0.28
New England	0.43	0.42	0.50
Middle Atlantic	0.15	0.29	0.34
Midwest	0.08	0.17	0.19
East North Central	0.09	0.21	0.23
West North Central	0.19	0.24	0.32
South	0.09	0.20	0.21
South Atlantic	0.15	0.28	0.31
East South Central	0.14	0.50	0.53
West South Central	0.13	0.34	0.35
West	0.22	0.27	0.32
Mountain	0.26	0.34	0.34
Pacific	0.29	0.36	0.43
<b>COUNTY TYPE</b>			
Large Metro	0.12	0.18	0.20
Small Metro	0.09	0.16	0.18
250K - 1 Mil. Pop.	0.11	0.20	0.22
<250K Pop.	0.20	0.26	0.32
Nonmetro	0.11	0.23	0.25
Urbanized	0.15	0.30	0.32
Less Urbanized	0.17	0.31	0.34
Completely Rural	0.19	0.96	0.97

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.



**Table 5.13C Standard Errors of Estimated Numbers (in Thousands) of Persons Reporting Past Year Substance Abuse, by Age Group: 2000**

Past Year Abuse	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Any Illicit Drug <sup>1</sup>	86	27	35	73
Marijuana and Hashish	74	23	30	64
Cocaine	33	6	12	31
Heroin	5	3	4	*
Hallucinogens	26	11	16	15
Inhalants	11	9	6	*
Nonmedical Use of Any Psychotherapeutic <sup>2</sup>	47	14	15	42
Pain Relievers	41	11	13	37
Tranquilizers	28	6	8	26
Stimulants	13	8	7	7
Sedatives	13	4	5	11
Alcohol	230	32	70	212
Alcohol or Any Illicit Drug <sup>1</sup>	240	39	74	224
Alcohol and Any Illicit Drug <sup>1</sup>	44	14	21	36

\*Low precision; no estimate reported.

NOTE: Abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

<sup>1</sup> Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>2</sup> Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.13D Standard Errors of Percentages Reporting Past Year Substance Abuse, by Age Group: 2000**

Past Year Abuse	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Any Illicit Drug <sup>1</sup>	0.04	0.11	0.12	0.04
Marijuana and Hashish	0.03	0.10	0.10	0.04
Cocaine	0.01	0.03	0.04	0.02
Heroin	0.00	0.01	0.01	*
Hallucinogens	0.01	0.05	0.06	0.01
Inhalants	0.00	0.04	0.02	*
Nonmedical Use of Any Psychotherapeutic <sup>2</sup>	0.02	0.06	0.05	0.02
Pain Relievers	0.02	0.05	0.04	0.02
Tranquilizers	0.01	0.02	0.03	0.02
Stimulants	0.01	0.04	0.02	0.00
Sedatives	0.01	0.02	0.02	0.01
Alcohol	0.10	0.14	0.24	0.12
Alcohol or Any Illicit Drug <sup>1</sup>	0.11	0.17	0.25	0.13
Alcohol and Any Illicit Drug <sup>1</sup>	0.02	0.06	0.07	0.02

\*Low precision; no estimate reported.

NOTE: Abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

<sup>1</sup> Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>2</sup> Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.14C Standard Errors of Estimated Numbers (in Thousands) of Persons Reporting Past Year Any Illicit Drug or Alcohol Abuse, by Detailed Age Categories: 2000**

Age Category	TYPE OF PAST YEAR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
Total	86	230	240
12	5	4	7
13	8	8	10
14	9	10	13
15	12	14	17
16	13	17	20
17	14	18	22
18	17	24	28
19	18	27	29
20	12	28	29
21	11	22	24
22	11	23	24
23	8	23	24
24	9	21	22
25	6	21	21
26-29	24	57	58
30-34	27	59	64
35-39	35	98	100
40-44	21	89	91
45-49	20	61	63
50-54	42	64	76
55-59	16	58	60
60-64	*	37	37
65 or Older	*	92	92

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.14D Standard Errors of Percentages Reporting Past Year Any Illicit Drug or Alcohol Abuse, by Detailed Age Categories: 2000**

Age Category	TYPE OF PAST YEAR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
Total	0.04	0.10	0.11
12	0.14	0.11	0.17
13	0.21	0.19	0.26
14	0.22	0.25	0.32
15	0.31	0.35	0.43
16	0.32	0.42	0.50
17	0.40	0.50	0.60
18	0.39	0.57	0.64
19	0.45	0.69	0.74
20	0.31	0.72	0.76
21	0.32	0.64	0.68
22	0.31	0.67	0.70
23	0.22	0.68	0.70
24	0.28	0.67	0.70
25	0.19	0.62	0.63
26-29	0.17	0.40	0.41
30-34	0.14	0.32	0.34
35-39	0.16	0.46	0.46
40-44	0.09	0.40	0.41
45-49	0.10	0.30	0.31
50-54	0.22	0.34	0.40
55-59	0.12	0.43	0.44
60-64	*	0.36	0.36
65 or Older	*	0.29	0.29

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.15C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 12 or Older Reporting Past Year Any Illicit Drug or Alcohol Abuse, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	86	230	240
<b>AGE</b>			
12-17	27	32	39
18-25	35	70	74
26 or Older	73	212	224
<b>GENDER</b>			
Male	63	204	208
Female	58	112	127
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	59	202	207
Black Only	26	54	59
American Indian or Alaska Native Only	10	12	15
Native Hawaiian or Other Pacific Islander	0	3	3
Asian Only	7	21	22
More Than One Race	*	16	41
Hispanic	41	85	90

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.15D Standard Errors of Percentages of Persons Aged 12 or Older Reporting Past Year Any Illicit Drug or Alcohol Abuse, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.04	0.10	0.11
<b>AGE</b>			
12-17	0.11	0.14	0.17
18-25	0.12	0.24	0.25
26 or Older	0.04	0.12	0.13
<b>GENDER</b>			
Male	0.06	0.19	0.19
Female	0.05	0.10	0.11
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	0.04	0.12	0.13
Black Only	0.10	0.21	0.23
American Indian or Alaska Native Only	0.95	1.14	1.38
Native Hawaiian or Other Pacific Islander	0.05	0.60	0.60
Asian Only	0.09	0.27	0.29
More Than One Race	*	0.85	2.20
Hispanic	0.17	0.36	0.38

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.16C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 12 to 17 Reporting Past Year Any Illicit Drug or Alcohol Abuse, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	27	32	39
<b>GENDER</b>			
Male	20	23	28
Female	17	21	25
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	20	29	34
Black Only	10	10	13
American Indian or Alaska Native Only	3	1	4
Native Hawaiian or Other Pacific Islander	*	*	*
Asian Only	4	4	5
More Than One Race	2	5	5
Hispanic	13	11	16
<b>GENDER/RACE/HISPANIC ORIGIN</b>			
Male - White	15	21	24
Female - White	14	18	22
Male - Black	9	8	10
Female - Black	5	7	8
Male - Hispanic	10	9	13
Female - Hispanic	7	7	9

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.16D Standard Errors of Percentages of Persons Aged 12 to 17 Reporting Past Year Any Illicit Drug or Alcohol Abuse, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.11	0.14	0.17
<b>GENDER</b>			
Male	0.17	0.20	0.23
Female	0.15	0.19	0.22
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	0.13	0.19	0.22
Black Only	0.30	0.31	0.40
American Indian or Alaska Native Only	2.32	0.98	2.47
Native Hawaiian or Other Pacific Islander	*	*	*
Asian Only	0.47	0.47	0.60
More Than One Race	0.49	1.24	1.31
Hispanic	0.39	0.32	0.48
<b>GENDER/RACE/HISPANIC ORIGIN</b>			
Male - White	0.19	0.26	0.31
Female - White	0.19	0.25	0.29
Male - Black	0.50	0.46	0.61
Female - Black	0.32	0.41	0.52
Male - Hispanic	0.60	0.51	0.75
Female - Hispanic	0.42	0.41	0.55

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.



**Table 5.17C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 18 to 25 Reporting Past Year Any Illicit Drug or Alcohol Abuse, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	35	70	74
<b>GENDER</b>			
Male	29	57	59
Female	19	38	41
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	30	59	62
Black Only	10	22	23
American Indian or Alaska Native Only	*	8	8
Native Hawaiian or Other Pacific Islander	0	1	1
Asian Only	3	8	8
More Than One Race	2	8	8
Hispanic	13	26	30
<b>ADULT EDUCATION</b>			
< High School	20	29	33
High School Graduate	20	37	40
Some College	19	43	45
College Graduate	5	26	26
<b>CURRENT EMPLOYMENT</b>			
Full-Time	25	52	55
Part-Time	16	33	35
Unemployed	11	17	18
Other <sup>1</sup>	15	25	27

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

<sup>1</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.17D Standard Errors of Percentages of Persons Aged 18 to 25 Reporting Past Year Any Illicit Drug or Alcohol Abuse, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.12	0.24	0.25
<b>GENDER</b>			
Male	0.20	0.39	0.41
Female	0.13	0.26	0.28
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	0.16	0.31	0.32
Black Only	0.25	0.56	0.60
American Indian or Alaska Native Only	*	4.27	4.27
Native Hawaiian or Other Pacific Islander	0.17	1.44	1.44
Asian Only	0.26	0.79	0.82
More Than One Race	0.80	2.75	2.79
Hispanic	0.30	0.61	0.68
<b>ADULT EDUCATION</b>			
< High School	0.33	0.48	0.53
High School Graduate	0.19	0.36	0.38
Some College	0.21	0.47	0.50
College Graduate	0.16	0.76	0.77
<b>CURRENT EMPLOYMENT</b>			
Full-Time	0.16	0.34	0.36
Part-Time	0.23	0.49	0.53
Unemployed	0.70	1.05	1.11
Other <sup>1</sup>	0.28	0.47	0.51

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

<sup>1</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.18C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 26 or Older Reporting Past Year Any Illicit Drug or Alcohol Abuse, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	73	212	224
<b>GENDER</b>			
Male	51	191	195
Female	52	103	115
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	45	188	193
Black Only	21	45	49
American Indian or Alaska Native Only	8	9	12
Native Hawaiian or Other Pacific Islander	*	*	*
Asian Only	5	20	20
More Than One Race	*	12	*
Hispanic	36	77	79
<b>ADULT EDUCATION</b>			
< High School	37	86	89
High School Graduate	31	108	112
Some College	27	92	95
College Graduate	47	127	135
<b>CURRENT EMPLOYMENT</b>			
Full-Time	52	174	179
Part-Time	19	54	55
Unemployed	19	35	38
Other <sup>1</sup>	44	106	114

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

<sup>1</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.18D Standard Errors of Percentages of Persons Aged 26 or Older Reporting Past Year Any Illicit Drug or Alcohol Abuse, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.04	0.12	0.13
<b>GENDER</b>			
Male	0.06	0.24	0.24
Female	0.06	0.11	0.13
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	0.03	0.15	0.15
Black Only	0.12	0.25	0.27
American Indian or Alaska Native Only	1.06	1.18	1.57
Native Hawaiian or Other Pacific Islander	*	*	*
Asian Only	0.09	0.34	0.35
More Than One Race	*	0.99	*
Hispanic	0.22	0.47	0.49
<b>ADULT EDUCATION</b>			
< High School	0.13	0.29	0.30
High School Graduate	0.05	0.19	0.20
Some College	0.07	0.23	0.24
College Graduate	0.10	0.28	0.30
<b>CURRENT EMPLOYMENT</b>			
Full-Time	0.05	0.17	0.18
Part-Time	0.11	0.32	0.33
Unemployed	0.63	1.14	1.25
Other <sup>1</sup>	0.09	0.21	0.22

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

<sup>1</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.19C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 18 or Older Reporting Past Year Any Illicit Drug or Alcohol Abuse, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	82	227	238
<b>GENDER</b>			
Male	59	201	205
Female	55	111	125
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	55	198	203
Black Only	24	51	55
American Indian or Alaska Native Only	10	12	15
Native Hawaiian or Other Pacific Islander	0	3	3
Asian Only	6	21	21
More Than One Race	*	15	41
Hispanic	38	84	88
<b>ADULT EDUCATION</b>			
< High School	44	91	97
High School Graduate	38	117	121
Some College	34	103	107
College Graduate	47	129	137
<b>CURRENT EMPLOYMENT</b>			
Full-Time	58	184	191
Part-Time	25	65	67
Unemployed	23	40	45
Other <sup>1</sup>	46	110	118

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

<sup>1</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.19D Standard Errors of Percentages of Persons Aged 18 or Older Reporting Past Year Any Illicit Drug or Alcohol Abuse, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.04	0.11	0.12
<b>GENDER</b>			
Male	0.06	0.21	0.22
Female	0.05	0.11	0.12
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	0.04	0.13	0.14
Black Only	0.11	0.23	0.25
American Indian or Alaska Native Only	1.04	1.32	1.56
Native Hawaiian or Other Pacific Islander	0.04	0.60	0.60
Asian Only	0.08	0.30	0.31
More Than One Race	*	1.00	2.71
Hispanic	0.19	0.41	0.43
<b>ADULT EDUCATION</b>			
< High School	0.12	0.26	0.27
High School Graduate	0.06	0.17	0.18
Some College	0.07	0.21	0.22
College Graduate	0.10	0.26	0.28
<b>CURRENT EMPLOYMENT</b>			
Full-Time	0.05	0.16	0.16
Part-Time	0.11	0.28	0.29
Unemployed	0.50	0.87	0.96
Other <sup>1</sup>	0.08	0.20	0.21

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

<sup>1</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.20C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 12 or Older Reporting Past Year Any Illicit Drug or Alcohol Abuse, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	86	230	240
<b>GEOGRAPHIC DIVISION</b>			
Northeast	38	99	101
New England	16	73	75
Middle Atlantic	34	67	67
Midwest	43	84	95
East North Central	34	69	76
West North Central	26	47	56
South	36	130	128
South Atlantic	26	77	78
East South Central	14	42	44
West South Central	21	96	92
West	54	138	149
Mountain	21	53	63
Pacific	49	127	135
<b>COUNTY TYPE</b>			
Large Metro	74	187	200
Small Metro	32	117	115
250K - 1 Mil. Pop.	28	101	99
<250K Pop.	15	58	58
Nonmetro	29	75	79
Urbanized	12	45	46
Less Urbanized	26	54	59
Completely Rural	8	26	27

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.20D Standard Errors of Percentages of Persons Aged 12 or Older Reporting Past Year Any Illicit Drug or Alcohol Abuse, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.04	0.10	0.11
<b>GEOGRAPHIC DIVISION</b>			
Northeast	0.09	0.23	0.24
New England	0.14	0.65	0.67
Middle Atlantic	0.11	0.21	0.21
Midwest	0.08	0.16	0.18
East North Central	0.09	0.19	0.21
West North Central	0.17	0.30	0.36
South	0.04	0.16	0.16
South Atlantic	0.06	0.19	0.19
East South Central	0.10	0.30	0.32
West South Central	0.08	0.39	0.38
West	0.11	0.28	0.30
Mountain	0.15	0.37	0.44
Pacific	0.14	0.37	0.39
<b>COUNTY TYPE</b>			
Large Metro	0.07	0.17	0.18
Small Metro	0.05	0.17	0.17
250K - 1 Mil. Pop.	0.06	0.20	0.19
<250K Pop.	0.08	0.33	0.33
Nonmetro	0.06	0.17	0.17
Urbanized	0.09	0.34	0.35
Less Urbanized	0.10	0.20	0.22
Completely Rural	0.14	0.50	0.50

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.



**Table 5.21C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 12 to 17 Reporting Past Year Any Illicit Drug or Alcohol Abuse, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	27	32	39
<b>GEOGRAPHIC DIVISION</b>			
Northeast	10	13	16
New England	6	8	8
Middle Atlantic	8	10	14
Midwest	11	17	20
East North Central	9	15	16
West North Central	7	9	12
South	18	20	24
South Atlantic	11	13	16
East South Central	6	11	12
West South Central	13	12	13
West	13	13	17
Mountain	6	7	9
Pacific	11	11	14
<b>COUNTY TYPE</b>			
Large Metro	18	22	27
Small Metro	15	17	21
250K - 1 Mil. Pop.	14	15	18
<250K Pop.	6	9	11
Nonmetro	12	16	19
Urbanized	5	8	9
Less Urbanized	9	10	12
Completely Rural	6	10	10

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.21D Standard Errors of Percentages of Persons Aged 12 to 17 Reporting Past Year Any Illicit Drug or Alcohol Abuse, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.11	0.14	0.17
<b>GEOGRAPHIC DIVISION</b>			
Northeast	0.24	0.31	0.37
New England	0.56	0.74	0.71
Middle Atlantic	0.26	0.31	0.44
Midwest	0.20	0.32	0.37
East North Central	0.24	0.39	0.43
West North Central	0.40	0.55	0.70
South	0.21	0.24	0.29
South Atlantic	0.26	0.31	0.40
East South Central	0.44	0.74	0.83
West South Central	0.45	0.42	0.45
West	0.24	0.24	0.31
Mountain	0.37	0.42	0.54
Pacific	0.30	0.30	0.37
<b>COUNTY TYPE</b>			
Large Metro	0.16	0.20	0.24
Small Metro	0.21	0.24	0.29
250K - 1 Mil. Pop.	0.26	0.28	0.33
<250K Pop.	0.35	0.51	0.63
Nonmetro	0.25	0.33	0.39
Urbanized	0.38	0.57	0.69
Less Urbanized	0.31	0.35	0.43
Completely Rural	1.07	1.64	1.75

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.22C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 18 to 25 Reporting Past Year Any Illicit Drug or Alcohol Abuse, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	35	70	74
<b>GEOGRAPHIC DIVISION</b>			
Northeast	15	26	28
New England	6	13	14
Middle Atlantic	14	23	25
Midwest	15	37	37
East North Central	13	33	34
West North Central	7	16	16
South	21	42	43
South Atlantic	16	34	35
East South Central	6	14	16
West South Central	12	18	20
West	18	33	36
Mountain	8	15	16
Pacific	16	29	33
<b>COUNTY TYPE</b>			
Large Metro	25	45	50
Small Metro	21	42	43
250K - 1 Mil. Pop.	18	31	33
<250K Pop.	11	29	29
Nonmetro	11	26	27
Urbanized	9	16	17
Less Urbanized	7	20	20
Completely Rural	1	6	6

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.22D Standard Errors of Percentages of Persons Aged 18 to 25 Reporting Past Year Any Illicit Drug or Alcohol Abuse, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.12	0.24	0.25
<b>GEOGRAPHIC DIVISION</b>			
Northeast	0.29	0.51	0.56
New England	0.42	1.00	1.07
Middle Atlantic	0.37	0.61	0.67
Midwest	0.22	0.53	0.54
East North Central	0.28	0.69	0.70
West North Central	0.34	0.76	0.78
South	0.20	0.40	0.42
South Atlantic	0.32	0.69	0.70
East South Central	0.33	0.78	0.87
West South Central	0.34	0.52	0.56
West	0.26	0.49	0.55
Mountain	0.38	0.74	0.80
Pacific	0.34	0.62	0.70
<b>COUNTY TYPE</b>			
Large Metro	0.18	0.32	0.36
Small Metro	0.22	0.45	0.46
250K - 1 Mil. Pop.	0.26	0.47	0.50
<250K Pop.	0.37	1.01	1.03
Nonmetro	0.20	0.47	0.49
Urbanized	0.49	0.84	0.89
Less Urbanized	0.22	0.61	0.63
Completely Rural	0.22	1.26	1.26

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.23C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 26 or Older Reporting Past Year Any Illicit Drug or Alcohol Abuse, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	73	212	224
<b>GEOGRAPHIC DIVISION</b>			
Northeast	33	97	97
New England	12	71	71
Middle Atlantic	31	65	64
Midwest	37	71	80
East North Central	29	56	62
West North Central	23	43	52
South	25	115	116
South Atlantic	19	66	68
East South Central	9	34	35
West South Central	14	88	87
West	48	131	144
Mountain	17	49	56
Pacific	44	122	132
<b>COUNTY TYPE</b>			
Large Metro	66	179	191
Small Metro	20	96	97
250K - 1 Mil. Pop.	18	85	85
<250K Pop.	7	45	45
Nonmetro	23	66	69
Urbanized	7	41	42
Less Urbanized	22	45	50
Completely Rural	3	22	22

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.23D Standard Errors of Percentages of Persons Aged 26 or Older Reporting Past Year Any Illicit Drug or Alcohol Abuse, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.04	0.12	0.13
<b>GEOGRAPHIC DIVISION</b>			
Northeast	0.10	0.29	0.29
New England	0.14	0.81	0.81
Middle Atlantic	0.13	0.26	0.26
Midwest	0.09	0.18	0.20
East North Central	0.10	0.20	0.22
West North Central	0.19	0.36	0.44
South	0.04	0.19	0.19
South Atlantic	0.06	0.21	0.21
East South Central	0.08	0.32	0.33
West South Central	0.07	0.48	0.48
West	0.13	0.35	0.39
Mountain	0.16	0.46	0.52
Pacific	0.17	0.46	0.50
<b>COUNTY TYPE</b>			
Large Metro	0.08	0.21	0.23
Small Metro	0.04	0.19	0.19
250K - 1 Mil. Pop.	0.05	0.22	0.22
<250K Pop.	0.05	0.35	0.35
Nonmetro	0.07	0.19	0.20
Urbanized	0.07	0.41	0.41
Less Urbanized	0.10	0.22	0.24
Completely Rural	0.06	0.52	0.52

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.24C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 18 or Older Reporting Past Year Any Illicit Drug or Alcohol Abuse, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	82	227	238
<b>GEOGRAPHIC DIVISION</b>			
Northeast	37	98	99
New England	14	72	74
Middle Atlantic	34	66	66
Midwest	41	81	91
East North Central	33	66	73
West North Central	25	46	54
South	31	127	128
South Atlantic	23	76	75
East South Central	11	41	43
West South Central	17	94	94
West	51	138	149
Mountain	20	53	61
Pacific	47	128	136
<b>COUNTY TYPE</b>			
Large Metro	72	186	199
Small Metro	28	113	113
250K - 1 Mil. Pop.	25	96	97
<250K Pop.	12	59	58
Nonmetro	24	73	76
Urbanized	11	45	46
Less Urbanized	23	53	57
Completely Rural	3	22	22

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.24D Standard Errors of Percentages of Persons Aged 18 or Older Reporting Past Year Any Illicit Drug or Alcohol Abuse, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.04	0.11	0.12
<b>GEOGRAPHIC DIVISION</b>			
Northeast	0.10	0.25	0.26
New England	0.14	0.72	0.73
Middle Atlantic	0.12	0.23	0.23
Midwest	0.09	0.17	0.20
East North Central	0.10	0.20	0.22
West North Central	0.18	0.33	0.39
South	0.04	0.18	0.18
South Atlantic	0.06	0.20	0.20
East South Central	0.09	0.33	0.35
West South Central	0.08	0.43	0.43
West	0.12	0.32	0.34
Mountain	0.16	0.42	0.48
Pacific	0.15	0.41	0.44
<b>COUNTY TYPE</b>			
Large Metro	0.07	0.19	0.20
Small Metro	0.05	0.18	0.18
250K - 1 Mil. Pop.	0.05	0.21	0.21
<250K Pop.	0.08	0.37	0.37
Nonmetro	0.06	0.18	0.19
Urbanized	0.10	0.38	0.38
Less Urbanized	0.10	0.22	0.24
Completely Rural	0.06	0.48	0.48

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.



**Table 5.25C Standard Errors of Estimated Numbers (in Thousands) of Persons Reporting Past Year Substance Dependence or Abuse, by Age Group: 2000**

Past Year Dependence or Abuse	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Any Illicit Drug <sup>1</sup>	168	38	55	152
Marijuana and Hashish	119	34	48	103
Cocaine	85	10	22	80
Heroin	43	4	10	42
Hallucinogens	32	13	21	19
Inhalants	31	11	7	27
Nonmedical Use of Any Psychotherapeutic <sup>2</sup>	101	18	24	98
Pain Relievers	73	15	20	68
Tranquilizers	52	8	11	52
Stimulants	64	11	13	62
Sedatives	33	5	7	33
Alcohol	307	41	83	292
Alcohol or Any Illicit Drug <sup>1</sup>	338	50	92	320
Alcohol and Any Illicit Drug <sup>1</sup>	109	25	38	99

\*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

<sup>1</sup> Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>2</sup> Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.25D Standard Errors of Percentages Reporting Past Year Substance Dependence or Abuse, by Age Group: 2000**

Past Year Dependence or Abuse	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Any Illicit Drug <sup>1</sup>	0.08	0.16	0.19	0.09
Marijuana and Hashish	0.05	0.14	0.16	0.06
Cocaine	0.04	0.04	0.08	0.05
Heroin	0.02	0.02	0.03	0.02
Hallucinogens	0.01	0.06	0.07	0.01
Inhalants	0.01	0.05	0.03	0.02
Nonmedical Use of Any Psychotherapeutic <sup>2</sup>	0.05	0.08	0.08	0.06
Pain Relievers	0.03	0.06	0.07	0.04
Tranquilizers	0.02	0.03	0.04	0.03
Stimulants	0.03	0.05	0.04	0.04
Sedatives	0.01	0.02	0.02	0.02
Alcohol	0.14	0.18	0.29	0.17
Alcohol or Any Illicit Drug <sup>1</sup>	0.15	0.21	0.32	0.19
Alcohol and Any Illicit Drug <sup>1</sup>	0.05	0.11	0.13	0.06

\*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

<sup>1</sup> Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>2</sup> Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.26C Standard Errors of Estimated Numbers (in Thousands) of Persons Reporting Past Year Dependence or Abuse for Any Illicit Drug or Alcohol, by Detailed Age Categories: 2000**

Age Category	TYPE OF PAST YEAR DEPENDENCE OR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
Total	168	307	338
12	7	5	8
13	11	9	13
14	12	12	15
15	18	18	22
16	19	21	25
17	20	22	27
18	26	30	34
19	25	30	33
20	23	33	37
21	19	28	30
22	18	29	31
23	15	30	31
24	15	25	27
25	13	24	25
26-29	38	74	77
30-34	42	78	87
35-39	66	135	145
40-44	56	125	135
45-49	68	105	116
50-54	72	112	120
55-59	24	86	89
60-64	*	49	49
65 or Older	24	113	115

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.26D Standard Errors of Percentages Reporting Past Year Dependence or Abuse for Any Illicit Drug or Alcohol, by Detailed Age Categories: 2000**

Age Category	TYPE OF PAST YEAR DEPENDENCE OR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
Total	0.08	0.14	0.15
12	0.18	0.14	0.22
13	0.28	0.22	0.32
14	0.29	0.30	0.38
15	0.46	0.45	0.56
16	0.49	0.53	0.63
17	0.57	0.61	0.76
18	0.59	0.69	0.78
19	0.64	0.77	0.85
20	0.60	0.85	0.95
21	0.55	0.80	0.86
22	0.52	0.85	0.90
23	0.43	0.86	0.90
24	0.45	0.79	0.82
25	0.39	0.72	0.74
26-29	0.27	0.52	0.54
30-34	0.22	0.42	0.46
35-39	0.30	0.63	0.67
40-44	0.25	0.56	0.61
45-49	0.34	0.52	0.58
50-54	0.38	0.59	0.63
55-59	0.18	0.64	0.66
60-64	*	0.48	0.48
65 or Older	0.08	0.36	0.37

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.27C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 12 or Older Reporting Past Year Dependence or Abuse for Any Illicit Drug or Alcohol, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR DEPENDENCE OR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	168	307	338
<b>AGE</b>			
12-17	38	41	50
18-25	55	83	92
26 or Older	152	292	320
<b>GENDER</b>			
Male	119	271	281
Female	123	163	200
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	132	269	295
Black Only	61	101	108
American Indian or Alaska Native Only	11	17	19
Native Hawaiian or Other Pacific Islander	4	5	6
Asian Only	13	54	55
More Than One Race	40	22	43
Hispanic	59	103	109

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.27D Standard Errors of Percentages of Persons Aged 12 or Older Reporting Past Year Dependence or Abuse for Any Illicit Drug or Alcohol, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR DEPENDENCE OR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.08	0.14	0.15
<b>AGE</b>			
12-17	0.16	0.18	0.21
18-25	0.19	0.29	0.32
26 or Older	0.09	0.17	0.19
<b>GENDER</b>			
Male	0.11	0.25	0.26
Female	0.11	0.14	0.17
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	0.08	0.17	0.18
Black Only	0.24	0.40	0.43
American Indian or Alaska Native Only	1.02	1.53	1.76
Native Hawaiian or Other Pacific Islander	0.78	0.88	1.11
Asian Only	0.16	0.70	0.72
More Than One Race	2.13	1.17	2.29
Hispanic	0.25	0.43	0.46

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.28C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 12 to 17 Reporting Past Year Dependence or Abuse for Any Illicit Drug or Alcohol, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR DEPENDENCE OR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	38	41	50
<b>GENDER</b>			
Male	29	29	35
Female	25	26	32
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	31	36	43
Black Only	12	11	15
American Indian or Alaska Native Only	4	2	4
Native Hawaiian or Other Pacific Islander	*	*	*
Asian Only	6	6	8
More Than One Race	3	6	7
Hispanic	17	14	19
<b>GENDER/RACE/HISPANIC ORIGIN</b>			
Male - White	24	25	31
Female - White	20	23	27
Male - Black	11	9	12
Female - Black	7	7	10
Male - Hispanic	12	11	14
Female - Hispanic	10	9	12

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.28D Standard Errors of Percentages of Persons Aged 12 to 17 Reporting Past Year Dependence or Abuse for Any Illicit Drug or Alcohol, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR DEPENDENCE OR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.16	0.18	0.21
<b>GENDER</b>			
Male	0.24	0.24	0.29
Female	0.22	0.23	0.28
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	0.21	0.24	0.28
Black Only	0.37	0.34	0.46
American Indian or Alaska Native Only	2.78	1.69	3.12
Native Hawaiian or Other Pacific Islander	*	*	*
Asian Only	0.75	0.74	0.95
More Than One Race	0.81	1.69	1.79
Hispanic	0.51	0.41	0.56
<b>GENDER/RACE/HISPANIC ORIGIN</b>			
Male - White	0.31	0.32	0.40
Female - White	0.27	0.31	0.36
Male - Black	0.63	0.51	0.71
Female - Black	0.44	0.44	0.63
Male - Hispanic	0.72	0.64	0.81
Female - Hispanic	0.66	0.53	0.75

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.



**Table 5.29C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 18 to 25 Reporting Past Year Dependence or Abuse for Any Illicit Drug or Alcohol, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR DEPENDENCE OR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	55	83	92
<b>GENDER</b>			
Male	44	67	73
Female	34	48	53
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	48	69	76
Black Only	19	26	29
American Indian or Alaska Native Only	*	8	8
Native Hawaiian or Other Pacific Islander	*	3	3
Asian Only	5	11	11
More Than One Race	6	10	11
Hispanic	21	31	35
<b>ADULT EDUCATION</b>			
< High School	31	38	43
High School Graduate	37	43	49
Some College	31	51	55
College Graduate	12	30	30
<b>CURRENT EMPLOYMENT</b>			
Full-Time	43	62	69
Part-Time	28	38	44
Unemployed	19	21	23
Other <sup>1</sup>	21	34	37

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

<sup>1</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.29D Standard Errors of Percentages of Persons Aged 18 to 25 Reporting Past Year Dependence or Abuse for Any Illicit Drug or Alcohol, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR DEPENDENCE OR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.19	0.29	0.32
<b>GENDER</b>			
Male	0.31	0.47	0.51
Female	0.23	0.33	0.36
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	0.25	0.36	0.40
Black Only	0.48	0.66	0.73
American Indian or Alaska Native Only	*	4.61	4.57
Native Hawaiian or Other Pacific Islander	*	3.14	3.14
Asian Only	0.54	1.06	1.09
More Than One Race	2.07	3.32	3.49
Hispanic	0.48	0.71	0.80
<b>ADULT EDUCATION</b>			
< High School	0.51	0.62	0.70
High School Graduate	0.35	0.42	0.47
Some College	0.34	0.56	0.60
College Graduate	0.35	0.87	0.89
<b>CURRENT EMPLOYMENT</b>			
Full-Time	0.28	0.40	0.45
Part-Time	0.41	0.58	0.66
Unemployed	1.19	1.30	1.43
Other <sup>1</sup>	0.38	0.64	0.69

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

<sup>1</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.30C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 26 or Older Reporting Past Year Dependence or Abuse for Any Illicit Drug or Alcohol, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR DEPENDENCE OR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	152	292	320
<b>GENDER</b>			
Male	103	257	268
Female	116	156	188
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	119	251	274
Black Only	57	96	103
American Indian or Alaska Native Only	8	14	17
Native Hawaiian or Other Pacific Islander	*	*	*
Asian Only	9	53	54
More Than One Race	*	17	*
Hispanic	51	98	101
<b>ADULT EDUCATION</b>			
< High School	80	147	150
High School Graduate	63	166	174
Some College	62	139	150
College Graduate	84	144	161
<b>CURRENT EMPLOYMENT</b>			
Full-Time	97	242	257
Part-Time	66	88	96
Unemployed	35	50	58
Other <sup>1</sup>	78	151	167

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

<sup>1</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.30D Standard Errors of Percentages of Persons Aged 26 or Older Reporting Past Year Dependence or Abuse for Any Illicit Drug or Alcohol, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR DEPENDENCE OR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.09	0.17	0.19
<b>GENDER</b>			
Male	0.13	0.32	0.33
Female	0.13	0.17	0.21
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	0.09	0.20	0.21
Black Only	0.32	0.53	0.57
American Indian or Alaska Native Only	1.09	1.82	2.18
Native Hawaiian or Other Pacific Islander	*	*	*
Asian Only	0.16	0.92	0.93
More Than One Race	*	1.46	*
Hispanic	0.31	0.60	0.62
<b>ADULT EDUCATION</b>			
< High School	0.27	0.50	0.51
High School Graduate	0.11	0.29	0.31
Some College	0.16	0.35	0.38
College Graduate	0.18	0.32	0.35
<b>CURRENT EMPLOYMENT</b>			
Full-Time	0.10	0.24	0.25
Part-Time	0.39	0.53	0.57
Unemployed	1.15	1.63	1.89
Other <sup>1</sup>	0.15	0.30	0.33

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

<sup>1</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.31C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 18 or Older Reporting Past Year Dependence or Abuse for Any Illicit Drug or Alcohol, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR DEPENDENCE OR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	166	305	338
<b>GENDER</b>			
Male	114	269	280
Female	122	162	199
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	129	265	292
Black Only	61	100	108
American Indian or Alaska Native Only	10	16	19
Native Hawaiian or Other Pacific Islander	4	4	6
Asian Only	10	53	54
More Than One Race	39	21	42
Hispanic	55	103	108
<b>ADULT EDUCATION</b>			
< High School	88	152	157
High School Graduate	77	172	185
Some College	68	151	165
College Graduate	85	147	164
<b>CURRENT EMPLOYMENT</b>			
Full-Time	111	252	268
Part-Time	71	98	107
Unemployed	42	57	64
Other <sup>1</sup>	82	154	171

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

<sup>1</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.31D Standard Errors of Percentages of Persons Aged 18 or Older Reporting Past Year Dependence or Abuse for Any Illicit Drug or Alcohol, by Demographic Characteristics: 2000**

Demographic Characteristic	TYPE OF PAST YEAR DEPENDENCE OR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.08	0.15	0.17
<b>GENDER</b>			
Male	0.12	0.28	0.29
Female	0.12	0.15	0.19
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic			
White Only	0.09	0.18	0.20
Black Only	0.28	0.46	0.49
American Indian or Alaska Native Only	1.10	1.74	1.99
Native Hawaiian or Other Pacific Islander	0.82	0.92	1.17
Asian Only	0.15	0.78	0.79
More Than One Race	2.62	1.40	2.80
Hispanic	0.27	0.50	0.53
<b>ADULT EDUCATION</b>			
< High School	0.25	0.43	0.44
High School Graduate	0.12	0.26	0.28
Some College	0.14	0.31	0.34
College Graduate	0.17	0.30	0.33
<b>CURRENT EMPLOYMENT</b>			
Full-Time	0.10	0.22	0.23
Part-Time	0.30	0.42	0.46
Unemployed	0.90	1.22	1.37
Other <sup>1</sup>	0.15	0.28	0.31

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

<sup>1</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.32C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 12 or Older Reporting Past Year Dependence or Abuse for Any Illicit Drug or Alcohol, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR DEPENDENCE OR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	168	307	338
<b>GEOGRAPHIC DIVISION</b>			
Northeast	77	127	139
New England	48	80	89
Middle Atlantic	59	99	106
Midwest	66	118	138
East North Central	50	102	114
West North Central	42	59	78
South	74	197	201
South Atlantic	60	128	138
East South Central	24	81	82
West South Central	35	125	122
West	113	159	189
Mountain	42	58	63
Pacific	104	148	178
<b>COUNTY TYPE</b>			
Large Metro	141	242	277
Small Metro	70	149	154
250K - 1 Mil. Pop.	59	128	131
<250K Pop.	37	77	83
Nonmetro	55	126	135
Urbanized	23	61	63
Less Urbanized	49	96	105
Completely Rural	12	58	59

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.32D Standard Errors of Percentages of Persons Aged 12 or Older Reporting Past Year Dependence or Abuse for Any Illicit Drug or Alcohol, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR DEPENDENCE OR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.08	0.14	0.15
<b>GEOGRAPHIC DIVISION</b>			
Northeast	0.18	0.30	0.32
New England	0.43	0.71	0.79
Middle Atlantic	0.19	0.31	0.34
Midwest	0.13	0.23	0.27
East North Central	0.14	0.28	0.31
West North Central	0.27	0.38	0.50
South	0.09	0.25	0.25
South Atlantic	0.15	0.31	0.33
East South Central	0.18	0.58	0.59
West South Central	0.14	0.51	0.50
West	0.23	0.32	0.38
Mountain	0.29	0.40	0.44
Pacific	0.30	0.43	0.51
<b>COUNTY TYPE</b>			
Large Metro	0.13	0.22	0.25
Small Metro	0.10	0.22	0.22
250K - 1 Mil. Pop.	0.12	0.25	0.26
<250K Pop.	0.21	0.43	0.47
Nonmetro	0.12	0.28	0.30
Urbanized	0.17	0.46	0.47
Less Urbanized	0.18	0.36	0.39
Completely Rural	0.24	1.10	1.12

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.



**Table 5.33C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 12 to 17 Reporting Past Year Dependence or Abuse for Any Illicit Drug or Alcohol, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR DEPENDENCE OR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	38	41	50
<b>GEOGRAPHIC DIVISION</b>			
Northeast	16	15	18
New England	12	9	10
Middle Atlantic	11	12	15
Midwest	17	20	24
East North Central	14	17	20
West North Central	10	10	12
South	24	26	32
South Atlantic	15	15	21
East South Central	9	12	15
West South Central	16	18	18
West	18	20	24
Mountain	9	8	11
Pacific	16	19	21
<b>COUNTY TYPE</b>			
Large Metro	26	27	33
Small Metro	23	24	28
250K - 1 Mil. Pop.	19	19	22
<250K Pop.	13	14	18
Nonmetro	16	19	23
Urbanized	7	9	12
Less Urbanized	12	13	16
Completely Rural	7	10	11

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.33D Standard Errors of Percentages of Persons Aged 12 to 17 Reporting Past Year Dependence or Abuse for Any Illicit Drug or Alcohol, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR DEPENDENCE OR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.16	0.18	0.21
<b>GEOGRAPHIC DIVISION</b>			
Northeast	0.38	0.35	0.43
New England	1.08	0.80	0.88
Middle Atlantic	0.34	0.38	0.49
Midwest	0.31	0.36	0.44
East North Central	0.38	0.45	0.54
West North Central	0.57	0.57	0.74
South	0.29	0.31	0.38
South Atlantic	0.38	0.36	0.53
East South Central	0.65	0.84	1.08
West South Central	0.58	0.62	0.64
West	0.34	0.38	0.45
Mountain	0.56	0.50	0.70
Pacific	0.42	0.49	0.56
<b>COUNTY TYPE</b>			
Large Metro	0.23	0.24	0.29
Small Metro	0.32	0.33	0.39
250K - 1 Mil. Pop.	0.35	0.36	0.41
<250K Pop.	0.71	0.77	0.97
Nonmetro	0.33	0.40	0.48
Urbanized	0.56	0.70	0.87
Less Urbanized	0.40	0.45	0.55
Completely Rural	1.17	1.67	1.82

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.34C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 18 to 25 Reporting Past Year Dependence or Abuse for Any Illicit Drug or Alcohol, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR DEPENDENCE OR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	55	83	92
<b>GEOGRAPHIC DIVISION</b>			
Northeast	25	32	35
New England	14	15	16
Middle Atlantic	21	29	33
Midwest	21	44	46
East North Central	18	40	42
West North Central	10	19	19
South	34	48	53
South Atlantic	22	37	40
East South Central	13	19	19
West South Central	22	23	28
West	29	41	47
Mountain	14	20	21
Pacific	25	36	42
<b>COUNTY TYPE</b>			
Large Metro	41	55	64
Small Metro	29	49	50
250K - 1 Mil. Pop.	24	38	38
<250K Pop.	16	32	34
Nonmetro	21	34	36
Urbanized	13	20	21
Less Urbanized	17	26	27
Completely Rural	3	7	7

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.34D Standard Errors of Percentages of Persons Aged 18 to 25 Reporting Past Year Dependence or Abuse for Any Illicit Drug or Alcohol, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR DEPENDENCE OR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.19	0.29	0.32
<b>GEOGRAPHIC DIVISION</b>			
Northeast	0.49	0.62	0.69
New England	1.03	1.15	1.19
Middle Atlantic	0.55	0.76	0.86
Midwest	0.30	0.64	0.66
East North Central	0.38	0.84	0.87
West North Central	0.47	0.90	0.90
South	0.33	0.46	0.51
South Atlantic	0.45	0.75	0.82
East South Central	0.70	1.04	1.02
West South Central	0.62	0.64	0.80
West	0.43	0.62	0.70
Mountain	0.70	0.97	1.05
Pacific	0.53	0.78	0.89
<b>COUNTY TYPE</b>			
Large Metro	0.30	0.40	0.46
Small Metro	0.30	0.52	0.53
250K - 1 Mil. Pop.	0.37	0.57	0.58
<250K Pop.	0.57	1.11	1.18
Nonmetro	0.38	0.61	0.64
Urbanized	0.69	1.05	1.11
Less Urbanized	0.52	0.81	0.85
Completely Rural	0.54	1.42	1.53

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.35C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 26 or Older Reporting Past Year Dependence or Abuse for Any Illicit Drug or Alcohol, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR DEPENDENCE OR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	152	292	320
<b>GEOGRAPHIC DIVISION</b>			
Northeast	69	129	136
New England	44	82	90
Middle Atlantic	52	99	101
Midwest	54	103	120
East North Central	39	87	97
West North Central	38	54	71
South	58	183	188
South Atlantic	52	123	130
East South Central	14	71	73
West South Central	20	114	114
West	109	156	185
Mountain	39	50	56
Pacific	102	147	176
<b>COUNTY TYPE</b>			
Large Metro	131	235	263
Small Metro	56	129	137
250K - 1 Mil. Pop.	48	112	117
<250K Pop.	28	62	70
Nonmetro	48	114	123
Urbanized	18	53	55
Less Urbanized	43	86	94
Completely Rural	9	55	55

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.35D Standard Errors of Percentages of Persons Aged 26 or Older Reporting Past Year Dependence or Abuse for Any Illicit Drug or Alcohol, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR DEPENDENCE OR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.09	0.17	0.19
<b>GEOGRAPHIC DIVISION</b>			
Northeast	0.21	0.38	0.41
New England	0.50	0.93	1.02
Middle Atlantic	0.21	0.40	0.41
Midwest	0.14	0.26	0.30
East North Central	0.14	0.31	0.35
West North Central	0.32	0.46	0.60
South	0.09	0.30	0.31
South Atlantic	0.16	0.38	0.40
East South Central	0.13	0.67	0.69
West South Central	0.11	0.63	0.63
West	0.29	0.42	0.50
Mountain	0.36	0.47	0.53
Pacific	0.39	0.56	0.67
<b>COUNTY TYPE</b>			
Large Metro	0.16	0.28	0.31
Small Metro	0.11	0.25	0.26
250K - 1 Mil. Pop.	0.12	0.29	0.30
<250K Pop.	0.22	0.48	0.54
Nonmetro	0.14	0.33	0.35
Urbanized	0.18	0.53	0.54
Less Urbanized	0.20	0.41	0.45
Completely Rural	0.22	1.32	1.32

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.36C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 18 or Older Reporting Past Year Dependence or Abuse for Any Illicit Drug or Alcohol, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR DEPENDENCE OR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	166	305	338
<b>GEOGRAPHIC DIVISION</b>			
Northeast	75	126	138
New England	47	79	89
Middle Atlantic	59	98	104
Midwest	62	117	136
East North Central	48	101	112
West North Central	40	58	77
South	71	193	200
South Atlantic	61	127	137
East South Central	21	80	81
West South Central	31	121	122
West	112	161	192
Mountain	42	57	63
Pacific	104	151	181
<b>COUNTY TYPE</b>			
Large Metro	140	242	276
Small Metro	67	144	152
250K - 1 Mil. Pop.	58	124	129
<250K Pop.	34	74	82
Nonmetro	53	122	131
Urbanized	22	61	62
Less Urbanized	47	94	102
Completely Rural	9	55	55

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.36D Standard Errors of Percentages of Persons Aged 18 or Older Reporting Past Year Dependence or Abuse for Any Illicit Drug or Alcohol, by Geographic Characteristics: 2000**

Geographic Characteristic	TYPE OF PAST YEAR DEPENDENCE OR ABUSE		
	Any Illicit Drug	Alcohol	Any Illicit Drug or Alcohol
<b>TOTAL</b>	0.08	0.15	0.17
<b>GEOGRAPHIC DIVISION</b>			
Northeast	0.19	0.33	0.36
New England	0.46	0.78	0.88
Middle Atlantic	0.21	0.34	0.37
Midwest	0.13	0.25	0.29
East North Central	0.15	0.31	0.34
West North Central	0.29	0.42	0.55
South	0.10	0.27	0.28
South Atlantic	0.16	0.34	0.37
East South Central	0.17	0.64	0.65
West South Central	0.14	0.56	0.56
West	0.26	0.37	0.44
Mountain	0.33	0.45	0.49
Pacific	0.34	0.49	0.58
<b>COUNTY TYPE</b>			
Large Metro	0.14	0.25	0.28
Small Metro	0.11	0.23	0.25
250K - 1 Mil. Pop.	0.13	0.27	0.28
<250K Pop.	0.21	0.47	0.52
Nonmetro	0.13	0.30	0.32
Urbanized	0.18	0.51	0.52
Less Urbanized	0.20	0.39	0.42
Completely Rural	0.20	1.17	1.17

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on the definition found in the 4th ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.



**Table 5.37C Standard Errors of Estimated Numbers (in Thousands) of Persons Who Received Substance Abuse Treatment in the Past Year, by Detailed Age Categories: 2000**

Age Category	RECEIVED TREATMENT IN THE PAST YEAR <sup>1</sup>			
	Any Illicit Drug	Alcohol	Both Any Illicit Drug and Alcohol	Any Illicit Drug or Alcohol <sup>2</sup>
Total	108	151	96	169
12	1	2	1	3
13	4	5	4	6
14	5	5	4	7
15	8	8	7	10
16	11	9	9	12
17	11	11	9	13
18	10	11	9	13
19	8	11	6	13
20	12	13	11	15
21	8	12	7	13
22	8	7	4	11
23	8	14	6	15
24	7	8	5	9
25	4	5	3	6
26-29	24	34	20	39
30-34	23	33	16	38
35-39	35	52	32	67
40-44	45	58	42	62
45-49	29	38	21	45
50-54	65	84	63	86
55-59	10	52	*	53
60-64	*	17	*	24
65 or Older	2	2	*	3

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> "Received treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>2</sup> Estimates include persons who received treatment specifically for alcohol or any illicit drug, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.37D Standard Errors of Percentages of Persons Who Received Substance Abuse Treatment in the Past Year, by Detailed Age Categories: 2000**

Age Category	RECEIVED TREATMENT IN THE PAST YEAR <sup>1</sup>			
	Any Illicit Drug	Alcohol	Both Any Illicit Drug and Alcohol	Any Illicit Drug or Alcohol <sup>2</sup>
Total	0.05	0.07	0.04	0.08
12	0.04	0.05	0.04	0.07
13	0.10	0.12	0.10	0.16
14	0.13	0.12	0.10	0.17
15	0.20	0.19	0.18	0.24
16	0.27	0.23	0.22	0.29
17	0.31	0.30	0.26	0.37
18	0.22	0.24	0.20	0.29
19	0.20	0.28	0.16	0.33
20	0.32	0.34	0.29	0.38
21	0.22	0.34	0.20	0.36
22	0.24	0.21	0.11	0.32
23	0.22	0.40	0.18	0.44
24	0.21	0.25	0.16	0.29
25	0.12	0.15	0.09	0.17
26-29	0.17	0.24	0.14	0.28
30-34	0.12	0.17	0.08	0.20
35-39	0.16	0.24	0.15	0.31
40-44	0.20	0.26	0.19	0.28
45-49	0.15	0.19	0.10	0.22
50-54	0.34	0.44	0.33	0.45
55-59	0.07	0.38	*	0.39
60-64	*	0.17	*	0.24
65 or Older	0.01	0.01	*	0.01

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> "Received treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>2</sup> Estimates include persons who received treatment specifically for alcohol or any illicit drug, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.38C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 12 or Older Who Received Substance Abuse Treatment in the Past Year, by Demographic Characteristics: 2000**

Demographic Characteristic	RECEIVED TREATMENT IN THE PAST YEAR <sup>1</sup>			
	Any Illicit Drug	Alcohol	Both Any Illicit Drug and Alcohol	Any Illicit Drug or Alcohol <sup>2</sup>
<b>TOTAL</b>	108	151	96	169
<b>AGE</b>				
12-17	20	18	16	24
18-25	24	29	20	35
26 or Older	99	142	88	159
<b>GENDER</b>				
Male	61	120	54	134
Female	85	83	75	94
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic				
White Only	94	132	81	148
Black Only	42	54	41	59
American Indian or Alaska Native Only	5	15	4	15
Native Hawaiian or Other Pacific Islander	2	2	2	4
Asian Only	3	11	3	11
More Than One Race	13	13	13	13
Hispanic	20	38	18	41

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> "Received treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>2</sup> Estimates include persons who received treatment specifically for alcohol or any illicit drug, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.38D Standard Errors of Percentages of Persons Aged 12 or Older Who Received Substance Abuse Treatment in the Past Year, by Demographic Characteristics: 2000**

Demographic Characteristic	RECEIVED TREATMENT IN THE PAST YEAR <sup>1</sup>			
	Any Illicit Drug	Alcohol	Both Any Illicit Drug and Alcohol	Any Illicit Drug or Alcohol <sup>2</sup>
<b>TOTAL</b>	0.05	0.07	0.04	0.08
<b>AGE</b>				
12-17	0.09	0.08	0.07	0.10
18-25	0.08	0.10	0.07	0.12
26 or Older	0.06	0.08	0.05	0.09
<b>GENDER</b>				
Male	0.06	0.11	0.05	0.12
Female	0.07	0.07	0.06	0.08
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic				
White Only	0.06	0.08	0.05	0.09
Black Only	0.17	0.21	0.16	0.23
American Indian or Alaska Native Only	0.45	1.38	0.37	1.41
Native Hawaiian or Other Pacific Islander	0.30	0.31	0.30	0.67
Asian Only	0.04	0.14	0.04	0.14
More Than One Race	0.68	0.69	0.68	0.69
Hispanic	0.08	0.16	0.07	0.17

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> "Received treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>2</sup> Estimates include persons who received treatment specifically for alcohol or any illicit drug, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.39C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 12 to 17 Who Received Substance Abuse Treatment in the Past Year, by Demographic Characteristics: 2000**

Demographic Characteristic	RECEIVED TREATMENT IN THE PAST YEAR <sup>1</sup>			
	Any Illicit Drug	Alcohol	Both Any Illicit Drug and Alcohol	Any Illicit Drug or Alcohol <sup>2</sup>
<b>TOTAL</b>	20	18	16	24
<b>GENDER</b>				
Male	16	14	13	19
Female	11	11	10	13
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic				
White Only	16	15	13	19
Black Only	6	5	4	7
American Indian or Alaska Native Only	*	*	*	4
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian Only	2	2	2	2
More Than One Race	1	2	1	2
Hispanic	7	8	7	9
<b>GENDER/RACE/HISPANIC ORIGIN</b>				
Male - White	12	10	9	14
Female - White	10	10	9	12
Male - Black	5	4	4	6
Female - Black	2	2	2	3
Male - Hispanic	7	8	6	8
Female - Hispanic	4	3	3	5

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> "Received treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>2</sup> Estimates include persons who received treatment specifically for alcohol or any illicit drug, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.39D Standard Errors of Percentages of Persons Aged 12 to 17 Who Received Substance Abuse Treatment in the Past Year, by Demographic Characteristics: 2000**

Demographic Characteristic	RECEIVED TREATMENT IN THE PAST YEAR <sup>1</sup>			
	Any Illicit Drug	Alcohol	Both Any Illicit Drug and Alcohol	Any Illicit Drug or Alcohol <sup>2</sup>
<b>TOTAL</b>	0.09	0.08	0.07	0.10
<b>GENDER</b>				
Male	0.13	0.12	0.10	0.16
Female	0.10	0.10	0.09	0.12
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic				
White Only	0.10	0.10	0.09	0.12
Black Only	0.17	0.14	0.13	0.21
American Indian or Alaska Native Only	*	*	*	2.46
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian Only	0.21	0.21	0.21	0.26
More Than One Race	0.21	0.48	0.21	0.50
Hispanic	0.23	0.24	0.21	0.28
<b>GENDER/RACE/HISPANIC ORIGIN</b>				
Male - White	0.15	0.13	0.12	0.18
Female - White	0.13	0.14	0.12	0.16
Male - Black	0.31	0.25	0.24	0.36
Female - Black	0.12	0.11	0.10	0.20
Male - Hispanic	0.38	0.44	0.37	0.47
Female - Hispanic	0.25	0.21	0.20	0.32

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> "Received treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>2</sup> Estimates include persons who received treatment specifically for alcohol or any illicit drug, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.40C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 18 to 25 Who Received Substance Abuse Treatment in the Past Year, by Demographic Characteristics: 2000**

Demographic Characteristic	RECEIVED TREATMENT IN THE PAST YEAR <sup>1</sup>			
	Any Illicit Drug	Alcohol	Both Any Illicit Drug and Alcohol	Any Illicit Drug or Alcohol <sup>2</sup>
<b>TOTAL</b>	24	29	20	35
<b>GENDER</b>				
Male	19	24	16	28
Female	15	17	12	20
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic				
White Only	21	24	17	30
Black Only	8	10	8	11
American Indian or Alaska Native Only	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian Only	2	3	2	3
More Than One Race	2	3	2	3
Hispanic	6	8	5	9
<b>ADULT EDUCATION</b>				
< High School	14	15	12	19
High School Graduate	14	19	11	22
Some College	14	16	12	19
College Graduate	3	4	2	5
<b>CURRENT EMPLOYMENT</b>				
Full-Time	18	23	15	26
Part-Time	11	12	8	14
Unemployed	8	9	8	10
Other <sup>3</sup>	8	9	5	14

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> "Received treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>2</sup> Estimates include persons who received treatment specifically for alcohol or any illicit drug, as well as persons who received treatment but did not specify for what substance(s).

<sup>3</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.40D Standard Errors of Percentages of Persons Aged 18 to 25 Who Received Substance Abuse Treatment in the Past Year, by Demographic Characteristics: 2000**

Demographic Characteristic	RECEIVED TREATMENT IN THE PAST YEAR <sup>1</sup>			
	Any Illicit Drug	Alcohol	Both Any Illicit Drug and Alcohol	Any Illicit Drug or Alcohol <sup>2</sup>
<b>TOTAL</b>	0.08	0.10	0.07	0.12
<b>GENDER</b>				
Male	0.13	0.16	0.11	0.19
Female	0.10	0.12	0.08	0.14
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic				
White Only	0.11	0.13	0.09	0.16
Black Only	0.21	0.26	0.21	0.29
American Indian or Alaska Native Only	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian Only	0.23	0.31	0.23	0.31
More Than One Race	0.81	0.86	0.81	0.86
Hispanic	0.13	0.18	0.11	0.20
<b>ADULT EDUCATION</b>				
< High School	0.22	0.25	0.19	0.32
High School Graduate	0.13	0.19	0.11	0.21
Some College	0.15	0.18	0.13	0.21
College Graduate	0.10	0.12	0.07	0.14
<b>CURRENT EMPLOYMENT</b>				
Full-Time	0.12	0.15	0.10	0.17
Part-Time	0.16	0.17	0.11	0.21
Unemployed	0.49	0.57	0.48	0.64
Other <sup>3</sup>	0.16	0.17	0.10	0.25

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> "Received treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>2</sup> Estimates include persons who received treatment specifically for alcohol or any illicit drug, as well as persons who received treatment but did not specify for what substance(s).

<sup>3</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.



**Table 5.41C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 26 or Older Who Received Substance Abuse Treatment in the Past Year, by Demographic Characteristics: 2000**

Demographic Characteristic	RECEIVED TREATMENT IN THE PAST YEAR <sup>1</sup>			
	Any Illicit Drug	Alcohol	Both Any Illicit Drug and Alcohol	Any Illicit Drug or Alcohol <sup>2</sup>
<b>TOTAL</b>	99	142	88	159
<b>GENDER</b>				
Male	55	116	50	129
Female	83	82	72	91
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic				
White Only	89	125	76	139
Black Only	41	54	40	58
American Indian or Alaska Native Only	3	11	3	11
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian Only	*	10	*	10
More Than One Race	12	13	12	13
Hispanic	16	36	15	39
<b>ADULT EDUCATION</b>				
< High School	62	75	58	83
High School Graduate	47	92	43	101
Some College	48	64	42	71
College Graduate	36	45	30	50
<b>CURRENT EMPLOYMENT</b>				
Full-Time	63	110	56	116
Part-Time	52	59	52	60
Unemployed	22	25	21	46
Other <sup>3</sup>	46	63	39	71

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> "Received treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>2</sup> Estimates include persons who received treatment specifically for alcohol or any illicit drug, as well as persons who received treatment but did not specify for what substance(s).

<sup>3</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.41D Standard Errors of Percentages of Persons Aged 26 or Older Who Received Substance Abuse Treatment in the Past Year, by Demographic Characteristics: 2000**

Demographic Characteristic	RECEIVED TREATMENT IN THE PAST YEAR <sup>1</sup>			
	Any Illicit Drug	Alcohol	Both Any Illicit Drug and Alcohol	Any Illicit Drug or Alcohol <sup>2</sup>
<b>TOTAL</b>	0.06	0.08	0.05	0.09
<b>GENDER</b>				
Male	0.07	0.14	0.06	0.16
Female	0.09	0.09	0.08	0.10
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic				
White Only	0.07	0.10	0.06	0.11
Black Only	0.22	0.30	0.22	0.32
American Indian or Alaska Native Only	0.42	1.37	0.42	1.39
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian Only	*	0.17	*	0.17
More Than One Race	1.04	1.05	1.04	1.05
Hispanic	0.10	0.22	0.09	0.24
<b>ADULT EDUCATION</b>				
< High School	0.21	0.26	0.20	0.28
High School Graduate	0.08	0.16	0.08	0.18
Some College	0.12	0.16	0.11	0.18
College Graduate	0.08	0.10	0.06	0.11
<b>CURRENT EMPLOYMENT</b>				
Full-Time	0.06	0.11	0.06	0.12
Part-Time	0.31	0.35	0.31	0.36
Unemployed	0.72	0.81	0.67	1.52
Other <sup>3</sup>	0.09	0.13	0.08	0.14

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> "Received treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>2</sup> Estimates include persons who received treatment specifically for alcohol or any illicit drug, as well as persons who received treatment but did not specify for what substance(s).

<sup>3</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.42C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 18 or Older Who Received Substance Abuse Treatment in the Past Year, by Demographic Characteristics: 2000**

Demographic Characteristic	RECEIVED TREATMENT IN THE PAST YEAR <sup>1</sup>			
	Any Illicit Drug	Alcohol	Both Any Illicit Drug and Alcohol	Any Illicit Drug or Alcohol <sup>2</sup>
<b>TOTAL</b>	105	149	94	167
<b>GENDER</b>				
Male	59	120	53	134
Female	84	82	73	93
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic				
White Only	92	131	79	147
Black Only	42	54	41	58
American Indian or Alaska Native Only	4	15	4	15
Native Hawaiian or Other Pacific Islander	2	2	2	4
Asian Only	2	10	2	10
More Than One Race	13	13	13	13
Hispanic	17	37	16	39
<b>ADULT EDUCATION</b>				
< High School	64	77	59	86
High School Graduate	49	94	44	105
Some College	51	66	44	74
College Graduate	36	45	30	51
<b>CURRENT EMPLOYMENT</b>				
Full-Time	68	115	62	122
Part-Time	53	60	53	63
Unemployed	23	27	22	48
Other <sup>3</sup>	47	64	39	72

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> "Received treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>2</sup> Estimates include persons who received treatment specifically for alcohol or any illicit drug, as well as persons who received treatment but did not specify for what substance(s).

<sup>3</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.42D Standard Errors of Percentages of Persons Aged 18 or Older Who Received Substance Abuse Treatment in the Past Year, by Demographic Characteristics: 2000**

Demographic Characteristic	RECEIVED TREATMENT IN THE PAST YEAR <sup>1</sup>			
	Any Illicit Drug	Alcohol	Both Any Illicit Drug and Alcohol	Any Illicit Drug or Alcohol <sup>2</sup>
<b>TOTAL</b>	0.05	0.07	0.05	0.08
<b>GENDER</b>				
Male	0.06	0.13	0.06	0.14
Female	0.08	0.08	0.07	0.09
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic				
White Only	0.06	0.09	0.05	0.10
Black Only	0.19	0.25	0.19	0.26
American Indian or Alaska Native Only	0.38	1.58	0.38	1.59
Native Hawaiian or Other Pacific Islander	0.34	0.34	0.34	0.73
Asian Only	0.03	0.15	0.03	0.15
More Than One Race	0.84	0.85	0.84	0.85
Hispanic	0.08	0.18	0.08	0.19
<b>ADULT EDUCATION</b>				
< High School	0.18	0.22	0.17	0.24
High School Graduate	0.07	0.14	0.07	0.16
Some College	0.10	0.14	0.09	0.15
College Graduate	0.07	0.09	0.06	0.10
<b>CURRENT EMPLOYMENT</b>				
Full-Time	0.06	0.10	0.05	0.11
Part-Time	0.23	0.26	0.23	0.27
Unemployed	0.50	0.57	0.47	1.02
Other <sup>3</sup>	0.08	0.11	0.07	0.13

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> "Received treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>2</sup> Estimates include persons who received treatment specifically for alcohol or any illicit drug, as well as persons who received treatment but did not specify for what substance(s).

<sup>3</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.43C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 12 or Older Who Received Substance Abuse Treatment in the Past Year, by Geographic Characteristics: 2000**

Geographic Characterist	RECEIVED TREATMENT IN THE PAST YEAR <sup>1</sup>			
	Any Illicit Drug	Alcohol	Both Any Illicit Drug and Alcohol	Any Illicit Drug or Alcohol <sup>2</sup>
<b>TOTAL</b>	108	151	96	169
<b>GEOGRAPHIC DIVISION</b>				
Northeast	31	66	29	72
New England	24	22	17	28
Middle Atlantic	20	63	23	66
Midwest	53	67	41	82
East North Central	30	48	24	61
West North Central	43	47	34	55
South	50	76	40	83
South Atlantic	34	65	25	69
East South Central	12	19	9	21
West South Central	34	36	30	42
West	73	89	71	98
Mountain	23	33	22	33
Pacific	69	82	68	93
<b>COUNTY TYPE</b>				
Large Metro	91	125	86	132
Small Metro	40	59	35	75
250K - 1 Mil. Pop.	33	50	29	67
<250K Pop.	22	31	19	36
Nonmetro	38	59	25	71
Urbanized	23	25	21	35
Less Urbanized	29	52	13	60
Completely Rural	6	11	6	11

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> "Receiving treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>2</sup> Estimates include persons who received treatment specifically for alcohol or any illicit drug, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.43D Standard Errors of Percentages of Persons Aged 12 or Older Who Received Substance Abuse Treatment in the Past Year, by Geographic Characteristics: 2000**

Geographic Characteristic	RECEIVED TREATMENT IN THE PAST YEAR <sup>1</sup>			
	Any Illicit Drug	Alcohol	Both Any Illicit Drug and Alcohol	Any Illicit Drug or Alcohol <sup>2</sup>
<b>TOTAL</b>	0.05	0.07	0.04	0.08
<b>GEOGRAPHIC DIVISION</b>				
Northeast	0.07	0.15	0.07	0.17
New England	0.21	0.19	0.15	0.25
Middle Atlantic	0.06	0.20	0.07	0.21
Midwest	0.10	0.13	0.08	0.16
East North Central	0.08	0.13	0.07	0.17
West North Central	0.28	0.30	0.22	0.35
South	0.06	0.10	0.05	0.10
South Atlantic	0.08	0.16	0.06	0.17
East South Central	0.09	0.14	0.07	0.15
West South Central	0.14	0.15	0.12	0.17
West	0.15	0.18	0.14	0.20
Mountain	0.16	0.23	0.15	0.23
Pacific	0.20	0.24	0.19	0.27
<b>COUNTY TYPE</b>				
Large Metro	0.08	0.11	0.08	0.12
Small Metro	0.06	0.09	0.05	0.11
250K - 1 Mil. Pop.	0.06	0.10	0.06	0.13
<250K Pop.	0.13	0.17	0.11	0.20
Nonmetro	0.08	0.13	0.06	0.16
Urbanized	0.17	0.19	0.16	0.26
Less Urbanized	0.11	0.19	0.05	0.22
Completely Rural	0.11	0.20	0.11	0.21

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> "Receiving treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>2</sup> Estimates include persons who received treatment specifically for alcohol or any illicit drug, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.44C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 12 to 17 Who Received Substance Abuse Treatment in the Past Year, by Geographic Characteristics: 2000**

Geographic Characterist	RECEIVED TREATMENT IN THE PAST YEAR <sup>1</sup>			
	Any Illicit Drug	Alcohol	Both Any Illicit Drug and Alcohol	Any Illicit Drug or Alcohol <sup>2</sup>
<b>TOTAL</b>	20	18	16	24
<b>GEOGRAPHIC DIVISION</b>				
Northeast	9	7	7	11
New England	7	5	5	9
Middle Atlantic	5	5	5	7
Midwest	9	7	6	11
East North Central	7	6	6	8
West North Central	6	3	2	7
South	11	11	10	13
South Atlantic	9	8	7	10
East South Central	4	4	3	4
West South Central	6	6	6	7
West	11	10	9	13
Mountain	7	5	5	7
Pacific	9	9	8	11
<b>COUNTY TYPE</b>				
Large Metro	14	13	11	16
Small Metro	11	10	9	14
250K - 1 Mil. Pop.	7	7	6	9
<250K Pop.	8	7	7	10
Nonmetro	10	9	8	11
Urbanized	6	6	6	7
Less Urbanized	7	6	6	9
Completely Rural	1	1	1	2

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> "Receiving treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>2</sup> Estimates include persons who received treatment specifically for alcohol or any illicit drug, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.44D Standard Errors of Percentages of Persons Aged 12 to 17 Who Received Substance Abuse Treatment in the Past Year, by Geographic Characteristics: 2000**

Geographic Characteristic	RECEIVED TREATMENT IN THE PAST YEAR <sup>1</sup>			
	Any Illicit Drug	Alcohol	Both Any Illicit Drug and Alcohol	Any Illicit Drug or Alcohol <sup>2</sup>
<b>TOTAL</b>	0.09	0.08	0.07	0.10
<b>GEOGRAPHIC DIVISION</b>				
Northeast	0.20	0.18	0.16	0.26
New England	0.62	0.45	0.42	0.83
Middle Atlantic	0.17	0.18	0.15	0.21
Midwest	0.17	0.13	0.12	0.20
East North Central	0.18	0.16	0.16	0.22
West North Central	0.38	0.20	0.14	0.41
South	0.13	0.13	0.12	0.16
South Atlantic	0.21	0.20	0.18	0.24
East South Central	0.27	0.25	0.24	0.30
West South Central	0.21	0.22	0.20	0.26
West	0.21	0.19	0.17	0.24
Mountain	0.40	0.32	0.31	0.44
Pacific	0.24	0.23	0.20	0.29
<b>COUNTY TYPE</b>				
Large Metro	0.12	0.11	0.10	0.14
Small Metro	0.15	0.13	0.13	0.19
250K - 1 Mil. Pop.	0.14	0.13	0.11	0.17
<250K Pop.	0.44	0.37	0.37	0.56
Nonmetro	0.20	0.18	0.17	0.24
Urbanized	0.45	0.45	0.42	0.50
Less Urbanized	0.26	0.22	0.20	0.30
Completely Rural	0.11	0.11	0.11	0.31

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> "Receiving treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>2</sup> Estimates include persons who received treatment specifically for alcohol or any illicit drug, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.



**Table 5.45C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 18 to 25 Who Received Substance Abuse Treatment in the Past Year, by Geographic Characteristics: 2000**

Geographic Characterist	RECEIVED TREATMENT IN THE PAST YEAR <sup>1</sup>			
	Any Illicit Drug	Alcohol	Both Any Illicit Drug and Alcohol	Any Illicit Drug or Alcohol <sup>2</sup>
<b>TOTAL</b>	24	29	20	35
<b>GEOGRAPHIC DIVISION</b>				
Northeast	10	9	8	10
New England	4	5	3	5
Middle Atlantic	9	8	7	9
Midwest	9	16	9	18
East North Central	7	14	6	16
West North Central	6	8	6	9
South	13	13	9	18
South Atlantic	8	9	6	13
East South Central	6	7	5	8
West South Central	8	6	5	9
West	15	18	13	20
Mountain	10	11	9	12
Pacific	11	14	10	17
<b>COUNTY TYPE</b>				
Large Metro	19	22	16	26
Small Metro	11	14	9	17
250K - 1 Mil. Pop.	9	11	8	13
<250K Pop.	7	8	4	11
Nonmetro	9	14	8	15
Urbanized	5	6	5	8
Less Urbanized	7	11	6	13
Completely Rural	2	4	2	4

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> "Receiving treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>2</sup> Estimates include persons who received treatment specifically for alcohol or any illicit drug, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.45D Standard Errors of Percentages of Persons Aged 18 to 25 Who Received Substance Abuse Treatment in the Past Year, by Geographic Characteristics: 2000**

Geographic Characteristic	RECEIVED TREATMENT IN THE PAST YEAR <sup>1</sup>			
	Any Illicit Drug	Alcohol	Both Any Illicit Drug and Alcohol	Any Illicit Drug or Alcohol <sup>2</sup>
<b>TOTAL</b>	0.08	0.10	0.07	0.12
<b>GEOGRAPHIC DIVISION</b>				
Northeast	0.20	0.18	0.15	0.21
New England	0.32	0.34	0.25	0.39
Middle Atlantic	0.24	0.20	0.19	0.24
Midwest	0.13	0.24	0.12	0.27
East North Central	0.14	0.30	0.12	0.34
West North Central	0.29	0.38	0.29	0.41
South	0.12	0.12	0.09	0.17
South Atlantic	0.16	0.18	0.11	0.26
East South Central	0.33	0.38	0.29	0.45
West South Central	0.22	0.16	0.13	0.27
West	0.22	0.27	0.20	0.31
Mountain	0.49	0.56	0.46	0.59
Pacific	0.23	0.30	0.21	0.36
<b>COUNTY TYPE</b>				
Large Metro	0.14	0.16	0.12	0.19
Small Metro	0.12	0.14	0.09	0.18
250K - 1 Mil. Pop.	0.14	0.17	0.12	0.20
<250K Pop.	0.23	0.28	0.15	0.40
Nonmetro	0.16	0.24	0.14	0.28
Urbanized	0.28	0.32	0.27	0.41
Less Urbanized	0.21	0.33	0.18	0.39
Completely Rural	0.39	0.89	0.34	0.93

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> "Receiving treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>2</sup> Estimates include persons who received treatment specifically for alcohol or any illicit drug, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.46C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 26 or Older Who Received Substance Abuse Treatment in the Past Year, by Geographic Characteristics: 2000**

Geographic Characterist	RECEIVED TREATMENT IN THE PAST YEAR <sup>1</sup>			
	Any Illicit Drug	Alcohol	Both Any Illicit Drug and Alcohol	Any Illicit Drug or Alcohol <sup>2</sup>
<b>TOTAL</b>	99	142	88	159
<b>GEOGRAPHIC DIVISION</b>				
Northeast	28	66	26	69
New England	22	20	16	25
Middle Atlantic	17	63	20	65
Midwest	48	62	37	76
East North Central	28	44	22	56
West North Central	39	43	30	51
South	45	72	37	79
South Atlantic	31	60	23	64
East South Central	9	17	7	18
West South Central	32	37	29	41
West	68	82	66	92
Mountain	17	29	16	30
Pacific	66	77	64	87
<b>COUNTY TYPE</b>				
Large Metro	85	117	79	122
Small Metro	35	57	31	73
250K - 1 Mil. Pop.	30	49	26	66
<250K Pop.	19	29	17	31
Nonmetro	35	55	22	66
Urbanized	22	24	19	33
Less Urbanized	26	49	8	57
Completely Rural	5	10	6	10

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> "Receiving treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>2</sup> Estimates include persons who received treatment specifically for alcohol or any illicit drug, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.46D Standard Errors of Percentages of Persons Aged 26 or Older Who Received Substance Abuse Treatment in the Past Year, by Geographic Characteristics: 2000**

Geographic Characteristic	RECEIVED TREATMENT IN THE PAST YEAR <sup>1</sup>			
	Any Illicit Drug	Alcohol	Both Any Illicit Drug and Alcohol	Any Illicit Drug or Alcohol <sup>2</sup>
<b>TOTAL</b>	0.06	0.08	0.05	0.09
<b>GEOGRAPHIC DIVISION</b>				
Northeast	0.08	0.20	0.08	0.21
New England	0.25	0.23	0.19	0.28
Middle Atlantic	0.07	0.26	0.08	0.26
Midwest	0.12	0.16	0.09	0.19
East North Central	0.10	0.16	0.08	0.20
West North Central	0.33	0.37	0.25	0.43
South	0.07	0.12	0.06	0.13
South Atlantic	0.10	0.19	0.07	0.20
East South Central	0.08	0.16	0.07	0.17
West South Central	0.18	0.21	0.16	0.23
West	0.18	0.22	0.18	0.25
Mountain	0.16	0.27	0.15	0.28
Pacific	0.25	0.29	0.24	0.33
<b>COUNTY TYPE</b>				
Large Metro	0.10	0.14	0.09	0.15
Small Metro	0.07	0.11	0.06	0.14
250K - 1 Mil. Pop.	0.08	0.13	0.07	0.17
<250K Pop.	0.15	0.22	0.13	0.24
Nonmetro	0.10	0.16	0.06	0.19
Urbanized	0.21	0.23	0.19	0.32
Less Urbanized	0.13	0.24	0.04	0.27
Completely Rural	0.13	0.25	0.14	0.25

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> "Receiving treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>2</sup> Estimates include persons who received treatment specifically for alcohol or any illicit drug, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.47C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 18 or Older Who Received Substance Abuse Treatment in the Past Year, by Geographic Characteristics: 2000**

Geographic Characterist	RECEIVED TREATMENT IN THE PAST YEAR <sup>1</sup>			
	Any Illicit Drug	Alcohol	Both Any Illicit Drug and Alcohol	Any Illicit Drug or Alcohol <sup>2</sup>
<b>TOTAL</b>	105	149	94	167
<b>GEOGRAPHIC DIVISION</b>				
Northeast	30	66	27	73
New England	22	21	17	25
Middle Atlantic	20	63	22	68
Midwest	52	67	41	80
East North Central	29	48	23	59
West North Central	43	47	34	54
South	48	73	38	81
South Atlantic	32	61	23	66
East South Central	11	19	9	20
West South Central	33	36	29	41
West	72	89	70	98
Mountain	23	34	22	35
Pacific	68	82	66	91
<b>COUNTY TYPE</b>				
Large Metro	90	124	84	131
Small Metro	38	58	33	74
250K - 1 Mil. Pop.	32	50	28	67
<250K Pop.	20	30	18	33
Nonmetro	36	57	23	68
Urbanized	22	25	20	34
Less Urbanized	27	50	10	58
Completely Rural	6	11	6	11

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> "Receiving treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>2</sup> Estimates include persons who received treatment specifically for alcohol or any illicit drug, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.47D Standard Errors of Percentages of Persons Aged 18 or Older Who Received Substance Abuse Treatment in the Past Year, by Geographic Characteristics: 2000**

Geographic Characteristic	RECEIVED TREATMENT IN THE PAST YEAR <sup>1</sup>			
	Any Illicit Drug	Alcohol	Both Any Illicit Drug and Alcohol	Any Illicit Drug or Alcohol <sup>2</sup>
<b>TOTAL</b>	0.05	0.07	0.05	0.08
<b>GEOGRAPHIC DIVISION</b>				
Northeast	0.08	0.17	0.07	0.19
New England	0.22	0.20	0.17	0.25
Middle Atlantic	0.07	0.22	0.08	0.24
Midwest	0.11	0.14	0.09	0.17
East North Central	0.09	0.15	0.07	0.18
West North Central	0.31	0.34	0.24	0.39
South	0.07	0.10	0.05	0.11
South Atlantic	0.09	0.17	0.06	0.18
East South Central	0.09	0.15	0.07	0.16
West South Central	0.15	0.17	0.13	0.19
West	0.16	0.20	0.16	0.22
Mountain	0.18	0.27	0.17	0.28
Pacific	0.22	0.26	0.21	0.29
<b>COUNTY TYPE</b>				
Large Metro	0.09	0.13	0.09	0.13
Small Metro	0.06	0.09	0.05	0.12
250K - 1 Mil. Pop.	0.07	0.11	0.06	0.15
<250K Pop.	0.13	0.19	0.11	0.21
Nonmetro	0.09	0.14	0.06	0.17
Urbanized	0.19	0.21	0.17	0.29
Less Urbanized	0.11	0.21	0.04	0.24
Completely Rural	0.12	0.23	0.13	0.23

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> "Receiving treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>2</sup> Estimates include persons who received treatment specifically for alcohol or any illicit drug, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.48C Standard Errors of Estimated Numbers (in Thousands) of Persons Who Needed and Received Treatment for an Illicit Drug Problem in the Past Year, by Detailed Age Categories: 2000**

Age Category	NEEDED TREATMENT FOR AN ILLICIT DRUG PROBLEM IN THE PAST YEAR			Percentage Who Received Treatment at a Specialty Facility Among Persons Who Needed Treatment
	Total	Received Treatment at a Specialty Facility	Did Not Receive Treatment at a Specialty Facility	
Total	174	86	154	1.65
12	7	0	7	*
13	11	3	11	*
14	12	4	12	2.85
15	18	5	18	2.02
16	19	7	18	2.17
17	21	8	20	2.43
18	26	7	25	1.81
19	25	3	25	1.07
20	24	8	22	2.79
21	19	5	19	2.45
22	18	6	17	3.30
23	16	6	15	*
24	15	5	14	*
25	13	3	13	*
26-29	41	21	35	*
30-34	45	20	38	4.53
35-39	67	25	63	*
40-44	62	31	53	*
45-49	68	25	64	*
50-54	75	60	45	*
55-59	26	10	24	*
60-64	*	*	*	*
65 or Older	24	2	24	*

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on any illicit drug; (2) abuse of any illicit drug; or (3) received treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities (inpatient or outpatient), hospitals (inpatient only), and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type psychotherapeutic (nonmedical use).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.48D Standard Errors of Percentages of Persons Who Needed and Received Treatment for an Illicit Drug Problem in the Past Year, by Detailed Age Categories: 2000**

Age Category	NEEDED TREATMENT FOR AN ILLICIT DRUG PROBLEM IN THE PAST YEAR			Percentage Who Received Treatment at a Specialty Facility Among Persons Who Needed Treatment
	Total	Received Treatment at a Specialty Facility	Did Not Receive Treatment at a Specialty Facility	
Total	0.08	0.04	0.07	1.65
12	0.18	0.01	0.18	*
13	0.28	0.08	0.27	*
14	0.30	0.09	0.29	2.85
15	0.46	0.12	0.44	2.02
16	0.49	0.17	0.45	2.17
17	0.59	0.22	0.55	2.43
18	0.59	0.15	0.58	1.81
19	0.64	0.09	0.63	1.07
20	0.61	0.21	0.58	2.79
21	0.55	0.14	0.54	2.45
22	0.53	0.18	0.50	3.30
23	0.46	0.17	0.43	*
24	0.46	0.15	0.44	*
25	0.39	0.08	0.38	*
26-29	0.29	0.14	0.25	*
30-34	0.24	0.11	0.20	4.53
35-39	0.31	0.12	0.29	*
40-44	0.28	0.14	0.24	*
45-49	0.34	0.12	0.32	*
50-54	0.39	0.32	0.24	*
55-59	0.19	0.07	0.18	*
60-64	*	*	*	*
65 or Older	0.08	0.01	0.08	*

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on any illicit drug; (2) abuse of any illicit drug; or (3) received treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities (inpatient or outpatient), hospitals (inpatient only), and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type psychotherapeutic (nonmedical use).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.



10913

**Table 5.49C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 12 or Older Who Needed and Received Treatment for an Illicit Drug Problem in the Past Year, by Demographic Characteristics: 2000**

Demographic Characteristic	NEEDED TREATMENT FOR AN ILLICIT DRUG PROBLEM IN THE PAST YEAR			Percentage Who Received Treatment at a Specialty Facility Among Persons Who Needed Treatment
	Total	Received Treatment at a Specialty Facility	Did Not Receive Treatment at a Specialty Facility	
<b>TOTAL</b>	174	86	154	1.65
<b>AGE</b>				
12-17	39	13	36	1.12
18-25	56	16	54	0.94
26 or Older	157	81	136	3.64
<b>GENDER</b>				
Male	124	48	115	1.61
Female	127	71	101	3.11
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic				
White Only	140	74	122	2.04
Black Only	65	37	53	*
American Indian or Alaska Native Only	11	3	11	*
Native Hawaiian or Other Pacific Islander	5	2	4	*
Asian Only	13	1	13	*
More Than One Race	40	12	38	*
Hispanic	60	15	56	2.34

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on any illicit drug; (2) abuse of any illicit drug; or (3) received treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities (inpatient or outpatient), hospitals (inpatient only), and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type psychotherapeutic (nonmedical use).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.49D Standard Errors of Percentages of Persons Aged 12 or Older Who Needed and Received Treatment for an Illicit Drug Problem in the Past Year, by Demographic Characteristics: 2000**

Demographic Characteristic	NEEDED TREATMENT FOR AN ILLICIT DRUG PROBLEM IN THE PAST YEAR			Percentage Who Received Treatment at a Specialty Facility Among Persons Who Needed Treatment
	Total	Received Treatment at a Specialty Facility	Did Not Receive Treatment at a Specialty Facility	
<b>TOTAL</b>	0.08	0.04	0.07	1.65
<b>AGE</b>				
12-17	0.17	0.06	0.15	1.12
18-25	0.19	0.06	0.19	0.94
26 or Older	0.09	0.05	0.08	3.64
<b>GENDER</b>				
Male	0.12	0.04	0.11	1.61
Female	0.11	0.06	0.09	3.11
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic				
White Only	0.09	0.05	0.08	2.04
Black Only	0.26	0.15	0.21	*
American Indian or Alaska Native Only	1.03	0.26	0.99	*
Native Hawaiian or Other Pacific Islander	0.85	0.30	0.76	*
Asian Only	0.16	0.01	0.16	*
More Than One Race	2.13	0.67	2.06	*
Hispanic	0.25	0.06	0.23	2.34

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on any illicit drug; (2) abuse of any illicit drug; or (3) received treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities (inpatient or outpatient), hospitals (inpatient only), and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type psychotherapeutic (nonmedical use).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.50C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 12 to 17 Who Needed and Received Treatment for an Illicit Drug Problem in the Past Year, by Demographic Characteristics: 2000**

Demographic Characteristic	NEEDED TREATMENT FOR AN ILLICIT DRUG PROBLEM IN THE PAST YEAR			Percentage Who Received Treatment at a Specialty Facility Among Persons Who Needed Treatment
	Total	Received Treatment at a Specialty Facility	Did Not Receive Treatment at a Specialty Facility	
<b>TOTAL</b>	39	13	36	1.12
<b>GENDER</b>				
Male	29	11	27	1.67
Female	25	7	24	1.45
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic				
White Only	32	11	30	1.36
Black Only	13	5	12	3.62
American Indian or Alaska Native Only	4	*	3	*
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian Only	6	*	6	*
More Than One Race	3	*	3	*
Hispanic	17	4	16	2.05
<b>GENDER/RACE/HISPANIC ORIGIN</b>				
Male - White	25	8	23	2.00
Female - White	20	7	19	1.83
Male - Black	11	4	10	*
Female - Black	7	1	7	*
Male - Hispanic	12	3	12	*
Female - Hispanic	10	2	10	*

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on any illicit drug; (2) abuse of any illicit drug; or (3) received treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities (inpatient or outpatient), hospitals (inpatient only), and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type psychotherapeutic (nonmedical use).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.50D Standard Errors of Percentages of Persons Aged 12 to 17 Who Needed and Received Treatment for an Illicit Drug Problem in the Past Year, by Demographic Characteristics: 2000**

Demographic Characteristic	NEEDED TREATMENT FOR AN ILLICIT DRUG PROBLEM IN THE PAST YEAR			Percentage Who Received Treatment at a Specialty Facility Among Persons Who Needed Treatment
	Total	Received Treatment at a Specialty Facility	Did Not Receive Treatment at a Specialty Facility	
<b>TOTAL</b>	0.17	0.06	0.15	1.12
<b>GENDER</b>				
Male	0.24	0.09	0.23	1.67
Female	0.22	0.06	0.21	1.45
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic				
White Only	0.21	0.07	0.20	1.36
Black Only	0.38	0.14	0.35	3.62
American Indian or Alaska Native Only	2.78	*	2.29	*
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian Only	0.75	*	0.75	*
More Than One Race	0.81	*	0.81	*
Hispanic	0.50	0.11	0.49	2.05
<b>GENDER/RACE/HISPANIC ORIGIN</b>				
Male - White	0.31	0.11	0.30	2.00
Female - White	0.27	0.09	0.25	1.83
Male - Black	0.64	0.26	0.59	*
Female - Black	0.45	0.09	0.43	*
Male - Hispanic	0.71	0.16	0.72	*
Female - Hispanic	0.66	0.15	0.65	*

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on any illicit drug; (2) abuse of any illicit drug; or (3) received treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities (inpatient or outpatient), hospitals (inpatient only), and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type psychotherapeutic (nonmedical use).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.51C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 18 to 25 Who Needed and Received Treatment for an Illicit Drug Problem in the Past Year, by Demographic Characteristics: 2000**

Demographic Characteristic	NEEDED TREATMENT FOR AN ILLICIT DRUG PROBLEM IN THE PAST YEAR			Percentage Who Received Treatment at a Specialty Facility Among Persons Who Needed Treatment
	Total	Received Treatment at a Specialty Facility	Did Not Receive Treatment at a Specialty Facility	
<b>TOTAL</b>	56	16	54	0.94
<b>GENDER</b>				
Male	45	13	44	1.19
Female	34	10	32	1.56
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic				
White Only	49	14	47	1.13
Black Only	20	5	19	2.40
American Indian or Alaska Native Only	*	0	*	*
Native Hawaiian or Other Pacific Islander	3	*	*	*
Asian Only	5	1	5	*
More Than One Race	6	1	6	*
Hispanic	21	4	20	2.43
<b>ADULT EDUCATION</b>				
< High School	31	11	30	1.96
High School Graduate	37	9	36	1.38
Some College	31	7	30	1.68
College Graduate	12	2	12	*
<b>CURRENT EMPLOYMENT</b>				
Full-Time	44	12	42	1.37
Part-Time	28	5	27	1.45
Unemployed	20	7	19	3.36
Other <sup>1</sup>	21	6	20	2.31

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on any illicit drug; (2) abuse of any illicit drug; or (3) received treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities (inpatient or outpatient), hospitals (inpatient only), and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type psychotherapeutic (nonmedical use).

<sup>1</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.51D Standard Errors of Percentages of Persons Aged 18 to 25 Who Needed and Received Treatment for an Illicit Drug Problem in the Past Year, by Demographic Characteristics: 2000**

Demographic Characteristic	NEEDED TREATMENT FOR AN ILLICIT DRUG PROBLEM IN THE PAST YEAR			Percentage Who Received Treatment at a Specialty Facility Among Persons Who Needed Treatment
	Total	Received Treatment at a Specialty Facility	Did Not Receive Treatment at a Specialty Facility	
<b>TOTAL</b>	0.19	0.06	0.19	0.94
<b>GENDER</b>				
Male	0.31	0.09	0.30	1.19
Female	0.24	0.07	0.22	1.56
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic				
White Only	0.25	0.07	0.24	1.13
Black Only	0.51	0.14	0.48	2.40
American Indian or Alaska Native Only	*	0.10	*	*
Native Hawaiian or Other Pacific Islander	2.45	*	*	*
Asian Only	0.54	0.07	0.53	*
More Than One Race	2.07	0.38	2.04	*
Hispanic	0.48	0.10	0.46	2.43
<b>ADULT EDUCATION</b>				
< High School	0.51	0.18	0.49	1.96
High School Graduate	0.36	0.08	0.35	1.38
Some College	0.34	0.08	0.34	1.68
College Graduate	0.35	0.05	0.34	*
<b>CURRENT EMPLOYMENT</b>				
Full-Time	0.28	0.08	0.27	1.37
Part-Time	0.42	0.08	0.41	1.45
Unemployed	1.23	0.43	1.16	3.36
Other <sup>1</sup>	0.39	0.11	0.38	2.31

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on any illicit drug; (2) abuse of any illicit drug; or (3) received treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities (inpatient or outpatient), hospitals (inpatient only), and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type psychotherapeutic (nonmedical use).

<sup>1</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.52C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 26 or Older Who Needed and Received Treatment for an Illicit Drug Problem in the Past Year, by Demographic Characteristics: 2000**

Demographic Characteristic	NEEDED TREATMENT FOR AN ILLICIT DRUG PROBLEM IN THE PAST YEAR			Percentage Who Received Treatment at a Specialty Facility Among Persons Who Needed Treatment
	Total	Received Treatment at a Specialty Facility	Did Not Receive Treatment at a Specialty Facility	
<b>TOTAL</b>	157	81	136	3.64
<b>GENDER</b>				
Male	108	44	99	3.68
Female	120	70	94	*
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic				
White Only	127	72	104	4.50
Black Only	60	36	49	*
American Indian or Alaska Native Only	8	1	8	*
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian Only	9	*	9	*
More Than One Race	*	12	*	*
Hispanic	52	13	48	*
<b>ADULT EDUCATION</b>				
< High School	85	61	59	*
High School Graduate	67	42	53	*
Some College	66	31	61	*
College Graduate	84	20	82	*
<b>CURRENT EMPLOYMENT</b>				
Full-Time	98	37	93	3.43
Part-Time	66	51	44	*
Unemployed	40	21	34	*
Other <sup>1</sup>	83	46	70	*

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on any illicit drug; (2) abuse of any illicit drug; or (3) received treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities (inpatient or outpatient), hospitals (inpatient only), and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type psychotherapeutic (nonmedical use).

<sup>1</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.52D Standard Errors of Percentages of Persons Aged 26 or Older Who Needed and Received Treatment for an Illicit Drug Problem in the Past Year, by Demographic Characteristics: 2000**

Demographic Characteristic	NEEDED TREATMENT FOR AN ILLICIT DRUG PROBLEM IN THE PAST YEAR			Percentage Who Received Treatment at a Specialty Facility Among Persons Who Needed Treatment
	Total	Received Treatment at a Specialty Facility	Did Not Receive Treatment at a Specialty Facility	
<b>TOTAL</b>	0.09	0.05	0.08	3.64
<b>GENDER</b>				
Male	0.13	0.05	0.12	3.68
Female	0.13	0.08	0.10	*
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic				
White Only	0.10	0.06	0.08	4.50
Black Only	0.33	0.20	0.27	*
American Indian or Alaska Native Only	1.10	0.15	1.09	*
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian Only	0.16	*	0.16	*
More Than One Race	*	1.04	*	*
Hispanic	0.32	0.08	0.30	*
<b>ADULT EDUCATION</b>				
< High School	0.29	0.21	0.20	*
High School Graduate	0.12	0.07	0.09	*
Some College	0.17	0.08	0.16	*
College Graduate	0.18	0.04	0.18	*
<b>CURRENT EMPLOYMENT</b>				
Full-Time	0.10	0.04	0.09	3.43
Part-Time	0.40	0.31	0.26	*
Unemployed	1.32	0.70	1.12	*
Other <sup>1</sup>	0.16	0.09	0.14	*

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on any illicit drug; (2) abuse of any illicit drug; or (3) received treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities (inpatient or outpatient), hospitals (inpatient only), and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type psychotherapeutic (nonmedical use).

<sup>1</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.



**Table 5.53C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 18 or Older Who Needed and Received Treatment for an Illicit Drug Problem in the Past Year, by Demographic Characteristics: 2000**

Demographic Characteristic	NEEDED TREATMENT FOR AN ILLICIT DRUG PROBLEM IN THE PAST YEAR			Percentage Who Received Treatment at a Specialty Facility Among Persons Who Needed Treatment
	Total	Received Treatment at a Specialty Facility	Did Not Receive Treatment at a Specialty Facility	
<b>TOTAL</b>	171	85	150	2.09
<b>GENDER</b>				
Male	120	47	110	1.99
Female	126	71	100	4.05
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic				
White Only	137	74	117	2.55
Black Only	65	36	53	*
American Indian or Alaska Native Only	10	1	10	*
Native Hawaiian or Other Pacific Islander	4	2	4	*
Asian Only	10	1	10	*
More Than One Race	39	12	38	*
Hispanic	56	14	52	3.21
<b>ADULT EDUCATION</b>				
< High School	94	62	68	4.78
High School Graduate	81	43	68	3.25
Some College	73	33	68	3.39
College Graduate	85	20	83	*
<b>CURRENT EMPLOYMENT</b>				
Full-Time	112	39	107	2.00
Part-Time	72	51	52	*
Unemployed	46	22	40	*
Other <sup>1</sup>	86	46	73	5.37

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on any illicit drug; (2) abuse of any illicit drug; or (3) received treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities (inpatient or outpatient), hospitals (inpatient only), and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type psychotherapeutic (nonmedical use).

<sup>1</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.53D Standard Errors of Percentages of Persons Aged 18 or Older Who Needed and Received Treatment for an Illicit Drug Problem in the Past Year, by Demographic Characteristics: 2000**

Demographic Characteristic	NEEDED TREATMENT FOR AN ILLICIT DRUG PROBLEM IN THE PAST YEAR			Percentage Who Received Treatment at a Specialty Facility Among Persons Who Needed Treatment
	Total	Received Treatment at a Specialty Facility	Did Not Receive Treatment at a Specialty Facility	
<b>TOTAL</b>	0.09	0.04	0.08	2.09
<b>GENDER</b>				
Male	0.13	0.05	0.12	1.99
Female	0.12	0.07	0.10	4.05
<b>HISPANIC ORIGIN AND RACE</b>				
Not Hispanic				
White Only	0.09	0.05	0.08	2.55
Black Only	0.30	0.17	0.24	*
American Indian or Alaska Native Only	1.11	0.12	1.10	*
Native Hawaiian or Other Pacific Islander	0.89	0.34	0.82	*
Asian Only	0.15	0.01	0.15	*
More Than One Race	2.62	0.83	2.54	*
Hispanic	0.27	0.07	0.25	3.21
<b>ADULT EDUCATION</b>				
< High School	0.26	0.18	0.19	4.78
High School Graduate	0.12	0.06	0.10	3.25
Some College	0.15	0.07	0.14	3.39
College Graduate	0.17	0.04	0.17	*
<b>CURRENT EMPLOYMENT</b>				
Full-Time	0.10	0.03	0.09	2.00
Part-Time	0.31	0.22	0.22	*
Unemployed	0.99	0.48	0.86	*
Other <sup>1</sup>	0.15	0.08	0.13	5.37

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on any illicit drug; (2) abuse of any illicit drug; or (3) received treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities (inpatient or outpatient), hospitals (inpatient only), and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type psychotherapeutic (nonmedical use).

<sup>1</sup> Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.54C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 12 or Older Who Needed and Received Treatment for an Illicit Drug Problem in the Past Year, by Geographic Characteristics: 2000**

Geographic Characteristic	NEEDED TREATMENT FOR AN ILLICIT DRUG PROBLEM IN THE PAST YEAR			Percentage Who Received Treatment at a Specialty Facility Among Persons Who Needed Treatment
	Total	Received Treatment at a Specialty Facility	Did Not Receive Treatment at a Specialty Facility	
<b>TOTAL</b>	174	86	154	1.65
<b>GEOGRAPHIC DIVISION</b>				
Northeast	78	28	73	2.60
New England	49	22	44	*
Middle Atlantic	60	18	58	2.67
Midwest	69	32	61	3.02
East North Central	54	24	47	3.11
West North Central	44	22	39	*
South	78	40	69	2.62
South Atlantic	62	32	58	4.18
East South Central	26	9	22	3.59
West South Central	40	23	30	4.29
West	115	63	99	4.23
Mountain	43	17	39	3.94
Pacific	106	60	91	*
<b>COUNTY TYPE</b>				
Large Metro	142	70	128	2.55
Small Metro	76	33	66	2.10
250K - 1 Mil. Pop.	63	27	56	2.38
<250K Pop.	43	19	36	4.35
Nonmetro	62	34	52	3.73
Urbanized	27	21	20	*
Less Urbanized	54	26	47	*
Completely Rural	13	6	11	*

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on any illicit drug; (2) abuse of any illicit drug; or (3) received treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities (inpatient or outpatient), hospitals (inpatient only), and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type psychotherapeutic (nonmedical use).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.54D Standard Errors of Percentages of Persons Aged 12 or Older Who Needed and Received Treatment for an Illicit Drug Problem in the Past Year, by Geographic Characteristics: 2000**

Geographic Characteristic	NEEDED TREATMENT FOR AN ILLICIT DRUG PROBLEM IN THE PAST YEAR			Percentage Who Received Treatment at a Specialty Facility Among Persons Who Needed Treatment
	Total	Received Treatment at a Specialty Facility	Did Not Receive Treatment at a Specialty Facility	
<b>TOTAL</b>	0.08	0.04	0.07	1.65
<b>GEOGRAPHIC DIVISION</b>				
Northeast	0.18	0.07	0.17	2.60
New England	0.44	0.19	0.40	*
Middle Atlantic	0.19	0.06	0.18	2.67
Midwest	0.13	0.06	0.12	3.02
East North Central	0.15	0.06	0.13	3.11
West North Central	0.28	0.14	0.25	*
South	0.10	0.05	0.09	2.62
South Atlantic	0.15	0.08	0.14	4.18
East South Central	0.19	0.06	0.16	3.59
West South Central	0.16	0.09	0.12	4.29
West	0.23	0.13	0.20	4.23
Mountain	0.30	0.12	0.27	3.94
Pacific	0.30	0.17	0.26	*
<b>COUNTY TYPE</b>				
Large Metro	0.13	0.06	0.12	2.55
Small Metro	0.11	0.05	0.10	2.10
250K - 1 Mil. Pop.	0.12	0.05	0.11	2.38
<250K Pop.	0.24	0.10	0.20	4.35
Nonmetro	0.14	0.08	0.11	3.73
Urbanized	0.20	0.16	0.15	*
Less Urbanized	0.20	0.10	0.17	*
Completely Rural	0.24	0.11	0.21	*

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on any illicit drug; (2) abuse of any illicit drug; or (3) received treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities (inpatient or outpatient), hospitals (inpatient only), and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type psychotherapeutic (nonmedical use).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.55C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 12 to 17 Who Needed and Received Treatment for an Illicit Drug Problem in the Past Year, by Geographic Characteristics: 2000**

Geographic Characteristic	NEEDED TREATMENT FOR AN ILLICIT DRUG PROBLEM IN THE PAST YEAR			Percentage Who Received Treatment at a Specialty Facility Among Persons Who Needed Treatment
	Total	Received Treatment at a Specialty Facility	Did Not Receive Treatment at a Specialty Facility	
<b>TOTAL</b>	39	13	36	1.12
<b>GEOGRAPHIC DIVISION</b>				
Northeast	16	5	15	2.62
New England	12	3	11	*
Middle Atlantic	11	4	10	3.62
Midwest	18	7	16	2.56
East North Central	14	4	13	2.46
West North Central	10	5	8	*
South	25	8	23	2.01
South Atlantic	16	6	15	3.56
East South Central	9	2	9	*
West South Central	16	5	15	2.84
West	18	6	17	1.87
Mountain	9	4	8	4.11
Pacific	16	4	15	1.98
<b>COUNTY TYPE</b>				
Large Metro	26	9	25	1.76
Small Metro	23	7	22	1.81
250K - 1 Mil. Pop.	19	5	18	1.93
<250K Pop.	13	4	12	*
Nonmetro	16	6	15	2.43
Urbanized	7	2	7	3.22
Less Urbanized	12	5	11	3.90
Completely Rural	7	1	7	*

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on any illicit drug; (2) abuse of any illicit drug; or (3) received treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities (inpatient or outpatient), hospitals (inpatient only), and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type psychotherapeutic (nonmedical use).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.55D Standard Errors of Percentages of Persons Aged 12 to 17 Who Needed and Received Treatment for an Illicit Drug Problem in the Past Year, by Geographic Characteristics: 2000**

Geographic Characteristic	NEEDED TREATMENT FOR AN ILLICIT DRUG PROBLEM IN THE PAST YEAR			Percentage Who Received Treatment at a Specialty Facility Among Persons Who Needed Treatment
	Total	Received Treatment at a Specialty Facility	Did Not Receive Treatment at a Specialty Facility	
<b>TOTAL</b>	0.17	0.06	0.15	1.12
<b>GEOGRAPHIC DIVISION</b>				
Northeast	0.38	0.12	0.36	2.62
New England	1.07	0.29	1.02	*
Middle Atlantic	0.35	0.13	0.33	3.62
Midwest	0.32	0.12	0.29	2.56
East North Central	0.38	0.12	0.35	2.46
West North Central	0.61	0.29	0.50	*
South	0.30	0.10	0.28	2.01
South Atlantic	0.39	0.16	0.37	3.56
East South Central	0.65	0.13	0.62	*
West South Central	0.58	0.16	0.53	2.84
West	0.34	0.11	0.32	1.87
Mountain	0.57	0.24	0.49	4.11
Pacific	0.41	0.11	0.41	1.98
<b>COUNTY TYPE</b>				
Large Metro	0.23	0.08	0.22	1.76
Small Metro	0.32	0.10	0.30	1.81
250K - 1 Mil. Pop.	0.36	0.10	0.33	1.93
<250K Pop.	0.71	0.24	0.67	*
Nonmetro	0.34	0.12	0.32	2.43
Urbanized	0.55	0.17	0.53	3.22
Less Urbanized	0.41	0.17	0.39	3.90
Completely Rural	1.17	0.11	1.17	*

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on any illicit drug; (2) abuse of any illicit drug; or (3) received treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities (inpatient or outpatient), hospitals (inpatient only), and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type psychotherapeutic (nonmedical use).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.56C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 18 to 25 Who Needed and Received Treatment for an Illicit Drug Problem in the Past Year, by Geographic Characteristics: 2000**

Geographic Characteristic	NEEDED TREATMENT FOR AN ILLICIT DRUG PROBLEM IN THE PAST YEAR			Percentage Who Received Treatment at a Specialty Facility Among Persons Who Needed Treatment
	Total	Received Treatment at a Specialty Facility	Did Not Receive Treatment at a Specialty Facility	
<b>TOTAL</b>	56	16	54	0.94
<b>GEOGRAPHIC DIVISION</b>				
Northeast	25	6	25	1.59
New England	14	4	13	3.24
Middle Atlantic	21	5	21	1.82
Midwest	21	6	21	1.76
East North Central	19	4	19	1.60
West North Central	11	5	10	*
South	34	10	33	1.84
South Atlantic	22	6	22	2.64
East South Central	15	6	12	*
West South Central	21	5	22	*
West	29	8	27	2.18
Mountain	14	5	12	3.79
Pacific	25	7	24	2.63
<b>COUNTY TYPE</b>				
Large Metro	41	13	40	1.47
Small Metro	29	6	28	1.10
250K - 1 Mil. Pop.	25	5	24	1.25
<250K Pop.	16	3	16	2.35
Nonmetro	23	8	20	2.68
Urbanized	14	5	13	*
Less Urbanized	17	6	15	3.57
Completely Rural	3	2	2	*

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on any illicit drug; (2) abuse of any illicit drug; or (3) received treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities (inpatient or outpatient), hospitals (inpatient only), and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type psychotherapeutic (nonmedical use).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.56D Standard Errors of Percentages of Persons Aged 18 to 25 Who Needed and Received Treatment for an Illicit Drug Problem in the Past Year, by Geographic Characteristics: 2000**

Geographic Characteristic	NEEDED TREATMENT FOR AN ILLICIT DRUG PROBLEM IN THE PAST YEAR			Percentage Who Received Treatment at a Specialty Facility Among Persons Who Needed Treatment
	Total	Received Treatment at a Specialty Facility	Did Not Receive Treatment at a Specialty Facility	
<b>TOTAL</b>	0.19	0.06	0.19	0.94
<b>GEOGRAPHIC DIVISION</b>				
Northeast	0.50	0.13	0.48	1.59
New England	1.09	0.27	1.00	3.24
Middle Atlantic	0.55	0.14	0.55	1.82
Midwest	0.31	0.09	0.31	1.76
East North Central	0.39	0.09	0.39	1.60
West North Central	0.52	0.23	0.47	*
South	0.33	0.10	0.32	1.84
South Atlantic	0.45	0.13	0.45	2.64
East South Central	0.82	0.33	0.67	*
West South Central	0.59	0.15	0.61	*
West	0.43	0.13	0.40	2.18
Mountain	0.70	0.26	0.62	3.79
Pacific	0.54	0.14	0.51	2.63
<b>COUNTY TYPE</b>				
Large Metro	0.30	0.09	0.29	1.47
Small Metro	0.31	0.07	0.30	1.10
250K - 1 Mil. Pop.	0.37	0.08	0.36	1.25
<250K Pop.	0.57	0.12	0.56	2.35
Nonmetro	0.40	0.14	0.36	2.68
Urbanized	0.72	0.27	0.70	*
Less Urbanized	0.52	0.18	0.46	3.57
Completely Rural	0.65	0.31	0.50	*

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on any illicit drug; (2) abuse of any illicit drug; or (3) received treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities (inpatient or outpatient), hospitals (inpatient only), and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type psychotherapeutic (nonmedical use).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.



**Table 5.57C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 26 or Older Who Needed and Received Treatment for an Illicit Drug Problem in the Past Year, by Geographic Characteristics: 2000**

Geographic Characteristic	NEEDED TREATMENT FOR AN ILLICIT DRUG PROBLEM IN THE PAST YEAR			Percentage Who Received Treatment at a Specialty Facility Among Persons Who Needed Treatment
	Total	Received Treatment at a Specialty Facility	Did Not Receive Treatment at a Specialty Facility	
<b>TOTAL</b>	157	81	136	3.64
<b>GEOGRAPHIC DIVISION</b>				
Northeast	71	27	65	*
New England	46	21	42	*
Middle Atlantic	53	16	50	*
Midwest	58	29	49	*
East North Central	43	22	36	*
West North Central	39	19	34	*
South	61	37	55	*
South Atlantic	53	30	51	*
East South Central	14	6	13	*
West South Central	27	21	16	*
West	110	60	93	*
Mountain	40	15	36	*
Pacific	103	59	85	*
<b>COUNTY TYPE</b>				
Large Metro	131	66	115	5.00
Small Metro	61	31	53	*
250K - 1 Mil. Pop.	51	26	45	*
<250K Pop.	33	18	28	*
Nonmetro	55	32	45	*
Urbanized	23	21	15	*
Less Urbanized	48	24	41	*
Completely Rural	9	6	7	*

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on any illicit drug; (2) abuse of any illicit drug; or (3) received treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities (inpatient or outpatient), hospitals (inpatient only), and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type psychotherapeutic (nonmedical use).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.57D Standard Errors of Percentages of Persons Aged 26 or Older Who Needed and Received Treatment for an Illicit Drug Problem in the Past Year, by Geographic Characteristics: 2000**

Geographic Characteristic	NEEDED TREATMENT FOR AN ILLICIT DRUG PROBLEM IN THE PAST YEAR			Percentage Who Received Treatment at a Specialty Facility Among Persons Who Needed Treatment
	Total	Received Treatment at a Specialty Facility	Did Not Receive Treatment at a Specialty Facility	
<b>TOTAL</b>	0.09	0.05	0.08	3.64
<b>GEOGRAPHIC DIVISION</b>				
Northeast	0.21	0.08	0.19	*
New England	0.52	0.24	0.48	*
Middle Atlantic	0.22	0.07	0.20	*
Midwest	0.15	0.07	0.12	*
East North Central	0.15	0.08	0.13	*
West North Central	0.33	0.16	0.28	*
South	0.10	0.06	0.09	*
South Atlantic	0.17	0.09	0.16	*
East South Central	0.13	0.06	0.12	*
West South Central	0.15	0.12	0.09	*
West	0.30	0.16	0.25	*
Mountain	0.37	0.14	0.34	*
Pacific	0.39	0.22	0.32	*
<b>COUNTY TYPE</b>				
Large Metro	0.16	0.08	0.14	5.00
Small Metro	0.12	0.06	0.10	*
250K - 1 Mil. Pop.	0.13	0.07	0.12	*
<250K Pop.	0.25	0.14	0.21	*
Nonmetro	0.16	0.09	0.13	*
Urbanized	0.23	0.21	0.15	*
Less Urbanized	0.23	0.12	0.20	*
Completely Rural	0.22	0.14	0.17	*

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on any illicit drug; (2) abuse of any illicit drug; or (3) received treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities (inpatient or outpatient), hospitals (inpatient only), and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type psychotherapeutic (nonmedical use).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.58C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 18 or Older Who Needed and Received Treatment for an Illicit Drug Problem in the Past Year, by Geographic Characteristics: 2000**

Geographic Characteristic	NEEDED TREATMENT FOR AN ILLICIT DRUG PROBLEM IN THE PAST YEAR			Percentage Who Received Treatment at a Specialty Facility Among Persons Who Needed Treatment
	Total	Received Treatment at a Specialty Facility	Did Not Receive Treatment at a Specialty Facility	
<b>TOTAL</b>	171	85	150	2.09
<b>GEOGRAPHIC DIVISION</b>				
Northeast	77	28	71	3.10
New England	48	22	43	*
Middle Atlantic	60	17	57	3.03
Midwest	66	31	58	3.78
East North Central	51	22	45	3.88
West North Central	42	21	37	*
South	76	39	67	3.51
South Atlantic	63	31	58	*
East South Central	22	9	19	*
West South Central	36	22	28	*
West	114	62	98	*
Mountain	43	17	40	*
Pacific	106	60	89	*
<b>COUNTY TYPE</b>				
Large Metro	141	69	125	3.12
Small Metro	73	32	62	2.71
250K - 1 Mil. Pop.	62	26	53	3.04
<250K Pop.	39	18	33	*
Nonmetro	60	33	50	4.96
Urbanized	26	21	19	*
Less Urbanized	52	25	45	*
Completely Rural	10	6	7	*

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on any illicit drug; (2) abuse of any illicit drug; or (3) received treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities (inpatient or outpatient), hospitals (inpatient only), and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type psychotherapeutic (nonmedical use).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.58D Standard Errors of Percentages of Persons Aged 18 or Older Who Needed and Received Treatment for an Illicit Drug Problem in the Past Year, by Geographic Characteristics: 2000**

Geographic Characteristic	NEEDED TREATMENT FOR AN ILLICIT DRUG PROBLEM IN THE PAST YEAR			Percentage Who Received Treatment at a Specialty Facility Among Persons Who Needed Treatment
	Total	Received Treatment at a Specialty Facility	Did Not Receive Treatment at a Specialty Facility	
<b>TOTAL</b>	0.09	0.04	0.08	2.09
<b>GEOGRAPHIC DIVISION</b>				
Northeast	0.20	0.07	0.18	3.10
New England	0.48	0.21	0.43	*
Middle Atlantic	0.21	0.06	0.20	3.03
Midwest	0.14	0.07	0.12	3.78
East North Central	0.16	0.07	0.14	3.88
West North Central	0.31	0.15	0.26	*
South	0.11	0.05	0.09	3.51
South Atlantic	0.17	0.08	0.16	*
East South Central	0.18	0.07	0.15	*
West South Central	0.17	0.10	0.13	*
West	0.26	0.14	0.22	*
Mountain	0.34	0.13	0.31	*
Pacific	0.34	0.19	0.29	*
<b>COUNTY TYPE</b>				
Large Metro	0.14	0.07	0.13	3.12
Small Metro	0.12	0.05	0.10	2.71
250K - 1 Mil. Pop.	0.14	0.06	0.12	3.04
<250K Pop.	0.24	0.11	0.21	*
Nonmetro	0.15	0.08	0.12	4.96
Urbanized	0.22	0.18	0.16	*
Less Urbanized	0.22	0.10	0.19	*
Completely Rural	0.20	0.13	0.16	*

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug problem if they met at least one of three criteria during the past year: (1) dependent on any illicit drug; (2) abuse of any illicit drug; or (3) received treatment for an illicit drug problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities (inpatient or outpatient), hospitals (inpatient only), and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), inhalants, hallucinogens, heroin, or prescription-type psychotherapeutic (nonmedical use).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.59C Standard Errors of Estimated Numbers (in Thousands) of Persons Who Reported the Specific Substances for Which They Received Their Last or Current Treatment Among Persons Who Received Treatment in the Past Year, by Age Group: 2000**

Last or Current Treatment <sup>1</sup>	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Marijuana	71	18	21	62
Cocaine	81	6	12	78
Heroin	33	3	8	32
Hallucinogens	49	6	12	46
Inhalants	38	6	9	36
Pain Relievers	33	7	12	29
Tranquilizers	34	5	9	32
Stimulants	49	4	11	48
Sedatives	35	3	5	34
Alcohol	145	16	28	137

\*Low precision; no estimate reported.

NOTE: "Received Treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If a respondent reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

<sup>1</sup> Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.59D Standard Errors of Percentages of Persons Who Reported the Specific Substances for Which They Received Their Last or Current Treatment Among Persons Who Received Treatment in the Past Year, by Age Group: 2000**

Last or Current Treatment <sup>1</sup>	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Marijuana	2.28	3.01	2.94	3.02
Cocaine	2.46	1.55	1.88	3.57
Heroin	1.13	0.71	1.40	1.64
Hallucinogens	1.62	1.69	1.95	2.33
Inhalants	1.30	1.56	1.53	1.84
Pain Relievers	1.13	1.81	1.86	1.52
Tranquilizers	1.16	1.29	1.57	1.65
Stimulants	1.65	1.16	1.76	2.42
Sedatives	1.20	0.87	0.79	1.77
Alcohol	2.56	3.13	2.85	3.60

\*Low precision; no estimate reported.

NOTE: "Received Treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If a respondent reported that they were currently receiving treatment, they were asked questions regarding specific substances for their current treatment; otherwise, questions pertained to the last treatment they received.

<sup>1</sup> Respondents could indicate multiple substances for which they received their last or current treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.60C Standard Errors of Estimated Numbers (in Thousands) of Persons Who Reported Source of Payment for Last or Current Any Illicit Drug Treatment Among Persons Who Received Any Illicit Drug Treatment in the Past Year, by Age Group: 2000**

Source of Payment <sup>1</sup>	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Private Health Insurance	46	11	11	*
Medicare	66	9	8	*
Medicaid	46	9	7	*
Public Assistance Other Than Medicaid	47	7	9	*
Own Savings or Earnings	78	7	16	*
Family Members	46	9	12	*
Courts	40	8	9	*
CHAMPUS, TRICARE, CHAMPVA, the VA or Some Other Military Health Care	32	3	5	*
Employer	39	2	3	*

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: "Received Any Illicit Drug Treatment" refers to treatment received in order to reduce or stop drug use, or for medical problems associated with drug use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If a respondent reported that they were currently receiving treatment, they were asked questions regarding the sources of payment for their current treatment; otherwise, questions pertained to the last treatment they received.

<sup>1</sup> Respondents could indicate multiple sources of payment for their last or current treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.60D Standard Errors of Percentages of Persons Who Reported Source of Payment for Last or Current Any Illicit Drug Treatment Among Persons Who Received Any Illicit Drug Treatment in the Past Year, by Age Group: 2000**

Source of Payment <sup>1</sup>	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Private Health Insurance	3.29	3.56	3.65	*
Medicare	4.34	3.14	2.60	*
Medicaid	3.14	3.14	2.58	*
Public Assistance Other Than Medicaid	3.25	2.51	3.06	*
Own Savings or Earnings	4.09	2.70	4.40	*
Family Members	3.16	3.15	3.61	*
Courts	2.85	2.96	3.02	*
CHAMPUS, TRICARE, CHAMPVA, the VA or Some Other Military Health Care	2.35	1.07	1.65	*
Employer	2.85	0.65	1.25	*

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: "Received Any Illicit Drug Treatment" refers to treatment received in order to reduce or stop drug use, or for medical problems associated with drug use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If a respondent reported that they were currently receiving treatment, they were asked questions regarding the sources of payment for their current treatment; otherwise, questions pertained to the last treatment they received.

<sup>1</sup> Respondents could indicate multiple sources of payment for their last or current treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.



**Table 5.61C Standard Errors of Estimated Numbers (in Thousands) of Persons Who Reported Source of Payment for Last or Current Alcohol Treatment Among Persons Who Received Alcohol Treatment in the Past Year, by Age Group: 2000**

Source of Payment <sup>1</sup>	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Private Health Insurance	92	10	15	90
Medicare	71	8	6	70
Medicaid	43	8	7	42
Public Assistance Other Than Medicaid	57	7	10	55
Own Savings or Earnings	120	8	23	115
Family Members	47	8	12	45
Courts	32	8	10	30
CHAMPUS, TRICARE, CHAMPVA, the VA or Some Other Military Health Care	37	2	5	36
Employer	43	2	4	43

\*Low precision; no estimate reported.

NOTE: "Received Alcohol Treatment" refers to treatment received in order to reduce or stop alcohol use, or for medical problems associated with alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If a respondent reported that they were currently receiving treatment, they were asked questions regarding the sources of payment for their current treatment; otherwise, questions pertained to the last treatment they received.

<sup>1</sup> Respondents could indicate multiple sources of payment for their last or current treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.61D Standard Errors of Percentages of Persons Who Reported Source of Payment for Last or Current Alcohol Treatment Among Persons Who Received Alcohol Treatment in the Past Year, by Age Group: 2000**

Source of Payment <sup>1</sup>	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Private Health Insurance	3.51	3.44	3.22	4.75
Medicare	2.95	2.97	1.53	4.20
Medicaid	1.96	3.06	1.74	2.74
Public Assistance Other Than Medicaid	2.48	2.61	2.30	3.49
Own Savings or Earnings	3.49	3.14	4.02	4.64
Family Members	2.11	2.93	2.69	2.91
Courts	1.49	3.00	2.36	1.97
CHAMPUS, TRICARE, CHAMPVA, the VA or Some Other Military Health Care	1.68	0.99	1.24	2.37
Employer	1.97	0.76	1.03	2.80

\*Low precision; no estimate reported.

NOTE: "Received Alcohol Treatment" refers to treatment received in order to reduce or stop alcohol use, or for medical problems associated with alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If a respondent reported that they were currently receiving treatment, they were asked questions regarding the sources of payment for their current treatment; otherwise, questions pertained to the last treatment they received.

<sup>1</sup> Respondents could indicate multiple sources of payment for their last or current treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.62C Standard Errors of Estimated Numbers (in Thousands) of Persons Who Reported Source of Payment for Last or Current Any Illicit Drug or Alcohol Treatment Among Persons Who Received Any Illicit Drug or Alcohol Treatment in the Past Year, by Age Group: 2000**

Source of Payment <sup>1</sup>	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Private Health Insurance	96	13	17	93
Medicare	79	10	9	78
Medicaid	58	10	9	56
Public Assistance Other Than Medicaid	67	9	11	65
Own Savings or Earnings	130	10	25	123
Family Members	63	11	14	61
Courts	49	9	11	47
CHAMPUS, TRICARE, CHAMPVA, the VA or Some Other Military Health Care	46	3	6	46
Employer	56	2	5	56

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: "Received Any Illicit Drug or Alcohol Treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If a respondent reported that they were currently receiving treatment, they were asked questions regarding the sources of payment for their current treatment; otherwise, questions pertained to the last treatment they received.

<sup>1</sup> Respondents could indicate multiple sources of payment for their last or current treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.62D Standard Errors of Percentages of Persons Who Reported Source of Payment for Last or Current Any Illicit Drug or Alcohol Treatment Among Persons Who Received Any Illicit Drug or Alcohol Treatment in the Past Year, by Age Group: 2000**

Source of Payment <sup>1</sup>	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Private Health Insurance	2.77	2.72	2.64	4.00
Medicare	2.44	2.34	1.47	3.64
Medicaid	1.88	2.45	1.59	2.77
Public Assistance Other Than Medicaid	2.15	2.05	1.88	3.14
Own Savings or Earnings	2.91	2.47	3.13	4.03
Family Members	2.03	2.48	2.22	2.94
Courts	1.64	2.23	1.97	2.34
CHAMPUS, TRICARE, CHAMPVA, the VA or Some Other Military Health Care	1.56	0.76	1.10	2.31
Employer	1.85	0.64	0.82	2.75

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: "Received Any Illicit Drug or Alcohol Treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

NOTE: If a respondent reported that they were currently receiving treatment, they were asked questions regarding the sources of payment for their current treatment; otherwise, questions pertained to the last treatment they received.

<sup>1</sup> Respondents could indicate multiple sources of payment for their last or current treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

50601

**Table 5.63C (REVISED) Standard Errors of Estimated Numbers (in Thousands) of Persons Who Reported Specific Locations Where They Received Any Illicit Drug Treatment in the Past Year Among Persons Who Received Any Illicit Drug Treatment in the Past Year, by Age Group: 2000**

Location of Treatment <sup>1</sup>	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Hospital as an Inpatient	63	8	9	*
Rehabilitation Facility - Inpatient	63	9	10	*
Rehabilitation Facility - Outpatient	69	9	12	*
Mental Health Center - Outpatient	71	7	10	*
Private Doctor's Office	63	8	10	*
Self-Help Group	80	10	15	*
Prison/Jail	19	5	7	16
Emergency Room	*	5	8	*

\*Low precision; no estimate reported.

NOTE: Estimates of past year illicit drug treatment at specific locations reported in previously published versions of this table were under-reported by excluding respondents who reported having received only illicit drug treatment in the past year.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: "Received Any Illicit Drug Treatment" refers to treatment received in order to reduce or stop drug use, or for medical problems associated with drug use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>1</sup> Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.63D (REVISED) Standard Errors of Percentages of Persons Who Reported Specific Locations Where They Received Any Illicit Drug Treatment in the Past Year Among Persons Who Received Any Illicit Drug Treatment in the Past Year, by Age Group: 2000**

Location of Treatment <sup>1</sup>	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Hospital as an Inpatient	4.03	2.88	3.01	*
Rehabilitation Facility - Inpatient	3.87	3.15	3.33	*
Rehabilitation Facility - Outpatient	4.01	3.22	3.90	*
Mental Health Center - Outpatient	4.26	2.61	3.41	*
Private Doctor's Office	4.14	2.99	3.17	*
Self-Help Group	3.97	3.43	4.29	*
Prison/Jail	1.50	2.01	2.50	2.25
Emergency Room	*	2.00	2.75	*

\*Low precision; no estimate reported.

NOTE: Estimates of past year illicit drug treatment at specific locations reported in previously published versions of this table were under-reported by excluding respondents who reported having received only illicit drug treatment in the past year.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: "Received Any Illicit Drug Treatment" refers to treatment received in order to reduce or stop drug use, or for medical problems associated with drug use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>1</sup> Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

50601

**Table 5.64C (REVISED) Standard Errors of Estimated Numbers (in Thousands) of Persons Who Reported Specific Locations Where They Received Alcohol Treatment in the Past Year Among Persons Who Received Alcohol Treatment in the Past Year, by Age Group: 2000**

Location of Treatment <sup>1</sup>	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Hospital as an Inpatient	77	8	11	76
Rehabilitation Facility - Inpatient	78	7	12	77
Rehabilitation Facility - Outpatient	99	10	16	97
Mental Health Center - Outpatient	98	8	13	98
Private Doctor's Office	90	7	9	89
Self-Help Group	120	10	23	113
Prison/Jail	24	6	7	22
Emergency Room	69	5	11	*

\*Low precision; no estimate reported.

NOTE: Estimates of past year alcohol treatment at specific locations reported in previously published versions of this table were under-reported by excluding respondents who reported having received only alcohol treatment in the past year.

NOTE: "Received Alcohol Treatment" refers to treatment received in order to reduce or stop alcohol use, or for medical problems associated with alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>1</sup> Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.64D (REVISED) Standard Errors of Percentages of Persons Who Reported Specific Locations Where They Received Alcohol Treatment in the Past Year Among Persons Who Received Alcohol Treatment in the Past Year, by Age Group: 2000**

Location of Treatment <sup>1</sup>	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Hospital as an Inpatient	3.08	3.21	2.47	4.39
Rehabilitation Facility - Inpatient	3.10	2.76	2.71	4.46
Rehabilitation Facility - Outpatient	3.37	3.52	3.24	4.78
Mental Health Center - Outpatient	3.58	3.08	2.89	5.01
Private Doctor's Office	3.56	2.67	2.23	5.01
Self-Help Group	3.38	3.51	3.79	4.64
Prison/Jail	1.15	2.30	1.73	1.50
Emergency Room	2.95	1.92	2.58	*

\*Low precision; no estimate reported.

NOTE: Estimates of past year alcohol treatment at specific locations reported in previously published versions of this table were under-reported by excluding respondents who reported having received only alcohol treatment in the past year.

NOTE: "Received Alcohol Treatment" refers to treatment received in order to reduce or stop alcohol use, or for medical problems associated with alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>1</sup> Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.



**Table 5.65C Standard Errors of Estimated Numbers (in Thousands) of Persons Who Reported Specific Locations Where They Received Any Illicit Drug or Alcohol Treatment in the Past Year Among Persons Who Received Any Illicit Drug or Alcohol Treatment in the Past Year, by Age Group: 2000**

Location of Treatment <sup>1</sup>	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Hospital as an Inpatient	84	11	12	80
Rehabilitation Facility - Inpatient	83	10	15	79
Rehabilitation Facility - Outpatient	105	11	19	101
Mental Health Center - Outpatient	109	9	14	107
Private Doctor's Office	96	10	13	95
Self-Help Group	125	13	25	117
Prison/Jail	25	7	8	23
Emergency Room	70	7	13	68

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: "Received Any Illicit Drug or Alcohol Treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>1</sup> Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.65D Standard Errors of Percentages of Persons Who Reported Specific Locations Where They Received Any Illicit Drug or Alcohol Treatment in the Past Year Among Persons Who Received Any Illicit Drug or Alcohol Treatment in the Past Year, by Age Group: 2000**

Location of Treatment <sup>1</sup>	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Hospital as an Inpatient	2.52	2.51	1.89	3.67
Rehabilitation Facility - Inpatient	2.50	2.38	2.37	3.63
Rehabilitation Facility - Outpatient	2.77	2.47	2.74	4.04
Mental Health Center - Outpatient	2.99	2.26	2.22	4.34
Private Doctor's Office	2.85	2.26	2.09	4.17
Self-Help Group	2.85	2.73	3.00	4.05
Prison/Jail	0.90	1.84	1.41	1.19
Emergency Room	2.26	1.72	2.19	3.34

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: "Received Any Illicit Drug or Alcohol Treatment" refers to treatment received in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use. It includes treatment received at any location, such as a hospital, at a rehabilitation facility (inpatient or outpatient), mental health center, emergency room, private doctor's office, self-help group, or prison/jail.

<sup>1</sup> Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

10913

**Table 5.66C Standard Errors of Estimated Numbers (in Thousands) of Persons Aged 12 or Older Reporting Past Year Any Illicit Drug and/or Alcohol Dependence and/or Abuse: 2000**

Any Illicit Drug	Alcohol				
	Total	No Dependence or Abuse	Dependence or Abuse		
			Total	Dependence	Abuse
Total	0	307	307	222	230
No Dependence or Abuse	168	338	294	210	223
Dependence or Abuse	168	134	109	90	58
Dependence	135	104	90	85	41
Abuse	86	70	51	24	44

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on the definition found in the 4<sup>th</sup> ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.

**Table 5.66D Standard Errors of Percentages of Persons Aged 12 or Older Reporting Past Year Any Illicit Drug and/or Alcohol Dependence and/or Abuse: 2000**

Any Illicit Drug	Alcohol				
	Total	No Dependence or Abuse	Dependence or Abuse		
			Total	Dependence	Abuse
Total	0.00	0.14	0.14	0.10	0.10
No Dependence or Abuse	0.08	0.15	0.13	0.09	0.10
Dependence or Abuse	0.08	0.06	0.05	0.04	0.03
Dependence	0.06	0.05	0.04	0.04	0.02
Abuse	0.04	0.03	0.02	0.01	0.02

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on the definition found in the 4<sup>th</sup> ed. of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 2000.