

APR08-2008-000592

Abstract Submitted  
for the APR08 Meeting of  
The American Physical Society

Sorting Category: A19. (E)

**Top quark mass measurement in the dilepton channel at CDF using a neural network event selection and matrix element method** RAVI SHEKHAR, Duke University, CDF COLLABORATION — We present a measurement of the top quark mass using events in the dilepton decay channel. The events used in this analysis are selected using a genetically-evolved artificial neural network that is optimized directly for precision in the top quark mass measurements. This is the first application of this method in high energy physics. We extract the top quark mass from a probability that a given event is consistent with  $t\bar{t}$  decay in the dilepton channel. The probability is evaluated using the differential cross-section for  $t\bar{t}$  production and decay. The effect of background events in the sample is accounted for by evaluating differential cross sections for major background processes. Using  $2\text{ fb}^{-1}$  of  $p\bar{p}$  data collected at the CDF II detector, we measure  $m_t = 171.2 \pm 2.7(\text{stat.}) \pm 2.9(\text{syst.}) \text{ GeV}/c^2$ . We discuss the gain in sensitivity due to the use of the evolving neural network.

- Prefer Oral Session  
 Prefer Poster Session

Florencia Canelli  
canelli@fnal.gov

Special instructions: Membership pending

Date submitted: 10 Jan 2008

Electronic form version 1.4