

# Search for High-Mass $e^+e^-$ Resonances in $p\bar{p}$ Collisions at $\sqrt{s}=1.96$ TeV

T. Aaltonen,<sup>24</sup> J. Adelman,<sup>14</sup> T. Akimoto,<sup>56</sup> M.G. Albrow,<sup>18</sup> B. Álvarez González,<sup>12</sup> S. Amerio<sup>x</sup>,<sup>44</sup> D. Amidei,<sup>35</sup>  
 A. Anastassov,<sup>39</sup> A. Annovi,<sup>20</sup> J. Antos,<sup>15</sup> G. Apollinari,<sup>18</sup> A. Apresyan,<sup>49</sup> T. Arisawa,<sup>58</sup> A. Artikov,<sup>16</sup>  
 W. Ashmanskas,<sup>18</sup> A. Attal,<sup>4</sup> A. Aurisano,<sup>54</sup> F. Azfar,<sup>43</sup> P. Azzurri<sup>aa</sup>,<sup>47</sup> W. Badgett,<sup>18</sup> A. Barbaro-Galtieri,<sup>29</sup>  
 V.E. Barnes,<sup>49</sup> B.A. Barnett,<sup>26</sup> V. Bartsch,<sup>31</sup> G. Bauer,<sup>33</sup> P.-H. Beauchemin,<sup>34</sup> F. Bedeschi,<sup>47</sup> D. Beecher,<sup>31</sup>  
 S. Behari,<sup>26</sup> G. Bellettini<sup>y</sup>,<sup>47</sup> J. Bellinger,<sup>60</sup> D. Benjamin,<sup>17</sup> A. Beretvas,<sup>18</sup> J. Beringer,<sup>29</sup> A. Bhatti,<sup>51</sup> M. Binkley,<sup>18</sup>  
 D. Bisello<sup>x</sup>,<sup>44</sup> I. Bizjak<sup>dd</sup>,<sup>31</sup> R.E. Blair,<sup>2</sup> C. Blocker,<sup>7</sup> B. Blumenfeld,<sup>26</sup> A. Bocci,<sup>17</sup> A. Bodek,<sup>50</sup> V. Boisvert,<sup>50</sup>  
 G. Bolla,<sup>49</sup> D. Bortoletto,<sup>49</sup> J. Boudreau,<sup>48</sup> A. Boveia,<sup>11</sup> B. Brau<sup>a</sup>,<sup>11</sup> A. Bridgeman,<sup>25</sup> L. Brigliadori,<sup>44</sup>  
 C. Bromberg,<sup>36</sup> E. Brubaker,<sup>14</sup> J. Budagov,<sup>16</sup> H.S. Budd,<sup>50</sup> S. Budd,<sup>25</sup> S. Burke,<sup>18</sup> K. Burkett,<sup>18</sup> G. Busetto<sup>x</sup>,<sup>44</sup>  
 P. Bussey<sup>k</sup>,<sup>22</sup> A. Buzatu,<sup>34</sup> K. L. Byrum,<sup>2</sup> S. Cabrera<sup>u</sup>,<sup>17</sup> C. Calancha,<sup>32</sup> M. Campanelli,<sup>36</sup> M. Campbell,<sup>35</sup>  
 F. Canelli,<sup>18</sup> A. Canepa,<sup>46</sup> B. Carls,<sup>25</sup> D. Carlsmith,<sup>60</sup> R. Carosi,<sup>47</sup> S. Carrillo<sup>m</sup>,<sup>19</sup> S. Carron,<sup>34</sup> B. Casal,<sup>12</sup>  
 M. Casarsa,<sup>18</sup> A. Castro<sup>w</sup>,<sup>6</sup> P. Catastini<sup>z</sup>,<sup>47</sup> D. Cauz<sup>cc</sup>,<sup>55</sup> V. Cavaliere<sup>z</sup>,<sup>47</sup> M. Cavalli-Sforza,<sup>4</sup> A. Cerri,<sup>29</sup>  
 L. Cerrito<sup>n</sup>,<sup>31</sup> S.H. Chang,<sup>28</sup> Y.C. Chen,<sup>1</sup> M. Chertok,<sup>8</sup> G. Chiarelli,<sup>47</sup> G. Chlachidze,<sup>18</sup> F. Chlebana,<sup>18</sup> K. Cho,<sup>28</sup>  
 D. Chokheli,<sup>16</sup> J.P. Chou,<sup>23</sup> G. Choudalakis,<sup>33</sup> S.H. Chuang,<sup>53</sup> K. Chung,<sup>13</sup> W.H. Chung,<sup>60</sup> Y.S. Chung,<sup>50</sup>  
 T. Chwalek,<sup>27</sup> C.I. Ciobanu,<sup>45</sup> M.A. Ciocci<sup>z</sup>,<sup>47</sup> A. Clark,<sup>21</sup> D. Clark,<sup>7</sup> G. Compostella,<sup>44</sup> M.E. Convery,<sup>18</sup>  
 J. Conway,<sup>8</sup> M. Cordelli,<sup>20</sup> G. Cortiana<sup>x</sup>,<sup>44</sup> C.A. Cox,<sup>8</sup> D.J. Cox,<sup>8</sup> F. Crescioli<sup>y</sup>,<sup>47</sup> C. Cuenca Almenar<sup>u</sup>,<sup>8</sup>  
 J. Cuevas<sup>r</sup>,<sup>12</sup> R. Culbertson,<sup>18</sup> J.C. Cully,<sup>35</sup> D. Dagenhart,<sup>18</sup> M. Datta,<sup>18</sup> T. Davies,<sup>22</sup> P. de Barbaro,<sup>50</sup>  
 S. De Cecco,<sup>52</sup> A. Deisher,<sup>29</sup> G. De Lorenzo,<sup>4</sup> M. Dell’Orso<sup>y</sup>,<sup>47</sup> C. Deluca,<sup>4</sup> L. Demortier,<sup>51</sup> J. Deng,<sup>17</sup> M. Deninno,<sup>6</sup>  
 P.F. Derwent,<sup>18</sup> G.P. di Giovanni,<sup>45</sup> C. Dionisi<sup>bb</sup>,<sup>52</sup> B. Di Ruzza<sup>cc</sup>,<sup>55</sup> J.R. Dittmann,<sup>5</sup> M. D’Onofrio,<sup>4</sup> S. Donati<sup>y</sup>,<sup>47</sup>  
 P. Dong,<sup>9</sup> J. Donini,<sup>44</sup> T. Dorigo,<sup>44</sup> S. Dube,<sup>53</sup> J. Efron,<sup>40</sup> A. Elagin,<sup>54</sup> R. Erbacher,<sup>8</sup> D. Errede,<sup>25</sup> S. Errede,<sup>25</sup>  
 R. Eusebi,<sup>18</sup> H.C. Fang,<sup>29</sup> S. Farrington,<sup>43</sup> W.T. Fedorko,<sup>14</sup> R.G. Feild,<sup>61</sup> M. Feindt,<sup>27</sup> J.P. Fernandez,<sup>32</sup>  
 C. Ferrazza<sup>aa</sup>,<sup>47</sup> R. Field,<sup>19</sup> G. Flanagan,<sup>49</sup> R. Forrest,<sup>8</sup> M.J. Frank,<sup>5</sup> M. Franklin,<sup>23</sup> J.C. Freeman,<sup>18</sup> I. Furic,<sup>19</sup>  
 M. Gallinaro,<sup>52</sup> J. Galyardt,<sup>13</sup> F. Garberson,<sup>11</sup> J.E. Garcia,<sup>21</sup> A.F. Garfinkel,<sup>49</sup> K. Genser,<sup>18</sup> H. Gerberich,<sup>25</sup>  
 D. Gerdes,<sup>35</sup> A. Gessler,<sup>27</sup> S. Giagu<sup>bb</sup>,<sup>52</sup> V. Giakoumopoulou,<sup>3</sup> P. Giannetti,<sup>47</sup> K. Gibson,<sup>48</sup> J.L. Gimmell,<sup>50</sup>  
 C.M. Ginsburg,<sup>18</sup> N. Giokaris,<sup>3</sup> M. Giordani<sup>cc</sup>,<sup>55</sup> P. Giromini,<sup>20</sup> M. Giunta<sup>y</sup>,<sup>47</sup> G. Giurgiu,<sup>26</sup> V. Glagolev,<sup>16</sup>  
 D. Glenzinski,<sup>18</sup> M. Gold,<sup>38</sup> N. Goldschmidt,<sup>19</sup> A. Golossanov,<sup>18</sup> G. Gomez,<sup>12</sup> G. Gomez-Ceballos,<sup>33</sup>  
 M. Goncharov,<sup>54</sup> O. González,<sup>32</sup> I. Gorelov,<sup>38</sup> A.T. Goshaw,<sup>17</sup> K. Goulios,<sup>51</sup> A. Gresele<sup>x</sup>,<sup>44</sup> S. Grinstein,<sup>23</sup>  
 C. Grosso-Pilcher,<sup>14</sup> R.C. Group,<sup>18</sup> U. Grundler,<sup>25</sup> J. Guimaraes da Costa,<sup>23</sup> Z. Gunay-Unalan,<sup>36</sup> C. Haber,<sup>29</sup>  
 K. Hahn,<sup>33</sup> S.R. Hahn,<sup>18</sup> E. Halkiadakis,<sup>53</sup> B.-Y. Han,<sup>50</sup> J.Y. Han,<sup>50</sup> F. Happacher,<sup>20</sup> K. Hara,<sup>56</sup> D. Hare,<sup>53</sup>  
 M. Hare,<sup>57</sup> S. Harper,<sup>43</sup> R.F. Harr,<sup>59</sup> R.M. Harris,<sup>18</sup> M. Hartz,<sup>48</sup> K. Hatakeyama,<sup>51</sup> C. Hays,<sup>43</sup> M. Heck,<sup>27</sup>  
 A. Heijboer,<sup>46</sup> J. Heinrich,<sup>46</sup> C. Henderson,<sup>33</sup> M. Herndon,<sup>60</sup> J. Heuser,<sup>27</sup> S. Hewamanage,<sup>5</sup> D. Hidas,<sup>17</sup>  
 C.S. Hill<sup>c</sup>,<sup>11</sup> D. Hirschbuehl,<sup>27</sup> A. Hocker,<sup>18</sup> S. Hou,<sup>1</sup> M. Houlden,<sup>30</sup> S.-C. Hsu,<sup>29</sup> B.T. Huffman,<sup>43</sup> R.E. Hughes,<sup>40</sup>  
 U. Husemann,<sup>61</sup> J. Huston,<sup>36</sup> J. Incandela,<sup>11</sup> G. Introzzi,<sup>47</sup> M. Iori<sup>bb</sup>,<sup>52</sup> A. Ivanov,<sup>8</sup> E. James,<sup>18</sup> B. Jayatilaka,<sup>17</sup>  
 E.-J. Jeon,<sup>28</sup> M.K. Jha,<sup>6</sup> S. Jindariani,<sup>18</sup> W. Johnson,<sup>8</sup> M. Jones,<sup>49</sup> K.K. Joo,<sup>28</sup> S.Y. Jun,<sup>13</sup> J.E. Jung,<sup>28</sup>  
 T.R. Junk,<sup>18</sup> T. Kamon,<sup>54</sup> D. Kar,<sup>19</sup> P.E. Karchin,<sup>59</sup> Y. Kato,<sup>42</sup> R. Kephart,<sup>18</sup> J. Keung,<sup>46</sup> V. Khotilovich,<sup>54</sup>  
 B. Kilminster,<sup>18</sup> D.H. Kim,<sup>28</sup> H.S. Kim,<sup>28</sup> H.W. Kim,<sup>28</sup> J.E. Kim,<sup>28</sup> M.J. Kim,<sup>20</sup> S.B. Kim,<sup>28</sup> S.H. Kim,<sup>56</sup>  
 Y.K. Kim,<sup>14</sup> N. Kimura,<sup>56</sup> L. Kirsch,<sup>7</sup> S. Klimentenko,<sup>19</sup> B. Knuteson,<sup>33</sup> B.R. Ko,<sup>17</sup> K. Kondo,<sup>58</sup> D.J. Kong,<sup>28</sup>  
 J. Konigsberg,<sup>19</sup> A. Korytov,<sup>19</sup> A.V. Kotwal,<sup>17</sup> M. Kreps,<sup>27</sup> J. Kroll,<sup>46</sup> D. Krop,<sup>14</sup> N. Krumnack,<sup>5</sup> M. Kruse,<sup>17</sup>  
 V. Krutelyov,<sup>11</sup> T. Kubo,<sup>56</sup> T. Kuhr,<sup>27</sup> N.P. Kulkarni,<sup>59</sup> M. Kurata,<sup>56</sup> Y. Kusakabe,<sup>58</sup> S. Kwang,<sup>14</sup> A.T. Laasanen,<sup>49</sup>  
 S. Lami,<sup>47</sup> S. Lammel,<sup>18</sup> M. Lancaster,<sup>31</sup> R.L. Lander,<sup>8</sup> K. Lannon<sup>q</sup>,<sup>40</sup> A. Lath,<sup>53</sup> G. Latino<sup>z</sup>,<sup>47</sup> I. Lazzizzera<sup>x</sup>,<sup>44</sup>  
 T. LeCompte,<sup>2</sup> E. Lee,<sup>54</sup> H.S. Lee,<sup>14</sup> S.W. Lee<sup>t</sup>,<sup>54</sup> S. Leone,<sup>47</sup> J.D. Lewis,<sup>18</sup> C.-S. Lin,<sup>29</sup> J. Linacre,<sup>43</sup> M. Lindgren,<sup>18</sup>  
 E. Lipeles,<sup>46</sup> A. Lister,<sup>8</sup> D.O. Litvintsev,<sup>18</sup> C. Liu,<sup>48</sup> T. Liu,<sup>18</sup> N.S. Lockyer,<sup>46</sup> A. Loginov,<sup>61</sup> M. Loretix<sup>x</sup>,<sup>44</sup>  
 L. Lovas,<sup>15</sup> D. Lucchesi<sup>x</sup>,<sup>44</sup> C. Luci<sup>bb</sup>,<sup>52</sup> J. Lueck,<sup>27</sup> P. Lujan,<sup>29</sup> P. Lukens,<sup>18</sup> G. Lungu,<sup>51</sup> L. Lyons,<sup>43</sup> J. Lys,<sup>29</sup>  
 R. Lysak,<sup>15</sup> D. MacQueen,<sup>34</sup> R. Madrak,<sup>18</sup> K. Maeshima,<sup>18</sup> K. Makhoul,<sup>33</sup> T. Maki,<sup>24</sup> P. Maksimovic,<sup>26</sup> S. Malde,<sup>43</sup>  
 S. Malik,<sup>31</sup> G. Manca<sup>e</sup>,<sup>30</sup> A. Manousakis-Katsikakis,<sup>3</sup> F. Margaroli,<sup>49</sup> C. Marino,<sup>27</sup> C.P. Marino,<sup>25</sup> A. Martin,<sup>61</sup>  
 V. Martin<sup>l</sup>,<sup>22</sup> M. Martínez,<sup>4</sup> R. Martínez-Ballarín,<sup>32</sup> T. Maruyama,<sup>56</sup> P. Mastrandrea,<sup>52</sup> T. Masubuchi,<sup>56</sup>  
 M. Mathis,<sup>26</sup> M.E. Mattson,<sup>59</sup> P. Mazzanti,<sup>6</sup> K.S. McFarland,<sup>50</sup> P. McIntyre,<sup>54</sup> R. McNulty<sup>j</sup>,<sup>30</sup> A. Mehta,<sup>30</sup>  
 P. Mehtala,<sup>24</sup> A. Menzione,<sup>47</sup> P. Merkel,<sup>49</sup> C. Mesropian,<sup>51</sup> T. Miao,<sup>18</sup> N. Miladinovic,<sup>7</sup> R. Miller,<sup>36</sup> C. Mills,<sup>23</sup>  
 M. Milnik,<sup>27</sup> A. Mitra,<sup>1</sup> G. Mitselmakher,<sup>19</sup> H. Miyake,<sup>56</sup> N. Moggi,<sup>6</sup> C.S. Moon,<sup>28</sup> R. Moore,<sup>18</sup> M.J. Morello<sup>y</sup>,<sup>47</sup>  
 J. Morlok,<sup>27</sup> P. Movilla Fernandez,<sup>18</sup> J. Mülmenstädt,<sup>29</sup> A. Mukherjee,<sup>18</sup> Th. Muller,<sup>27</sup> R. Mumford,<sup>26</sup> P. Murat,<sup>18</sup>  
 M. Mussini<sup>w</sup>,<sup>6</sup> J. Nachtman,<sup>18</sup> Y. Nagai,<sup>56</sup> A. Nagano,<sup>56</sup> J. Naganoma,<sup>56</sup> K. Nakamura,<sup>56</sup> I. Nakano,<sup>41</sup> A. Napier,<sup>57</sup>

V. Neula,<sup>17</sup> J. Nett,<sup>60</sup> C. Neu<sup>v</sup>,<sup>46</sup> M.S. Neubauer,<sup>25</sup> S. Neubauer,<sup>27</sup> J. Nielsen<sup>9</sup>,<sup>29</sup> L. Nodulman,<sup>2</sup> M. Norman,<sup>10</sup> O. Normiella,<sup>25</sup> E. Nurse,<sup>31</sup> L. Oakes,<sup>43</sup> S.H. Oh,<sup>17</sup> Y.D. Oh,<sup>28</sup> I. Oksuzian,<sup>19</sup> T. Okusawa,<sup>42</sup> R. Orava,<sup>24</sup> S. Pagan Griso<sup>x</sup>,<sup>44</sup> E. Palencia,<sup>18</sup> V. Papadimitriou,<sup>18</sup> A. Papaikonomou,<sup>27</sup> A.A. Paramonov,<sup>14</sup> B. Parks,<sup>40</sup> S. Pashapour,<sup>34</sup> J. Patrick,<sup>18</sup> G. Pauletta<sup>cc</sup>,<sup>55</sup> M. Paulini,<sup>13</sup> C. Paus,<sup>33</sup> T. Peiffer,<sup>27</sup> D.E. Pellett,<sup>8</sup> A. Penzo,<sup>55</sup> T.J. Phillips,<sup>17</sup> G. Piacentino,<sup>47</sup> E. Pianori,<sup>46</sup> L. Pinera,<sup>19</sup> K. Pitts,<sup>25</sup> C. Plager,<sup>9</sup> L. Pondrom,<sup>60</sup> O. Poukhov<sup>\*</sup>,<sup>16</sup> N. Pounder,<sup>43</sup> F. Prakoshyn,<sup>16</sup> A. Pronko,<sup>18</sup> J. Proudfoot,<sup>2</sup> F. Ptohos<sup>i</sup>,<sup>18</sup> E. Pueschel,<sup>13</sup> G. Punzi<sup>y</sup>,<sup>47</sup> J. Pursley,<sup>60</sup> J. Rademacker<sup>c</sup>,<sup>43</sup> A. Rahaman,<sup>48</sup> V. Ramakrishnan,<sup>60</sup> N. Ranjan,<sup>49</sup> I. Redondo,<sup>32</sup> V. Rekovic,<sup>38</sup> P. Renton,<sup>43</sup> M. Renz,<sup>27</sup> M. Rescigno,<sup>52</sup> S. Richter,<sup>27</sup> F. Rimondi<sup>w</sup>,<sup>6</sup> L. Ristori,<sup>47</sup> A. Robson,<sup>22</sup> T. Rodrigo,<sup>12</sup> T. Rodriguez,<sup>46</sup> E. Rogers,<sup>25</sup> S. Rolli,<sup>57</sup> R. Roser,<sup>18</sup> M. Rossi,<sup>55</sup> R. Rossin,<sup>11</sup> P. Roy,<sup>34</sup> A. Ruiz,<sup>12</sup> J. Russ,<sup>13</sup> V. Rusu,<sup>18</sup> A. Safonov,<sup>54</sup> W.K. Sakumoto,<sup>50</sup> O. Saltó,<sup>4</sup> L. Santi<sup>cc</sup>,<sup>55</sup> S. Sarkar<sup>bb</sup>,<sup>52</sup> L. Sartori,<sup>47</sup> K. Sato,<sup>18</sup> A. Savoy-Navarro,<sup>45</sup> P. Schlabach,<sup>18</sup> A. Schmidt,<sup>27</sup> E.E. Schmidt,<sup>18</sup> M.A. Schmidt,<sup>14</sup> M.P. Schmidt<sup>\*</sup>,<sup>61</sup> M. Schmitt,<sup>39</sup> T. Schwarz,<sup>8</sup> L. Scodellaro,<sup>12</sup> A. Scribano<sup>z</sup>,<sup>47</sup> F. Scuri,<sup>47</sup> A. Sedov,<sup>49</sup> S. Seidel,<sup>38</sup> Y. Seiya,<sup>42</sup> A. Semenov,<sup>16</sup> L. Sexton-Kennedy,<sup>18</sup> F. Sforza,<sup>47</sup> A. Sfyrila,<sup>25</sup> S.Z. Shalhout,<sup>59</sup> T. Shears,<sup>30</sup> P.F. Shepard,<sup>48</sup> M. Shimojima<sup>p</sup>,<sup>56</sup> S. Shiraishi,<sup>14</sup> M. Shochet,<sup>14</sup> Y. Shon,<sup>60</sup> I. Shreyber,<sup>37</sup> A. Sidoti,<sup>47</sup> P. Sinervo,<sup>34</sup> A. Sisakyan,<sup>16</sup> A.J. Slaughter,<sup>18</sup> J. Slaunwhite,<sup>40</sup> K. Sliwa,<sup>57</sup> J.R. Smith,<sup>8</sup> F.D. Snider,<sup>18</sup> R. Snihur,<sup>34</sup> A. Soha,<sup>8</sup> S. Somalwar,<sup>53</sup> V. Sorin,<sup>36</sup> J. Spalding,<sup>18</sup> T. Spreitzer,<sup>34</sup> P. Squillacioti<sup>z</sup>,<sup>47</sup> M. Stanitzki,<sup>61</sup> R. St. Denis,<sup>22</sup> B. Stelzer<sup>s</sup>,<sup>9</sup> O. Stelzer-Chilton,<sup>17</sup> D. Stentz,<sup>39</sup> J. Strogas,<sup>38</sup> G.L. Strycker,<sup>35</sup> D. Stuart,<sup>11</sup> J.S. Suh,<sup>28</sup> A. Sukhanov,<sup>19</sup> I. Suslov,<sup>16</sup> T. Suzuki,<sup>56</sup> A. Taffard<sup>f</sup>,<sup>25</sup> R. Takashima,<sup>41</sup> Y. Takeuchi,<sup>56</sup> R. Tanaka,<sup>41</sup> M. Tecchio,<sup>35</sup> P.K. Teng,<sup>1</sup> K. Terashi,<sup>51</sup> J. Thom<sup>h</sup>,<sup>18</sup> A.S. Thompson,<sup>22</sup> G.A. Thompson,<sup>25</sup> E. Thomson,<sup>46</sup> P. Tipton,<sup>61</sup> P. Ttito-Guzmán,<sup>32</sup> S. Tkaczyk,<sup>18</sup> D. Toback,<sup>54</sup> S. Tokar,<sup>15</sup> K. Tollefson,<sup>36</sup> T. Tomura,<sup>56</sup> D. Tonelli,<sup>18</sup> S. Torre,<sup>20</sup> D. Torretta,<sup>18</sup> P. Totaro<sup>cc</sup>,<sup>55</sup> S. Tourneur,<sup>45</sup> M. Trovato,<sup>47</sup> S.-Y. Tsai,<sup>1</sup> Y. Tu,<sup>46</sup> N. Turini<sup>z</sup>,<sup>47</sup> F. Ukegawa,<sup>56</sup> S. Vallecorsa,<sup>21</sup> N. van Remortel<sup>b</sup>,<sup>24</sup> A. Varganov,<sup>35</sup> E. Vataga<sup>aa</sup>,<sup>47</sup> F. Vázquez<sup>m</sup>,<sup>19</sup> G. Velev,<sup>18</sup> C. Vellidis,<sup>3</sup> V. Veszpremi,<sup>49</sup> M. Vidal,<sup>32</sup> R. Vidal,<sup>18</sup> I. Vila,<sup>12</sup> R. Vilar,<sup>12</sup> T. Vine,<sup>31</sup> M. Vogel,<sup>38</sup> I. Volobouev<sup>t</sup>,<sup>29</sup> G. Volpi<sup>y</sup>,<sup>47</sup> P. Wagner,<sup>46</sup> R.G. Wagner,<sup>2</sup> R.L. Wagner,<sup>18</sup> W. Wagner,<sup>27</sup> J. Wagner-Kuhr,<sup>27</sup> T. Wakisaka,<sup>42</sup> R. Wallny,<sup>9</sup> C. Wang,<sup>17</sup> S.M. Wang,<sup>1</sup> A. Warburton,<sup>34</sup> D. Waters,<sup>31</sup> M. Weinberger,<sup>54</sup> J. Weinelt,<sup>27</sup> W.C. Wester III,<sup>18</sup> B. Whitehouse,<sup>57</sup> D. Whiteson<sup>f</sup>,<sup>46</sup> A.B. Wicklund,<sup>2</sup> E. Wicklund,<sup>18</sup> S. Wilbur,<sup>14</sup> G. Williams,<sup>34</sup> H.H. Williams,<sup>46</sup> P. Wilson,<sup>18</sup> B.L. Winer,<sup>40</sup> P. Wittich<sup>h</sup>,<sup>18</sup> S. Wolbers,<sup>18</sup> C. Wolfe,<sup>14</sup> T. Wright,<sup>35</sup> X. Wu,<sup>21</sup> F. Würthwein,<sup>10</sup> S.M. Wynne,<sup>30</sup> S. Xie,<sup>33</sup> A. Yagil,<sup>10</sup> K. Yamamoto,<sup>42</sup> J. Yamaoka,<sup>53</sup> U.K. Yang<sup>o</sup>,<sup>14</sup> Y.C. Yang,<sup>28</sup> W.M. Yao,<sup>29</sup> G.P. Yeh,<sup>18</sup> J. Yoh,<sup>18</sup> K. Yorita,<sup>14</sup> T. Yoshida,<sup>42</sup> G.B. Yu,<sup>50</sup> I. Yu,<sup>28</sup> S.S. Yu,<sup>18</sup> J.C. Yun,<sup>18</sup> L. Zanello<sup>bb</sup>,<sup>52</sup> A. Zanetti,<sup>55</sup> X. Zhang,<sup>25</sup> Y. Zheng<sup>d</sup>,<sup>9</sup> and S. Zucchelli<sup>w</sup>,<sup>6</sup>

(CDF Collaboration<sup>†</sup>)

<sup>1</sup>*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*

<sup>2</sup>*Argonne National Laboratory, Argonne, Illinois 60439*

<sup>3</sup>*University of Athens, 157 71 Athens, Greece*

<sup>4</sup>*Institut de Fisica d'Altes Energies, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain*

<sup>5</sup>*Baylor University, Waco, Texas 76798*

<sup>6</sup>*Istituto Nazionale di Fisica Nucleare Bologna, <sup>w</sup>University of Bologna, I-40127 Bologna, Italy*

<sup>7</sup>*Brandeis University, Waltham, Massachusetts 02254*

<sup>8</sup>*University of California, Davis, Davis, California 95616*

<sup>9</sup>*University of California, Los Angeles, Los Angeles, California 90024*

<sup>10</sup>*University of California, San Diego, La Jolla, California 92093*

<sup>11</sup>*University of California, Santa Barbara, Santa Barbara, California 93106*

<sup>12</sup>*Instituto de Fisica de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*

<sup>13</sup>*Carnegie Mellon University, Pittsburgh, PA 15213*

<sup>14</sup>*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637*

<sup>15</sup>*Comenius University, 842 48 Bratislava, Slovakia; Institute of Experimental Physics, 040 01 Kosice, Slovakia*

<sup>16</sup>*Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*

<sup>17</sup>*Duke University, Durham, North Carolina 27708*

<sup>18</sup>*Fermi National Accelerator Laboratory, Batavia, Illinois 60510*

<sup>19</sup>*University of Florida, Gainesville, Florida 32611*

<sup>20</sup>*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*

<sup>21</sup>*University of Geneva, CH-1211 Geneva 4, Switzerland*

<sup>22</sup>*Glasgow University, Glasgow G12 8QQ, United Kingdom*

<sup>23</sup>*Harvard University, Cambridge, Massachusetts 02138*

<sup>24</sup>*Division of High Energy Physics, Department of Physics,*

*University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland*

<sup>25</sup>*University of Illinois, Urbana, Illinois 61801*

- <sup>26</sup>The Johns Hopkins University, Baltimore, Maryland 21218
- <sup>27</sup>Institut für Experimentelle Kernphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany
- <sup>28</sup>Center for High Energy Physics: Kyungpook National University, Daegu 702-701, Korea; Seoul National University, Seoul 151-742, Korea; Sungkyunkwan University, Suwon 440-746, Korea; Korea Institute of Science and Technology Information, Daejeon, 305-806, Korea; Chonnam National University, Gwangju, 500-757, Korea
- <sup>29</sup>Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720
- <sup>30</sup>University of Liverpool, Liverpool L69 7ZE, United Kingdom
- <sup>31</sup>University College London, London WC1E 6BT, United Kingdom
- <sup>32</sup>Centro de Investigaciones Energeticas Medioambientales y Tecnologicas, E-28040 Madrid, Spain
- <sup>33</sup>Massachusetts Institute of Technology, Cambridge, Massachusetts 02139
- <sup>34</sup>Institute of Particle Physics: McGill University, Montréal, Canada H3A 2T8; and University of Toronto, Toronto, Canada M5S 1A7
- <sup>35</sup>University of Michigan, Ann Arbor, Michigan 48109
- <sup>36</sup>Michigan State University, East Lansing, Michigan 48824
- <sup>37</sup>Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia
- <sup>38</sup>University of New Mexico, Albuquerque, New Mexico 87131
- <sup>39</sup>Northwestern University, Evanston, Illinois 60208
- <sup>40</sup>The Ohio State University, Columbus, Ohio 43210
- <sup>41</sup>Okayama University, Okayama 700-8530, Japan
- <sup>42</sup>Osaka City University, Osaka 588, Japan
- <sup>43</sup>University of Oxford, Oxford OX1 3RH, United Kingdom
- <sup>44</sup>Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, <sup>x</sup>University of Padova, I-35131 Padova, Italy
- <sup>45</sup>LPNHE, Université Pierre et Marie Curie/IN2P3-CNRS, UMR7585, Paris, F-75252 France
- <sup>46</sup>University of Pennsylvania, Philadelphia, Pennsylvania 19104
- <sup>47</sup>Istituto Nazionale di Fisica Nucleare Pisa, <sup>y</sup>University of Pisa, <sup>z</sup>University of Siena and <sup>aa</sup>Scuola Normale Superiore, I-56127 Pisa, Italy
- <sup>48</sup>University of Pittsburgh, Pittsburgh, Pennsylvania 15260
- <sup>49</sup>Purdue University, West Lafayette, Indiana 47907
- <sup>50</sup>University of Rochester, Rochester, New York 14627
- <sup>51</sup>The Rockefeller University, New York, New York 10021
- <sup>52</sup>Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1, <sup>bb</sup>Sapienza Università di Roma, I-00185 Roma, Italy
- <sup>53</sup>Rutgers University, Piscataway, New Jersey 08855
- <sup>54</sup>Texas A&M University, College Station, Texas 77843
- <sup>55</sup>Istituto Nazionale di Fisica Nucleare Trieste/Udine, <sup>cc</sup>University of Trieste/Udine, Italy
- <sup>56</sup>University of Tsukuba, Tsukuba, Ibaraki 305, Japan
- <sup>57</sup>Tufts University, Medford, Massachusetts 02155
- <sup>58</sup>Waseda University, Tokyo 169, Japan
- <sup>59</sup>Wayne State University, Detroit, Michigan 48201
- <sup>60</sup>University of Wisconsin, Madison, Wisconsin 53706
- <sup>61</sup>Yale University, New Haven, Connecticut 06520

A search for high-mass resonances in the  $e^+e^-$  final state is presented based on  $\sqrt{s}=1.96$  TeV  $p\bar{p}$  collision data from the CDF II detector at the Fermilab Tevatron from an integrated luminosity of  $2.5 \text{ fb}^{-1}$ . The largest excess over the standard model prediction is at an  $e^+e^-$  invariant mass of 240 GeV/ $c^2$ . The probability of observing such an excess arising from fluctuations in the standard model anywhere in the mass range of 150–1,000 GeV/ $c^2$  is 0.6% (equivalent to  $2.5 \sigma$ ). We set Bayesian upper limits on  $\sigma(p\bar{p} \rightarrow X) \cdot \mathcal{B}(X \rightarrow e^+e^-)$  at the 95% credibility level, where  $X$  is a spin 1 or spin 2 particle, and we exclude the standard model coupling  $Z'$  and the Randall-Sundrum graviton for  $k/\overline{M}_{Pl} = 0.1$  with masses below 963 and 848 GeV/ $c^2$ , respectively.

PACS numbers: 13.85.Rm, 13.85.Qk, 12.60.Cn, 14.70.Pw, 04.50.-h

---

\*Deceased

†With visitors from <sup>a</sup>University of Massachusetts Amherst, Amherst, Massachusetts 01003, <sup>b</sup>Universiteit Antwerpen, B-2610 Antwerp, Belgium, <sup>c</sup>University of Bristol, Bristol BS8 1TL, United Kingdom, <sup>d</sup>Chinese Academy of Sciences, Beijing 100864,

China, <sup>e</sup>Istituto Nazionale di Fisica Nucleare, Sezione di Cagliari, 09042 Monserrato (Cagliari), Italy, <sup>f</sup>University of California Irvine, Irvine, CA 92697, <sup>g</sup>University of California Santa Cruz, Santa Cruz, CA 95064, <sup>h</sup>Cornell University, Ithaca, NY 14853, <sup>i</sup>University of Cyprus, Nicosia CY-1678, Cyprus, <sup>j</sup>University

The charged lepton-antilepton pair signature, in particular  $e^+e^-$  and  $\mu^+\mu^-$ , has been a leading discovery channel for new particles such as the  $J/\psi$  and  $\Upsilon$  mesons and the  $Z$  boson. Even though leptonic decay channels generally have lower branching ratios than hadronic channels, these channels are preferred for particle searches since they have lower backgrounds than hadronic channels. Furthermore, leptons have cleaner experimental signatures, and their energies and momenta can be measured with better resolution than those of hadrons.

Many models beyond the standard model (SM) predict the existence of new particles decaying to lepton-antilepton pairs. The  $E_6$   $Z'$ 's [1] and the Randall-Sundrum (RS) graviton [2] are an example of specific new particles decaying to a lepton-antilepton final state. The  $E_6$  model unifies the interactions of the SM into the  $E_6$  gauge group with two additional neutral massive spin 1 gauge bosons that can mix with an arbitrary mixing angle. The  $Z'_\psi$ ,  $Z'_\chi$ ,  $Z'_\eta$ ,  $Z'_I$ ,  $Z'_{sec}$ , and  $Z'_N$  correspond to specific values of the mixing angle if there is just one  $Z'$  at low energy. The RS model predicts a series of neutral spin 2 resonances, the lightest of which is the RS graviton. We test this model assuming one extra dimension, in the  $k/\overline{M}_{Pl}$  range between 0.01 and 0.1 [3], where  $k$  is the curvature of the extra dimension and  $\overline{M}_{Pl}$  is the reduced effective Planck scale.

In a recent publication, the CDF Collaboration set limits on these models by analyzing the  $e^+e^-$  invariant mass distribution with  $1.3 \text{ fb}^{-1}$  of integrated luminosity [4]. Using a data set twice as large ( $2.5 \text{ fb}^{-1}$ ), this Letter describes a search for  $e^+e^-$  resonances in the invariant mass range of  $150\text{--}1,000 \text{ GeV}/c^2$ , and we set upper limits on  $\sigma(p\bar{p} \rightarrow X) \cdot \mathcal{B}(X \rightarrow e^+e^-)$  at the 95% credibility level (C.L.) where  $X$  is a spin 1 or spin 2 particle. We also set lower mass bounds on the  $Z'$  with SM coupling, the  $Z'$ 's in the  $E_6$  model, and the RS graviton.

This analysis is based on data collected with the CDF II detector, for which a full description can be found elsewhere [5]. The relevant components of the detector for this analysis are the tracking system and the calorimeters. The tracking system consists of a 96 layer

drift chamber, called the central outer tracker (COT), surrounding an eight-layer silicon tracker. Both are inside a 1.4 T solenoidal magnet. The COT covers the range of pseudorapidity  $|\eta| < 1.1$  [6], and the silicon tracker covers  $|\eta|$  up to 2.0. The electromagnetic (EM) and hadronic calorimeters, which are sandwiches of lead (EM) or iron (hadronic) absorber and plastic scintillator. They are outside the magnet, and are divided into a central calorimeter ( $|\eta| < 1.1$ ) and two plug calorimeters ( $1.1 < |\eta| < 3.6$ ). Both the central and the plug EM calorimeters have fine-grained shower profile detectors at EM shower maximum.

Three parallel on-line event selection criteria (triggers) are used to select the data. The first trigger requires any two EM clusters with  $E_T$  [6] greater than 18 GeV in the calorimeter, the second one requires a central EM cluster with  $E_T$  greater than 70 GeV, and the last one requires one central EM cluster that has  $E_T$  greater than 18 GeV and passes “loose electron” [7] selection criteria. For the last two triggers, a well-measured track based on the COT pointing to an energy deposit in the calorimeter is required. With these three triggers, the combined efficiency to collect events that pass the off-line selections is 100%.

Off-line events are required to have two isolated electrons, one in the central EM calorimeter and the other one in either the central (CC) or the plug (CP) EM calorimeters. Only electrons with  $E_T$  greater than 25 GeV and  $|\eta| < 2$  are used in order to ensure 100% trigger efficiency and coverage by the the silicon tracker. Electrons in the central EM calorimeter are required to have a well-measured track in the COT system pointing at an energy deposit in the calorimeter. For electrons in the plug EM calorimeter, the track association uses a calorimeter-seeded silicon-tracking algorithm [8]. An opposite-charge requirement is applied to electron-objects pairs detected in the central EM calorimeter. No such requirement is applied when one electron is detected in the plug, where  $\eta$ -dependent charge misidentification occurs. The 28 cm radial range of silicon at  $|\eta| > 1.1$ , where the COT coverage is incomplete, is insufficient to determine accurately the curvature for high  $p_T$  tracks, as predicted by simulation. Events with both electrons in the plug EM calorimeter are not considered in this Letter since adding them gains little sensitivity.

There are three sources of background. One is Drell-Yan production of  $e^+e^-$  pairs (DY), which is the dominant source of background and is irreducible. Another is dijets and  $W$ +jets production (referred to as “QCD” background) where one or more jets is misidentified as electron. Other contributions include  $Z/\gamma^* \rightarrow \tau^+\tau^-$ ,  $t\bar{t}$ , and diboson ( $W\gamma, WW, WZ, ZZ, \gamma\gamma$ ) production that collectively are referred to as “other SM” backgrounds.

The Monte Carlo (MC) event generator PYTHIA [9] Tune A [10] that performs a leading-order calculation is used to model the DY background. The simulated

---

College Dublin, Dublin 4, Ireland, <sup>k</sup>Royal Society of Edinburgh/Scottish Executive Support Research Fellow, <sup>l</sup>University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom, <sup>m</sup>Universidad Iberoamericana, Mexico D.F., Mexico, <sup>n</sup>Queen Mary, University of London, London, E1 4NS, England, <sup>o</sup>University of Manchester, Manchester M13 9PL, England, <sup>p</sup>Nagasaki Institute of Applied Science, Nagasaki, Japan, <sup>q</sup>University of Notre Dame, Notre Dame, IN 46556, <sup>r</sup>University de Oviedo, E-33007 Oviedo, Spain, <sup>s</sup>Simon Fraser University, Vancouver, British Columbia, Canada V6B 5K3, <sup>t</sup>Texas Tech University, Lubbock, TX 79409, <sup>u</sup>IFIC(CSIC-Universitat de Valencia), 46071 Valencia, Spain, <sup>v</sup>University of Virginia, Charlottesville, VA 22904, <sup>dd</sup>On leave from J. Stefan Institute, Ljubljana, Slovenia,

DY prediction is normalized to the data after subtracting other SM and QCD backgrounds in an invariant mass window from 76 to 106  $\text{GeV}/c^2$  for CC events and from 81 to 101  $\text{GeV}/c^2$  for CP events. Different mass windows are used because the QCD background in the CP events is higher than in the CC events and large uncertainty on QCD background estimation. We assign a 3.6% systematic uncertainty in the DY prediction to take into account the invariant-mass dependence of the  $k$ -factor [11] that is the difference between the leading and the next-to-next-to-leading order DY cross sections. The uncertainty in the DY prediction due to the choice of the parton distribution function set CTEQ6M [12] is evaluated using the Hessian method [13] and is found to be 3.7–6.4–13% (200–600–1,000  $\text{GeV}/c^2$ ) depending on the invariant mass.

The QCD background estimation is determined from the experimental data. The estimate is obtained using the probability for a jet to be misidentified as an electron. We measure this probability with a jet-triggered data sample. We then apply the misidentification probability to each jet in events with one good electron candidate and one or more jets. To estimate the dijet background contribution, events with  $W$  or  $Z$  candidates are removed from the sample before applying the jet misidentification probability (MP). The events with  $W$  candidate are identified with one good electron and a large missing transverse energy  $\cancel{E}_T$  [14] and the events with  $Z$  candidate are identified with two “loose electrons”. To estimate the  $W$ +jet background, events with  $Z$  candidate are removed and events with  $W$  candidate are retained. The dominant systematic uncertainty in the predicted QCD background is due to the 20% uncertainty in the jet MP.

Other SM contributions to the background are estimated with simulation samples generated with PYTHIA Tune A, except for the  $W+\gamma$  process. The  $W+\gamma$  process is generated with the matrix element generator WGAMMA [15]. These simulated samples are normalized to the product of the theoretical cross sections and the integrated luminosity. The systematic uncertainty for these other SM backgrounds is dominated by the 6% uncertainty in the integrated luminosity measurement [16] and 8% uncertainty in the theoretical cross sections [17].

The QCD and other SM backgrounds are small compared to the DY rate. Fig. 1 shows the observed  $e^+e^-$  invariant mass spectrum from 2.5  $\text{fb}^{-1}$  of data together with the expected backgrounds.

The dominant sources of systematic uncertainty in this analysis are the DY prediction, the luminosity, and the theoretical cross sections of other SM processes discussed above. Other systematic sources are the uncertainty on the scale factor of electron identification efficiency that comes from the difference between data and simulated events (1.3% for CC and 2.3% for CP events), the energy scale (1.0%), and the energy resolution (0.6% for CC and 0.3% for CP events), which affects the shape of the  $e^+e^-$




FIG. 1: Invariant mass distribution of  $e^+e^-$  events compared to the expected backgrounds. Dots with error bars are data. The dark shaded region represents “other SM” background, the light shaded region shows “QCD” background, and the white region corresponds to  $Z/\gamma^* \rightarrow e^+e^-$  background. The inset shows the same for the 240  $\text{GeV}/c^2$  region. The hatched histogram shows the shape of the expected signal from a 240  $\text{GeV}/c^2$  spin 1 particle (of negligible intrinsic width) on top of the total background. The hatched region is normalized to the number of excess events seen in the data.

invariant mass distribution. The uncertainty on the acceptance due to parton-distribution-function uncertainties is evaluated using the same method that was used for the DY prediction, and found to be 1.9% for CC and 0.6% for CP events.

The search for  $e^+e^-$  resonances in the high-mass range of 150–1,000  $\text{GeV}/c^2$  uses an unbinned likelihood ratio statistic,  $\lambda$ , defined in Eqs. 1–3 [18]:

$$\lambda = \frac{\max_{n_b \geq 0} \mathcal{L}_b}{\max_{n_b \geq 0, n_s \geq 0} \mathcal{L}_{s+b}}, \quad 0 \leq \lambda \leq 1, \quad 0 \leq -2 \ln \lambda \leq \infty \quad (1)$$

$$\mathcal{L}_{s+b} = \frac{(n_s + n_b)^N e^{-(n_s + n_b)}}{N!} \prod_i \frac{n_s S(x_i | \mu) + n_b B(x_i)}{n_s + n_b} \quad (2)$$

$$\mathcal{L}_b = \frac{n_b^N e^{-n_b}}{N!} \prod_i B(x_i). \quad (3)$$

where  $\mathcal{L}_b$  is the likelihood for a null hypothesis that is described by the SM only,  $\mathcal{L}_{s+b}$  is the likelihood for a test hypothesis that is described by physics beyond the SM together with the SM. The quantities  $n_s$  and  $n_b$  are the number of signal and background candidates which are determined by the fit and  $N$  is the number of candidates observed in data, each represented by a vector  $\{x_i\}$  of observables. The signal probability density function (PDF),  $S(x|\mu)$ , is a Gaussian with a floating mean  $\mu$  and a fixed

width, and  $B(x)$  is a background PDF obtained from the total background template. The widths of the signal PDF are determined from simulation ( $\sigma_{M_{ee}} = 0.8565 \text{ GeV}/c^2 + 0.0192 \cdot M_{ee}$  for  $M_{ee} > 150 \text{ GeV}/c^2$ ). The quantities  $\mathcal{L}_{s+b}$  and the  $\mathcal{L}_b$  are maximized separately without external background constraints. The function  $-2 \ln \lambda$  is calculated over the search range of 150–1,000  $\text{GeV}/c^2$  and the most prominent local maxima are listed in Table I. The most significant deviation between data and the SM prediction occurs at an invariant mass of 241.3  $\text{GeV}/c^2$  where  $-2 \ln \lambda$  is 14.4. The  $(\text{data} - \text{background})/\sigma_B$  [19] corresponding to the region of maximum  $-2 \ln \lambda$  is calculated by counting the number of observed events and estimated backgrounds within  $\pm 2 \sigma_{M_{ee}}$  of the maximum, and it is 3.8.

To estimate the probability of observing an excess equal to or greater than the maximum observed excess anywhere in the search range of 150–1,000  $\text{GeV}/c^2$ , we simulated 100,000 experiments assuming background only (null hypothesis). The distribution of maximum  $-2 \ln \lambda$  on these simulated experiments is shown in Fig. 2. Assuming only SM physics, the probability of observing a number of events equal to or greater than the observed excess is defined as the fraction of simulated experiments with maximum  $-2 \ln \lambda$  equal to or greater than 14.4, and is 0.6% which corresponds to the  $2.5 \sigma$  level of excess over the background.

TABLE I: The prominent local maxima in the search range of 150–1,000  $\text{GeV}/c^2$ .

$M_X$ ( $\text{GeV}/c^2$ )	241.3	272.7	478.9	725.2
$-2 \ln \lambda$	14.4	3.7	2.6	4.1




FIG. 2: Distribution of maximum  $-2 \ln \lambda$  in simulated experiments that assume only background. The arrow indicates the value observed in data:  $-2 \ln \lambda = 14.4$ .

Upper limits on  $\sigma(p\bar{p} \rightarrow X) \cdot \mathcal{B}(X \rightarrow e^+e^-)$  at the 95% C.L. are calculated as a function of mass using a Bayesian

binned likelihood method [20]. The likelihood is a function of the signal cross section and is given by a Poisson distribution. The likelihood is marginalized with gamma priors to allow for the uncertainty on the signal cross section due to uncertainty on the total signal efficiency and the background estimation. Then the posterior probability density function is formed with a constant prior for the signal cross section together with the likelihood. The limits are obtained by integrating the posterior probability density function for the signal cross section until we achieve the required fraction (95% for this analysis) of the total integral from zero to infinity. Fig. 3 (a) shows the observed upper limits from data and the expected limits from background-only simulated events for spin 1 particles as a function of the  $e^+e^-$  invariant mass, together with the expected cross sections for  $Z$ 's [21]. Fig. 3 (b) shows the same but for spin 2 particles, together with the expected cross sections for RS gravitons. The cross section lines for  $Z$ 's and RS gravitons are calculated at leading order with PYTHIA and then multiplied by a factor of 1.3 in order to approximate a next-to-leading-order prediction as done in reports of earlier results.

Table II shows the lower mass limits of the SM coupling and  $E_6$   $Z$ 's and Fig. 4 shows the excluded RS graviton mass region with respect to  $k/\overline{M}_{Pl}$ .

TABLE II: Expected and observed 95% C.L. lower limits on  $Z$ 's masses.

$Z'$ Model	$Z'_{SM}$	$Z'_\psi$	$Z'_X$	$Z'_\eta$	$Z'_I$	$Z'_{sec}$	$Z'_N$
Expected Limit ( $\text{GeV}/c^2$ )	961	846	857	928	755	788	831
Observed Limit ( $\text{GeV}/c^2$ )	963	851	862	930	735	792	837

To conclude, we have searched for  $e^+e^-$  resonances with  $2.5 \text{ fb}^{-1}$  of data collected by the CDF II detector. The largest excess over the standard model prediction is at an  $e^+e^-$  invariant mass of 240  $\text{GeV}/c^2$ . The probability of observing such an excess arising from fluctuation in the standard model anywhere in the mass range of 150–1,000  $\text{GeV}/c^2$  is 0.6%. We also set upper limits on  $\sigma(p\bar{p} \rightarrow X) \cdot \mathcal{B}$  at the 95% C.L. for spin 1 and spin 2 particles. The SM coupling  $Z'$  with mass below 963  $\text{GeV}/c^2$  and the  $E_6$   $Z$ 's with masses below 735/930 (lightest/heaviest)  $\text{GeV}/c^2$  are excluded at the 95% C.L. RS gravitons with masses below 848  $\text{GeV}/c^2$  are excluded at the 95% C.L. for  $k/\overline{M}_{Pl} = 0.1$ .

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium




FIG. 3: The upper limits on  $\sigma(p\bar{p} \rightarrow X) \cdot \mathcal{B}(X \rightarrow e^+e^-)$  as function of the mass of an  $X$  particle at the 95% C.L. where  $X$  is a spin 1 particle (a) or a spin 2 particle (b) together with model predictions.

für Bildung und Forschung, Germany; the Korean Science and Engineering Foundation and the Korean Research Foundation; the Science and Technology Facilities Council and the Royal Society, UK; the Institut National de Physique Nucleaire et Physique des Particules/CNRS; the Russian Foundation for Basic Research; the Ministerio de Ciencia e Innovación, and Programa Consolider-Ingenio 2010, Spain; the Slovak R&D Agency; and the Academy of Finland.

- 
- [1] D. London and J. L. Rosner, Phys. Rev. D **34**, 1530 (1986); F. del Aguila, M. Quiros, and F. Zwirner, Nucl. Phys. B **287**, 419 (1987); J. L. Rosner, Phys. Rev. D **35**, 2244 (1987); J. L. Hewett and T. G. Rizzo, Phys. Rept. **183**, 193 (1989); J. Erler, P. Langacker, and T. Li, Phys. Rev. D **66**, 015002 (2002); T. Han, P. Langacker, and B. McElrath, Phys. Rev. D **70**, 115006 (2004); J. Kang and P. Langacker, Phys. Rev. D **71**, 035014 (2005).  
 [2] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999).




FIG. 4:  $k/\overline{M}_{PI}$  as a function of RS graviton mass. The shading indicates the region excluded at the 95% credibility level.

- [3] H. Davoudiasl, J. L. Hewett, and T. G. Rizzo, Phys. Rev. Lett. **84**, 2080 (2000); B. C. Allanach *et al.*, J. High Energy Phys. **0212**, 039 (2002).  
 [4] T. Aaltonen *et al.*, (CDF Collaboration), Phys. Rev. Lett. **99**, 171802 (2007).  
 [5] A. Abulencia *et al.*, (CDF Collaboration), J. Phys. G: Nucl. Part. Phys. **34** 2457-2544 (2007).  
 [6] We use a cylindrical coordinate system where  $\theta$  is the polar angle with respect to the proton beam axis (positive  $z$  direction) and  $\phi$  is the azimuthal angle. The pseudorapidity is  $\eta = -\ln[\tan(\theta/2)]$ . We define transverse energy as  $E_T = E \sin \theta$  and transverse momentum as  $p_T = p \sin \theta$ , where  $E$  is the energy measured in the calorimeter and  $p$  is the magnitude of the momentum measured by the spectrometer.  
 [7] Throughout this Letter the charge conjugate state is implied.  
 [8] C. Issever, AIP Conf. Proc. **670**, 371 (2003).  
 [9] T. Sjöstrand *et al.*, Comput. Phys. Commun. **135**, 238 (2001).  
 [10] R. Field and R. C. Group, arXiv:hep-ph/0510198 (2005).  
 [11] R. Hamberg, W. L. van Neerven, and T. Matsuura, Nucl. Phys. B **359**, 343 (1991).  
 [12] H. L. Lai *et al.*, Phys. Rev. D **51**, 4763 (1995); J. Pumplin *et al.*, J. High Energy Phys. **0207**, 012 (2002).  
 [13] J. Pumplin *et al.*, Phys. Rev. D **65**, 014013 (2002).  
 [14] Missing transverse energy ( $\cancel{E}_T = |\vec{\cancel{E}}_T|$ ) is defined as  $\vec{\cancel{E}}_T = -\sum_i E_T^i \hat{n}_i$  where  $\hat{n}_i$  is a unit vector in the transverse plane that points from the beam-line to the  $i^{\text{th}}$  calorimeter tower.  
 [15] U. Baur and E. L. Berger, Phys. Rev. D **41**, 1476 (1990).  
 [16] D. Acosta *et al.*, Nucl. Instrum. Methods A **494**, 57 (2002);  
 [17] U. Baur, T. Han, and J. Ohnemus, Phys. Rev. D **48**, 5140 (1993); J. M. Campbell and R. K. Ellis, Phys. Rev. D **60**, 113006 (1999).  
 [18] L. Demortier, CERN Yellow Report, CERN-2008-001, 23 (2008); Y. Gao, L. Lu, and X. Wang, Eur. Phys. J. C **45** 659-667 (2006).  
 [19]  $\sigma_B$  is  $\sqrt{N_B + \sigma_{N_B}^2}$ , where  $N_B$  is the number of expected backgrounds and  $\sigma_{N_B}$  is the systematic uncertainty of the  $N_B$ .

[20] J. Heinrich *et al.*, arXiv:physics/0409129 (2004).

[21] C. Ciobanu *et al.*, Fermilab, Report No. FERMILAB-

FN-0773-E, (2005).