

Table I-23
AGE-ADJUSTED SEER INCIDENCE RATES AND TRENDS FOR THE TOP 15 CANCER SITES^a BY RACE/ETHNICITY

Both Sexes

	All Races		White		Black	
	Rate ^b 2000-2004	APC ^c 1995-2004	Rate ^b 2000-2004	APC ^c 1995-2004	Rate ^b 2000-2004	APC ^c 1995-2004
All Sites	470.1	-0.6*	477.5	-0.5	504.1	-1.0*
Prostate ^f	73.9	0.3	71.6	0.4	105.5	-1.1
Breast	69.6	-1.1*	71.5	-1.1	76.6	-1.6*
Lung and Bronchus	64.5	-1.4*	65.7	-1.3*	68.0	-0.6*
Colon and Rectum	51.6	-1.5*	51.2	-1.6*	62.1	-0.4
Urinary Bladder	21.1	0.0	23.0	0.1	15.0	-1.6*
Non-Hodgkin Lymphoma	19.3	0.1	21.6	1.8*	14.6	0.2
Melanoma of the Skin	18.5	1.2*	20.2	0.2	14.3	1.4
Kidney and Renal Pelvis	12.8	2.1*	13.3	2.3*	12.6	0.7
Corpus and Uterus, NOS ^f	12.6	-1.0*	13.1	-1.2*	12.5	-2.9*
Leukemia	12.3	-1.0*	12.8	-1.1*	11.3	1.3
Pancreas	11.4	0.0	11.2	0.3	11.3	-0.5
Oral Cavity and Pharynx	10.5	-1.5*	10.6	-1.3*	11.1	-3.0*
Thyroid	8.5	5.3*	8.9	5.7*	10.2	-0.2
Stomach	8.1	-1.5*	7.7	-1.4*	7.6	3.3*
Ovary ^{fh}	7.4	-1.3*	7.1	-1.4*	6.3	-5.0*
	Asian/Pacific Islander		American Indian/Alaska Native ^d		Hispanic ^e	
	Rate ^b 2000-2004	APC ^c 1995-2004	Rate ^b 2000-2004	APC ^c 1995-2004	Rate ^b 2000-2004	APC ^c 1995-2004
All Sites	314.9	-0.6*	297.6	-1.0	356.0	-0.7*
Breast	48.6	0.1	44.0	-1.2	60.8	0.0
Prostate ^f	41.6	-0.1	40.8	-0.5	48.1	-1.0
Colon and Rectum	41.6	-1.2*	37.6	-2.0	39.3	-0.9
Lung and Bronchus	39.4	-0.7*	30.1	-2.7	33.3	-2.1*
Stomach	14.3	-3.0*	14.7	1.4	16.5	-0.4
Liver & IBD ^g	13.9	0.0	11.5	-0.2	12.4	2.0*
Non-Hodgkin Lymphoma	13.2	0.2	10.4	0.4	12.3	-1.9*
Urinary Bladder	9.2	0.2	9.7	0.2	11.6	0.1
Pancreas	9.0	0.0	9.2	-1.1	10.5	-1.1
Corpus and Uterus, NOS ^f	8.8	0.2	8.1	-	9.7	1.0
Thyroid	8.5	2.3*	7.1	-3.5	9.6	-1.6*
Oral Cavity and Pharynx	7.9	-1.8	7.0	-	9.4	0.6
Leukemia	7.4	-1.1	6.4	1.1	7.5	4.4*
Kidney and Renal Pelvis	6.3	1.1	6.0	-3.4	7.1	-3.8*
Ovary ^{fh}	5.3	-0.2	5.3	0.5	6.2	-0.6

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Top 15 cancer sites selected based on 2000-2004 age-adjusted rates for the race/ethnic group.

^b Incidence data used in calculating the rates are from the 17 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^c The APC is the Annual Percent Change over the time interval. Incidence data used in calculating the trends are from the 13 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^d Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

^e Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

^f Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky.

^g The rates for sex-specific cancer sites are calculated using the population for both sexes combined.

^h IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

* Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

* The APC is significantly different from zero (p<.05).