

Table I-28  
AGE-ADJUSTED U.S. DEATH RATES AND TRENDS FOR THE TOP 15 CANCER SITES<sup>a</sup> BY RACE/ETHNICITY

	Females					
	All Races		White		Black	
	Rate <sup>b</sup> 2000-2004	APC <sup>c</sup> 1995-2004	Rate <sup>b</sup> 2000-2004	APC <sup>c</sup> 1995-2004	Rate <sup>b</sup> 2000-2004	APC <sup>c</sup> 1995-2004
All Sites	162.2	-1.0*	161.4	-0.9*	189.3	-1.1*
Lung and Bronchus	41.1	0.2*	42.1	0.3*	39.8	0.2
Breast	25.5	-2.3*	25.0	-2.4*	33.8	-1.6*
Colon and Rectum	16.4	-2.2*	15.9	-2.3*	22.9	-1.7*
Pancreas	9.2	0.1	9.2	-0.1	12.4	-0.6*
Ovary	8.9	-0.1	9.0	0.2	7.4	0.1
Non-Hodgkin Lymphoma	6.2	-2.7*	6.4	-2.8*	7.1	0.6
Leukemia	5.7	-0.9*	5.8	-0.8*	6.3	-1.3*
Corpus and Uterus, NOS	4.1	0.1	3.9	-1.0*	5.8	-3.2*
Brain and ONS <sup>f</sup>	3.6	-1.2*	3.9	0.0	5.3	-0.7
Myeloma	3.1	-1.0*	2.8	-0.9*	4.9	-4.7*
Liver & IBD <sup>f</sup>	3.1	0.9*	2.8	0.8*	4.3	-1.4*
Stomach	3.0	-2.7*	2.8	-0.7*	3.9	0.3
Kidney and Renal Pelvis	2.8	-0.8*	2.6	-2.8*	3.0	-2.9*
Cervix Uteri	2.6	-3.5*	2.3	-3.2*	2.8	-1.3*
Urinary Bladder	2.3	-0.5	2.3	-0.3	2.8	-0.9
	Asian/Pacific Islander		American Indian/Alaska Native <sup>d</sup>		Hispanic <sup>e</sup>	
	Rate <sup>b</sup> 2000-2004	APC <sup>c</sup> 1995-2004	Rate <sup>b</sup> 2000-2004	APC <sup>c</sup> 1995-2004	Rate <sup>b</sup> 2000-2004	APC <sup>c</sup> 1995-2004
All Sites	96.7	-1.4*	141.2	0.1	106.7	-0.9*
Lung and Bronchus	18.5	-1.2*	32.7	3.1*	16.1	-2.4*
Breast	12.6	-0.3	16.1	0.2	14.6	-0.3
Colon and Rectum	10.3	-1.9*	14.3	-2.4	11.1	-0.6
Pancreas	6.9	0.6	7.3	0.7	7.5	0.6
Liver & IBD <sup>f</sup>	6.7	0.4	7.1	2.2	6.1	0.5
Stomach	6.2	-4.6*	6.4	2.7	5.1	-1.8*
Ovary	4.8	0.7	5.5	-1.2	5.0	1.2
Non-Hodgkin Lymphoma	3.9	-1.6	5.0	2.2	4.8	-2.4*
Leukemia	3.0	-2.9*	4.3	-1.7	4.1	-1.1
Cervix Uteri	2.4	-4.5*	4.1	-	3.3	-3.3*
Corpus and Uterus, NOS	2.4	1.1	4.0	-1.6	3.1	-0.5
Brain and ONS <sup>f</sup>	1.5	-1.6	3.5	-4.1	2.6	-0.6
Myeloma	1.5	-1.4	2.9	-	2.4	0.3
Oral Cavity and Pharynx	1.3	-2.3	2.3	-	2.3	-0.3
Kidney and Renal Pelvis	1.1	-1.5	1.8	-	1.4	-5.2*

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

<sup>a</sup> Top 15 cancer sites selected based on 2000-2004 age-adjusted rates for the race/ethnic group.

<sup>b</sup> Mortality data used in calculating the rates are analyzed from a public use file provided by the National Center for Health Statistics (NCHS). Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>c</sup> The APC is the Annual Percent Change over the time interval. Mortality data used in calculating the trends are analyzed from a public use file provided by the National Center for Health Statistics (NCHS). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>d</sup> Rates for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

<sup>e</sup> Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. The 2000-2004 Hispanic death rates do not include deaths from Minnesota, New Hampshire and North Dakota. The 1995-2004 Hispanic mortality trends do not include deaths from Maine, Minnesota, New Hampshire, North Dakota, and Oklahoma.

<sup>f</sup> IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

\* The APC is significantly different from zero ( $p < .05$ ).