Table 173. Cancer—Estimated New Cases, 2008, and Survival Rates: 1987–1989 to 1996–2004

[1,437 represents 1,437,000. The 5-year relative survival rate, which is derived by adjusting the observed survival rate for expected mortality, represents the likelihood that a person will not die from causes directly related to their cancer within 5 years. Survival data shown are based on those patients diagnosed while residents of an area listed below during the time periods shown. Data are based on information collected as part of the National Cancer Institute's Surveillance, Epidemiology, and End Results (SEER) program, a collection of population-based registries in five states (Connecticut, Hawaii, Iowa, New Mexico, Utah) and four metropolitian areas (Atlanta. Detroit. San Francisco-Oakland. and Seattle-Puoet Soundi)

Site	Estimated new cases, 1 2008 (1,000)			5-year relative survival rates (percent)							
				White				Black			
	Total	Male	Female	1987– 1989	1990- 1992	1993– 1995	1996- 2004	1987– 1989	1990– 1992	1993- 1995	1996- 2004
All sites ² Lung ³ Breast ³ Colon and rectum. Colon Rectum Prostate Bladder Corpus uteri Non-Hodgkin's lymphoma ⁴	1,437 215 184 149 108 41 186 69 40 66	745 115 2 77 54 23 186 51 (X) 35	692 100 182 72 54 17 (X) 18 40 31	57.8 13.8 85.3 61.1 61.7 59.7 85.4 81.4 85.6 52.8	62.4 14.5 86.7 63.2 64.0 61.3 95.3 81.9 87.2 52.9	63.4 15.1 88.0 61.5 61.4 61.7 96.2 82.2 86.5 54.5	67.5 15.9 90.5 66.3 66.0 67.2 99.4 81.6 86.9 65.9	43.7 11.2 71.3 53.4 53.3 53.6 72.2 63.3 59.2 47.4	48.2 10.8 71.8 53.8 54.2 52.2 85.5 64.7 57.0 42.0	52.9 13.0 72.8 52.9 52.3 54.8 91.5 61.7 61.7 42.0	57.5 12.7 77.8 56.0 55.0 58.9 95.9 66.2 62.7 57.6
Oral cavity,4 and pharynx. Leukemia Melanoma of skin Pancreas Kidney Stomach Ovary. Cervix uteri 5	35 44 62 38 54 22 22	25 25 35 19 33 13 (X) (X)	10 19 28 19 21 8 22	56.6 45.5 88.3 3.4 58.5 19.1 39.9 73.6	58.7 48.0 89.6 4.7 62.1 19.3 42.4 71.9	60.9 49.5 89.7 4.2 63.1 20.6 42.7 74.7	62.4 52.2 92.1 4.9 66.6 23.1 45.3 73.7	34.4 36.7 79.9 5.9 55.8 20.0 35.2 58.2	33.3 37.4 60.3 3.7 57.7 23.8 37.8 58.8	38.2 42.3 66.8 3.7 58.6 19.8 43.3 64.0	41.7 42.4 77.9 4.6 66.1 24.5 38.3 65.3

X Not applicable. ¹ Estimates provided by American Cancer Society (<www.cancer.org>) are based on rates from the National Cancer Institute's SEER program. ² Includes other sites not shown separately. ³ Survival rates for female only. ⁴ All types combined. ⁵ Invasive cancer only.

Source: U.S. National Institutes of Health, National Cancer Institute; http://seer.cancer.gov/csr/1975_2004/>.