

III. Appendices

C. Food

4. Permissible Crop Translations for Pesticide Monitoring Data

Table III.C.4. Permissible Crop Translations for Pesticide Monitoring Data¹

Commodity Analyzed	Commodity translated to...	Comments
Potato	Subgroup 1-C	
Carrot	Subgroup 1-A or 1-C	
Head Lettuce	Cabbage, Chinese cabbage napa (tight headed varieties), Brussels sprouts, radicchio	All have a head morphology best represented by lettuce. All are in Subgroup 5-A except radicchio (4-A).
Broccoli	Cauliflower, Chinese broccoli, Chinese cabbage bok choy, Chinese mustard, kohlrabi	Broccoli better represents these heading, thickly stemmed and/or more branching cole crops than spinach does.
Spinach	Subgroup 4-A, Subgroup 5-B and Subgroup 4-B (except celery and fennel unless a strong case can be made)	Celery and fennel typically are excluded since residues may be higher in these crops due to the whorled, overlapping petioles which may retain spray residues.
Green Bean	Subgroups 6-A and 6-B	
Soybean	Subgroup 6-C	
Tomato or bell pepper	Group 8	All are fruiting vegetables.
Cucumber	Subgroup 9-B	All are cucurbit vegetables; residues in melon and pumpkin expected to be lower because of removal of rind.
Cantaloupe or Winter squash	Subgroup 9-A and pumpkin	
Orange	Group 10	Fruit will be peeled before analysis by PDP.
Apple or Pear	Group 11	All are pome fruits.
Peach	Group 12, except cherries (sweet and tart)	All are stone fruits.

Grape	Kiwifruit	Based on similar cultural practices.
Wheat	Group 15, except corn, rice, or wild rice	All are small grain crops or closely related thereto.
Milk	Meat	Metabolism study must indicate that residues in meat, fat, and meat-by-products will likely be equal to or lower than residues in milk. If dermal use is allowed on beef cattle, then it must be permitted and used on dairy cattle as well.
¹ Extracted from OPP/HED SOP 99.3		