


WHAT IS TICK-BORNE RELAPSING FEVER (TBRF)?

Relapsing Fever is a disease of relapsing episodes of fever caused by a spiral-shaped bacteria (spirochetes).

WHO CAN GET TBRF?

STATES WHERE TBRF OCCURS



Anybody of any age can develop relapsing fever. The disease occurs in the Western United States and is relatively uncommon. Most people become infected while vacationing in cabins in rural, wooded, mountainous areas. A few people have developed relapsing fever after being in caves.



HOW DOES SOMEONE GET TBRF?

In the United States, people get relapsing fever through the bite of an infected soft tick. These ticks, *Ornithodoros*, usually feed on small animals such as squirrels, mice, chipmunks, or rabbits. The ticks live in rodent nests under flooring and between walls. If these rodents are scarce, the ticks will take a meal from other warm-blooded animals including humans.




Soft ticks, unlike hard ticks, usually feed at night. Their bite is painless, and they feed between 15 and 30 minutes before they drop off. Most people are unaware that they have been bitten.

Relapsing fever can not be transmitted from person to person except in the case of pregnant women, who sometimes transmit the infection to their fetus.

WHAT ARE THE SYMPTOMS?

Symptoms usually develop within 7 days after being bitten. They include a sudden high fever with chills, sweats, headache, and body aches. Other possible symptoms include nausea, vomiting, loss of appetite (anorexia), dry cough, inability to tolerate bright lights (photophobia), rash, neck pain, eye pain, confusion and dizziness.



HOW IS TBRF DIAGNOSED AND TREATED?

When a person has fever it is often possible to see the spirochetes in a sample of the person's blood. Antibody tests are also available, but these are not useful in the early stages of the disease.

Although symptoms usually resolve on their own, treating relapsing fever with antibiotics will make the symptoms subside much more quickly.

If you think you may have relapsing fever consult a medical professional for further testing and treatment.

HOW CAN I PREVENT GETTING TBRF?

The best protection against relapsing fever is knowing where and how people get it and avoiding that exposure.

To prevent infection:

- Avoid sleeping in rodent infected buildings
- Use insect repellent containing DEET
- Rodent-proof buildings in areas where the disease is known to occur
- Identify and remove any rodent nesting material from walls, ceiling and floors
- In combination with removing the rodent material, fumigate the building with preparations containing pyrethrins and permethrins. More than one treatment is often needed to rid the building of the soft-ticks.

For more information contact either the CDC at 800-311-3435 or your local or state health department

Department of Health & Human Services
Centers for Disease Control and Prevention (CDC)
Division of Vector-Borne Infectious Diseases
PO Box 2087
Fort Collins, CO 80522-2087

TICK-BORNE RELAPSING FEVER



WHAT IS IT AND HOW TO PREVENT IT