

ROS signaling: the new wave?

Ron Mittler^{1,2}, Sandy Vanderauwera^{3,4}, Nobuhiro Suzuki¹, Gad Miller⁵,
Vanessa B. Tognetti^{3,4}, Klaas Vandepoele^{3,4}, Marty Gollery⁶, Vladimir Shulaev¹ and
Frank Van Breusegem^{3,4}

¹ Department of Biological Sciences, College of Arts and Sciences, University of North Texas, 1155 Union Circle #305220, Denton, TX 76203, USA

² Department of Plant Sciences, Hebrew University of Jerusalem, Jerusalem 91904, Israel

³ Department of Plant Systems Biology, VIB, 9052 Ghent, Belgium

⁴ Department of Plant Biotechnology and Genetics, Ghent University, 9052 Ghent, Belgium

⁵ Faculty of Life Sciences, Bar-Ilan University, Ramat-Gan, 52900, Israel

⁶ Tahoe Informatics, 1230 Styria Way, Incline Village, NV 89451, USA

Reactive oxygen species (ROS) play a multitude of signaling roles in different organisms from bacteria to mammalian cells. They were initially thought to be toxic byproducts of aerobic metabolism, but have now been acknowledged as central players in the complex signaling network of cells. In this review, we will attempt to address several key questions related to the use of ROS as signaling molecules in cells, including the dynamics and specificity of ROS signaling, networking of ROS with other signaling pathways, ROS signaling within and across different cells, ROS waves and the evolution of the ROS gene network.

Origins of the reactive oxygen species (ROS) network

It is easy to imagine how cells had to acquire different antioxidants and ROS scavenging/detoxifying enzymes during evolution to cope with the increased levels of atmospheric oxygen that accompanied the appearance of oxygen-evolving microorganisms on Earth billions of years ago [1–3]. It is nevertheless harder to imagine how ROS with their toxic potential can play such a current key signaling role in cells. When considering the evolution of ROS as important signaling molecules we can assume that once cells learned to deal with ROS toxicity, they were able to utilize ROS for signaling purposes (Box 1). Moreover, we can also assume that there are numerous advantages for using ROS as signaling molecules. What is it therefore that makes ROS such good signaling molecules (after all if cells evolved to use them as such they must have their advantages)?

Several possible advantages come to mind when considering the use of ROS as signaling molecules. These include the capacity of the cell to rapidly produce and scavenge different forms of ROS in a simultaneous manner, enabling rapid and dynamic changes in ROS levels (caused by simply tilting the balance between cellular production and scavenging rates). Another advantage could be a tight control over the subcellular localization of ROS signals in cells. If we assume a significant capacity of cells to detoxify/scavenge or buffer ROS throughout the cell, then local increases in ROS production can be limited to particular

locations of the cell, such as a certain membrane patch, or an organelle, making the spatial control of ROS accumulation highly specific. Detection of oscillating ROS signals in root hairs clearly demonstrates such capability [4,5]. Another advantage of ROS is that they could be used as rapid long distance auto-propagating signals transferred throughout the plant. Each individual cell along the path of the signal could activate its own ROS producing mechanism(s) in an autonomous manner carrying a ROS signal over long distances. It was recently reported for example that such a signal could propagate at a rate of up to 8.4 cm/min in *Arabidopsis* (*Arabidopsis thaliana*) [6]. An additional signaling advantage of ROS is that different forms of ROS exist, with significantly different molecular properties. For example, superoxide is a charged molecule under most physiological conditions and could not passively transfer across a membrane. By contrast, superoxide could be easily converted into hydrogen peroxide (H₂O₂) that readily transfers across membranes passively or through water channels [7]. Superoxide and H₂O₂ can also mediate the formation of lipid peroxides that would be membrane soluble. Thus, ROS have the advantage of being versatile signaling molecules with regard to their properties and mobility within cells. Moreover, as part of a cellular signaling network, ROS could be integrated with several different signaling pathways. Links with calcium and protein phosphorylation networks have been extensively studied, for example in the case of the ROS-generating respiratory burst oxidase (RBOH) NADPH oxidase proteins that contain an EF-calcium binding as well as phosphorylation domain(s) [8,9]. In addition, ROS levels are linked with cellular redox networks, for example through thioredoxins, peroxiredoxins, glutaredoxins and/or NADPH [10–12].


Another key signaling advantage of ROS is their tight link to cellular homeostasis and metabolism. Almost any change in cellular homeostasis could lead to a change in the steady-state level of ROS in a particular compartment(s). Physiological conditions that favor photorespiration would for example cause enhanced production of ROS in peroxisomes [13]. It is easy to envision how a tight link between metabolism and ROS levels would make ROS good signals to monitor changes in cellular metabolism. It is

Corresponding author: Mittler, R. (ron.mittler@unt.edu).

Box 1. An evolutionary view of the ROS gene network

An insight into the evolution of ROS signaling can be obtained by studying the ROS gene network of *Arabidopsis* [2] in a phylogenetic context, comparing it to several other representatives of the green plant lineage. In this regard, genome data of four dicots [*Arabidopsis*, poplar (*Populus trichocarpa*), grapevine (*Vitis vinifera*) and soybean (*Glycine max*)], four monocots [maize (*Zea mays*), rice (*Oryza sativa*), millet (*Sorghum vulgare*) and *Brachypodium distachyon*], a vascular non-seed plant (*Selaginella moellendorffii*), a moss (*Physcomitrella patens*) and four green algae (*Chlamydomonas reinhardtii*, *Volvox carteri*, *Micromonas* sp. *RCC299* and *Ostreococcus tauri*) were compared using the PLAZA platform [57]. Although the genomes of moss and green algae contain a smaller number of genes compared to flowering plants, they provide an excellent starting point to: (i) reconstruct the ancestral set of ROS-related genes at a certain time point during evolution and (ii) trace back the origin of newly acquired genes. Figure 1a summarizes the distribution of genes within the ROS-producing and ROS-scavenging families [2] across various

species. (A complete list with gene identifier codes for each family and species can be viewed in Table S2 in the supplementary material online). This comparative analysis revealed that, except for catalase, all genomes of the green plant lineage encode members of each ROS-scavenging family. By contrast, a complete absence of the ROS-producing NADPH oxidase gene family was observed in the algal genomes. Only from mosses on, plants appeared to have acquired genes to encode this gene family which then strongly expanded within the vascular plants. This family expansion might have been associated with the need for a more complex signaling network to coordinate multicellular growth, morphological complexity and biotic/abiotic stress responses. The evolutionary analysis of the ROS gene network in plants could therefore suggest that, at least with regard to the NADPH oxidase family, ROS-scavenging mechanisms were acquired before ROS-producing mechanisms, and that plants first learned to control their intracellular ROS levels and only then started using ROS for signaling purposes (Figure 1b).



TRENDS in Plant Science

Figure 1. An evolutionary framework of the ROS gene network within the green plant lineage. (a) Representation of ROS gene network genes in different species. For each ROS gene network family in *Arabidopsis* [2], orthologous genes in other species were determined using PLAZA 2.0, a comparative genomics resource to study gene and genome evolution in plants [57]. Starting from the *Arabidopsis* Genome Initiative codes, homologous gene families and phylogenetic profiles, reporting the gene distribution over the different species, were identified. Putative family members in other species were evaluated using PLAZA similarity heat maps, multiple sequence alignments, shared protein domain organization and phylogenetic trees. The absence of gene families in certain species was manually validated using dedicated sequence similarity searches against the genome sequence or assembled cDNA sequences. Circle size represents absolute number of genes in each family and species, whereas colors indicate relative family size (calculated as the ratio of the absolute number of genes divided by the average gene family size of each species). Abbreviations: ath, *Arabidopsis thaliana*; bdi, *Brachypodium distachyon*; cre, *Chlamydomonas reinhardtii*; gma, *Glycine max*; mrcc299, *Micromonas* sp. *RCC299*; osa, *Oryza sativa*; ota, *Ostreococcus tauri*; ppa, *Physcomitrella patens*; ptr, *Populus trichocarpa*; smo, *Selaginella moellendorffii*; sbi, *Sorghum bicolor*; vca, *Volvox carteri*; vvi, *Vitis vinifera*; zma, *Zea mays*. (b) A hypothetical model showing the different driving forces behind the ordered network evolution of ROS signaling in plants. A cell at an early evolutionary stage is shown as a circle at the top of the model. An increase in atmospheric pressure [O₂] is shown to drive the development of constitutive ROS buffering/scavenging systems (⊥). This is followed by the acquisition of simple signal transduction systems that include a sensing component (small semicircles linked with a dashed line to the buffering/scavenging systems) and finally ROS producing mechanisms (red circles), culminating in an advanced stage cell (bottom of model) that contains multiple ROS scavenging and ROS producing systems linked into a ROS signaling network.




Figure 1. Dynamics and specificity in ROS signaling. **(a)** Time-lapse imaging of a ROS wave propagating through an *Arabidopsis* plant in response to wounding. Wounding site is indicated with a white arrow. Imaging of the ROS wave was obtained using a luciferase reporter gene fused to the promoter of the zinc finger protein ZAT12 as described in [6]. **(b)** Model for the extracellular propagation of the ROS wave [6]. Each cell along the path of the wave activates its NADPH oxidase (RbohD) and produces ROS in an autonomous manner resulting in an auto-propagating ROS wave. **(c)** Different models for ROS signaling specificity. Different signals (A, B, C, D) are transmitted between cells, or between different organelles within a cell, via ROS and other signaling pathways such as calcium, MAPK pathways and others.

contributed to the evolution of ROS as key signaling molecules (Box 1).

The dynamics of ROS signaling

The dynamic and rapid nature of ROS signaling in cells is a result of the contrasting processes of ROS production and scavenging [1–3,14]. Because these two processes always occur in cells in a simultaneous manner, tipping the balance between scavenging and production rates would

result in rapid alterations in ROS levels that will generate a signal. In many biological systems a burst of ROS, often occurring as two distinguished peaks, accompanies several different signaling events [15]. Nevertheless, new research using advanced imaging tools, such as a luciferase reporter gene expressed under the control of a rapid ROS-response promoter in plants [6], or a new H₂O₂/redox-GFP sensor in Zebrafish [16], revealed that the initial burst of ROS production could trigger a cascade of cell-to-cell communication

events that results in the formation of a ROS wave that propagates throughout the different tissues and carries the signal over long distances (Figure 1a; Movie S1 in the supplementary material online). Thus, the temporal concept of a ROS burst occurring in selected cells could now be modified into a temporal-spatial concept of a ROS wave. We can therefore envision ROS signaling as a dynamic process that occurs within cells between different organelles, as well as between cells over long distances.

Because plants have a high capacity to scavenge ROS, the long distance aspect of ROS signaling can only be explained by the continuous production of ROS in individual cells along the path of the ROS wave/signal. Such a mechanistic view will imply that the ROS wave is auto-propagating. It has recently been shown [6] that a ROS wave triggered by different stimuli can be blocked by the local application of catalase or an NADPH oxidase inhibitor, at distances that are up to 5–8 cm away from the signal initiation site. Moreover, the signal requires the presence of the NADPH oxidase *RbohD* gene and spreads throughout the plant in both the upper and lower directions [6] (Movie S1 in the supplementary material online). These experiments clearly demonstrate the auto-propagating nature of the ROS wave. The initiation of the wave in specific cells must therefore be associated with a long distance signal that causes individual cells along its path to activate ROS production via their own ROS producing mechanisms (Figure 1b).

The activation of systemic signals by local application of high light stress was recently shown to be accompanied by plasma membrane electrical signals in a light wavelength specific manner [17]. Because membrane potential could be directly affected by ROS and because the rate of certain electric signals in plants matches the rate of the ROS wave as previously reported [6], it is possible that the generation of a ROS wave affects the formation, amplitude and/or rate of the electrical signal. Further research is needed to address this intriguing possibility.

Examples of localized changes in ROS levels and/or ROS oscillation patterns were reported for root hairs, guard cells, pollen-stigma interactions and cells interacting with different pathogens or pests [4,15,18,19]. These examples, as well as the new reports of ROS functioning as long distance auto-propagating signals [6], strengthen our view of ROS as highly dynamic signaling molecules.

Specificity in ROS signaling: how can it be?

The question of ROS signal specificity has been the focus of considerable attention recently [20,21]. The main point of debate concerns how can a ROS signal, generated in a particular compartment or a particular cell, be specific for a particular stimulus? For example, an increase in ROS levels in the chloroplast or peroxisome could result from light intensity, heat or cold stresses. How then could this increase act as a specific signal to trigger an appropriate acclimation response? Likewise, a ROS signal can be generated in a group of cells in the plant in response to wounding, pathogen attack or a local abiotic stress, and be transferred to the entire plant, but how could it be specific? One possibility is that ROS are mainly used as a general signal to prime or activate the cellular signaling network of cells and that other signals function together

with ROS to convey specificity (Figure 1c, top panel). These other signals could be small peptides, hormones, lipids, cell wall fragments and others. A different possibility is that the ROS signal itself carries within it a decoded message, much like calcium signals that have specific oscillation patterns within defined cellular locations (Figure 1c, middle panel). The specific features of the signal (amplitude, frequency and/or localization) could then be perceived and decoded by specialized mechanisms to trigger specific gene expression patterns. A third possibility is that each cellular compartment or individual cell has its own set(s) of ROS receptors to decode ROS signals generated within it, which are then transferred by other networks such as calcium and/or protein phosphorylation (Figure 1c, lower panel). As is usually the case with different biological systems, it is most probable that a combination of the different mechanisms described above, as well as others, function in cells and that ROS signaling cannot be evaluated as a stand-alone network but as an integrated signaling pathway that functions together with many different signaling networks. In future studies it would be interesting to determine whether different subnetworks within the entire cellular signaling apparatus could be defined, and whether these could be linked in a hierarchic manner to provide detailed understanding of the ROS signaling process. A strong link is also likely to exist between ROS signaling, the redox network of cells and the different antioxidant pools in different cells, or in a different subcellular compartment of cells. Unraveling the different interactions between these interconnected networks would be the first step in linking these networks with other cellular signaling pathways.

An interesting example of high specificity in ROS signaling is the analysis of double mutants deficient in APX1 and CAT2 in *Arabidopsis*, and APX1 and CAT1 in tobacco (*Nicotiana tabacum*) [13,22]. Compared to wild type, and the single *apx* or *cat* mutants, these double mutants are surprisingly more tolerant to different environmental conditions [13,22]. It was recently found that the combined lack of APX1 and CAT2 in *Arabidopsis* causes a unique ROS signature in cells that triggers a novel acclimation response involving the activation of DNA repair, cell cycle control and antiprogrammed cell death mechanisms. A similar response is not found in the *apx1* or *cat2* single mutants demonstrating the need for a specific ROS signature for its activation, a signature that is only found in the double mutants [13].

Deciphering the complex mode of ROS signaling within cells would require the development and utilization of many more mutants deficient in ROS signaling and the combined use of these mutants with advanced subcellular-specific ROS imaging tools. Advances in the study of ROS gene network genes are summarized in Table S1 in the supplementary material online. A highly complex mode of interaction between ROS signaling and growth, development and stress responses in plants is revealed. Especially notable are the multiple mutants, lacking two or more ROS metabolizing/signaling enzymes, which often produce unexpected results [22,23] (Table S1 in the supplementary material online). These can provide new insights into how plants balance and mediate ROS signaling, as well as reveal new and yet unknown ROS producing or scavenging mechanisms.

Networking of ROS signaling with other signaling pathways

As indicated above, ROS signaling is integrated with many different signaling networks in plants. These include protein kinase networks, calcium signaling, cellular metabolic networks and redox responses. In some instances, ROS accumulation was found to precede the activation of signaling through these networks, whereas in other examples ROS accumulation was found to be a direct result of signaling through these networks. A good example for a ROS-activated signaling network is the mitogen-activated protein kinase (MAPK) cascade.

Many different MAPKs cascades can be activated following ROS accumulation. These include the ROS-responsive MAPKKK MEKK1, MPK4 and MPK6 [19,24,25]. The MEKK1 pathway is highly active during abiotic and oxidative stress conditions, and MEKK1 is an activator of two highly homologous MAPKKs (MKK1 and MKK2), which function upstream of the MAPKs MPK4 and MPK6 [24,25]. MEKK1 was suggested to be specifically required for the activation of MPK4 by H₂O₂ [26]. Transcriptome analysis of MEKK1, MPK4 and MKK1/2 deficient mutants identified 20 different transcription factors previously classified by Gadjev *et al.* [27] as highly responsive to multiple ROS-inducing conditions [28]. Two other ROS-responsive MAP kinases MPK3 and MPK6 that are activated by H₂O₂ were recently shown to depend on MKK9 for their activity [24,29]. The guard cell highly expressed MPK9 and MPK12 was recently implicated as required for abscisic acid (ABA)-induced stomata closure functioning downstream to ROS signals and necessary for the activation of anion channels. MPK12 activity was also shown to increase in response to both ABA and H₂O₂ application [19]. MAPK pathways are also implicated in the induction of nitric oxide (NO) and ROS bursts and signaling, which synergistically function in defense responses and were recently shown to confer resistance to pathogens [30]. Both NO and ROS were reported to be produced simultaneously through the MAPK cascade MEK2–SIPK (salicylic acid-induced protein kinase) [30,31].

In contrast to MAPKs, the NADPH oxidase complex is a good example for ROS production that is driven by a compilation of different cellular signaling pathways. Pathways that proceed NADPH oxidase activation include calcium signaling and protein phosphorylation [8,9,32]. *Solanum tuberosum* StCDPK4 and StCDPK5, have been shown to phosphorylate StrbohB in potato (*Solanum tuberosum*) [8], and the ABA-activated SnRK2 protein kinase open stomata 1 (OST1) (SRK2E/SnRK2.6) was recently shown to act upstream of ROS in guard cell ABA signaling. OST1 physically interact with AtRbohF and phosphorylates it on Ser13 and Ser174 [33]. More recently, SLAC1, a protein essential for guard cell anion channel functioning was shown to be a substrate for OST1 phosphorylation. SLAC1 is essential for stomatal closure in response to ABA, CO₂, O₃, light–dark transitions and humidity change, and by Ca²⁺, H₂O₂ and NO [34]. AtRBOHD was shown to be directly phosphorylated *in vivo*, which together with Ca²⁺ binding synergistically activates its enzymatic activity [9]. Two MAPK cascades, MEK2–SIPK and MEK1–NTF6, were suggested to be involved in the induction of *NbRbohB* gene and protein activities in *Nicotiana benthamiana* [30].

Links between ROS signaling and cellular redox were proposed to be mediated by peroxiredoxins, NADPH, kinase activity or the detection of oxidized proteins and peptides [10–12,20]. Oxidation of methionine (Met) residues to Met sulfoxide (MetSO) in kinase substrate proteins, such as nitrate reductase, can inhibit the phosphorylation of nearby sites and thereby couple oxidative signals to changes in protein phosphorylation [35]. Direct coupling of ROS signaling with primary cellular metabolism is a key feature of ROS signaling in cells. ROS accumulation for example can cause an inhibition of the tricarboxylic acid cycle (TCA) in mitochondria and upregulation of glycolysis and oxidative pentose phosphate pathways [36–38]. In chloroplasts, ROS signaling is coupled to the redox state of the plastoquinone (PQ) pool and plays an important role in the response of plants to changes in environmental conditions [39–41]. Plants optimize their photosynthetic activity by regulating the association of light harvesting complexes with thylakoids, and by adjusting photosystem stoichiometry to rearrange the balance of excitation energy [42]. The STN7 kinase was recently identified as a key regulator of these responses, linking them to the PQ redox state [42]. A chloroplast sensor kinase was also recently shown to be required for the regulation of gene expression in chloroplasts in response to changes in redox state of electron carriers connecting the two photosystems [43]. These studies expand our understanding of how redox and ROS levels balance some of the key metabolic pathways in plants.

Links between ROS and hormonal signaling are summarized in Box 2. These demonstrate a complex interaction between ROS signaling, environmental conditions and plant development. Overall it appears as if the more we study ROS signaling, the more we discover how intertwined it is with almost all aspects of plant signaling, physiology and development. It is probable that unraveling the different interactions that tie the different signaling networks and pathways of plants will be a challenge for many years to come.

Imaging of ROS signaling

Recent developments in cellular imaging and real-time detection tools have advanced our understanding of ROS metabolism and cellular dynamics in human, animal and microbial systems. Among these are two-photon fluorescence microscopy, matrix-assisted laser desorption/ionization mass spectrometry (MALDI-MS) imaging, atomic force microscopy and optical tweezers force spectroscopy [44]. Although these techniques have proved valuable in measuring the level of ROS and other metabolites in different organisms, their use in plants is very limited. The main imaging tools that have been used in plants are the green fluorescent protein (GFP)-based redox probe roGFP, fluorescent dyes and luciferase [6,45–49]. Fluorescence-based probes provide an important means for non-invasive quantitative monitoring of redox changes and include redox-sensitive yellow fluorescent protein (rxYFP), roGFPs and the YFP-based probe HyPer [50–52], and have been successfully used to follow the cellular redox status in multiple biological systems. Application of some of the newer tools described above as well as development of


Box 2. Integration of ROS and hormonal signaling

ROS signaling is highly integrated with hormonal signaling networks (Figure 1), thereby allowing plants to regulate developmental processes, as well as adaptive responses to environmental cues. Similar to ROS, salicylic acid (SA) is involved in defense and cell death responses [58] and increased ROS levels can cause SA accumulation [59]. Interestingly, ablation of SA biosynthesis could prevent ROS-induced defense responses and cell death without affecting redox state [60]. A ROS burst is involved in SA-induced stomatal closure [61] and the crosstalk between ROS and ABA was extensively reported [62]. Interactions between ROS, NO and ethylene signaling were described in both a stress responsive and a developmental context [63,64]. Gibberellin (GA) signaling is linked with ROS contributing to the fine-tuning of ROS levels by stimulating the destruction of the nuclear growth-repressing DELLA proteins that regulate transcript levels of antioxidant enzymes [65].

Root apical meristems (RAMs) are rapidly reorganized in response to environmental triggers. This plasticity is linked to a complex signaling module in which ROS, NOS and antioxidants operate in strict association with hormonal signaling pathways such as auxin, GA and cytokinin [66]. RAM responses are modulated by localized ROS production in quiescent centers and the GA-DELLA signaling pathway [66]. GA-induced genes include the plant-specific GAST1-like genes which encode small proteins with a conserved cysteine-rich

domain possibly involved in redox regulation [67]. The stress response of C19-GA2ox genes, a major GA inactivating pathway, is another example of ROS effect on hormone homeostasis integrating extrinsic signals with developmental programs [68].

The integration of ROS with auxin signaling networks, triggered by environmental factors, is known as the stress-induced morphogenic response. In this response, ROS and auxin metabolism interfere and lead to morphological changes that help avoid deleterious effects of environmental stress [69,70]. These morphological changes can be partly explained by ROS affecting auxin homeostasis at different levels including oxidative degradation by stress-induced peroxidases, increased auxin catabolism, auxin transport and redistribution caused by altered expression and cellular location of PIN proteins [71–74]. A cohort of auxin responsive genes are differentially expressed during various stresses, and oxidative stress related genes are regulated differentially by auxin [70,75]. Auxins are also known to induce a programmed and cell-specific ROS generation or to regulate the level of antioxidants [76–78]. Two key regulators of redox homeostasis: NADP-linked thioredoxin (NTRX) and glutathione were recently found to alter both auxin transport and metabolism [79]. It would be interesting in future studies to identify additional convergence points between ROS and hormone signaling pathways.



TRENDS in Plant Science

Figure 1. ROS–auxin crosstalk in relation to other hormone signals. A schematic model showing coregulation and crosstalk between ROS and different hormonal signaling pathways in response to different environmental stimuli or developmental signals. Black and purple arrows indicate activation by abiotic or environmental and internal triggers, respectively, whereas red lines indicate repression. ROS can affect biosynthesis, degradation, metabolism, transport or perception of auxin. Abbreviations: ABA, abscisic acid; GA, gibberellins; IAA, auxins; MAPK, mitogen-activated protein kinase; ROS, reactive oxygen species; SA, salicylic acid.

new imaging tools in plants is highly needed to advance the study of ROS signaling in plants.

Future directions in ROS signaling research

Several possible research avenues come to mind when considering ROS signaling. The new concept of ROS waves, compared to ROS bursts, requires further research. How are these waves being propagated within and/or across

different cells? What is the degree of specificity communicated by these waves? How are they linked to changes in membrane potential? How are they even possible in light of the high capacity of cells to scavenge ROS?

The question of how ROS signals travel within or across different cells is also an intriguing one. It was previously reported that ROS can be formed within enclosed vesicles [53]. Is this a possible way to transfer ROS from one

subcellular location to another? Would these vesicles or other mechanisms transfer ROS signals along cytoskeleton fibers within cells? Are there yet unknown ROS producing enzymes/complexes that function within or on the surface of cells that could be associated with cytoskeleton components? Is the apoplasmic space an easier conducting media for transferring ROS signals, compared with the cytosol or vacuole?

The view of ROS as being produced by specific complexes, such as the NADPH complex, is an acknowledged concept that suggests a high degree of cellular regulation by different proteins that associate with the complex. Signal-specific sensing and transducing complexes were reported in plant and animal cells. However, a sensor and/or signal transducer complex for ROS related proteins has not yet been identified in plants. The identification of such a complex would be of immense importance to the field of ROS signaling research.

What can we learn from animal systems? Several interesting papers have recently been published describing the dynamic nature of mitochondrial-to-mitochondrial communication in heart muscle cells. In heart cells mitochondria are found in a lattice-like arrangement and communicate with each other via waves of ROS-induced-ROS-release [54]. This ROS-induced-ROS-release mechanism might be similar to the mechanism(s) that propagates cell-to-cell ROS signaling across long distances in plants [6]. Similarities between animal and plant models for wound-induced ROS waves can also be found between Zebrafish and *Arabidopsis* [6,16]. ROS signaling could therefore have broad similarities across different kingdoms, as is expected from an evolutionary conserved signaling pathway.

A mostly unanswered question in plant ROS research has been: How do plant cells sense ROS? Several different systems involving 'single-Cys' and 'two-Cys' redox sensors, as well as different variations on metal sensing proteins that involve His oxidation, or two-component receptors have been proposed in bacteria, algae and animal cells [2,55]. It is unclear, however, whether similar systems function in plants, and how are they integrated into the ROS signaling network of plant cells. It is probable that genetic screens for mutants impaired in ROS sensing will reveal some of these mechanisms.

The development of advanced imaging tools, such as MALDI-MS imaging, would enable the detection of ROS and antioxidant pools in specific tissues and/or subcellular compartments. These would be of key importance to our understanding of ROS signaling. Dynamic changes in the nuclear pool of glutathione were recently reported in plants [56]. Subcellular changes in glutathione, as well as other antioxidants and/or ROS, could explain many different regulatory events in plants linking redox status with ROS signaling and enabling rapid responses to changes in environmental conditions or different developmental cues.

Many unknown players in ROS signaling still exist. The vacuole, for example, could have a key buffering/signaling role in ROS metabolism. Organelle movement and the cytoskeleton could also play a central role in distributing ROS signals. Unknown ROS producing mechanisms could exist within plant cells as well as novel antioxidants. It is

probable that the increased effort to sequence many different plant genomes, coupled with a more direct research into gene function and ROS metabolism, will unravel many of these mechanisms. Because ROS are linked to many biotic and abiotic responses, deciphering ROS signaling is likely to have a significant impact on agriculture and biotechnology in many countries and could lead to the development of crops with enhanced yield under suboptimal conditions. The future of ROS research is very promising!

Acknowledgments

Supported by funding from The National Science Foundation (IBN-0420033, NSF-0431327, IOS-0639964 and IOS-0743954), University of North Texas College of Arts and Sciences, and EU grant FP7 – MARIE CURIE 447, Ghent University (Multidisciplinary Research Partnership 'Biotechnology for a Sustainable Economy' project no. 01MRB510W). S.V. is a postdoctoral fellow of the Research Foundation-Flanders. V.B.T. is the recipient of a Marie Curie Intra-European Fellowship for Career Development (PIEF-GA-2008-221427). Work in the laboratory of V.S. was supported in part by the National Science Foundation (NSF) Integrative Organismal Systems (IOS) grant 0820126, NFS Molecular and Cellular Biosciences (MCB) grant 0820823, National Institute of Health (NIH)-National Institute of Allergy and Infectious Diseases (NIAID) grant 2R01AI045774 and NIH National Cancer Institute (NCI) grant R01CA120170. KV is a Postdoctoral Fellow of the Research Foundation-Flanders (FWO) and acknowledges the support of Ghent University (Multidisciplinary Research Partnership "Bioinformatics: from nucleotides to networks").

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.tplants.2011.03.007](https://doi.org/10.1016/j.tplants.2011.03.007).

References

- Halliwell, B. and Gutteridge, J.M.C., eds (2007) *Free Radicals in Biology and Medicine* (4th edn), Clarendon
- Mittler, R. *et al.* (2004) The reactive oxygen gene network of plants. *Trends Plant Sci.* 9, 490–498
- Foyer, C.H. and Noctor, G. (2005) Redox homeostasis and antioxidant signaling: a metabolic interface between stress perception and physiological responses. *Plant Cell* 17, 1866–1875
- Monshausen, G.B. *et al.* (2007) Oscillations in extracellular pH and reactive oxygen species modulate tip growth of *Arabidopsis* root hairs. *Proc. Natl. Acad. Sci. U.S.A.* 104, 20996–21001
- Takeda, S. *et al.* (2008) Local positive feedback regulation determines cell shape in root hair cells. *Science* 319, 1241–1244
- Miller, G. *et al.* (2009) The plant NADPH oxidase RBOHD mediates rapid systemic signaling in response to diverse stimuli. *Sci. Signal.* 2, ra45
- Miller, E.W. *et al.* (2010) Aquaporin-3 mediates hydrogen peroxide uptake to regulate downstream intracellular signaling. *Proc. Natl. Acad. Sci. U.S.A.* 107, 15681–15686
- Kobayashi, M. *et al.* (2007) Calcium-dependent protein kinases regulate the production of reactive oxygen species by potato NADPH oxidase. *Plant Cell* 19, 1065–1080
- Ogasawara, Y. *et al.* (2008) Synergistic activation of the *Arabidopsis* NADPH oxidase AtrbohD by Ca²⁺ and phosphorylation. *J. Biol. Chem.* 283, 8885–8892
- Moon, H. *et al.* (2003) NDP kinase 2 interacts with two oxidative stress-activated MAPKs to regulate cellular redox state and enhances multiple stress tolerance in transgenic plants. *Proc. Natl. Acad. Sci. U.S.A.* 100, 358–363
- Dietz, K.J. *et al.* (2010) Hubs and bottlenecks in plant molecular signalling networks. *New Phytol.* 188, 919–938
- Rouhier, N. (2011) Plant glutaredoxins: pivotal players in redox biology and iron-sulphur centre assembly. *New Phytol.* 186, 365–372
- Vanderauwera, S. *et al.* (2011) Extracellular protection of chromosomal DNA from oxidative stress. *Proc. Natl. Acad. Sci. U.S.A.* 108, 1711–1716

- 14 Mittler, R. (2002) Oxidative stress, antioxidants, and stress tolerance. *Trends Plant Sci.* 7, 405–410
- 15 Nishimura, M.T. and Dangl, J.L. (2010) Arabidopsis and the plant immune system. *Plant J.* 61, 1053–1066
- 16 Niethammer, P. *et al.* (2009) A tissue-scale gradient of hydrogen peroxide mediates rapid wound detection in zebrafish. *Nature* 459, 996–999
- 17 Szechyńska-Hebda, M. *et al.* (2010) Evidence for light wavelength-specific photoelectrophysiological signaling and memory of excess light episodes in Arabidopsis. *Plant Cell* 22, 2201–2218
- 18 McInnis, S.M. *et al.* (2006) Production of reactive oxygen species and reactive nitrogen species by angiosperm stigmas and pollen: potential signalling cross-talk. *New Phytol.* 172, 221–228
- 19 Jammes, F. *et al.* (2009) MAP kinases MPK9 and MPK12 are preferentially expressed in guard cells and positively regulate ROS-mediated ABA signaling. *Proc. Natl. Acad. Sci. U.S.A.* 106, 20520–20525
- 20 Møller, I.M. and Sweetlove, L.J. (2010) ROS signaling–specificity is required. *Trends Plant Sci.* 15, 370–374
- 21 Triantaphylidès, C. and Havaux, M. (2009) Singlet oxygen in plants: production, detoxification and signaling. *Trends Plant Sci.* 14, 219–228
- 22 Rizhsky, L. *et al.* (2002) Double antisense plants lacking ascorbate peroxidase and catalase are less sensitive to oxidative stress than single antisense plants lacking ascorbate peroxidase or catalase. *Plant J.* 32, 329–342
- 23 Miller, G. *et al.* (2007) Double mutants deficient in cytosolic and thylakoid ascorbate peroxidase reveal a complex mode of interaction between reactive oxygen species, plant development, and response to abiotic stresses. *Plant Physiol.* 144, 1777–1785
- 24 Xing, Y. *et al.* (2008) AtMKK1 mediates ABA-induced CAT1 expression and H₂O₂ production via AtMPK6-coupled signaling in Arabidopsis. *Plant J.* 54, 440–451
- 25 Teige, M. *et al.* (2004) The MKK2 pathway mediates cold and salt stress signaling in Arabidopsis. *Mol. Cell* 15, 141–152
- 26 Nakagami, H. *et al.* (2006) A mitogen-activated protein kinase kinase mediates reactive oxygen species homeostasis in Arabidopsis. *J. Biol. Chem.* 281, 38697–38704
- 27 Gadjev, I. *et al.* (2006) Transcriptomic footprints disclose specificity of reactive oxygen species signaling in Arabidopsis. *Plant Physiol.* 141, 436–445
- 28 Pitzschke, H. *et al.* (2009) A major role of the MEKK1–MKK1/2–MPK4 pathway in ROS signalling. *Mol. Plant* 2, 120–137
- 29 Xu, J. *et al.* (2008) Activation of MAPK kinase 9 induces ethylene and camalexin biosynthesis and enhances sensitivity to salt stress in Arabidopsis. *J. Biol. Chem.* 283, 26996–27006
- 30 Asai, S. *et al.* (2008) MAPK signaling regulates nitric oxide and NADPH oxidase-dependent oxidative bursts in *Nicotiana benthamiana*. *Plant Cell* 20, 1390–1406
- 31 Asai, S. *et al.* (2010) A key enzyme for flavin synthesis is required for nitric oxide and reactive oxygen species production in disease resistance. *Plant J.* 62, 911–924
- 32 Demidchik, V. *et al.* (2009) Plant extracellular ATP signalling by plasma membrane NADPH oxidase and Ca²⁺ channels. *Plant J.* 58, 903–913
- 33 Sirichandra, C. *et al.* (2009) Phosphorylation of the Arabidopsis AtrbohF NADPH oxidase by OST1 protein kinase. *FEBS Lett.* 583, 2982–2986
- 34 Vahisalu, T. *et al.* (2010) Ozone-triggered rapid stomatal response involves the production of reactive oxygen species, and is controlled by SLAC1 and OST1. *Plant J.* 62, 442–453
- 35 Hardin, S. *et al.* (2009) Coupling oxidative signals to protein phosphorylation via methionine oxidation in Arabidopsis. *Biochem. J.* 422, 305–312
- 36 Umbach, A.L. *et al.* (2005) Characterization of transformed Arabidopsis with altered alternative oxidase levels and analysis of effects on reactive oxygen species in tissue. *Plant Physiol.* 139, 1806–1820
- 37 Baxter, C.J. *et al.* (2007) The metabolic response of heterotrophic Arabidopsis cells to oxidative stress. *Plant Physiol.* 143, 312–325
- 38 Rasmusson, A.G. and Wallström, S.V. (2010) Involvement of mitochondria in the control of plant cell NAD(P)H reduction levels. *Biochem. Soc. Trans.* 38, 661–666
- 39 Pfannschmidt, T. *et al.* (2009) Potential regulation of gene expression in photosynthetic cells by redox and energy state: approaches towards better understanding. *Ann. Bot.* 103, 599–607
- 40 Mühlenbock, P. *et al.* (2008) Chloroplast signaling and LESION SIMULATING DISEASE1 regulate crosstalk between light acclimation and immunity in Arabidopsis. *Plant Cell* 20, 2339–2356
- 41 Li, Z. *et al.* (2009) Sensing and responding to excess light. *Annu. Rev. Plant Biol.* 60, 239–260
- 42 Pesaresi, P. *et al.* (2009) Arabidopsis STN7 kinase provides a link between short- and long-term photosynthetic acclimation. *Plant Cell* 21, 2402–2423
- 43 Puthiyaveetil, S. *et al.* (2008) The ancestral symbiont sensor kinase CSK links photosynthesis with gene expression in chloroplasts. *Proc. Natl. Acad. Sci. U.S.A.* 105, 10061–10066
- 44 Anselmetti, D. *et al.* (2007) Analysis of subcellular surface structure, function and dynamics. *Anal. Bioanal. Chem.* 387, 83–89
- 45 Meyer, A.J. and Brach, T. (2009) Dynamic redox measurements with redox-sensitive GFP in plants by confocal laser scanning microscopy. *Methods Mol. Biol.* 479, 93–107
- 46 Meyer, A.J. and Dick, T.P. (2010) Fluorescent protein-based redox probes. *Antioxid. Redox Signal.* 13, 621–650
- 47 Mullineaux, P.M. and Lawson, T. (2009) Measuring redox changes *in vivo* in leaves: prospects and technical challenges. *Methods Mol. Biol.* 476, 65–75
- 48 Maughan, S.C. *et al.* (2010) Plant homologs of the Plasmodium falciparum chloroquine-resistance transporter, PfCRT, are required for glutathione homeostasis and stress responses. *Proc. Natl. Acad. Sci. U.S.A.* 107, 2331–2336
- 49 Rosenwasser, S. *et al.* (2010) A fluorometer-based method for monitoring oxidation of redox-sensitive GFP (roGFP) during development and extended dark stress. *Physiol. Plant.* 138, 493–502
- 50 Belousov, V.V. *et al.* (2006) Genetically encoded fluorescent indicator for intracellular hydrogen peroxide. *Nat. Methods* 3, 281–286
- 51 Lohman, J.R. and Remington, S.J. (2008) Development of a family of redox-sensitive green fluorescent protein indicators for use in relatively oxidizing subcellular environments. *Biochemistry* 47, 8678–8688
- 52 Maulucci, G. *et al.* (2008) High-resolution imaging of redox signaling in live cells through an oxidation-sensitive yellow fluorescent protein. *Sci. Signal.* 1, pl3
- 53 Leshem, Y. *et al.* (2006) Suppression of Arabidopsis vesicle-SNARE expression inhibited fusion of H₂O₂-containing vesicles with tonoplast and increased salt tolerance. *Proc. Natl. Acad. Sci. U.S.A.* 103, 18008–18013
- 54 Zhou, L. *et al.* (2010) A reaction-diffusion model of ROS-induced ROS release in a mitochondrial network. *PLoS Comput. Biol.* 6, e1000657
- 55 Giedroc, D.P. (2009) Hydrogen peroxide sensing in *Bacillus subtilis*: it's all about the (metal)regulator. *Mol. Microbiol.* 73, 1–4
- 56 Diaz Vivanco, P. *et al.* (2010) A nuclear glutathione cycle within the cell cycle. *Biochem. J.* 431, 169–178
- 57 Proost, S. *et al.* (2009) PLAZA, a comparative genomics resource to study gene and genome evolution in plants. *Plant Cell* 21, 3718–3731
- 58 Vlot, A.C. *et al.* (2009) Salicylic acid, a multifaceted hormone to combat disease. *Annu. Rev. Phytopathol.* 47, 177–206
- 59 Chamnongpol, S. *et al.* (1998) Defense activation and enhanced pathogen tolerance induced by H₂O₂ in transgenic tobacco. *Proc. Natl. Acad. Sci. U.S.A.* 95, 5818–5823
- 60 Chaouch, S. and Noctor, G. (2010) Myo-inositol abolishes salicylic acid-dependent cell death and pathogen defence responses triggered by peroxisomal hydrogen peroxide. *New Phytol.* 188, 711–718
- 61 Khokon, M.A. *et al.* (2011) Involvement of extracellular oxidative burst in salicylic acid-induced stomatal closure in Arabidopsis. *Plant Cell Environ.* 34, 434–443.
- 62 Wang, P. and Song, C.P. (2008) Guard-cell signalling for hydrogen peroxide and abscisic acid. *New Phytol.* 178, 703–718
- 63 Zago, E. *et al.* (2006) Nitric oxide- and hydrogen peroxide-responsive gene regulation during cell death induction in tobacco. *Plant Physiol.* 141, 404–411
- 64 Mühlenbock, P. *et al.* (2007) Lysigenous aerenchyma formation in Arabidopsis is controlled by LESION SIMULATING DISEASE1. *Plant Cell* 19, 3819–3830
- 65 Achard, P. *et al.* (2008) Plant DELLAs restrain growth and promote survival of adversity by reducing the levels of reactive oxygen species. *Curr. Biol.* 18, 656–660

- 66 De Tullio, M.C. *et al.* (2010) Redox regulation of root apical meristem organization: connecting root development to its environment. *Plant Physiol. Biochem.* 48, 328–336
- 67 Rubinovich, L. and Weiss, D. (2010) The Arabidopsis cysteine-rich protein GASA4 promotes GA responses and exhibits redox activity in bacteria and in planta. *Plant J.* 64, 1018–1027
- 68 Rieu, I. *et al.* (2008) Genetic analysis reveals that C19-GA 2-oxidation is a major gibberellin inactivation pathway in Arabidopsis. *Plant Cell* 20, 2420–2436
- 69 Potters, G. *et al.* (2009) Different stresses, similar morphogenic responses: integrating a plethora of pathways. *Plant Cell Environ.* 32, 158–169
- 70 Tognetti, V.B. *et al.* (2010) Perturbation of indole-3-butyric acid homeostasis by the UDP-glucosyltransferase UGT74E2 modulates Arabidopsis architecture and water stress tolerance. *Plant Cell* 22, 2660–2679
- 71 Gazarian, I.G. *et al.* (1998) Identification of skatolyl hydroperoxide and its role in the peroxidase catalysed oxidation of indol-3-yl acetic acid. *Biochem. J.* 333, 223–232
- 72 Jansen, M.A.K. *et al.* (2001) Phenol-oxidizing peroxidases contribute to the protection of plants from ultraviolet radiation stress. *Plant Physiol.* 126, 1012–1023
- 73 Santelia, D. *et al.* (2008) Flavonoids redirect PIN-mediated polar auxin fluxes during root gravitropic responses. *J. Biol. Chem.* 283, 31218–31226
- 74 Grunewald, W. and Friml, J. (2010) The march of the PINs: developmental plasticity by dynamic polar targeting in plant cells. *EMBO J.* 29, 2700–2714
- 75 Huang, Y.C. *et al.* (2008) Transcriptome analysis of auxin-regulated genes of Arabidopsis thaliana. *Gene* 420, 118–124
- 76 Joo, J.H. *et al.* (2001) Role of auxin-induced reactive oxygen species in root gravitropism. *Plant Physiol.* 126, 1055–1060
- 77 Pignocchi, C. *et al.* (2006) Ascorbate oxidase-dependent changes in the redox state of the apoplast modulate gene transcript accumulation leading to modified hormone signaling and orchestration of defense processes in tobacco. *Plant Physiol.* 141, 423–435
- 78 Tyburski, J. *et al.* (2010) A role for redox factors in shaping root architecture under phosphorus deficiency. *Plant Signal. Behav.* 5, 64–66
- 79 Bashandy, T. *et al.* (2010) Interplay between the NADP-linked thioredoxin and glutathione systems in Arabidopsis auxin signaling. *Plant Cell* 22, 376–391

Plant Science Conferences in 2011

ICAR 2011

22–25 June, 2011

Madison, USA

<http://www.union.wisc.edu/arabidopsis/>

New and Emerging Fungal Diseases of Animals and Plants: evolutionary aspects in the context of global changes

25–29 June, 2011

Roscoff, France

http://www.cnrs.fr/insb/cjm/cjmprog_e.html

IBC2011 XVIII International Botanical Congress

23–30 July, 2011

Melbourne, Australia

<http://www.ibc2011.com/>