


**TEXAS DEPARTMENT
OF
MENTAL HEALTH
AND
MENTAL RETARDATION**



**DRUG FORMULARY
2004**


<http://www.mhmr.state.tx.us/CentralOffice/MedicalDirector/EFC.html>

HOW TO USE THIS FORMULARY

The TDMHMR Drug Formulary is the publication that outlines the medications that have been approved for use in the TDMHMR system. The formulary is published annually in the last quarter of the calendar year. This document is divided into three sections (Alphabetical Listing, Therapeutic Classification/Cost Index, Alphabetical Index) to facilitate usage.


1. ALPHABETICAL LISTING:

This section lists all the medications alphabetically by generic (nonproprietary) name. Trade (proprietary) names and abbreviations are listed in parenthesis after the generic name. In this section, all approved dosage forms and strengths are also listed.



2. THERAPEUTIC CLASSIFICATION/COST INDEX:

This section groups the medications by therapeutic usage and lists them alphabetically by generic (nonproprietary) name. Medications may be listed in multiple categories. For trade names, consult the alphabetical index. Also in this section is a relative cost index so that medications used in similar situations can be evaluated keeping cost of care in mind. Relative cost is based on cost per day of the average dose. The cost of bulk items is listed per unit.



* Index = \$ < \$\$ < \$ < \$\$ < \$\$\$ < \$\$\$ < \$\$\$\$\$ < \$\$\$\$\$\$ < \$\$\$\$\$\$ < \$\$\$\$\$\$

3. ALPHABETICAL INDEX:

This section lists all the medications alphabetically by trade name, generic name, and abbreviations. Page numbers for each of the listings is provided.

INTRODUCTION

The purpose of the TDMHMR Executive Formulary Committee is to maintain and update the *Formulary* and to recommend standards for drug use within the TDMHMR system. The Committee is concerned with maintaining the highest standards for drug use. To assist the clinician in prescribing the most cost effective agents a Therapeutic Classification with relative cost data is provided. Because of their frequency of use, the highest level of concern and vigilance centers on the psychotropic medications. For the clinician's reference, the *Formulary* provides tables summarizing the recommended dosage ranges for the psychotropic drugs; however, these guidelines are not intended to replace other references or the clinician's clinical judgment. The clinician is urged to check with other references, including the TDMHMR Rule Governing Prescribing of Medications - Mental Health, Chapter 405, Subchapter A, *The American Hospital Formulary Service Drug Information, Facts & Comparisons*, and *The AMA Drug Evaluations* among other reliable sources.

This *Formulary*, is not intended to be a reference for drug use. Rather it is to serve the clinician as a listing of drugs, which are approved for use within the Department. Approval of a drug entity for inclusion in the *Formulary* does not imply approval of all formulations containing that entity. The Executive Formulary Committee will decide which formulations are approved for use within the department. Not all drugs listed in the *Formulary* will be stocked at each facility's pharmacy. If a physician, dentist, nurse or pharmacist desires to have a drug added to the *Formulary*, he or she should complete and submit the appropriate form (New Drug Application, DF-1, see Appendix 1) to the facility's Pharmacy and Therapeutics Committee. If it is approved at the facility level but is not included in the Departmental Formulary, the facility's Director of Pharmacy must then forward the request to the Departmental Executive Formulary Committee in care of the TDMHMR Office of the Medical Director. Requests received at least 30 days prior to the quarterly meeting will be considered at that meeting. Requests received less than 30 days before the meeting will not be considered until the following meeting. Clinicians and facilities are encouraged to submit supporting documents with their formulary request. The Executive Formulary Committee will evaluate the submission's appropriateness based upon the efficacy and safety of the proposed drug compared with existing formulary items and cost effectiveness of the new agent. When appropriate, the Executive Formulary committee will add the new drug or replace old agents in the same pharmacological/therapeutic category with the new agent. The TDMHMR *Formulary* and Interim Formulary Updates are available at <http://www.mhmr.state.tx.us/CentralOffice/MedicalDirector/Formulary.html>

The TDMHMR Formulary consists of routine and reserve drugs. Drugs in the reserve class have specific guidelines for use printed in the *Formulary*. These guidelines will be used to audit the appropriate use of reserve drugs. Based on the audit results, continuing education will be developed and targeted as needed. The purpose of reserve status is to stimulate thought and promote care in prescribing. Discussions about the use of reserve status drugs with peers are encouraged. Use of reserve status drugs requires documentation of justification in the patient's progress notes.

As a means of preventing medication errors, the *Formulary* has incorporated TALL

MAN characters to assist in distinguishing look-alike drug names. TALL MAN characters are being implemented in various parts of the pharmaceutical industry in order to prevent medication errors. Even though the *Formulary* may not play a major role in preventing medication errors, hopefully this change will stimulate the awareness of TALL MAN characters and assist in implementing this print style in other areas of our medication use process.

The Texas Department of Mental Health and Mental Retardation utilizes a closed formulary system. Only drugs listed in the *Formulary* are to be stocked, prescribed and dispensed in TDMHMR facilities including pharmaceutical products recommended by consultants for specialized treatments. When a patient's condition requires a drug not listed in the *Formulary*, limited quantities can be obtained for use in that particular patient. However, documentation (Non-Formulary Drug Justification Form, DF-2, see Appendix 2) should be submitted to the facility Clinical Director and the Office of the Medical Director where it will be reviewed by the Executive Formulary Committee to assure reasonable compliance with the *Formulary*.

When requested by the Commissioner, the Executive Formulary Committee will make other recommendations regarding drug use. If facility clinicians have topics for committee consideration, these requests should be sent to the Chairperson, TDMHMR Executive Formulary Committee in care of the TDMHMR Office of the Medical Director. The Committee also appreciates your comments concerning the printing of the *Formulary* and Committee deliberations and decisions. To facilitate your involvement, a schedule of the review process has been added to this edition of the *Formulary*. Please feel free to provide input.

Victoria B. Morgan, MD
Chairperson
TDMHMR Executive Formulary Committee

Steven P. Shon, MD
Medical Director
TDMHMR

December 2003
Date

MEMBERS OF THE TDMHMR EXECUTIVE FORMULARY COMMITTEE

Victoria B. Morgan, Chair Rusk State Hospital	Jack McCoy, MD San Angelo State School
Janet Adams, MSN, RN, CNS San Antonio State School	Dan Still, PharmD San Antonio State Hospital
Rebecca Alexander, RPh North Texas State Hospital	Ann L. Richards, PharmD, BCPP Pharmacy Discipline Head San Antonio State Hospital
Rosah Chadwick, RPh Denton State School	Cindy Sturdivant, BSN, RNC Big Spring State Hospital
Erlinda R. Devera, MD Corpus Christi State School	Bernardo C. Tarin-Godoy, MD El Paso Community MHMR
Emilio Dominguez, MD Center for Health Care Services San Antonio	James Van Norman, MD Austin-Travis County MHMR
Robin Mallett, MD Gulf Coast Center	Robert L. Ward, DO Kerrville State Hospital
<i>Ex-Officio Members</i>	
Stephen P. Shon, MD Medical Director	
Kenny Dudley Director, State Mental Health Facilities	
Robert Kifowit Director, State Mental Retardation Facilities	
Gerry McKimmy Deputy Commissioner, Community Programs	
Patricia Martin Director, Contracting & Procurement Support, State Operations	
Camille Hemlock, MD Associate Medical Director, CO TDMHMR	
Nina Jo Muse, MD Associate Medical Director, CO TDMHMR	
Earl B. Matthew, MD Consultant, Office of the Medical Director	
TDMHMR DRUG FORMULARY EDITOR	
Sharon M. Tramonte, PharmD San Antonio State School	

Table of Contents

<u>Formulary Review Schedule</u>	9
<u>Procedure for Addition of Drugs to the Formulary</u>	10
<u>Psychotropic Dosage Guidelines</u>	11
Antipsychotics.....	13
Antidepressants.....	14
Foods Containing Tyramine	15
Mood Stabilizers.....	16
Stimulants.....	16
Miscellaneous Drugs Used for Psychotropic Purposes.....	16
Anxiolytics.....	17
Sedatives and Hypnotics	17
<u>Reserve Drugs</u>	18
<u>Therapeutic Serum Concentrations of Some Anticonvulsants</u>	21
<u>Antibiotic Sensitivities</u>	22
<u>Alphabetical Listing</u>	24
<u>Therapeutic Classification/Cost Index</u>	72
<u>Antidiabetic Agents</u>	72
Insulins, Human.....	72
Sulfonylureas.....	72
Miscellaneous Antidiabetics	72
Glucose Elevating Agents	72
<u>Antidotes/Deterrents/Poison Control Agents</u>	73
<u>Antihistamines</u>	73
<u>Antineoplastic Agents</u>	73
<u>Blood Modifying Agents</u>	73
Antiplatelet.....	73
Anticoagulant.....	74
Anticoagulation Antagonist	74
Miscellaneous Blood Modifying Agents	74

Cardiovascular Agents 74

Diuretics	74
Cardiac Glycosides.....	74
Antianginals.....	75
Antiarrhythmics.....	75
Calcium Channel Blockers	75
Beta-Adrenergic Blockers.....	75
Antihyperlipidemics.....	75
Angiotensin Converting Enzyme Inhibitors	75
Vasopressors	76
Miscellaneous Antihypertensives	76
Miscellaneous Cardiovasculars	76

Central Nervous System Agents 76

Analgesics	76
Antiemetic/Antivertigo Agents.....	77
Psychotropic Agents.....	78
Sedatives and Hypnotics	80
Anticonvulsants	81
Muscle Relaxants	81
AntiParkinson Agents	82
Agents for Migraine	82
Miscellaneous Central Nervous System Agents.....	82

Endocrine Agents 82

Estrogens	82
Progesterones	82
Combination Products	83
Androgens	83
Adrenal Cortical Steroids.....	83
Thyroid Agents	83
Miscellaneous Endocrine Agents	83

Gastrointestinal Agents 84

Antacids.....	84
Antispasmodics/Anticholinergic Agents	84
Histamine (H ₂) Antagonists	84
Proton Pump Inhibitors	84
Antiflatulents	84
Stimulants.....	84
Laxatives	85
Antidiarrheals	85
Rectal Agents	86
Miscellaneous Gastrointestinal Agents.....	86

Genitourinary Agents	86
Interstitial Cystitis Agents	86
Genitourinary Irrigants	86
Urinary Alkalinizers	87
Urinary Anticholinergics	87
Urinary Cholinergics	87
Vaginal Antifungals.....	87
Miscellaneous Genitourinary Agents	87
Immunological Agents	87
Immune Serums	87
Bacterial Vaccines.....	87
Viral Vaccines.....	88
Toxoids.....	88
<i>In-Vivo</i> Diagnostic Biologicals	88
Infectious Disease Agents	88
Antibiotics	88
Antifungals.....	89
Antimalarials	89
Antituberculars	90
Antivirals	90
Antihelmintics	90
Urinary Anti-Infectives	90
Miscellaneous Anti-Infectives	90
Intravenous Solutions and Additives	91
Intravenous Solutions	91
Electrolyte Replacement Additives	91
Nutritional Agents	92
Vitamins.....	92
Minerals Trace Elements and Electrolytes	92
Combination Products	92

Respiratory Agents	93
Bronchodilators	93
Decongestants	93
Steroids	93
Antitussives	93
Expectorants	93
Cough and Cold Preparations	93
Miscellaneous Respiratory Drugs	94
Topical Agents	94
Ophthalmics	94
Otics	95
Nasal, Mouth and Throat Agents.....	96
Dermatologicals.....	96
Irrigation Solutions.....	99
Alphabetical Index	100
Appendix 1: New Drug Application Form	110
Appendix 2: Non-Formulary Drug Justification Form	112
Appendix 3: Adverse Drug Reaction Reporting (MedWatch) Form	113

FORMULARY REVIEW SCHEDULE

For FY04 – FY06 (9/1/03 – 8/31/06)

FY	Meeting Number	Category
04	1*	Dermatologicals, (Acne agents to Anti-Infectives Antifungals)
	2	Dermatologicals (Scabicides to Miscellaneous Dermatologicals)
	3	Psychotropic
	4	Cardiovascular
05	1*	Infectious Disease
	2	Antidiabetic Antidotes/Deterrents/Poison Control Antihistamines Antineoplastic Blood Modifying
	3	Analgesics Antiemetics/Antivertigo Sedative and Hypnotics Anticonvulsants
	4	Muscle Relaxants Antiparkinson Agents Migraine Miscellaneous CNS Endocrine
06	1*	Gastrointestinal Genitourinary
	2	Immunological Intravenous Solutions and Additives Nutritional
	3	Respiratory Ophthalmic
	4	Otics Nasal, Mouth and Throat Irrigation Solutions

* Reserve Drugs and Tables reviewed annually

PROCEDURE FOR ADDITION OF DRUGS TO THE FORMULARY

A physician, dentist, nurse or pharmacist desiring that a new drug be added to the *Formulary* should submit a new drug application (Form DF-1) to the facility's formulary committee. If approved, the new drug application should be forwarded to the Executive Formulary Committee. The following information should accompany the application:

- (1) published articles in the biomedical literature that substantiate the efficacy and safety of the proposed new drug;
- (2) advantages of the proposed new drug compared with similar therapeutic agents presently in the *Formulary*;
- (3) drugs in the *Formulary* which the proposed agent will replace or supplement;
- (4) cost effectiveness data.

New drug applications should be received by the secretary of the committee thirty days prior to the committee's scheduled meeting date. Applications received after this time will be considered at the next meeting.

The chairperson will assign a committee member to present an objective treatise and recommendations concerning the proposed to new drug to the Executive Formulary Committee. The committee may decide to approve or deny the drug's inclusion, approve the drug on a trial basis, or postpone a decision until the following meeting of the committee. The committee, at its discretion, may approve a drug's inclusion in the *Formulary* subject to specific limitations (e.g., recommendation by qualified specialists or consultants) or as a reserve drug.

The specific limitations or guidelines for use of a reserve drug are stated in the formulary. A credentialed clinician may prescribe reserve drugs outside *Formulary* guidelines; however, such exceptions must be justified in the patient record and must be reviewed in routine facility audits of reserve drug utilization.

PSYCHOTROPIC DOSAGE GUIDELINES

The following is a list of psychotropic drug dosages. These guidelines are not intended to establish rigid standards of treatment but to assist in monitoring the pharmacotherapy of the patient. Furthermore, guidelines for special patient populations are not intended to be absolute. For those medications that have a well established therapeutic serum range, the dosage should be based upon the desired serum range and response rather than a specific maximum administered daily dosage. These guidelines should be used in conjunction with sound clinical judgment and the prescriber's experience.

In children and adolescents, metabolic and physiologic differences from adults should be considered when prescribing. Dosing based on body weight may be more accurate when treating these patients.

Different dosage requirements are usually necessary in the geriatric population. Since there is no standard definition for "geriatric", the arbitrary age of 65 has been chosen to identify geriatric patients. In general, geriatric patient dosing guidelines should reflect a "go low, go slow" approach. Standard reference books should be consulted if needed for appropriate dosages when treating this population.

In general, when treating patients with developmental disabilities, a "go low, go slow" approach is recommended when increasing or decreasing psychotropic medication. The use of psychotropic medication can be therapeutic and empowering for a person with both mental retardation and a mental illness. The primary goal is to obtain an accurate diagnosis of behavioral and psychiatric symptoms so that the patients' treatment is appropriate. A functional analysis by a psychologist is vital prior to starting any psychotropic medication except in an emergency. The U. S. Health Care Financing Administration now states that the least intrusive and most positive intervention to treat behavioral or psychiatric symptoms in a person with mental retardation may be the use of a psychotropic medication.

Prescribing psychotropic medication should be based on the following resources:
TDMHMR Prescribing of Medications-Mental Health, Chapter 405, Subchapter A

TDMHMR Prescribing of Psychotropic Medication-Mental Retardation Facilities,
Chapter 405, Subchapter B

Other useful resources that reflect current Standards of Care for the mentally ill include but are not limited to the following:

Treatment Guidelines of Various Psychiatric Disorders

Examples include:

APA Practice Guideline for the treatment of Patients with Schizophrenia. *Am J Psychiatry* 1997; 154: 4 (April supplement)

APA Practice Guideline for Major Depressive Disorder in Adults. *Am J Psychiatry*

1993; 150 (supplement)
Consensus Guidelines for Bipolar Disorder. *J Clin Psychiatry* 1996; 57 (suppl 12a)

Psychotropic Medications and Developmental Disabilities: The International Consensus Handbook (Reiss and Aman, eds; AAMR, 202-387-1968)

Consensus Guidelines of Psychotropic Medication in Persons with Developmental Disabilities *AJMR*, May 2000 (special issue)

Texas Implementation of Medication Algorithms (TIMA):
www.mhmr.state.tx.us/CentralOffice/MedicalDirector/TIMA.html
(web site still under construction)

Revised 5 December 2000

ANTIPSYCHOTICS

Drug	Suggested Maximum Dose (mg/day)*
Aripiprazole (Abilify)	30
chlorproMAZINE (Thorazine)	2,000
Clozapine (Clozaril) - RESERVE USE	900
Fluphenazine ¹ (oral) (Prolixin)	60
Fluphenazine Decanoate ¹ (Prolixin, Permitil)	100 (q 1 - 4 weeks)
Haloperidol ² (oral) (Haldol)	100
Haloperidol Decanoate ² (Haldol)	450 mg per month
Loxapine (Loxitane)	250
Mesoridazine (Serentil) ³ - RESERVE USE	500
Molindone (Moban)	225
Olanzapine (Zyprexa)	30
Perphenazine (Trilafon)	64
Quetiapine (Seroquel)	800
Risperidone (Risperdal)	8 ⁴
Thioridazine (Mellaril) ³ - RESERVE USE	(ABSOLUTE) 800
Thiothixene (Navane)	60
Trifluoperazine (Stelazine)	80
Ziprasidone (Geodon)	240

*except where noted

¹ Fluphenazine Therapeutic Concentration = 1 - 3 ng/mL

² Haloperidol Therapeutic Concentration = 3 - 15 ng/mL

³ A boxed warning has been added to advise clinicians of prolongation of the QTc interval

⁴ Risperidone doses >6 mg/day have increased risk of EPS

Revised 24 October 2003

ANTIDEPRESSANTS

Drug	Suggested Maximum Dose (mg/day)
Amitriptyline (Elavil)	300
Amoxapine (Asendin)	600
buPROPion (Wellbutrin)	450/day (with no single dose > 150)
buPROPion SR (Wellbutrin SR)	400
buPROPion XL (Wellbutrin XL)	450
Citalopram (Celexa)	60
Desipramine (Norpramin)	300* ¹
Doxepin (Sinequan, Adapin)	300
Escitalopram (Lexapro)	20
Fluoxetine (Prozac)	80
Fluvoxamine (Luvox)	300
Imipramine (Tofranil)	300* ²
Maprotiline (Ludiomil)	225
Mirtazapine (Remeron)	45
Nefazodone (Serzone) - RESERVE USE	600
Nortriptyline (Pamelor, Aventyl)	200* ³
Paroxetine (Paxil)	50
Phenelzine (Nardil)	90
Protriptyline (Vivactil)	60
Sertraline (Zoloft)	200
Tranylcypromine (Parnate)	60
Trazodone (Desyrel)	600
Trimipramine (Surmontil)	300
Venlafaxine (Effexor)	375
Venlafaxine XR (Effexor XR)	

*Plasma concentration monitoring is recommended if these doses are exceeded.

¹Desipramine Therapeutic Concentration = 100-300 ng/mL

²Imipramine Therapeutic Concentration = 150-250 ng/mL

³Nortriptyline Therapeutic Concentration = 50-150 ng/mL

Revised 24 October 2003

FOODS CONTAINING TYRAMINE

***High Amounts of Tyramine**

Smoked, aged or pickled meat or fish
Sauerkraut
Aged Cheeses such as Swiss and Cheddar
Yeast extracts
Fava beans

****Moderate Amounts of Tyramine**

Beer
Avocados
Meat extracts
Red wines such as Chianti

*****Low Amounts of Tyramine**

Caffeine-containing beverages
Distilled spirits
Chocolate
Soy sauce
Cottage and cream cheeses
Yogurt and sour cream

*May not consume

**May consume in moderation

***May consume

Adapted from Shulman KI, et al. Dietary restriction, tyramine, and the use of monoamine oxidase inhibitors. J Clin Psychopharmacol. 1989; 9: 397.

MOOD STABILIZERS

Drug	Suggested Maximum Dose (mg/day)*	Therapeutic Serum Concentration
Carbamazepine (Tegretol, Tegretol XR, Carbatrol)	1600	4 - 12 mcg/mL
Lamotrigine (Lamictal)	400	nd
Lithium (Lithobid, Eskalith)	3600 (<65 years old) 1800 (\geq 65 years old)	0.6-1.5 mEq/L 0.6-1.0 mEq/L
Oxcarbazepine (Trileptal)	2400	nd
Topiramate (Topamax)	nd	nd
Valproic Acid/Valproate (Depakene), Divalproex (Depakote)	60 mg/kg	50 - 150 mcg/mL
Verapamil (Calan, Isoptin)	480	nd

* Plasma concentration monitoring is recommended if these doses are exceeded

nd = not yet determined

Revised 24 October 2003

STIMULANTS

Drug	Suggested Maximum Dose (mg/day)
Amphetamine Mixture (Adderall, Adderall XR)	60*
Dextroamphetamine (Dexedrine)	60*
Methylphenidate (Ritalin, Concerta)	60

* suggested maximum dosage in children

Revised 24 October 2003

MISCELLANEOUS DRUGS USED FOR PSYCHOTROPIC PURPOSES

Drug	Suggested Maximum Dose (mg/day)
clomiPRAMINE (Anafranil)	250
Clonidine (Catapres)	0.4
Propranolol (Inderal)	160 (anxiety)***

*** Maximum dose has not been determined for aggression or self-injurious behavior (SIB).

Revised 19 October 1998

ANXIOLYTICS

Drug	Suggested Maximum Dose (mg/day)	
	Under 65 years (mg/day)	Over 65 years (mg/day)
Alprazolam (Xanax) (exception: for panic disorder)	4 10	2 N/A
busPIRone (BuSpar)	60	60
Chlordiazepoxide (Librium)	100*	40*
Clonazepam (Klonopin)	20	10
Clorazepate (Tranxene)	60	30
Diazepam (Valium)	60	20
Lorazepam (Ativan)	10*	3*
Oxazepam (Serax)	90	60

*Larger doses may be necessary in some cases of alcohol/substance withdrawal.

Revised 19 October 1998

SEDATIVES AND HYPNOTICS

Drug	Suggested Maximum Dose (mg/day)	
	Under 65 years (mg/day)	Over 65 years (mg/day)
Amobarbital (Amytal) - RESERVE USE	500	500
Chloral Hydrate (Noctec)	1500*	1500*
diphenhydramINE (Benadryl)	300	300
hydrOXYzine (Atarax)	300	300
Mirtazapine (Remeron)	30	15
Temazepam (Restoril)	30	15
Trazodone (Desyrel)	150	150
Triazolam (Halcion)	0.25	0.125
Zaleplon (Sonata)	10	5
Zolpidem (Ambien)	10	5

* Individual dose usually 500 mg but doses up to 1 gram may be used to produce conscious sedation for certain procedures

Revised 25 October 2002

RESERVE DRUGS

The purpose of the reserve drug class is to stimulate thought and promote care in prescribing. Discussions and interaction with other prescribers is strongly encouraged. When reserve status drugs are utilized, documentation of justification in the patient's progress notes is mandated. To assist the prescriber, specific guidelines have been developed by the Executive Formulary Committee for use of each drug so classified. The guidelines will be used to audit the appropriate use of reserve drugs. Based on the audit results, continuing education will be developed and targeted as needed.

Drug	Guidelines for Use
Amobarbital (Amytal) C-II	For use with research protocols
Azithromycin (Zithromax)	<ol style="list-style-type: none"> 1) If a macrolide antibiotic is indicated for the treatment of an infection, then erythromycin is the drug of choice 2) The patient has prior side effects (not anaphylactic) to erythromycin which is documented in the chart 3) The prescriber is aware of the significant cost difference between erythromycin, clarithromycin, and azithromycin 4) The prescriber is aware that all macrolide antibiotics have the potential to have drug interactions
Cefuroxime Axetil (Ceftin) - Oral Form	<ol style="list-style-type: none"> 1) Streptococcus or Staphylococcus infections or any penicillin-sensitive organism infections in patients who are <ol style="list-style-type: none"> a) allergic to penicillin, and b) intolerant of erythromycin; or c) allergic or intolerant to sulfonamides; or d) therapeutic failures with penicillin, erythromycin, or sulfonamides; or 2) <i>H. influenza</i> or <i>Branhamella catarrhalis</i> infections or 3) Oral continuation of IV therapy for appropriate indications (not appropriate follow-up for coliforms); 4) Not to be used for oral therapy of gram negative infections
Clarithromycin (Biaxin)	<ol style="list-style-type: none"> 1) If a macrolide antibiotic is indicated for the treatment of an infection, then erythromycin is the drug of choice 2) The patient has prior side effects (not anaphylactic) to erythromycin which is documented in the chart 3) The prescriber is aware of the significant cost difference between erythromycin and clarithromycin 4) The prescriber is aware that all macrolide antibiotics have the potential to have drug interactions
Clobetasol (Temovate)	To be used under the direction of a dermatologist
Clopidogrel (Plavix)	Documented sensitivity to aspirin

Drug	Guidelines for Use
Clozapine (Clozaril)	<ol style="list-style-type: none"> 1) For use in patients with refractory schizophrenia or schizoaffective disorder; or 2) For use in schizophrenic or schizoaffective patients who cannot tolerate other antipsychotics; or 3) Psychosis associated with other organic conditions who have failed two antipsychotics or who cannot tolerate other antipsychotics; or 4) Manic disorders with psychosis in patients who have failed two antipsychotics
Divalproex ER (Depakote ER)	For the treatment of migraine headaches and epilepsy only.
Donepezil (Aricept)	<ol style="list-style-type: none"> 1) Preferred agent for patients who have been newly diagnosed with Alzheimer's disease 2) Patient's response needs to be monitored with a Mini-Mental Status Exam (MMSE) during the course of therapy 3) Initial dose is 5 mg/day for 4 to 6 weeks 4) Maximum dose is 10 mg/day and dose cannot be increased to 10 mg/day until the patient has been on 5 mg/day for 4 to 6 weeks
Felbamate (Felbatol)	Weekly progress note by physician monitoring for response and adverse effects for at least two weeks following initiation of therapy
Mesoridazine (Serentil)	<ol style="list-style-type: none"> 1) Indicated only for patients with refractory schizophrenia (failed other classes of antipsychotics) 2) EKG prior to initiating therapy: 7 – 14 days after dose change; 7 – 14 days after other medication changes that could significantly alter the cardiac effects of Thioridazine; every six months; and as clinically indicated 3) Potassium level prior to initiating therapy; every six months; and as clinically indicated 4) Magnesium level prior to initiating therapy; and as clinically indicated (especially if potassium level is low)
Nabumetone (Relafen)	For the treatment of chronic osteoarthritis or chronic rheumatoid arthritis
Ophthalmic agents containing a steroid	Consultation with an Ophthalmologist prior to initiation.
Pentamidine (Pentam)	<ol style="list-style-type: none"> 1) Prophylaxis post-treatment of documented <i>Pneumocystis carinii</i> pneumonia; or 2) Prophylaxis of <i>Pneumocystis carinii</i> in patients with T₄ counts of less than 200
Rivastigmine (Exelon)	Patient's response needs to be monitored with mini-mental status exam (MMSE) during the course of therapy

Drug	Guidelines for Use
Thioridazine (Mellaril)	<ul style="list-style-type: none"> 1) Indicated only for patients with refractory schizophrenia (failed other classes of antipsychotics) 2) EKG prior to initiating therapy; 7 – 14 days after dose change; 7 – 14 days after other medication changes that could significantly alter the cardiac effects of Thioridazine; every six months; and as clinically indicated 3) Potassium level prior to initiating therapy; every six months; and as clinically indicated 4) Magnesium level prior to initiating therapy; and as clinically indicated (especially if potassium level is low)
Tizanidine (Zanaflex)	<ul style="list-style-type: none"> 1) Documented failure or intolerance to baclofen 2) Liver enzymes obtained at baseline, 1, 3, and 6 months and at least annually 3) Documentation of efficacy after one month of therapy 4) The dose should be decreased in severe renal dysfunction ($\text{CrCl} < 25 \text{ mL/min}$)

THERAPEUTIC SERUM CONCENTRATIONS OF SOME ANTICONVULSANTS

Drug	(mcg/mL)
Carbamazepine (Tegretol, Tegretol XR, Carbatrol)	4-12
Ethosuximide (Zarontin)	40-100
Phenobarbital (Luminal)	10-40
Phenytoin (Dilantin)	10-20
Valproic Acid, Valproate, Divalproex (Depakene, Depakote)	Seizure Disorder: 50-150 Bipolar Disorder: 50-125 Aggression: not established

Serum concentrations are useful in the evaluation of therapy. However, they should only be considered as a guide to treatment, not the sole criterion for determination of dosage regimens.

The following anticonvulsants do not have an established therapeutic serum concentration range. Routine monitoring of these anticonvulsants is not warranted as serum concentrations cannot be correlated with clinical efficacy.

Benzodiazepines

Gabapentin
Lamotrigine
Levetiracetam
Oxcarbazepine
Tiagabine
Topiramate

*Source: USP Dispensing Information, 1998
The United States Pharmacopeial Convention, Inc.

Revised 25 October 2002

ANTIBIOTIC SENSITIVITIES

STREP/STAPH INFECTIONS

Oral Medication	Dose	Cost/Day*
Pen V Potassium (Strep only)	500 mg q6h	0.40
Cloxacillin/Dicloxacillin	500 mg q6h	1.40
Cloxacillin/Dicloxacillin	1 gm q6h	2.80
Cefuroxime Axetil - Reserve Use	500 mg q8h	15.90
Cephalexin	500 mg q6h	0.63
Erythromycin	500 mg q6h	0.48
Clarithromycin - Reserve Use	500 mg q6h	10.97
Trimethoprim/Sulfamethoxazole	DS q12h	0.12

STAPH INFECTIONS

IV Medication	Dose	Cost/Day*
Nafcillin	2 gm q4h	18.72
Vancomycin	500 mg q6h	13.18

SINUSITIS - OTITIS MEDIA

Oral Medication	Dose	Cost/Day*
Amoxicillin	500 mg q8h	0.38
Amoxicillin/Clavulanate	500 mg q8h	7.47
Cefuroxime Axetil - Reserve Use	500 mg q8h	15.90
Clarithromycin - Reserve Use	500 mg q8h	8.23
Trimethoprim/Sulfamethoxazole	DS q12h	0.12
Erythromycin	500 mg q6h	0.48
Clindamycin	300 mg q6h	7.11

* Cost may vary slightly between facilities

UTI - GRAM (-)
E. COLI, KLEBSIELLA, PROTEUS

Medication	Route	Dose	Cost/Day*
Ceftriaxone	IV	1 gm qD	25.44
Trimethoprim/Sulfamethoxazole	IV	1.2 gm sulfa q12h	1.32
Ciprofloxacin	IV	400 mg q12h	60.00
Ciprofloxacin	PO	500 mg q12h	6.20
Trimethoprim/Sulfamethoxazole	PO	DS q12h	0.12

PSEUDOMONAS INFECTIONS

Medication	Route	Dose	Cost/Day*
Ticarcillin/Clavulanate	IV	3.1 gm q6h	43.26
Ciprofloxacin	IV	400 mg q12h	60.00
Gentamicin	IV	1.8 mg/kg q8h	14.40
Ciprofloxacin	PO	500 mg q12h	6.20

ANAEROBES

Medication	Route	Dose	Cost/Day*
Ticarcillin/Clavulanate	IV	3.1 gm q6h	43.26
Metronidazole	IV	500 mg q12h	32.78
Clindamycin	IV	900 mg q8h	17.26
Amoxicillin/Clavulanate	PO	500 mg q8h	7.47
Metronidazole	PO	750 mg q8h	0.40
Clindamycin	PO	300 mg q6h	7.11

* Cost may vary slightly between facilities

Revised 30 November 1998

ALPHABETICAL LISTING

Abrasive Cleanser (Brasivol, Pernox)

Pernox Cleanser (contains sulfur, salicylic acid, EDTA): 56 g, 113 g

Brasivol Cleanser (contains aluminum oxide): fine (153 g), medium (180 g), rough (195 g) textures

Acetaminophen (Tylenol)

Liquid: 160 mg/5 mL, 500 mg/15 mL

Suppository, rectal: 120 mg, 125 mg, 325 mg, 650 mg

Tablet: 325 mg, 500 mg, 650 mg

Tablet, chewable: 80 mg

Acetaminophen/Codeine C-III

Suspension, oral: Acetaminophen 120 mg/Codeine 12 mg per 5 mL (C-V)

Tablet:

#2: Acetaminophen 300 mg/Codeine 15 mg

#3: Acetaminophen 300 mg/Codeine 30 mg

Acetaminophen/Hydrocodone (Lortab, Vicodin)

Elixir: Acetaminophen 167 mg/Hydrocodone 2.5 mg per 5 mL with 7% alcohol

Tablet: Acetaminophen 325 mg/Hydrocodone 5 mg, Acetaminophen 325

mg/Hydrocodone 10 mg, Acetaminophen 500 mg/Hydrocodone 2.5 mg,

Acetaminophen 500 mg/Hydrocodone 5 mg, Acetaminophen 500

mg/Hydrocodone 7.5 mg, Acetaminophen 500 mg/Hydrocodone 10 mg

acetaZOLAMIDE (Diamox)

Capsule, sustained release: 500 mg

Tablet: 125 mg, 250 mg

Acetic Acid (Acetasol, VoSol)

Solution, otic: 2%

Acetic Acid/Aluminum Acetate (Domeboro Otic)

Solution, otic: 2% acetic acid in aluminum acetate

Acetic Acid/Hydrocortisone/Propylene Glycol/Sodium Acetate/Benzethonium (VoSol HC)

Solution, otic: 1% Hydrocortisone, 2% Acetic Acid, 3% Propylene Glycol, 0.015% Sodium Acetate, and 0.02% Benzethonium

Acetylcysteine (Mucomyst)

Solution, inhalation: 10% [100 mg/mL], 20% [200 mg/mL]

Activated Charcoal

Liquid, oral, activated, with sorbitol: 25 g, 30 g, 50 g
Powder for oral suspension, activated: 15 g, 30 g, 40 g, 120 g, 240 g

Acyclovir (Zovirax)

Capsule: 200 mg
Powder for injection: 500 mg, 1000 mg
Ointment, topical 5% [50 mg/g]: 3 gm, 15 gm
Suspension, oral: 200 mg/5 mL
Tablet: 400 mg, 800 mg

Adapalene (Differin)

Gel, topical: 0.1%

Adenosine (Adenocard)

Injection: 3 mg/mL

Albuterol (Proventil, Ventolin)

Aerosol, inhalation, chlorofluorocarbon free: 90 mcg/dose (17g) [200 doses]
Solution, inhalation: 0.083% [83 mg/mL], 0.5% [50 mg/mL]
Syrup: 2 mg/5 mL
Tablet: 2 mg, 4 mg
Tablet, extended release: 4 mg, 8 mg

Alendronate (Fosamax)

Tablet: 5 mg, 10 mg, 35 mg, 40 mg, 70 mg

Allopurinol (Zyloprim)

Tablet: 100 mg, 300 mg

Alprazolam (Xanax) C-IV

Tablet: 0.25 mg, 0.5 mg, 1 mg, 2 mg

Aluminum Acetate (Burow's Solution,)

Solution, topical: 480 mL

Aluminum Hydroxide (Amphojel)

Suspension, oral: 320 mg/5 mL, 600 mg/5 mL
Tablet: 300 mg, 400 mg, 500 mg, 600 mg

Aluminum Hydroxide/Magnesium Trisilicate (Gaviscon)

Tablet, chewable: each tablet contains Aluminum Hydroxide/Magnesium Trisilicate

Aluminum Hydroxide/Magnesium Hydroxide (Maalox)

Suspension, oral: containing Aluminum Hydroxide/Magnesium Hydroxide
Tablet: each tablet contains Aluminum Hydroxide/Magnesium Hydroxide

Aluminum Hydroxide/Magnesium Hydroxide/Simethicone (Mylanta, Aludrox)

Liquid, oral: containing Aluminum Hydroxide/Magnesium Hydroxide/Simethicone
Tablet: each tablet contains Aluminum Hydroxide/Magnesium Hydroxide/Simethicone

Amantadine (Symmetrel)

Capsule: 100 mg
Syrup: 50 mg/5 mL

Amino Acid Injection (Aminosyn)

Infusion: 3.5%, 5%, 7%, 8.5%, 10%, 15%

Aminophylline (79% Theophylline)

Injection: 25 mg/mL
Suppository, rectal: 250 mg

Amitriptyline (Elavil)

Tablet: 10 mg, 25 mg, 50 mg, 75 mg, 100 mg, 150 mg

Amlodipine (Norvasc)

Tablet, extended release: 2.5 mg, 5 mg, 10 mg

Amobarbital (Amytal) C-II - RESERVE USE

Capsule: 65 mg, 200 mg 100 mg
Injection: 250 mg, 500 mg
Tablet: 30 mg, 50 mg, 100 mg, 500 mg

Amoxapine (Asendin)

Tablet: 25 mg, 50 mg, 100 mg, 150 mg

Amoxicillin (Amoxil, Polymox)

Capsule: 250 mg, 500 mg
Powder for oral suspension: 50 mg/mL, 125 mg/5 mL, 250 mg/5 mL
Tablet: 500 mg, 875 mg
Tablet, chewable: 125 mg, 250 mg

Amoxicillin/Clavulanate (Augmentin)

Tablet: 200 mg (contains 28.5 mg Clavulanate), 250 mg (contains 125 mg Clavulanate), 400 mg (contains 57 mg Clavulanate), 500 mg (contains 125 mg Clavulanate), 875 mg (contains 125 mg Clavulanate)
Tablet, chewable: 125 mg (contains 31.25 mg Clavulanate), 250 mg (contains 62.5 mg Clavulanate)

Amphetamine Mixture (Adderall, Adderall XR) CII

Capsule, extended release: 5 mg, 10 mg, 15 mg, 20 mg, 25 mg, 30 mg
Tablet: 5 mg, 10 mg

Ampicillin (Polycillin, Omnipen)

Capsule, as anhydrous: 250 mg, 500 mg

Capsule, as trihydrate: 250 mg, 500 mg

Powder for injection: 125 mg, 250 mg, 500 mg, 1 g, 2 g, 10 g

Powder for oral suspension, as trihydrate: 125 mg/5 mL, 250 mg/5 mL

Antipyrine/Benzocaine (Allergen, Auralgan)

Solution, otic

Aripiprazole (Abilify)

Tablet: 10 mg, 15 mg, 20 mg, 30 mg

Ascorbic Acid (Vitamin C)

Solution, oral: 100 mg/mL

Tablet: 250 mg, 500 mg

Tablet, chewable: 250 mg, 500 mg

Aspirin

Suppository, rectal: 300 mg, 600 mg

Tablet: 81 mg, 325 mg, 500 mg

Tablet, buffered: 325 mg with buffering agents

Tablet, chewable: 81 mg

Tablet, enteric coated: 81 mg, 325 mg, 500 mg, 650 mg, 975 mg

Atenolol (Tenormin)

Tablet: 25 mg, 50 mg, 100 mg

Atomoxetine (Strattera)

Capsule: 10 mg, 18 mg, 25 mg, 40 mg, 60 mg

Atorvastatin (Lipitor)

Tablet: 10 mg, 20 mg, 40 mg

Atropine Sulfate (Isopto Atropine)

Ointment, ophthalmic: 1%

Solution, ophthalmic: 1%

Attapulgite

Liquid, oral concentrate: 600 mg/15 mL, 750 mg/15 mL

Azithromycin (Zithromax)- RESERVE USE

Powder for oral solution: 200 mg/5 mL, 400 mg/5 mL

Tablet: 250 mg

Bacitracin (Baciguent)

Injection: 50,000 units

Ointment, ophthalmic: 500 units/g

Ointment, topical: 500 units/g

Bacitracin/Polymyxin B (Polysporin)

Ointment, ophthalmic: Bacitracin 500 units/Polymyxin B 10,000 units/g

Ointment, topical: Bacitracin 500 units/Polymyxin B 10,000 units/g

Powder, topical: Bacitracin 500 units/Polymyxin B 10,000 units/g

Baclofen (Lioresal)

Tablet: 10 mg, 20 mg

Beclomethasone (Vanceril, Beconase)

Spray, nasal: 0.084% (19 g) [120 metered doses]

Spray, nasal, aqueous: 42 mcg/inhalation (25 g) [\geq 200 metered doses], 84 mcg/inhalation (25 g) [\geq 200 metered doses]

Benazepril (Lotensin)

Tablet: 5 mg, 10 mg, 20 mg, 40 mg

Benzalkonium Chloride (Zephiran)

Concentrate, topical: 17%

Solution, topical, aqueous: 1:750

Spray, topical: 1:750

Benzocaine (Lanacaine)

Topical, for mucous membranes:

Gel: 6%, 20%

Liquid: 20%

Topical, dermatologic:

Cream, topical: 5%, 6%

Lotion: 8%

Ointment: 5%

Spray: 5%, 20%

Mouth/Throat preparations:

Gel: 6.3%, 7.5%, 10%, 15%, 20%

Liquid: 5%, 6.3%, 10%, 20%

Lozenges: 5 mg, 6 mg, 10 mg, 15 mg

Ointment: 20%

Benzoic and Salicylic Acids (Whitfield's)

Ointment, topical: 6% Benzoic acid/3% Salicylic acid

Benzoin, Compound Tincture

Tincture, topical (also contains aloe, storax, tolu balsam, 74% to 80% alcohol): 30 mL, 60 mL, 120 mL, 480 mL, 4000 mL

Benzoyl Peroxide

Bar: 5%, 10%
Cream, topical: 5%, 10%
Gel, topical: 2.5%, 5%, 10%, 20%
Liquid, topical: 5%, 10%
Lotion: 5%, 5.5%, 10%

Benzoyl Peroxide/Clindamycin (BenzaClin)

Gel, topical: Benzoyl Peroxide 5%/Clindamycin 1%

Benztropine (Cogentin)

Injection: 1 mg/mL
Tablet: 0.5 mg, 1 mg, 2 mg

Betamethasone Valerate (Valisone)

Cream, topical: 0.01%, 0.1%
Lotion: 0.1%
Ointment, topical: 0.1%

Betaxolol (Betoptic S)

Solution, ophthalmic: 0.5%
Suspension, ophthalmic: 0.25%

Bethanechol (Urecholine)

Injection: 5 mg/mL
Tablet: 5 mg, 10 mg, 25 mg, 50 mg

Bimatoprost (Lumigan)

Solution, ophthalmic: 0.03%

Biperiden (Akineton)

Injection: 5 mg/mL
Tablet: 2 mg

Bisacodyl (Dulcolax)

Suppository, rectal: 10 mg
Tablet, enteric coated: 5 mg

Bismuth Subsalicylate (Pepto-Bismol, Bismatrol, Kaopectate)

Liquid, oral: 262mg/15 mL
Tablet, chewable: 262 mg

Brimonidine (Alphagan)

Solution, ophthalmic: 0.15%, 0.2%

Bromocriptine (Parlodel)

Capsule: 5 mg
Tablet: 2.5 mg

Brompheniramine/Pseudoephedrine (Bromfed)

Capsule: 12 mg Brompheniramine/120 mg Pseudoephedrine
Elixir: 4 mg Brompheniramine/30 mg Pseudoephedrine
Syrup: 2 mg Brompheniramine/30 mg Pseudoephedrine
Tablet: 4 mg Brompheniramine/60 mg Pseudoephedrine

Bupivacaine (Marcaine)

Injection: 0.25%, 0.5%, 0.75%

buPROPion (Wellbutrin, Wellbutrin SR, Wellbutrin XL)

Tablet: 75 mg, 100 mg
Tablet, extended release: 150 mg, 300 mg
Tablet, sustained release: 100 mg, 150 mg

busPIRone (BuSpar)

Tablet: 5 mg, 10 mg

Calamine/Zinc Oxide/Glycerin (Calamine Lotion)

Lotion, topical: 120 mL, 240 mL, 480 mL

Calamine/Pramoxine (Caladryl)

Lotion, topical: 180 mL

Calcipotriene (Dovonex)

Cream, topical: 0.005%
Ointment, topical: 0.005%

Calcitonin-Salmon (Miacalcin)

Nasal spray: 200 IU/activation

Calcium Carbonate (Os-Cal, Titralac) [40% elemental calcium]

Liquid, oral: 500 mg/5 mL, 1000 mg/5 mL
Tablet: 600 mg, 1250 mg, 1500 mg
Tablet, chewable: 350 mg, 500 mg, 550 mg, 750 mg, 850 mg, 1000 mg

Calcium Citrate (Citracal) [21% elemental calcium]

Tablet: 200 mg, 250 mg

Calcium Carbonate/Vitamin D (Oscal + D)

Tablet: Calcium 250 mg/Vitamin D 125 IU, Calcium 500 mg/Vitamin D 125 IU

Calcium Glubionate (Neo-Calglucon) [6% elemental calcium]

Syrup: 1.8 g/5 mL

Calcium Gluconate [9% elemental calcium]

Injection: 10% [100 mg/mL]

Tablet: 500 mg, 650 mg, 975 mg, 1g

Calcium Undecylenate (Caldesene)

Powder, topical: 10%

Camphor-Phenol (Campho-Phenique)

Liquid, topical: 10.8% Camphor/4.7% phenol [with eucalyptus oil and mineral oil]

Captopril (Capoten)

Tablet: 12.5 mg, 25 mg, 50 mg, 100 mg

Carbamazepine (Tegretol, Tegretol XR, Carbatrol)

Capsule, extended release: 200 mg, 300 mg

Suspension, oral: 100 mg/5 mL

Tablet: 200 mg

Tablet, chewable: 100 mg

Tablet, extended release: 100 mg, 200 mg, 400 mg

Carbamide Peroxide/Glycerin/Propylene Glycol/Sodium Stannate (Debrox, Gly-Oxide)

Gel, oral: 10%

Solution, oral: 10%, 15%

Solution, otic: 6.5%

Carboxymethylcellulose/Electrolytes (Saliva Substitute, Moi-Stir, Salivart, MouthKote, Salix)

Solution, oral

Cascara Sagrada (Cascara Aromatic)

Aromatic fluid extract: 120 mL, 473 mL

Tablet: 325 mg

Cefazolin (Kefzol, Ancef)

Injection: 500 mg, 1 g

Powder for injection: 250 mg, 500 mg, 1 g, 5 g, 10 g, 20 g

Cefoperazone (Cefobid)

Infusion, premixed in dextrose: 1 g, 2 g

Powder for injection: 1 g, 2 g

Ceftriaxone (Rocephin)

Infusion, premixed in dextrose: 1 g, 2 g

Powder for injection: 250 mg, 500 mg, 1 g, 2 g, 10 g

Cefuroxime Axetil (Ceftin) - Oral form only - RESERVE USE

Powder for oral suspension: 125 mg/5 mL, 250 mg/5 mL
Tablet: 125 mg, 250 mg, 500 mg

Cellulose (Unifiber)

Powder, oral: 150 g, 270 g, 480 g

Cephalexin (Keflex)

Capsule: 250 mg, 500 mg
Powder for oral suspension: 100 mg/mL, 125 mg/5 mL, 250 mg/5 mL
Tablet: 250 mg, 500 mg, 1 g
Tablet: 500 mg

Cetylpyridinium (Cepacol)

Lozenges: 0.07% Cetylpyridinium/0.3% Benzyl Alcohol [with tartrazine]
Mouthwash: 0.05% Cetylpyridinium/14% Alcohol [with tartrazine]
Troches: 0.07% Cetylpyridinium/10 mg Benzocaine [with tartrazine]

Chloral Hydrate (Noctec) C-IV

Capsule: 500 mg
Suppository, rectal: 324 mg, 500 mg
Syrup: 250 mg/5 mL, 500 mg/5 mL

Chlordiazepoxide (Librium) - oral form only - C-IV

Capsule: 5 mg, 10 mg, 25 mg
Tablet: 5 mg, 10 mg, 25 mg

Chlorhexidine (Peridex, Hibiclens, Bactoshield)

Foam, topical, with 4% isopropyl alcohol: 4%
Liquid, topical, with 4% isopropyl alcohol: 2%, 4%
Rinse, oral, with 12% alcohol: 0.12%

Chloroquine (Aralen)

Tablet: 250 mg, 500 mg

Chlorpheniramine (Chlor-Trimeton, Teldrin)

Capsule: 12 mg
Syrup: 2 mg/5 mL
Tablet: 4 mg, 8 mg, 12 mg
Tablet, chewable: 2 mg
Tablet, timed release: 8 mg, 12 mg

chlorproMAZINE (Thorazine)

Concentrate, oral: 30 mg/mL, 100 mg/mL
Injection: 25 mg/mL
Syrup: 10 mg/5 mL
Tablet: 10 mg, 25 mg, 50 mg, 100 mg, 200 mg

chlorproPAMIDE (Diabinese)

Tablet: 100 mg, 250 mg

Chlorthalidone (Hygroton)

Tablet: 15 mg, 25 mg, 50 mg, 100 mg

Cholestyramine (Questran)

Powder, oral: 4 gm resin/9 gm powder

Powder for oral suspension (with aspartame): 4 gm resin/5 gm powder

Powder for oral suspension (with phenylalanine): 4 gm resin/5.5 gm powder

Tablet: 1 gm

Ciprofloxacin (Cipro, Ciloxan)

Injection: 200 mg, 400 mg

Solution, ophthalmic: 0.3%

Suspension, oral: 5 gm/100 mL, 10 gm/100 mL

Tablet: 100 mg, 250 mg, 500 mg, 750 mg

Ciprofloxacin/Hydrocortisone (Cipro Otic)

Solution, otic: Ciprofloxacin 2 mg/Hydrocortisone 10 mg per mL

Citalopram (Celexa)

Tablet: 20 mg, 40 mg

Clarithromycin (Biaxin) - RESERVE USE

Granules for oral suspension: 125 mg/5 mL, 250 mg/5 mL

Tablet, film coated: 250 mg, 500 mg

Clindamycin (Cleocin, Cleocin T)

Capsule: 75 mg, 150 mg, 300 mg

Gel, topical: 1% [10 mg/g]

Granules for oral solution: 75 mg/5 mL

Injection: 150 mg/mL

Lotion: 1% [10 mg/mL]

Solution, topical: 1% [10 mg/mL]

Clobetasol (Temovate) - RESERVE USE

Cream, topical: 0.05%

Cream, topical, in emollient base: 0.05%

Gel, topical: 0.05%

Ointment, topical: 0.05%

Scalp application: 0.05%

clomiPRAMINE (Anafranil) - for Obsessive Compulsive Disorder

Capsule: 25 mg, 50 mg, 75 mg

Clonazepam (Klonopin) C-IV

Tablet: 0.5 mg, 1 mg, 2 mg

Clonidine (Catapres)

Patch, transdermal: 1, 2, and 3 (0.1, 0.2, 0.3 mg/day, 7 day duration)

Tablet: 0.1 mg, 0.2 mg, 0.3 mg

Clopidogrel (Plavix) - RESERVE USE

Tablet: 75 mg

Clorazepate (Tranxene, Traxene SD) C-IV

Capsule: 3.75 mg, 7.5 mg, 15 mg

Tablet: 3.75 mg, 7.5 mg, 15 mg

Tablet, sustained release: 11.25 mg, 22.5 mg

Clotrimazole (Lotrimin, Mycelex, Gyne-Lotrimin, Fungoid)

Cream, topical: 1%

Cream, vaginal: 1%, 2%

Lotion: 1%

Solution, topical: 1%

Suppository, vaginal: 100 mg, 200 mg

Tablet, vaginal: 100 mg, 500 mg

Troche: 10 mg

Cloxacillin (Cloxapen, Tegopen)

Capsule: 250 mg, 500 mg

Powder for oral suspension: 125 mg/5 mL

Clozapine (Clozaril) - RESERVE USE

Tablet: 25 mg, 100 mg

Coal Tar (Ionil-T, Tegrin)

Liquid, topical: 30%

Shampoo: 1%

Solution, topical: 120 mL, 480 mL

Cod Liver Oil/Zinc Oxide/Talc (Desitin)

Ointment, topical: 40% Zinc Oxide [with Cod Liver Oil, Talc, Petrolatum, Lanolin, and Methylparaben]

Colchicine

Tablet: 0.5 mg, 0.6 mg

Collagenase (Santyl)

Ointment, topical: 250 units/g

Corticotropin (ACTH)

Injection, repository: 40 units/mL, 80 units/mL
Powder for injection: 25 units, 40 units

Cortisone

Injection: 50 mg/mL
Tablet: 5 mg, 10 mg, 25 mg

Cromolyn (Intal)

Inhalation, oral: 800 mcg/spray
Solution, nebulizing: 10 mg/mL
Solution, nasal: 40 mg/mL
Solution, ophthalmic: 4%

Crotamiton (Eurax)

Cream: 10%
Lotion: 10%

Cyanocobalamin (Vitamin B₁₂)

Injection: 1000 mcg/mL
Tablet: 100 mcg, 250 mcg, 500 mcg, 1000 mcg

Cyproheptadine (Periactin)

Syrup: 2 mg/5 mL with 5% alcohol
Tablet: 4 mg

Dantrolene (Dantrium)

Capsule: 25 mg, 50 mg, 100 mg
Powder for injection: 20 mg

Deferoxamine (Desferal)

Powder for injection: 500 mg

Delavirdine (DLV, Rescriptor)

Tablet: 100 mg

Desipramine (Norpramin, Pertofrane)

Capsule: 25 mg, 50 mg
Tablet: 10 mg, 25 mg, 50 mg, 75 mg, 100 mg, 150 mg

Desmopressin (DDAVP, Stimate)

Solution, nasal: 100 mcg/mL, 1.5 mg/mL

Dexamethasone (Decadron)

Injection, as sodium phosphate: 4 mg/mL, 10 mg/mL, 20 mg/mL, 24 mg/mL
Solution, oral: 0.5 mg/5 mL
Suspension, ophthalmic: 0.1% with methylcellulose 0.5% - **RESERVE USE**
Tablet: 0.25 mg, 0.75 mg, 1 mg, 1.5 mg, 2 mg, 4 mg, 6 mg

Dextran (Gentran, LMD, Macrodex, Rheomacrodex)

High molecular weight: 6% Dextran 75 in D5W, 6% Dextran 75 in NS, 6% Dextran 70 in NS
Low molecular weight: 10% Dextran 40 in D5W, 10% Dextran 40 in NS

Dextroamphetamine (Dexedrine)

Capsule, sustained release: 5 mg, 10 mg, 15 mg
Tablet: 5 mg, 10 mg (5 mg tablet contain tartrazine)

Dextromethorphan

Capsule: 30 mg
Liquid, oral: 3.5 mg/5 mL, 7.5 mg/5 mL, 10 mg/15 mL, 15 mg/5 mL
Liquid, oral, sustained release: 30 mg/5 mL
Lozenges: 2.5 mg, 5 mg, 7.5 mg

Dextrose/Sodium Chloride Intravenous Solution

Dextrose 5% in 0.2% Sodium Chloride
Dextrose 5% in 0.45% Sodium Chloride
Dextrose 5% in 0.9% Sodium Chloride

Dextrose 5%/Sodium Chloride/Potassium Chloride Intravenous Solution

Dextrose 5%/Sodium Chloride 0.2%/Potassium Chloride
Infusion with Potassium Chloride: 10 mEq, 20 mEq
Dextrose 5%/Sodium Chloride 0.45%/Potassium Chloride
Infusion with Potassium Chloride: 10 mEq, 20 mEq, 40 mEq
Dextrose 5%/Sodium Chloride 0.9%/Potassium Chloride
Infusion with Potassium Chloride: 20 mEq, 40 mEq

Dextrose 5% in Water

Infusion

Dextrose 5% in Ringer's Lactate

Infusion

Dextrose 5% with Multiple Electrolytes (D5 E75, Baxter)

Infusion

Dextrose 50% in Water

Infusion: 500 mL, 1000 mL, 2000 mL
Syringe: 50 mL
Vials: 50 mL

Diaper Rash Ointment (Desitin, Diaperene, Vitamin A&D)

see *Cod Liver Oil/Zinc Oxide/Talc (Desitin)*

see *Vitamin A&D Ointment*

see *Zinc Oxide/Petrolatum/Imidazolidinyl Urea (Diaperene)*

Diaper Rash Powder (Mexsana)

Powder: contains kaolin, eucalyptus oil, camphor, corn starch, lemon oil, zinc oxide

Diazepam (Valium, Diastat) C-IV

Gel, rectal: 2.5 mg, 5 mg, 10 mg, 15 mg, 20 mg

Injection: 5 mg/mL

Solution, oral: 1 mg/mL, 5 mg/mL

Tablet: 2 mg, 5 mg, 10 mg

Dibucaine (Nupercainal)

Cream, topical: 0.5%

Ointment, topical: 1%

Dicloxacillin (Dycill, Dynapen, Pathocil)

Capsule: 125 mg, 250 mg, 500 mg

Powder for oral suspension: 62.5 mg/mL

Dicyclomine (Bentyl)

Capsule: 10 mg, 20 mg

Injection: 10 mg/mL

Syrup: 10 mg/5 mL

Tablet: 10 mg

Didanosine (ddI, Videx)

Powder for oral solution: 100 mg, 167 mg, 250 mg, 375 mg, 2 gm, 4 gm

Tablet, chewable: 25 mg, 50 mg, 100 mg, 150 mg, 200 mg

Digoxin (Lanoxin)

Capsule: 50 mcg, 100 mcg, 200 mcg

Elixir: 50 mcg/mL with 10% alcohol

Injection: 100 mcg/mL, 250 mcg/mL

Tablet: 125 mcg, 250 mcg, 500 mcg

Diltiazem (Cardizem)

Capsule, sustained release:

Cardizem CD: 120 mg, 180 mg, 240 mg, 300 mg

Cardizem SR: 60 mg, 90 mg, 120 mg

Dilacor XR: 180 mg, 240 mg

Tiazac: 120 mg, 180 mg, 240 mg, 300 mg, 360 mg

Tablet: 30 mg, 60 mg, 90 mg, 120 mg

Tablet, sustained release: 120 mg, 180 mg, 240 mg

Dimercaprol (B.A.L.)

Injection: 100 mg/mL

diphenhydramine (Benadryl)

Capsule: 25 mg, 50 mg

Cream, topical: 1%, 2%

Injection: 50 mg/mL

Lotion: 1%

Syrup: 12.5 mg/5 mL

Tablet: 25 mg, 50 mg

Diphtheria & Tetanus Toxoids Adsorbed (DT)

Injection, single dose

Diphtheria & Tetanus Toxoids Adsorbed for Adult Use (Td)

Injection, single dose

Disulfiram (Antabuse)

Tablet: 250 mg, 500 mg

Divalproex (Depakote, Depakote ER, Divalproex ER)

Capsule, sprinkles: 125 mg

Tablet, delayed release: 125 mg, 250 mg, 500 mg

Tablet, extended release: 500 mg - **RESERVE USE****Docusate Calcium (Surfak)**

Capsule: 250 mg

Docusate Sodium (Colace, Doxinate)

Capsule: 100 mg, 250 mg

Liquid, oral: 150 mg/15 mL

Syrup: 60 mg/15 mL

Tablet: 100 mg

Docusate Sodium/Casanthrol (Peri-Colace)

Syrup, oral: Docusate 60 mg/Casanthrol 30 mg per 15 mL

Docusate Sodium/Sennosides (Peri-Colace)

Tablet: Docusate 50 mg/Sennosides 8.6 mg

Donepezil (Aricept) - **RESERVE USE**

Tablet: 5 mg, 10 mg

DOPamine (Intropin)

Infusion in D5W: 0.8 mg/mL, 1.6 mg/mL, 3.2 mg/mL

Injection: 40 mg/mL, 80 mg/mL, 160 mg/mL

Doxepin (Sinequan, Adapin)

Capsule: 10 mg, 25 mg, 50 mg, 75 mg, 100 mg, 150 mg
Concentrate, oral: 10 mg/mL

Doxycycline (Vibramycin, Periostat)

Capsule: 50 mg, 100 mg
Powder for injection: 100 mg, 200 mg
Powder for oral suspension: 25 mg/5 mL
Syrup: 50 mg/5 mL
Tablet: 20 mg, 50 mg, 100 mg

Emollient Lotion/Cream (Lubriderm, Allercreme, Keri Lotion, Cetaphil, Lac-Hydrin)

Lotion/Cream, topical

Emollient Ointment (Lanolin)

Ointment, topical

Enalapril (Vasotec)

Tablet: 2.5 mg, 5 mg, 10 mg, 20 mg

Enoxaparin (Lovenox)

Injection: 30 mg, 40 mg, 60 mg, 80 mg, 100 mg

Epinephrine (Adrenalin)

Auto-injector: 1:2000 [0.15 mg], 1:1000 [0.3 mg]
Injection: 1:100,000 [0.01 mg/mL], 1:10,000 [0.1 mg/mL], 1:1000 [1 mg/mL]

Ergocalciferol (Calciferol, Drisdol)

See *Vitamin D*

Erythromycin (Erythrocin)

Erythromycin base(Eryc, E-Mycin, Ery-Tab, E-Base, PCE):

Capsule, delayed release: 250 mg
Tablet, enteric coated: 250 mg, 333 mg, 500 mg
Tablet, film coated: 250 mg, 500 mg
Tablet, polymer coated particles: 333 mg, 500 mg

Erythromycin Ethylsuccinate (EryPed, E.E.S.):

Granules/Powder for oral suspension: 200 mg/5 mL, 400 mg/5 mL
Suspension, oral: 200 mg/5 mL, 400 mg/5 mL
Suspension, oral (drops): 100 mg/2.5 mL
Tablet: 400 mg
Tablet, chewable: 200 mg
Ointment, ophthalmic: 5%

Erythromycin/Benzoyl Peroxide (Benzamycin)

Gel, topical: Erythromycin 30 mg/Benzoyl Peroxide 50 mg per gram (with 16% alcohol)

Erythromycin Ethylsuccinate/Sulfisoxazole Suspension (Pediazole)

Suspension, oral: 200 mg/600 mg per 5 mL

Escitalopram (Lexapro)

Tablet: 10 mg, 20 mg

Estradiol (Estrace, Vivelle, Alora, Climara, Estraderm)

Cream, vaginal: 43 gm

Systems, transdermal: 0.025 mg, 0.0375 mg, 0.05 mg, 0.075 mg, 0.1 mg per 24 hr

Tablets: 0.5 mg, 1 mg, 2 mg

Estrogen/medroxyPROGESTERone (PremPro)

Tablet: Conjugated estrogen 0.625 mg/medroxyPROGESTERone 2.5 mg

Estrogens, Conjugated (Premarin)

Cream, vaginal: 0.625 mg/g

Injection: 25 mg

Tablet: 0.3 mg, 0.625 mg, 0.9 mg, 1.25 mg, 2.5 mg

Ethambutol (Myambutol)

Tablet: 100 mg, 400 mg

Ethinyl Estradiol/Norethindrone (Loestrin, Ortho-Novum 777)

Loestrin:

1/20: Ethinyl Estradiol 0.02 mg/Norethindrone 1 mg

1.5/30: Ethinyl Estradiol 0.03 mg/Norethindrone 1.5 mg

Ortho-Novum 777: Phase 1(Ethinyl Estradiol 0.035 mg/Norethindrone 0.5 mg),

Phase 2 (Ethinyl Estradiol 0.035 mg/Norethindrone 0.75 mg), Phase 3

(Ethinyl Estradiol 0.035 mg/Norethindrone 1 mg)

Ethinyl Estradiol/Norgestrel (Ovral, Lo-Ovral)

Lo-Ovral: Ethinyl Estradiol 0.03 mg/Norgestrel 0.3 mg

Ovral: Ethinyl Estradiol 0.05 mg/Norgestrel 0.5 mg

Ethionamide

Tablet, sugar coated: 250 mg

Ethosuximide (Zarontin)

Capsule: 250 mg

Syrup: 250 mg/5 mL

Ethyl Chloride

Spray: 100 g, 105 mL, 120 mL, 270 mL

Famotidine (Pepcid)

Injection: 10 mg/mL

Powder for oral suspension: 40 mg/5 mL

Tablet: 10 mg, 20 mg, 40 mg

Felbamate (Felbatol) - RESERVE USE

Suspension, oral: 600 mg/5 mL

Tablet: 400 mg, 600 mg

Felodipine (Plendil)

Tablet, extended release: 2.5 mg, 5 mg, 10 mg

Fentanyl (Duragesic) C-II

Patch, transdermal: 25 mcg/hr, 50 mcg/hr, 75 mcg/hr, 100 mcg/hr

Ferrous Fumarate/Docusate Sodium (Ferro-Sequels)[contains 33% elemental iron]

Tablet, timed released: Ferrous fumarate 150 mg [50 mg]/Docusate Sodium 100 mg

Ferrous Sulfate (Feosol, Fer-In-Sol) [contains 20% elemental iron]

Elixir with 5% alcohol: 220 mg/5 mL [18 mg/5 mL]

Tablet: 300 mg [60 mg], 325 mg [65 mg]

Fexofenadine (Allegra)

Capsule: 60 mg

Tablet: 30 mg, 60 mg, 180 mg

Fexofenadine/Pseudoephedrine (Allegra-D)

Tablet, extended release: 60 mg Fexofenadine/120 mg Pseudoephedrine

Flavoxate (Urispas)

Tablet, film coated: 100 mg

Fluconazole (Diflucan)

Tablet: 100 mg, 250 mg, 500 mg

Fludrocortisone (Florinef)

Tablet: 0.1 mg

Fluocinolone (Synalar)

Cream, topical: 0.01%, 0.025%

Ointment, topical: 0.025%

Shampoo: 0.01%

Solution, topical: 0.01%

Fluocinonide (Lidex)

Cream, topical: 0.05%
Gel, topical: 0.05%
Ointment, topical: 0.05%
Solution, topical: 0.05%

Fluorescein Sodium

Injection: 10%
Strip, ophthalmic: 1 mg

Fluoxetine (Prozac)

Capsule: 10 mg, 20 mg
Liquid, oral: 20 mg/5 mL
Tablet: 10 mg

Fluphenazine (Prolixin, Permitil)

Concentrate:
Permitil: 5 mg/mL with 1% alcohol
Prolixin: 5 mg/mL with 14% alcohol
Elixir: 2.5 mg/5 mL with 14% alcohol
Injection, as decanoate: 25 mg/mL
Injection, as hydrochloride: 2.5 mg/mL
Tablet: 1 mg, 2.5 mg, 5 mg, 10 mg

Fluticasone (Flonase, Flovent)

Aerosol, inhalation, oral: 44 mcg/actuation, 110 mcg/actuation, 220 mcg/actuation
Inhalation, nasal: 50 mcg/actuation

Fluvastatin (Lescol)

Capsule: 20 mg, 40 mg

Fluvoxamine (Luvox)

Tablet: 50 mg, 100 mg

Folic Acid (Folvite)

Tablet: 0.4 mg, 0.8 mg, 1 mg

Fosphenytoin (Cerebyx)

Injection: 100 Phenytoin Equivalents [PE]/2 mL, 500 PE/10 mL

Furosemide (Lasix)

Injection: 10 mg/mL
Solution, oral: 10 mg/mL, 40 mg/5 mL
Tablet: 20 mg, 40 mg, 80 mg

Gabapentin (Neurontin)

Capsule: 100 mg, 300 mg, 400 mg
Tablet: 600 mg, 800 mg

Galantamine (Reminyl)

Solution, oral: 4 mg/mL
Tablet, film coated: 4 mg, 8 mg, 12 mg

Gemfibrozil (Lopid)

Tablet, film coated: 600 mg

Gentamicin (Garamycin)

Infusion, premixed in D5W: 60 mg, 80 mg, 100 mg
Infusion, premixed in NS: 40 mg, 60 mg, 80 gm, 90 mg, 100 mg, 120 mg
Injection: 10 mg/mL, 40 mg/mL
Injection, intrathecal (preservative free): 2 mg/mL
Ointment, ophthalmic: 0.3% [3 mg/g]
Solution, ophthalmic: 0.3% [3 mg/mL]

glipiZIDE (Glucotrol)

Tablet: 5 mg, 10 mg
Tablet, extended release: 2.5 mg, 5 mg, 10 mg

Glucagon

Powder for injection: 1 mg

glyBURIDE (Micronase, DiaBeta)

Tablet: 1.25 mg, 2.5 mg, 5 mg
Tablet, micronized: 1.5 mg, 3 mg, 4.5 mg, 6 mg

Glycerin (Sani-Supp)

Suppository, rectal

Griseofulvin (Fulvicin)

Microsize:
 Capsule: 125 mg, 250 mg
 Suspension, oral: 125 mg/5 mL with 0.2% alcohol
 Tablet: 250 mg, 500 mg
Ultramicrosize:
 Tablet: 125 mg, 165 mg, 250 mg, 330 mg

Guaifenesin (Robitussin)

Caplet, sustained release: 600 mg
Liquid, oral: 100 mg/5 mL, 200 mg/5 mL
Tablet: 100 mg, 200 mg
Tablet, sustained release: 600 mg

Guaifenesin/Dextromethorphan (Robitussin DM)

Liquid, oral: Guaifenesin 100 mg/Dextromethorphan 100 mg per 5 mL

Guaifenesin/Pseudoephedrine (Entex PSE)

Tablet: Guaifenesin 600 mg/Pseudoephedrine 120 mg

Guanethidine (Ismelin)

Tablet: 10 mg, 25 mg

Haloperidol (Haldol)

Concentrate, oral: 2 mg/mL

Injection, as decanoate: 50 mg/mL, 100 mg/mL

Injection, as lactate: 5 mg/mL

Tablet: 0.5 mg, 1 mg, 2 mg, 5 mg, 10 mg, 20 mg

Heparin

Injection: 100 units/mL, 1,000 units/mL, 10,000 units/mL, 20,000 units/mL

Hepatitis A Vaccine (Vaqta)

Injection, single dose

Hepatitis B Immune Globulin (HBIG)

Injection, single dose

Hepatitis B Virus Vaccine, Recombinant (Recombivax HB, Engerix-B)

Injection: 5 mcg/mL, 10 mcg/mL, 20 mcg/mL

Hexachlorophene (pHisoHex)

Foam: 0.23% with 56% alcohol

Liquid, topical: 3%

Homatropine (Isopto Homatropine)

Solution, ophthalmic: 2%, 5%

hydrALAZINE (Apresoline)

Tablet: 10 mg, 25 mg, 50 mg, 100 mg

Hydrochlorothiazide (HydroDIURIL, Esidrix)

Solution, oral: 50 mg/5 mL

Tablet: 25 mg, 50 mg, 100 mg

Hydrocodone/Guaifenesin (Hycotuss, Kwelcof)

Liquid, oral: Hydrocodone 5 mg/Guaifenesin 100 mg per 5 mL

Hydrocortisone

Injection, as sodium succinate: 100 mg, 250 mg, 500 mg, 1000 mg

Suppositories, rectal, as acetate: 10 mg, 25 mg

Suspension, oral, as cypionate: 10 mg/5 mL

Hydrocortisone base:

Cream, rectal: 1%, 2.5%

Tablet, oral: 5 mg, 10 mg, 20 mg

Hydrocortisone, topical (Lanacort, Corticaine):

Cream, topical: 0.1%, 0.2%, 0.5%, 1%, 2.5%

Lotion, topical: 0.25%, 0.5%, 1%, 2%, 2.5%

Ointment, topical: 0.1%, 0.2%, 0.5%, 1%, 2.5%

Solution, topical: 0.1%

Hydrogen Peroxide

Solution, topical: 3%

hydrOXYzine (Atarax)

Injection, as hydrochloride: 25 mg/mL, 50 mg/mL

Syrup, as hydrochloride: 10 mg/5 mL

Tablet, as hydrochloride: 10 mg, 25 mg, 50 mg, 100 mg

Ibuprofen (Motrin)

Suspension, oral: 40 mg/mL, 100 mg/5 mL

Tablet: 200 mg, 400 mg, 600 mg, 800 mg

Tablet, chewable: 50 mg, 100 mg

Imipramine (Tofranil)

Capsule: 75 mg, 100 mg, 125 mg, 150 mg

Tablet: 10 mg, 25 mg, 50 mg

Indinavir (Crixivan)

Capsule: 400 mg

Influenza Virus Vaccine (Fluzone, Fluvirin)

Injection, single dose

Insulin, Combination (70/30)

Injection: 100 units/mL

Insulin, Glargine (Lantus)

Injection: 100 units/mL

Insulin, Lente

Injection: 100 units/mL

Insulin, Lispro (Humalog)

Injection: 100 units/mL

Insulin, Lispro/Insulin, Lispro Protamine (Humalog Mix 75/25)

Injection: 100 units/mL

Insulin, NPH

Injection: 100 units/mL

Insulin, Regular

Injection: 100 units/mL

Insulin, Ultralente

Injection: 100 units/mL

Ipecac Syrup

Syrup: 70 mg/mL

Ipratropium (Atrovent)

Inhalation: 18 mcg/actuation

Solution, nasal: 0.03%, 0.06%

Solution, nebulizing: 0.02%

Iron Dextran Complex (Imferon)

Injection: 50 mg/mL

Isoniazid (INH)

Injection: 100 mg/mL

Syrup: 50 mg/5 mL

Tablet: 50 mg, 100 mg, 300 mg

Isosorbide Dinitrate (Isordil, Sorbitrate)

Capsule, sustained release: 40 mg

Tablet, chewable: 5 mg, 10 mg

Tablet, oral: 5 mg, 10 mg, 20 mg, 30 mg, 40 mg

Tablet, sublingual: 2.5 mg, 5 mg, 10 mg

Tablet, sustained release: 40 mg

Isosorbide Mononitrate (Imdur, ISMO, Monoket)

Tablet: 10 mg, 20 mg

Tablet, extended release: 30 mg, 60 mg, 120 mg

Kaolin-Pectin

Suspension: 30 mL, 120 mL, 180 mL, 240 mL

Ketoconazole (Nizoral)

Cream, topical: 2%

Shampoo: 2%

Tablet: 200 mg

Ketorolac (Toradol)

Injection: 15 mg/mL, 30 mg/mL

Labetalol (Normodyne)

Tablet: 100 mg, 200 mg, 300 mg

Lactobacillus Acidophilus (Lactinex, Bacid)

Capsule

Granules: 1 g/packet

Tablet, chewable

Lactulose (Cephulac)

Syrup: 10 g/15 mL

Lamivudine (Epivir)

Solution, oral: 10 mg/mL

Tablet: 150 mg

Lamivudine/Zidovudine (Combivir)

Tablet: Lamivudine 150 mg/Zidovudine 300 mg

Lamotrigine (Lamictal)

Tablet: 25 mg, 100 mg, 150 mg, 200 mg

Lansoprazole (Prevacid)

Capsule, enteric coated granules: 15 mg, 30 mg

Granules for oral suspension: 15 mg, 30 mg

Latanoprost (Xalatan)

Solution, ophthalmic: 0.005%

Leucovorin (Wellcovorin)

Injection: 3 mg/mL

Powder for injection: 25 mg, 50 mg, 100 mg, 350 mg

Tablet: 5 mg, 10 mg, 15 mg, 25 mg

Levarterenol (Levophed)

see *Norepinephrine*

Levetiracetam (Keppra)

Tablets: 250 mg, 500 mg, 750 mg

Levodopa (Larodopa)

Capsule: 100 mg, 250 mg, 500 mg

Tablet: 100 mg, 250 mg, 500 mg

Levodopa/Carbidopa (Sinemet)

10/100: Carbidopa 10 mg/Levodopa 100 mg
25/100: Carbidopa 25 mg/Levodopa 100 mg
25/250: Carbidopa 25 mg/Levodopa 250 mg
Sustained release: Carbidopa 25 mg/Levodopa 100 mg, Carbidopa 50 mg/Levodopa 200 mg

Levofloxacin (Levaquin)

Infusion: 250 mg, 500 mg
Tablet: 250 mg, 500 mg

Levonorgestrel/Ethinyl Estradiol (Tri-Levlen, Triphasil)

Tablet: Phase I (Levonorgestrel 0.05 mg/Ethinyl Estradiol 30 mcg), Phase 2 (Levonorgestrel 0.075 mg/Ethinyl Estradiol 40 mg), Phase 3 (Levonorgestrel 0.125 mg/ Ethinyl Estradiol 30 mg)

Levothyroxine (Synthroid)

Powder for injection: 200 mcg/mL, 500 mcg/mL
Tablet: 25 mcg, 50 mcg, 75 mcg, 88 mcg, 100 mcg, 112 mcg, 112 mcg, 125 mcg, 150 mcg, 175 mcg, 200 mcg, 300 mcg

Lidocaine (Xylocaine)

Cream, topical: 2%
Injection: 10%
Gel, topical: 2%, 2.5%
Liquid, topical: 2.5%
Liquid, viscous: 2%
Ointment, topical: 2.5%, 5%
Solution, topical: 2%, 4%

Lindane (Gamma Benzene Hexachloride, Kwell)

Cream, topical: 1%
Lotion: 1%
Shampoo: 1%

Liotrix (Thyrolar, Euthroid)

Tablet: 15 mg, 30 mg, 60 mg, 120 mg, 180 mg [thyroid equivalent]

Lisinopril (Prinivil, Zestril)

Tablet: 2.5 mg, 5 mg, 10 mg, 20 mg, 40 mg

Lithium Carbonate (Eskalith, Lithonate, Lithobid)

Capsule: 150 mg, 300 mg, 600 mg
Tablet: 300 mg
Tablet, controlled release: 450 mg
Tablet, slow release: 300 mg

Lithium Citrate

Syrup: 300 mg/5 mL

Loperamide (Imodium)

Capsule: 2 mg

Liquid, oral: 1 mg/5 mL

Tablet: 2 mg

Loratadine (Claritin)

Tablet: 10 mg

Lorazepam (Ativan) C-IV

Injection: 2 mg/mL, 4 mg/mL

Solution, oral: 2 mg/mL

Tablet: 0.5 mg, 1 mg, 2 mg

Loxapine (Loxitane)

Capsule: 5 mg, 10 mg, 25 mg, 50 mg

Concentrate, oral: 25 mg/mL

Injection: 50 mg/mL

Magnesium Citrate

Solution, oral: 300 mL

Magnesium Hydroxide (Milk of Magnesia)

Liquid, oral: 400 mg/5 mL

Liquid, oral, concentrate: 800 mg/5 mL

Tablet, chewable: 311 mg

Magnesium Sulfate (Epsom Salt)

Granules: ~40 mEq magnesium/5 g

Injection: 100 mg/mL, 125 mg/mL, 250 mg/mL, 500 mg/mL

Maprotiline (Ludiomil)

Tablet: 25 mg, 50 mg, 75 mg

Measles, Mumps and Rubella Virus Vaccine, Live (MMR II)

Injection, single dose

Mebendazole (Vermox)

Tablet, chewable: 100 mg

Meclizine (Antivert, Bonine)

Tablet: 12.5 mg, 25 mg, 50 mg

Tablet, chewable: 25 mg

medroxyPROGESTERone (Provera)

Injection, suspension: 100 mg/mL, 150 mg/mL, 400 mg/mL
Tablet: 2.5 mg, 5 mg, 10 mg

Mephobarbital (Mebaral) C-IV

Tablet: 32 mg, 50 mg, 100 mg

Mesalamine (Asacol, Pentasa, Rowasa)

Capsule, controlled release: 250 mg
Suppository: 500 mg
Suspension, rectal: 4 gm/60 mL
Tablet, delayed release: 400 mg

Mesoridazine (Serentil) - RESERVE USE

Injection: 25 mg/mL
Liquid, oral: 25 mg/mL
Tablet: 10 mg, 25 mg, 50 mg, 100 mg

Metaproterenol (Alupent)

Aerosol, oral: 0.65 mg/metered dose
Solution for inhalation: 0.4%, 0.6%, 5%

Metformin (Glucophage, Glucophage XR)

Tablet: 500 mg, 850 mg, 1000 mg
Tablet, extended release: 500 mg

Methadone (Dolophine) C-II

Solution, oral: 1 mg/mL
Tablet: 5 mg, 10 mg, 40 mg

Methimazole (Tapazole)

Tablet: 5 mg, 10 mg

Methocarbamol (Robaxin)

Tablet: 500 mg, 750 mg

Methotrexate

Injection: 2.5 mg/mL, 25 mg/mL
Injection, preservative free: 25 mg/mL
Powder for injection: 20 mg, 25 mg, 50 mg, 100 mg, 250 mg, 1 g
Tablet: 2.5 mg

Methylcellulose (Citrucel)

Powder: Methylcellulose 2 g per tbsp [with sucrose]
Powder, sugar free: Methylcellulose 2 g per tbsp [with phylalanine]

Methyldopa (Aldomet)

Injection: 50 mg/mL

Suspension, oral: 250 mg/5 mL

Tablet: 125 mg, 250 mg, 500 mg

Methylphenidate (Ritalin, Concerta)

Tablet: 5 mg, 10 mg, 20 mg

Tablet, extended release: 18 mg, 36 mg

Tablet, sustained release: 20 mg

Methylprednisolone (Medrol)

Injection, as acetate: 20 mg/mL, 40 mg/mL, 80 mg/mL

Injection, as sodium succinate: 40 mg, 125 mg, 500 mg, 1000 mg, 2000 mg

Tablet: 2 mg, 4 mg, 8 mg, 16 mg, 24 mg, 32 mg

Methyl Salicylate (Ben-Gay)

Cream, topical: 30%

methylTESTOSTERone (Android, Oretón) C-IV

Capsule: 10 mg

Tablet: 10 mg, 25 mg

Tablet, buccal: 5 mg, 10 mg

Metoclopramide (Reglan)

Injection: 5 mg/mL

Syrup, sugar free: 5 mg/5 mL

Tablet: 5 mg, 10 mg

Metoprolol (Lopressor)

Tablet: 50 mg, 100 mg

Tablet, sustained release: 50 mg, 100 mg, 200 mg

Metronidazole (Flagyl, Noritate, MetroGel)

Capsule: 375 mg

Cream, topical: 1%

Gel, topical: 0.75% [7.5 mg/mL]

Gel, vaginal: 0.75%

Injection: 5 mg/mL

Powder for injection: 500 mg

Tablet: 250 mg, 500 mg

Miconazole (Monistat)

Cream, topical: 2%
Cream, vaginal: 2%
Injection: 10 mg/mL
Lotion: 2%
Powder, topical: 2%
Spray, topical: 2%
Suppository, vaginal: 100 mg, 200 mg

Midazolam (Versed) C-IV

Injection: 1 mg/mL, 5 mg/mL

Mineral Oil - for topical use only

Oil: 480 mL

Mirtazapine (Remeron)

Tablet: 15 mg, 30 mg

Misoprostol (Cytotec)

Tablet: 100 mcg, 200 mcg

Molindone (Moban)

Concentrate, oral: 20 mg/mL
Tablet: 5 mg, 10 mg, 25 mg, 50 mg, 100 mg

Mometasone (Nasonex)

Inhalation, nasal: 50 mcg/actuation

Morphine C-II

Injection: 1 mg/mL, 2 mg/mL, 4 mg/mL, 10 mg/mL
Solution, oral: 20 mg/mL
Tablet, controlled release: 15 mg, 30 mg

Multivitamin (Unicap, Hexavitamins)

Liquid, oral: each solution contains a minimum of USDA requirements
Tablet: each tablet contains a minimum of USDA requirements
Tablet, chew: each tablet contains a minimum of USDA requirements

Multivitamin, Prenatal (Filibon)

Tablet: each tablet contains a minimum of USDA requirements

Multivitamin/Minerals

Liquid, oral: each solution contains a minimum of USDA requirements
Tablet: each tablet contains a minimum of USDA requirements
Tablet, chew: each tablet contains a minimum of USDA requirements

Multivitamins, Pediatric (Poly-Vi-Sol)

Liquid, oral: each solution contains a minimum of USDA requirements

Mupirocin (Bactroban)

Ointment, intranasal: 2%

Ointment, topical: 2%

Nabumetone (Relafen) - RESERVE USE

Tablet: 500 mg, 750 mg

Nadolol (Corgard)

Tablet: 20 mg, 40 mg, 80 mg, 120 mg, 160 mg

Nafcillin (Unipen)

Capsule: 250 mg

Powder for injection: 500 mg, 1 g, 2 g, 4 g, 10 g

Solution: 250 mg/5 mL

Tablet: 500 mg

Naloxone (Narcan)

Injection: 0.4 mg/mL, 1 mg/mL

Naltrexone (Trexan, ReVia)

Tablet: 50 mg

Naphazoline (Naphcon, AK-Con)

Solution, ophthalmic: 0.012%, 0.1%

Naproxen (Naprosyn)

Tablet: 220 mg, 250 mg, 275 mg, 375 mg, 500 mg, 550 mg

Tablet, controlled release: 500 mg

Nefazodone (Serzone)

Tablet: 50 mg, 100 mg, 150 mg, 200 mg, 250 mg

Nelfinavir (Viracept)

Powder for oral solution: 50 mg/g

Tablet: 250 mg

Neomycin (Mycifradin)

Tablet: 500 mg

Neomycin/Polymyxin B/Bacitracin (Triple Antibiotic Ointment)

Ointment, topical: Neomycin 3.5 mg/Polymyxin B 5000 units/Bacitracin 400 units

Neomycin/Polymyxin B/Hydrocortisone (Cortisporin)

Solution, otic: Neomycin 5 mg/Polymyxin B 10,000 units/Hydrocortisone 10 mg per mL

Suspension, otic: Neomycin 5 mg/Polymyxin B 10,000 units/Hydrocortisone 10 mg per mL

Nevirapine (NVP, Viramune)

Tablet: 200 mg

Niacin/Nicotinamide (Nicobid)

Capsule, extended release: 250 mg, 500 mg

Tablet: 50 mg, 100 mg, 250 mg, 500 mg

Tablet, extended release: 250 mg, 500 mg, 750 mg, 1000 mg

Nicotine (Nicoderm, Habitrol, ProStep, Nicotrol, Nicorette)

Patch, transdermal:

Habitrol: 21 mg/day, 14 mg/day, 7 mg/day

Nicoderm: 21 mg/day, 14 mg/day, 7 mg/day

Nicotrol: 15 mg/day (gradual release over 16 hours)

Pieces, chewing gum, as polacrilex: 2 mg/square, 4 mg/square

NIFEdipine (Procardia)

Capsule, liquid-filled: 10 mg, 20 mg

Tablet, sustained release: 30 mg, 60 mg, 90 mg

Nitrofurantoin (Macrodantin)

Capsule: 50 mg, 100 mg

Capsule, extended release: 100 mg

Capsule, macrocrystal: 25 mg, 50 mg, 100 mg

Capsule, macrocrystal/monohydrate: 100 mg

Suspension, oral: 25 mg/mL

Nitroglycerin

Capsule, sustained release: 2.5 mg, 6.5 mg, 9 mg, 13 mg

Ointment, topical 2%: 30 gm, 60 gm

Patch, transdermal, topical: systems designed to deliver 2.5 mg, 5 mg, 7.5 mg, 10 mg, or 15 mg over 24 hours

Spray, translinqual: 0.4 mg/metered spray

Tablet, buccal, controlled release: 2 mg, 3 mg

Tablet, sublingual: 0.3 mg, 0.4 mg, 0.6 mg

Tablet, sustained release: 2.6 mg, 6.5 mg, 9 mg

Non-Soap Cleanser (Cetaphil)

Lotion

Norepinephrine or Levarterenol (Levophed)

Injection: 1 mg/mL

Norgestimate/Ethinyl Estradiol (Ortho Tri-Cyclen)

Tablet: 21 day, 28 day

Nortriptyline (Aventyl, Pamelor)

Capsule: 10 mg, 25 mg, 50 mg, 75 mg

Solution: 10 mg/5 mL

Nystatin (Mycostatin)

Cream, topical: 100,000 units/g

Ointment, topical: 100,000 units/g

Powder for oral suspension: 50 million units, 1 billion units, 2 billion units, 5 billion units

Powder, topical: 100,000 units/g

Suspension, oral: 100,000 units/mL

Tablet, oral: 500,000 units

Troche: 200,000 units

Olanzapine (Zyprexa, Zydis)

Tablet: 2.5 mg, 5 mg, 7.5 mg, 10 mg

Tablet, rapid dissolving: 5 mg, 10 mg

Olopatadine (Patanol)

Solution, ophthalmic: 0.1%

Omeprazole (Prilosec)

Capsule, delayed release: 10 mg, 20 mg, 40 mg

Ophthalmic Lubricant (HypoTears, HypoTears PF) [preservative-free, lanolin-free]

Ointment, ophthalmic: 3.5 gm

Solution, ophthalmic: 0.6 mL

Oxacillin (Prostaphlin)

Capsule: 250 mg, 500 mg

Powder for injection: 250 mg, 500 mg, 1 g, 2 g, 4 g, 10 g

Powder for oral solution: 250 mg/5 mL

Oxazepam (Serax) C-IV

Capsule: 10 mg, 15 mg, 30 mg

Tablet: 15 mg

Oxcarbazepine (Trileptal)

Tablet: 150 mg, 300 mg, 600 mg

Oxybutynin (Ditropan, Ditropan XL)

Syrup: 5 mg/5 mL

Tablet: 5 mg

Tablet, extended release: 5 mg, 10 mg, 15 mg

Oxycodone (OxyContin) C-II

Tablet, controlled release: 10 mg, 20 mg, 40 mg, 80 mg, 160 mg

Oxymetazoline (Afrin)

Solution, nasal, drops: 0.025%, 0.05%

Solution, nasal, spray: 0.05%

Pancrelipase (Pancrease, Creon)

Capsule: contains lipase, protease and amylase

Tablet: contains lipase, protease and amylase

Pantoprazole (Protonix)

Tablet: 40 mg, 20 mg

Paroxetine (Paxil)

Tablet: 10 mg, 20 mg, 30 mg, 40 mg

Tablet, controlled release: 12.5 mg, 25 mg, 37.5 mg, 50 mg

Penicillamine (Cuprimine)

Capsule: 125 mg, 250 mg

Tablet: 250 mg

Penicillin G Benzathine (Bicillin)

Injection: 300,000 units/mL, 600,000 units/mL

Penicillin G Benzathine/Penicillin G Procaine (Bicillin C-R)

Injection: Penicillin G Benzathine 150,000 units/Penicillin G Procaine 150,000 units,

Penicillin G Benzathine 900,000 units/Penicillin G Procaine 300,000 units

Penicillin G Procaine (Wycillin)

Injection (suspension): 300,000 units/mL, 500,000 units/mL, 600,000 units/mL

Penicillin G Sodium

Injection: 5 million units

Penicillin V Potassium (Pen-Vee K, V-Cillin K)

Powder for oral solution: 125 mg/5 mL, 250 mg/5 mL

Tablet: 125 mg, 250 mg, 500 mg

Pentamidine (Pentam) - RESERVE USE

Inhalation: 300 mg

Powder for injection: 300 mg

Permethrin (Elimite, NIX)

Cream, topical: 5%

Liquid, topical: 1%

Perphenazine (Trilafon)

Concentrate, oral: 16 mg/5 mL
Injection: 5 mg/mL
Tablet: 2 mg, 4 mg, 8 mg, 16 mg

Petrolatum, White (Vaseline)

Ointment, topical: 430 g

Phenazopyridine (Pyridium)

Tablet: 95 mg, 100 mg, 200 mg

Phenelzine (Nardil)

Tablet: 15 mg

Phenobarbital (Luminal) C-IV

Capsule: 16 mg
Elixir: 20 mg/5 mL
Injection: 30 mg/mL, 60 mg/mL, 65 mg/mL, 130 mg/mL
Tablet: 8 mg, 15 mg, 16 mg, 30 mg, 32 mg, 60 mg, 65 mg, 100 mg

Phenol (Chloraseptic)

Mouthwash/Gargle: 1.4%
Throat Spray: 0.5%

Phenylephrine (Neo-Synephrine)

Solution, nasal, drops: 0.125%, 0.25%, 0.5%
Solution, nasal, spray: 0.25%, 0.5%, 1%
Solution, ophthalmic: 2.5%, 10%

Phenytoin (Dilantin)

Capsule, extended release: 30 mg, 100 mg
Capsule, immediate release: 100 mg
Injection: 50 mg/mL
Suspension, oral: 125 mg/5 mL
Tablet, chewable: 50 mg

Physostigmine (Antilirium)

Injection: 1 mg/mL

Phytonadione (Vitamin K₁, Mephyton, Konakion)

Injection, aqueous colloidal: 2 mg/mL, 10 mg/mL
Injection, aqueous (IM only): 2 mg/mL, 10 mg/mL
Tablet: 5 mg

Pilocarpine (Isotopto Carpine)

Solution, ophthalmic, as hydrochloride: 1%, 2%, 4%

Pneumococcal Vaccine, Polysaccharide (Pneumovax)

Injection, single dose

Podophyllum Resin

Liquid, topical: 25% in benzoin

Poliovirus Vaccine, Inactivated (IPOL)

Injection, single dose

Polycarbophil (Fibercon, Fiber-Lax)

Tablet: 500 mg, 600 mg

Polyethylene Glycol (MiraLax)

Powder for oral solution

Polymyxin B/Bacitracin (Polysporin)

Ointment, ophthalmic: Polymyxin B 10,000 units/Bacitracin 500 units

Ointment, topical: Polymyxin B 10,000 units/Bacitracin 500 units

Powder, topical: Polymyxin B 10,000 units/Bacitracin 500 units

Polymyxin B/Neomycin (Neosporin)

Cream: Polymyxin B 10,000 units/Neomycin 3.5 mg

Polymyxin B/Trimethoprim (Polytrim)

Solution, ophthalmic: Polymyxin B 10,000 units/Trimethoprim 1 mg/mL

Potassium Chloride

Capsules: 8 mEq, 10 mEq

Crystals for oral suspension, extended release: 20 mEq/packet

Liquid, oral: 10% [20 mEq/15 mL], 15% [30 mEq/15 mL], 20% [40 mEq/15 mL]

Powder for oral suspension(per packet): 15 mEq, 20 mEq, 25 mEq

Injection, concentrate: 2 mEq/mL

Tablet, controlled release (microencapsulated): 750 mg [10 mEq], 1500 mg [20 mEq]

Tablet, controlled release (wax matrix): 500 mg [6.7 mEq], 600 mg [8 mEq], 750 mg [10 mEq]

Potassium Citrate (Urocit K)

Tablet: 5 mEq, 10 mEq

Potassium Citrate Combinations (Polycitra, Polycitra-LC, Polycitra K, Citrolith)

Solution, oral: containing Sodium Citrate /Potassium Citrate/Citric Acid

Solution, oral: containing Sodium Citrate/Potassium Citrate

Potassium Iodide (SSKI)

Solution, oral: 100 mg/mL, 1 g/mL

Povidone-Iodine (Betadine)

Cleanser, topical: 60 mL, 240 mL
Foam, topical: 10%
Liquid, topical: 473 mL
Ointment, topical: 10%
Solution, prep: 30 mL, 60 mL, 240 mL, 473 mL, 1000 mL, 4000 mL
Solution, swabsticks: 4"
Solution, topical: 10%

Pramoxine (Tronothane)

Cream, topical: 1%
Gel, topical: 1%
Lotion: 1%
Ointment, topical: 1%
Spray: 1%

Prazosin (Minipress)

Capsule: 1 mg, 2 mg, 5 mg

prednisoLONE (Delta-Cortef)

Injection, as sodium phosphate: 20 mg/mL
Liquid, oral: 5 mg/mL
Syrup: 15 mg/mL
Tablet: 5 mg

predniSONE (Meticorten, Deltasone)

Solution, oral, concentrate: 5 mg/mL with 30% alcohol
Syrup: 5 mg/5 mL
Tablet: 1 mg, 2.5 mg, 5 mg, 10 mg, 20 mg, 50 mg

Primidone (Mysoline)

Suspension, oral: 250 mg/5 mL
Tablet: 50 mg, 250 mg

Probenecid (Benemid)

Tablet: 500 mg

Procainamide (Pronestyl)

Capsule: 250 mg, 375 mg, 500 mg
Tablet: 250 mg, 375 mg, 500 mg
Tablet, sustained release: 250 mg, 500 mg, 750 mg, 1000 mg

Prochlorperazine (Compazine)

Injection: 5 mg/mL
Suppository, rectal: 2.5 mg, 5 mg, 25 mg
Syrup: 5 mg/5 mL
Tablet: 5 mg, 10 mg, 25 mg

Promethazine (Phenergan)

Injection: 25 mg/mL, 50 mg/mL
Suppository, rectal: 12.5 mg, 25 mg, 50 mg
Syrup: 6.25 mg/5 mL, 25 mg/5 mL
Tablet: 12.5 mg, 25 mg, 50 mg

Propantheline (Pro-Banthine)

Tablet: 7.5 mg, 15 mg

Proparacaine (Alcaine)

Solution, ophthalmic: 0.5%

Propranolol (Inderal)

Capsule, sustained release: 60 mg, 80 mg, 120 mg, 160 mg
Injection: 1 mg/mL
Solution, oral: 4 mg/mL, 8 mg/mL, 80 mg/mL
Tablet: 10 mg, 20 mg, 40 mg, 60 mg, 80 mg, 90 mg

Propylethylene Glycol Electrolyte Solution (PEG, Co-Lyte, GoLYTELY, OCL)

Powder for oral solution: 2000 mL, 4000 mL, 4800 mL, 6000 mL

Propylthiouracil

Tablet: 50 mg

Protamine

Injection: 10 mg/mL

Protriptyline (Vivactil)

Tablet: 5 mg, 10 mg

Pseudoephedrine (Sudafed)

Liquid, oral: 15 mg/5 mL, 30 mg/mL
Tablet, immediate release: 30 mg, 60 mg
Tablet, timed release: 120 mg
Tablet, extended release: 120 mg, 240 mg

Psyllium (Metamucil)

Granules: 4.03 g per rounded teaspoon, 2.5 g per rounded teaspoon
Powder, hydrophilic: 3.4 g per rounded teaspoon

Pyrantel (Antiminth)

Capsule: 180 mg
Liquid, oral: 50 mg/mL
Suspension, oral: 50 mg/mL

Pyrazinamide

Tablet: 500 mg

Pyrethrins/Piperonyl Butoxide (A-200, RID)

Gel, topical: 0.3%
Liquid, topical: 0.18%, 0.3%
Shampoo: 0.3%, 0.33%

Pyridoxine (Vitamin B₆)

Injection: 100 mg/mL
Tablet: 25 mg, 50 mg, 100 mg

Quetiapine (Seroquel) – RESERVE USE

Tablet: 25 mg, 100 mg, 200 mg, 300 mg

Quinidine Gluconate

Tablet, sustained release: 324 mg

Quinidine Sulfate

Tablet: 200 mg, 300 mg
Tablet, sustained action: 300 mg

Raloxifene (Evista)

Tablet: 60 mg

Ranitidine (Zantac)

Injection: 25 mg/mL
Granules, effervescent: 150 mg
Syrup: 15 mg/mL
Tablet: 75 mg, 150 mg, 300 mg
Tablet, effervescent: 150 mg

Rectal Hemorrhoidal Cream with Hydrocortisone

Cream: see Hydrocortisone

Rectal Hemorrhoidal Ointment (Anusol)

Ointment

Rectal Hemorrhoidal Suppositories (Wyanoids, Anusol)

Suppository, rectal:

Rectal Hemorrhoidal Suppositories with Hydrocortisone (Anusol-HC)

Suppository, rectal: see Hydrocortisone

Repaglinide (Prandin)

Tablet: 0.5 mg, 1 mg, 2 mg

Rifampin (Rifadin)

Capsule: 150 mg, 300 mg
Injection: 600 mg

Rifampin/Isoniazid (Rifamate)

Capsule: Rifampin 300 mg/Isoniazid 150 mg

Ringer's Lactate Solution (Hartmann's Solution)

Infusion: 150 mL, 250 mL, 500 mL, 1000 mL

Risperidone (Risperdal, Risperdal Consta)

Injection, long acting: 25 mg/2 mL, 37.5 mg/2 mL, 50 mg/2 mL

Solution, oral: 1 mg/mL

Tablet: 0.25 mg, 0.5 mg, 1 mg, 2 mg, 3 mg, 4 mg

Tablet, orally disintegrating: 0.5 mg, 1 mg, 2 mg

Ritonavir (Norvir)

Capsule: 100 mg

Solution, oral: 80 mg/mL

Rivastigmine (Exelon) - RESERVE USE

Capsule: 1.5 mg, 3 mg, 4.5 mg, 6 mg

Rosiglitazone (Avandia)

Tablet: 2 mg, 4 mg, 8 mg

Rubella Virus Vaccine Live (Meruvax II)

Injection, single dose

Salicylic Acid

Cream, topical: 2%

Gel, topical: 5%, 6%, 17%

Liquid, topical: 13.6%, 16.7%, 17%

Lotion: 3%

Ointment, topical: 3%

Soap: 2%

Salmeterol (Serevent)

Aerosol, inhalation: 25 mcg/dose

Powder, inhalation: 50 mcg

Saquinavir (Invirase, Fortovase)

Capsule: 200 mg

Scopolamine (Isopto Hyoscine)

Solution, ophthalmic: 0.25%

Selenium Sulfide (Selsun)

Shampoo: 1%, 2.5%

Senna (Senokot)

Tablet: 8.6 mg, 25 mg

Sertraline (Zoloft)

Tablet: 25 mg, 50 mg, 100 mg

Sevelamer (Renagel)

Tablet: 400 mg, 800 mg

Silver Nitrate

Applicator sticks: 75% with potassium nitrate 25%

Ointment, topical: 10%

Solution, topical: 10%, 25%, 50%

Silver Sulfadiazine (Silvadene)

Cream, topical: 1%

Simethicone (Mylicon)

Drops, oral: 40 mg/0.6 mL

Tablet, chewable: 40 mg, 80 mg, 125 mg

Simvastatin (Zocor)

Tablet: 5 mg, 10 mg, 20 mg, 40 mg, 80 mg

Sodium Bicarbonate

Injection: 4.2% [5 mEq/10 mL], 8.4% [10 mEq/10 mL]

Sodium Chloride

Drops, nasal: 0.9%

Infusion: 0.2%, 0.45%, 0.9%, 3%, 5%, 20%, 23.4%

Injection, bacteriostatic: 0.9%

Injection, for admixtures: 50 mEq, 100 mEq, 635 mEq

Ointment, ophthalmic: 5%

Solution, irrigation: 0.45%, 0.9%

Solution, nasal: 0.4%, 0.6%, 0.65%

Solution, nebulizing: 0.9%

Solution, ophthalmic: 2%, 5%

Tablet: 650 mg, 1 g

Tablet, enteric coated: 1 g

Tablet, slow release: 600 mg

Sodium Chloride Intravenous Solution

Sodium Chloride 0.2%

Sodium Chloride 0.45%

Sodium Chloride 0.9%

Sodium Citrate/Citric Acid (Bicitra, Oracit)

Solution, oral:

Bicitra: Sodium Citrate 500 mg/Citric Acid 334 mg per 5 mL

Oracit: Sodium Citrate 400 mg/Citric Acid 640 mg per 5 mL

Sodium Fluoride

Liquid, oral: 360 mL, 480 mL, 540 mL

Tablet, chewable: 0.25 mg, 0.5 mg, 1 mg, 1.1 mg, 2.2 mg

Sodium Lactate

Injection: Sodium 167 mEq/Lactate 168 mEq per liter

Sodium Phosphate/Biphosphate (Fleet Phospho-Soda, Fleet's Enema)

Enema: Sodium Phosphate 7 g/Sodium Biphosphate 19 g per 118 mL

Injection: Phosphate 3 mmol and Sodium 4 mEq per mL

Solution, oral: Sodium Phosphate 18 g/Sodium Biphosphate 48 g per 100 mL

Sodium Polystyrene Sulfonate (Kayexalate)

Powder for suspension: 454 gm

Suspension, oral: 1.25 gm/5 mL (with sorbitol and alcohol)

Sorbitol

Solution, oral: 70%

Spironolactone (Aldactone)

Tablet: 25 mg, 50 mg, 100 mg

Spironolactone/Hydrochlorothiazide (Aldactazide)

Tablet: Spironolactone 25 mg/Hydrochlorothiazide 25 mg, Spironolactone 50 mg/

Hydrochlorothiazide 50 mg

Stannous Fluoride (OmniMed, PerioMed)

Solution, oral: 0.4%, 0.63%

Stavudine (d4T, Zerit)

Capsule: 15 mg, 20 mg, 30 mg, 40 mg

Solution, oral: 1 mg/mL

Sucralfate (Carafate)

Suspension, oral: 1 g/10 mL

Tablet: 1 g

Sulfacetamide Sodium (Sulamyd, Sebizon)

Lotion: 10%

Ointment, ophthalmic: 10%

Solution, ophthalmic: 10%

Sulfasalazine (Azulfidine)

Tablet: 500 mg

Tablet, delayed release: 500 mg

Sulfur/Resorcinol (Sulforcin, Rezamid)

Lotion: Sulfur 5%/Resorcinol 2% [with up to 28% alcohol]

Sulindac (Clinoril)

Tablet: 150 mg, 200 mg

Sumatriptan (Imitrex)

Injection: 12 mg/mL

Nasal Spray: 5 mg, 20 mg

Tablet: 25 mg, 50 mg, 100 mg

Sunscreen/block

Cream/Lotion: contains a minimum SPF of 15

Tamoxifen (Nolvadex)

Tablet: 10 mg, 20 mg

Tazarotene (Tazorac, Avage)

Cream, topical: 0.05%, 0.1%

Gel, topical: 0.05%, 0.1%

Temazepam (Restoril) C-IV

Capsule: 7.5 mg, 15 mg, 30 mg

Terbinafine (Lamisil)

Cream, topical: 1%

Tablet: 250 mg

Terbutaline (Brethine)

Aerosol, oral: 0.2 mg/actuation

Injection: 1 mg/mL

Tablet: 2.5 mg, 5 mg

Testosterone (Androlan) C-IV

Injection, in oil, as cypionate: 100 mg/mL, 200 mg/mL

Injection, in oil, as enanthate: 100 mg/mL, 200 mg/mL

Tetracycline (Achromycin, Panmycin)

Capsule: 100 mg, 250 mg, 500 mg

Suspension, oral: 125 mg/5 mL

Tablet: 250 mg, 500 mg

Tetrahydrozoline (Visine Allergy Relief, Visine Moisturizing)

Solution, ophthalmic: 0.05%

Theophylline (Elixophyllin)

Capsule, timed release (12 hour): 130 mg, 260 mg

Capsule, timed release (24 hour): 100 mg, 200 mg, 300 mg

Solution, oral: 80 mg/15 mL, 150 mg/15 mL

Tablet, immediate release [Slo-phyllin]: 100 mg, 125 mg, 200 mg, 250 mg, 300 mg

Tablet, timed release:

Theolair SR (8-12 hour): 100 mg, 200 mg, 250 mg, 300 mg, 500 mg

Theo-Dur (8-24 hour): 100 mg, 200 mg, 300 mg, 450 mg

Theophylline SR (12-24 hour): 100 mg, 200 mg, 300 mg

Uniphyll (24 hour): 400 mg

Thiabendazole (Mintezol)

Suspension, oral: 500 mg/5 mL

Tablet, chewable: 500 mg

Thiamine (Vitamin B₁)

Injection: 100 mg/mL, 200 mg/mL

Tablet: 50 mg, 100 mg, 250 mg, 500 mg

Thioridazine (Mellaril) - RESERVE USE

Concentrate, oral: 30 mg/mL, 100 mg/mL

Suspension, oral: 25 mg/5 mL, 100 mg/mL

Tablet: 10 mg, 15 mg, 25 mg, 50 mg, 100 mg, 150 mg, 200 mg

Thiothixene (Navane)

Capsule: 1 mg, 2 mg, 5 mg, 10 mg, 20 mg

Concentrate, oral: 5 mg/mL

Thyroid, Desiccated (Thyroid)

Capsule (pork source): 60 mg, 120 mg, 180 mg, 300 mg

Tablet:

Armour: 15 mg, 30 mg, 60 mg, 90 mg, 120 mg, 180 mg, 240 mg, 300 mg

Thyrar (bovine source): 30 mg, 60 mg, 120 mg

Thyroid Strong (60 mg is equivalent to 90 mg thyroid, USP)

Thyroid, USP: 15 mg, 30 mg, 60 mg, 120 mg, 180 mg, 300 mg

Tiagabine (Gabatril)

Tablet: 4 mg, 12 mg, 16 mg, 20 mg

Ticarcillin (Ticar)

Powder for injection: 1 g, 3 g, 6 g, 20 g, 30 g

Ticarcillin/Clavulanate (Timentin)

Powder for injection: 3.1 g

Timolol (Timoptic)

Gel, ophthalmic: 0.25%, 0.5%

Solution, as maleate, ophthalmic: 0.25%, 0.5%

Solution, as maleate, ophthalmic, preservative free, single use: 0.25%, 0.5%

Tablet: 5 mg, 10 mg, 20 mg

Timolol/Dorzolamide (Cosopt)

Solution, ophthalmic: Timolol 0.5%/Dorzolamide 2%

Tioconazole (Vagistat-1)

Ointment, vaginal: 6.5%

Tizanidine (Zanaflex) - RESERVE USE

Tablet: 4 mg

Tobramycin (Nebcin, Tobrex)

Injection: 10 mg/mL, 40 mg/mL

Ointment, ophthalmic: 0.3%

Powder for injection: 40 mg/mL

Solution, ophthalmic: 0.3%

Tobramycin/Dexamethasone (TobraDex) [contains Benzalkonium] - RESERVE USE

Ointment, ophthalmic: Tobramycin 0.3%/Dexamethasone 0.1%

Suspension, ophthalmic: Tobramycin 0.3% / Dexamethasone 0.1%

TOLBUTamide (Orinase)

Injection, diagnostic: 1 g

Tablet: 250 mg, 500 mg

Tolnaftate (Tinactin)

Aerosol, topical, liquid: 1%

Aerosol, topical, powder: 1%

Cream, topical: 1%

Gel, topical: 1%

Powder, topical: 1%

Solution, topical: 1%

Tolterodine (Detrol, Detrol LA)

Capsule, extended release: 2 mg, 4 mg

Tablet: 1 mg, 2 mg

Topiramate (Topamax)

Tablet: 25 mg, 100 mg, 200 mg

Tramadol (Ultram)

Tablet: 50 mg

Tranylcypromine (Parnate)

Tablet: 10 mg

Travoprost (Travatan)

Solution, ophthalmic: 0.004%

Trazodone (Desyrel)

Tablet: 50 mg, 100 mg, 150 mg, 300 mg

Tretinoin Gel (Retin-A)

Cream, topical: 0.025%, 0.05%, 0.1%

Gel, topical: 0.01%, 0.025%, 0.1%

Liquid, topical: 0.05%

Triamcinolone (Aristocort, Kenacort, Azmacort, Nasacort)

Aerosol, oral, inhalation: 100 mcg/metered spray

Aerosol, topical: 0.2 mg/2 second spray

Cream, topical: 0.025%, 0.1%, 0.5%

Lotion, topical: 0.025%, 0.1%

Ointment, topical: 0.025%, 0.1%, 0.5%

Spray, intranasal: 55 mcg/actuation [100 sprays/canister]

Triamcinolone in Oral Adhesive Base (Kenalog in Orabase)

Paste: Triamcinolone 0.1%

Triamterene (Dyrenium)

Capsule: 50 mg, 100 mg

Triamterene/Hydrochlorothiazide (Dyazide, Maxzide)

Capsule (Dyazide): 37.5 mg Triamterene/25 mg Hydrochlorothiazide. 50 mg

Triamterene/25 mg Hydrochlorothiazide

Tablet (Maxzide): 37.5 mg Triamterene/25 mg Hydrochlorothiazide, 75 mg

Triamterene/50 mg Hydrochlorothiazide

Triazolam (Halcion) C-IV

Tablet: 0.125 mg, 0.25 mg

Trifluoperazine (Stelazine)

Concentrate, oral: 10 mg/mL

Injection: 2 mg/mL

Tablet: 1 mg, 2 mg, 5 mg, 10 mg

Trihexyphenidyl (Artane)

Capsule, sustained release: 5 mg

Elixir: 2 mg/5 mL

Tablet: 2 mg, 5 mg

Trimethobenzamide (Tigan)

Injection: 100 mg/mL

Suppository, rectal: 100 mg, 200 mg

Trimethoprim/Sulfamethoxazole (Co-Trimoxazole, Bactrim, Septra)

The 5:1 ratio of Sulfamethoxazole (SMX) to Trimethoprim (TMP) is constant in all dosage forms

Injection: 80 mg SMX/16 mg TMP per mL

Suspension, oral: 200 mg SMX/40 mg TMP per 5 mL

Tablet: 400 mg SMX/80 mg TMP, 800 mg SMX/160 mg TMP

Trimipramine (Surmontil)

Capsule: 25 mg, 50 mg, 100 mg

Triprolidine/Pseudoephedrine (Actifed)

Capsule, extended release: Triprolidine 5 mg/Pseudoephedrine 120 mg

Syrup: Triprolidine 1.25 mg/Pseudoephedrine 30 mg per 10 mL

Tablet: Triprolidine 2.5 mg/Pseudoephedrine 60 mg

Tropicamide (Mydriacyl)

Solution, ophthalmic: 0.5%, 1%

Trypsin/Balsam Peru/Castor Oil (Granulex)

Aerosol: Trypsin 0.1 mg/Balsam Peru 72.5 mg/Castor Oil 650 mg per 0.82 mL

Tuberculin, Purified Protein Derivative (P.P.D.)

Intermediate test strength: 5 TU/0.1 mL

Valproic Acid/Valproate (Depakene)

Capsule: 250 mg

Syrup: 250 mg/5 mL

Vancomycin (Vancocin)

Capsule: 125 mg, 250 mg

Powder for oral solution: 1 g, 10 g

Powder for injection: 500 mg, 1 g, 2 g, 5 g, 10 g

Varicella Virus Vaccine, Live (Varivax)

Injection, single dose

Venlafaxine (Effexor)

Capsule, sustained release: 37.5 mg, 75 mg, 150 mg

Tablet: 25 mg, 37.5 mg, 50 mg, 75 mg, 100 mg

Verapamil (Calan, Isoptin)

Capsule, sustained release: 120 mg, 180 mg, 240 mg, 360 mg
Injection: 2.5 mg/mL
Tablet: 40 mg, 80 mg, 120 mg
Tablet, sustained release: 120 mg, 180 mg, 240 mg

Vitamin A (Aquasol A)

Capsule: 10,000 units, 25,000 units, 50,000 units
Injection: 50,000 units/mL
Tablet: 5000 units

Vitamin A&D Ointment

Ointment, topical: 113 g

Vitamin B Complex/Vitamin C (Stresscaps, Allbee with C)

Capsule: each capsule contains a minimum of USDA requirements
Tablet: each tablet contains a minimum of USDA requirements

Vitamin B Complex/Vitamin C/Zinc

Tablet: each tablet contains a minimum of USDA requirements

Vitamin D (Ergocalciferol, Calciferol, Drisdol)

Drops, oral: 200 IU/drop

Vitamin E (Aquasol E)

Capsule: 100 units, 200 units, 400 units, 1000 units
Tablet: 200 units, 400 units

Warfarin (Coumadin)

Tablet: 1 mg, 2 mg, 2.5 mg, 3 mg, 4 mg, 5 mg, 7.5 mg, 10 mg

Water for Injection

Infusion: 250 mL, 500 mL, 1000 mL, 2000 mL, 3000 mL

Water for Irrigation

Solution, irrigation

Zafirlukast (Accolate)

Tablet: 20 mg

Zaleplon (Sonata)

Capsule: 5 mg, 10 mg

Zidovudine (AZT, Retrovir)

Capsule: 100 mg
Injection: 10 mg/mL
Syrup: 50 mg/5 mL
Tablet: 300 mg

Zinc Oxide

Ointment, topical: 20% in white ointment
Paste, topical: 25% in white petrolatum

Ziprasidone (Geodon)

Capsule: 20 mg, 40 mg, 60 mg, 80 mg
Injection: 20 mg

Zinc Oxide/Petrolatum/Imidazolidinyl Urea (Diaperene)

Ointment, topical

Zinc Sulfate

Capsule: 220 mg [50 mg zinc]
Injection: 1 mg/mL, 5 mg/mL

Zinc Undecylenate (Desenex)

Cream, topical: 20%
Foam, topical: 10% [with 35.2% alcohol]
Ointment, topical: 30 gm
Powder, topical: 19%

Zolpidem (Ambien) C-IV

Tablet: 5 mg, 10 mg

Zonisamide (Zonegran)

Capsule: 100 mg

THERAPEUTIC CLASSIFICATION/COST INDEX

ANTIDIABETIC AGENTS

INSULINS, HUMAN

Insulin, Combination (70/30)	\$\$\$\$
Insulin, Glargine (Lantus)	\$\$\$\$\$\$\$
Insulin, Lente	\$\$\$\$
Insulin, Lispro (Humalog)	\$\$\$\$\$\$\$
Insulin, Lispro/Insulin, Lispro Protamine (Humalog Mix 75/25)	\$\$\$\$\$\$\$
Insulin, NPH	\$\$\$\$
Insulin, Regular	\$\$\$\$
Insulin, Ultralente	\$\$\$\$\$

SULFONYLUREAS

chlorproPAMIDE (Diabinese)	\$
glipiZIDE (Glucotrol)	\$
glyBURIDE (Micronase, DiaBeta)	\$\$
TOLBUTamide (Orinase)	\$\$

MISCELLANEOUS ANTIDIABETICS

Metformin (Glucophage, Glucophage XR)	\$\$
Repaglinide (Prandin)	\$\$ - \$\$\$
Rosiglitazone (Avandia)	\$\$-\$\$\$

GLUCOSE ELEVATING AGENTS

Dextrose 50% in Water	\$\$
Glucagon	\$\$\$\$\$\$\$

ANTIDOTES/DETERRENTS/POISON CONTROL AGENTS

Acetylcysteine (Mucomyst)	
Activated Charcoal	
Deferoxamine (Desferal)	
Dimercaprol (B.A.L.)	
Disulfiram (Antabuse)	
Glucagon	
Ipecac Syrup	
Leucovorin (Wellcovorin)	\$\$ - \$\$
Naloxone (Narcan)	\$\$\$
Naltrexone (Trexan, ReVia)	\$\$ - \$\$\$\$\$
Nicotine Polacrilex (Nicorette)	\$\$\$
Nicotine Transdermal Patch (Nicoderm, Habitrol, ProStep, Nicotrol)	\$\$\$
Penicillamine (Cuprimine)	\$\$ - \$\$\$
Physostigmine (Antilirium)	\$
Phytonadione (Vitamin K ₁ , Mephyton)	\$\$ - \$\$\$
Protamine	\$

ANTIHISTAMINES

Brompheniramine/Pseudoephedrine (Bromfed)	\$\$
Chlorpheniramine (Chlor-Trimeton, Teldrin)	\$\$ - \$
Cyproheptadine (Periactin)	\$
diphenhydRAME (Benadryl)	\$
Fexofenadine (Allegra)	\$\$
Fexofenadine/Pseudoephedrine (Allegra-D)	\$\$
hydrOXYzine (Atarax)	\$\$ - \$\$
Loratadine (Claritin)	\$\$
Promethazine (Phenergan)	\$\$
Triprolidine/Pseudoephedrine (Actifed)	\$\$

ANTINEOPLASTIC AGENTS

Methotrexate
Tamoxifen (Nolvadex)

BLOOD MODIFYING AGENTS

ANTIPLATELET

Aspirin	\$
Clopidogrel (Plavix) – RESERVE USE	\$\$\$

ANTICOAGULANT

Heparin	\$\$
Enoxaparin (Lovenox)	\$\$\$\$\$\$\$
Warfarin (Coumadin)	\$

ANTICOAGULATION ANTAGONIST

Phytonadione (Vitamin K ₁ , Mephyton)	\$\$ - \$\$\$
Protamine	\$

MISCELLANEOUS BLOOD MODIFYING AGENTS

Ferrous Fumarate/Docusate Sodium (Ferro-Sequels) [33% elemental iron]	\$\$
Ferrous Sulfate (Feosol, Fer-In-Sol) [20% elemental iron]	\$
Iron Dextran Complex (Imferon)	\$\$\$\$\$\$\$

CARDIOVASCULAR AGENTS

DIURETICS

Thiazides & Related Diuretics	
Chlorthalidone (Hygroton)	\$
Hydrochlorothiazide (HydroDIURIL, Esidrix)	\$
Loop Diuretics	
Furosemide (Lasix)	\$
Potassium-Sparing Diuretics	
Spironolactone (Aldactone)	\$
Triamterene (Dyrenium)	\$ - \$
Carbonic Anhydrase Inhibitors	
acetaZOLAMIDE (Diamox)	\$
Combination Diuretics	
Spironolactone/Hydrochlorothiazide (Aldactazide)	\$ - \$\$
Triamterene/Hydrochlorothiazide (Dyazide, Maxzide)	\$ - \$

CARDIAC GLYCOSIDES

Digoxin (Lanoxin)	\$
-------------------	----

ANTIANGINALS

Isosorbide Dinitrate (Isordil, Sorbitrate)	\$
Isosorbide Mononitrate (Imdur, ISMO, Monoket)	\$\$
Nitroglycerin	\$\$ - \$

ANTIARRHYTHMICS

Adenosine (Adenocard)	\$\$\$\$\$\$\$
Procainamide (Pronestyl)	\$\$ - \$
Quinidine Gluconate	\$\$
Quinidine Sulfate	\$\$ - \$\$\$

CALCIUM CHANNEL BLOCKERS

Amlodipine (Norvasc)	\$ - \$\$
Diltiazem (Cardizem)	\$ - \$\$
Felodipine (Plendil)	\$ - \$\$
NIFEdipine (Procardia)	\$\$ - \$\$\$
Verapamil (Calan, Isoptin)	\$ - \$\$

BETA-ADRENERGIC BLOCKERS

Atenolol (Tenormin)	\$
Labetalol (Normodyne)	\$\$ - \$\$\$
Metoprolol (Lopressor)	\$\$ - \$\$
Nadolol (Corgard)	\$\$ - \$\$\$
Propranolol (Inderal)	\$\$ - \$\$\$

ANTIHYPERLIPIDEMICS

Atorvastatin (Lipitor)	\$\$
Cholestyramine (Questran)	\$ - \$\$\$
Fluvastatin (Lescol)	\$\$
Gemfibrozil (Lopid)	\$
Niacin/Nicotinamide (Nicobid)	\$\$ - \$
Simvastatin (Zocor)	\$ - \$\$

ANGIOTENSIN CONVERTING ENZYME INHIBITORS

Benazepril (Lotensin)	\$\$
Captopril (Capoten)	\$
Enalapril (Vasotec)	\$\$
Lisinopril (Prinivil, Zestril)	\$

VASOPRESSORS

DOPamine (Intropin)	\$\$\$\$ -\$\$\$\$\$
Epinephrine (Adrenalin)	\$ -\$\$\$\$\$\$\$
Norepinephrine or Levarterenol (Levophed)	\$\$\$ -\$\$\$\$\$\$\$\$

MISCELLANEOUS ANTIHYPERTENSIVES

Clonidine (Catapres)	\$ -\$\$\$\$\$\$\$
Guanethidine (Ismelin)	\$\$ - \$\$
hydrALAZINE (Apresoline)	\$
Methyldopa (Aldomet)	\$ -\$\$\$\$\$\$\$\$
Prazosin (Minipress)	\$ - \$\$

MISCELLANEOUS CARDIOVASCULARS

Sodium Polystyrene Sulfonate (Kayexalate)	\$\$ - \$\$\$
---	---------------

CENTRAL NERVOUS SYSTEM AGENTS

ANALGESICS

Nonsteroidal Anti-Inflammatory Agents

Aspirin	\$ - \$
Ibuprofen (Motrin)	\$ - \$\$
Ketorolac (Toradol)	\$\$\$\$ -\$\$\$\$\$\$\$\$
Nabumetone (Relafen) - RESERVE USE	\$\$
Naproxen (Naprosyn)	\$\$
Sulindac (Clinoril)	\$ - \$\$

Opiate Agonists

Fentanyl (Duragesic)	\$\$\$\$-\$\$\$\$\$
Methadone (Dolophine) C-II	\$ - \$\$\$
Morphine C-II	\$ - \$\$\$
Oxycodone (OxyContin)	\$\$ -\$\$\$\$\$

Miscellaneous Analgesics & Antipyretics

Acetaminophen (Tylenol)	\$ - \$\$
Acetaminophen/Codeine C-III	\$ - \$
Acetaminophen/Hydrocodone (Lortab, Vicodin)	\$\$ - \$\$
Tramadol (Ultram)	\$\$ - \$\$\$

ANTIEMETIC/ANTIVERTIGO AGENTS

diphenhydramine (Benadryl)	\$
hydrOXYzine (Atarax)	\$ - \$\$
Meclizine (Antivert, Bonine)	\$
Metoclopramide (Reglan)	\$ - \$\$\$
Prochlorperazine (Compazine)	\$ - \$\$\$
Promethazine (Phenergan)	\$ -\$\$\$\$
Trimethobenzamide (Tigan)	\$\$

SEE "PSYCHOTROPIC DOSAGE GUIDELINES", PAGE 11

PSYCHOTROPIC AGENTS

Benzodiazepines

Alprazolam (Xanax) C-IV	\$ - \$
Chlordiazepoxide (Librium) <u>oral only</u> C-IV	\$
Clonazepam (Klonopin)	\$\$ - \$\$\$\$
Clorazepate (Tranxene) C-IV	\$ - \$\$
Diazepam (Valium) C-IV	\$
Lorazepam (Ativan) C-IV	\$ - \$\$\$
Oxazepam (Serax) C-IV	\$ - \$\$
Temazepam (Restoril) C-IV	\$
Triazolam (Halcion) C-IV	\$ - \$

Antidepressants

Monoamine Oxidase Inhibitors

Phenelzine (Nardil)	\$\$ - \$
Tranylcypromine (Parnate)	\$\$

Serotonin Selective Reuptake Inhibitors (SSRIs)

Citalopram (Celexa)	\$\$
Escitalopram (Lexapro)	\$\$
Fluoxetine (Prozac)	\$\$ - \$\$\$\$
Fluvoxamine (Luvox)	\$\$ - \$\$\$
Paroxetine (Paxil)	\$\$ - \$\$\$\$
Sertraline (Zoloft)	\$ - \$\$\$

Serotonin Norepinephrine Reuptake Inhibitors (SNRIs)

Venlafaxine (Effexor)	\$ - \$\$\$\$
-----------------------	---------------

Tricyclic Agents

Amitriptyline (Elavil)	\$\$ - \$
clomiPRAMINE (Anafranil) - for OCD	\$ - \$\$\$
Desipramine (Norpramin, Pertofrane)	\$ - \$
Doxepin (Sinequan, Adapin)	\$ - \$\$
Imipramine (Tofranil)	\$
Nortriptyline (Aventyl, Pamelor)	\$ - \$\$
Protriptyline (Vivactil)	\$ - \$\$
Trimipramine (Surmontil)	\$ - \$\$\$

SEE "PSYCHOTROPIC DOSAGE GUIDELINES", PAGE 11

Miscellaneous Agents	
Amoxapine (Asendin)	\$ - \$\$
buPROPION (Wellbutrin, Wellbutrin SR, Wellbutrin XL)	\$\$ - \$\$\$
Maprotiline (Ludiomil)	\$ - \$
Mirtazapine (Remeron)	\$\$ - \$\$\$
Nefazodone (Serzone)	\$ - \$\$
Trazodone (Desyrel)	\$ - \$
Antipsychotics	
Aripiprazole (Abilify)	\$\$\$\$ -\$\$\$\$\$
chlorproMAZINE (Thorazine)	\$
Clozapine (Clozaril) - RESERVE USE	\$\$ -\$\$\$\$\$\$\$
Fluphenazine (Prolixin, Permitil)	\$ - \$\$\$
Fluphenazine Decanoate (Prolixin Decanoate)	\$\$ - \$\$\$
Haloperidol (Haldol)	\$
Haloperidol Decanoate (Haldol Decanoate)	\$\$ - \$\$\$
Loxapine (Loxitane)	\$ - \$\$\$
Mesoridazine (Serpentil) - RESERVE USE	\$ - \$\$\$
Molindone (Moban)	\$ - \$\$\$
Olanzapine (Zyprexa, Zydis)	\$\$\$ -\$\$\$\$\$
Perphenazine (Trilafon)	\$\$ - \$\$
Quetiapine (Seroquel) - RESERVE USE	\$\$ -\$\$\$\$\$
Risperidone (Risperdal, Risperdal Consta)	\$\$ -\$\$\$\$\$
Thioridazine (Mellaril) - RESERVE USE	\$ - \$
Thiothixene (Navane)	\$ - \$
Trifluoperazine (Stelazine)	\$\$ - \$
Ziprasidone (Geodon)	\$\$\$\$
Mood Stabilizers	
Carbamazepine (Tegretol)	\$ - \$\$\$
Divalproex (Depakote)	\$\$ - \$\$\$
Lithium Carbonate (Eskalith, Lithonate, Lithobid)	\$
Lithium Citrate	\$\$ - \$\$
Valproate (Depakene)	\$\$ - \$\$
Stimulants	
Amphetamine Mixture (Adderall, Adderall XR CII)	\$\$ - \$\$
Atomoxetine (Strattera)	\$\$ -\$\$\$\$\$
Dextroamphetamine (Dexedrine) C-II (<u>Rarely used in adults</u>)	\$ - \$\$
Methylphenidate (Ritalin, Concerta) C-II <u>(Rarely used in adults)</u>	\$ - \$\$

SEE "PSYCHOTROPIC DOSAGE GUIDELINES", PAGE 11

Miscellaneous Psychotropic Agents

busPIRone (BuSpar)	\$\$\$ - \$\$\$\$\$
hydrOXYzine (Atarax)	\$
Naloxone (Narcan)	\$ - \$\$
Naltrexone (Trexan, ReVia)	\$\$\$\$ - \$\$\$\$\$\$

SEDATIVES AND HYPNOTICS

Barbiturates

Amobarbital (Amytal) C-II - RESERVE USE	\$\$\$\$
---	----------

Benzodiazepines

Alprazolam (Xanax)	\$ - \$
Clonazepam (Klonopin)	\$\$ - \$\$\$\$\$
Diazepam (Valium)	\$
Lorazepam (Ativan)	\$ - \$\$\$
Midazolam (Versed)	\$\$\$ - \$\$\$\$\$\$
Oxazepam (Serax)	\$
Temazepam (Restoril)	\$
Triazolam (Halcion)	\$ - \$

Miscellaneous Sedative and Hypnotics

Chloral Hydrate (Noctec) C-IV	\$ - \$\$
diphenhydRamine (Benadryl)	\$
Midazolam (Versed)	\$\$\$ - \$\$\$\$\$\$
Trazodone (Desyrel)	\$ - \$
Zaleplon (Sonata)	\$\$
Zolpidem (Ambien) C-IV	\$\$

ANTICONVULSANTS

Barbiturates

Mephobarbital (Mebaral) C-IV	\$ - \$\$
Phenobarbital (Luminal) C-IV	\$ - \$\$
Primidone (Mysoline)	\$\$ - \$\$

Benzodiazepines

Clonazepam (Klonopin) C-IV	\$\$ - \$\$
Clorazepate (Tranxene, Tranxene SD) C-IV	\$ - \$\$
Diazepam (Valium, Diastat) C-IV	\$ -\$\$\$\$\$
Lorazepam (Ativan) C-IV	\$ - \$\$\$

Hydantoins

Fosphenytoin (Cerebyx)	\$\$\$\$\$\$\$
Phenytoin (Dilantin)	\$\$ - \$\$

Succinimides

Ethosuximide (Zarontin)	\$ - \$\$\$
-------------------------	-------------

Miscellaneous Anticonvulsants

Carbamazepine (Tegretol)	\$\$ - \$\$\$\$
Divalproex (Depakote)	\$\$ - \$\$\$\$
Felbamate (Felbatol) - RESERVE USE	\$\$ - \$\$\$\$
Gabapentin (Neurontin)	\$\$ - \$\$\$
Lamotrigine (Lamictal)	\$\$ - \$\$\$
Levetiracetam (Keppra)	\$\$-\$\$\$\$
Oxcarbazepine (Trileptal)	\$\$-\$\$\$\$
Tiagabine (Gabatril)	\$\$\$ - \$\$\$\$
Topiramate (Topamax)	\$\$\$ - \$\$\$\$
Valproate (Depakene)	\$ - \$\$\$
Zonisamide (Zonegran)	\$\$ - \$\$\$\$

MUSCLE RELAXANTS

Baclofen (Lioresal)	\$\$ - \$
Dantrolene (Dantrium)	\$\$ - \$\$\$\$
Diazepam (Valium) C-IV	\$ - \$\$
Methocarbamol (Robaxin)	\$ - \$\$\$
Tizanidine (Zanaflex) - RESERVE USE	\$\$

ANTIPARKINSON AGENTS

Amantadine (Symmetrel)	\$\$
Benztropine (Cogentin)	\$ - \$\$\$\$
Biperiden (Akineton)	\$
Bromocriptine (Parlodel)	\$\$ -\$\$\$\$\$
Levodopa (Larodopa)	\$ - \$\$
Levodopa/Carbidopa (Sinemet)	\$\$ - \$\$\$
Trihexyphenidyl (Artane)	\$ - \$\$

AGENTS FOR MIGRAINE

Atenolol (Tenormin)	\$
Divalproex (Depakote, Depakote ER- RESERVE USE)	\$\$ - \$\$\$
Metoprolol (Lopressor)	\$ - \$
Nadolol (Corgard)	\$ - \$\$\$
Naproxen (Naprosyn)	\$\$
Propranolol (Inderal)	\$ - \$\$\$
Sumatriptan (Imitrex)	\$\$\$\$ -\$\$\$\$\$
Valproate (Depakene)	\$ - \$\$\$
Verapamil (Calan, Isoptin)	\$ - \$\$

MISCELLANEOUS CENTRAL NERVOUS SYSTEM AGENTS

Donepezil (Aricept) - RESERVE USE	\$\$\$
Galantamine (Reminyl)	\$\$\$
Rivastigmine (Exelon) - RESERVE USE	\$\$\$

ENDOCRINE AGENTS

ESTROGENS

Estradiol (Estrace, Vivelle, Alora, Climara, Estraderm)	\$ -\$\$\$\$\$
Estrogens, Conjugated (Premarin)	\$\$

PROGESTERONES

medroxyPROGESTERone (Provera)	\$ - \$\$
-------------------------------	-----------

COMBINATION PRODUCTS

Estrogen/medroxyPROGESTERone (PremPro)	\$\$
Ethinyl Estradiol/Norethindrone (Loestrin, Ortho-Novum 777)	\$\$\$\$\$\$
Ethinyl Estradiol/Norgestrel (Ovral, Lo-Ovral)	\$\$\$\$\$\$
Levonorgestrel/Ethinyl Estradiol (Tri-Levlen, Triphasil)	\$\$\$\$\$\$
Norgestimate/Ethinyl Estradiol (Ortho Tri-Cyclen)	\$\$\$\$\$\$

ANDROGENS

methylTESTOSTERone (Android, Oretan) C-IV	\$\$ - \$\$\$
Testosterone (Androlan) C-IV	\$\$ - \$\$

ADRENAL CORTICAL STEROIDS

Cortisone	\$ - \$\$
Dexamethasone (Decadron)	\$ - \$\$
Fludrocortisone (Florinef)	\$ - \$\$
Hydrocortisone	\$\$ - \$\$\$\$\$
Methylprednisolone (Medrol)	\$ - \$\$\$\$
prednisolONE (Delta-Cortef)	\$\$ - \$\$
prednisONE (Meticorten, Deltasone)	\$ - \$\$\$\$
Triamcinolone (Aristocort, Kenacort)	\$ - \$\$\$\$\$

THYROID AGENTS

Levothyroxine (Synthroid)	\$
Liotrix (Thyrolar, Euthroid)	\$\$
Methimazole (Tapazole)	\$\$ - \$\$
Propylthiouracil	\$
Thyroid, Desiccated (Thyroid)	\$

MISCELLANEOUS ENDOCRINE AGENTS

Alendronate (Fosamax)	\$\$
Allopurinol (Zyloprim)	\$
Calcitonin-Salmon (Miacalcin)	\$\$\$
Colchicine	\$
Corticotropin (ACTH)	\$\$\$\$
Desmopressin (DDAVP, Stimate)	\$\$\$ - \$\$\$\$
Probenecid (Benemid)	\$
Raloxifene (Evista)	\$\$

GASTROINTESTINAL AGENTS

ANTACIDS

Aluminum Hydroxide (Amphojel)	\$ - \$\$
Aluminum Hydroxide/Magnesium Trisilicate (Gaviscon)	\$ - \$\$
Aluminum Hydroxide/Magnesium Hydroxide (Maalox)	\$ - \$
Aluminum Hydroxide/Magnesium Hydroxide/ Simethicone (Mylanta, Aludrox)	\$ - \$
Calcium Carbonate (Os-Cal, Titrалac) - 40% elemental calcium	\$

ANTISPASMODICS/ANTICHOLINERGIC AGENTS

Dicyclomine (Bentyl)	\$ - \$\$
Propantheline (Pro-Banthine)	\$ - \$\$\$

HISTAMINE (H₂) ANTAGONISTS

Preferred Agent:	
Famotidine (Pepcid)	\$ -\$\$\$\$
<i>Others:</i>	
Ranitidine (Zantac)	\$ -\$\$\$\$

PROTON PUMP INHIBITORS

Preferred Agent:	
Lansoprazole (Prevacid)	\$\$\$
Pantoprazole (Protonix) – price based on market share	\$\$ - \$\$\$
<i>Others:</i>	
Omeprazole (Prilosec)	\$\$\$

ANTIFLATULENTS

Simethicone (Mylicon)	\$ - \$\$
-----------------------	-----------

STIMULANTS

Metoclopramide (Reglan)	\$ - \$\$
-------------------------	-----------

LAXATIVES

Saline Laxatives

Magnesium Citrate	\$
Magnesium Hydroxide (Milk of Magnesia)	\$\$
Magnesium Sulfate (Epsom Salt)	\$\$
Sodium Phosphate/Biphosphate (Fleet)	\$\$ - \$\$
Phospho-Soda, Fleet's Enema)	

Irritant/Stimulant Laxatives

Bisacodyl (Dulcolax)	\$
Cascara Sagrada (Cascara Aromatic)	\$
Senna (Senokot)	\$ - \$

Bulk Laxatives

Cellulose (Unifiber)	\$
Methylcellulose (Citrucel)	\$\$
Polycarbophil (Fibercon, Fiber-Lax)	\$ -\$\$\$\$\$
Psyllium (Metamucil)	\$

Osmotic Laxatives

Lactulose (Cephulac)	\$ - \$\$
Polyethylene Glycol (MiraLax)	\$ - \$\$
Propylethylene Glycol Electrolyte Solution (PEG, Co-Lyte, GoLYTELY, OCL)	\$
Sorbitol	\$ - \$

Combination Laxatives

Docusate Sodium/Casanthrol (Peri-Colace)	\$
Docusate Sodium/Sennosides (Peri-Colace)	\$

Surfactants

Docusate Calcium (Surfak)	\$
Docusate Sodium (Colace, Doxinate)	\$

Miscellaneous Laxatives

Glycerin Suppository (Sani-Supp)	\$
----------------------------------	----

ANTIDIARRHEALS

Attapulgite	\$\$ - \$\$
Bismuth Subsalicylate (Pepto-Bismol, Bismatrol, Kaopectate)	\$ - \$
Kaolin-Pectin	\$\$ - \$\$
Lactobacillus Acidophilus (Lactinex, Bacid)	\$\$ - \$\$
Loperamide (Imodium)	\$ - \$\$

RECTAL AGENTS

Pramoxine (Tronothane)	\$ - \$\$
Rectal Hemorrhoidal Cream with Hydrocortisone	\$\$\$
Rectal Hemorrhoidal Ointment (Anusol)	\$\$
Rectal Hemorrhoidal Suppositories (Wyanoids, Anusol)	\$
Rectal Hemorrhoidal Suppositories with Hydrocortisone (Anusol-HC)	\$ - \$\$

MISCELLANEOUS GASTROINTESTINAL AGENTS

Activated Charcoal	\$ - \$
Bismuth Subsalicylate (Pepto-Bismol, Bismatrol)	\$\$\$ - \$\$\$\$\$
Ipecac Syrup	\$
Meclizine (Antivert, Bonine)	\$
Mesalamine (Asacol, Pentasa, Rowasa)	\$\$\$\$ - \$\$\$\$\$
Misoprostol (Cytotec)	\$\$
Pancrelipase (Pancrease, Creon)	\$\$ - \$\$\$\$
Prochlorperazine (Compazine)	\$ - \$\$\$
Sucralfate (Carafate)	\$ - \$\$\$\$
Sulfasalazine (Azulfidine)	\$\$
Trimethobenzamide (Tigan)	\$\$

GENITOURINARY AGENTS

INTERSTITIAL CYSTITIS AGENTS

Phenazopyridine (Pyridium)	\$ - \$\$
----------------------------	-----------

GENITOURINARY IRRIGANTS

Sodium Chloride	\$\$\$
Water for Irrigation	\$ - \$\$

URINARY ALKALINIZERS

Potassium Citrate (Urocit K)	\$ - \$\$\$\$\$
Potassium Citrate Combinations (Polycitra, Polycitra-LC, Polycitra K, Citrolith)	\$\$ - \$\$
Sodium Bicarbonate	\$ - \$\$
Sodium Citrate/Citric Acid (Bicitra, Oracit)	\$\$ - \$\$\$

URINARY ANTICHOLINERGICS

Flavoxate (Urispas)	\$\$\$
Oxybutynin (Ditropan, Ditropan XL)	\$\$ - \$\$
Tolterodine (Detrol, Detrol LA)	\$\$ - \$\$\$

URINARY CHOLINERGICS

Bethanechol (Urecholine)	\$ - \$\$\$
--------------------------	-------------

VAGINAL ANTIFUNGALS

Clotrimazole (Gyne-Lotrimin, Mycelex)	\$\$\$\$ - \$\$\$\$\$\$
Miconazole (Monistat)	\$\$\$\$ - \$\$\$\$\$\$
Tioconazole (Vagistat-1)	\$\$\$\$

MISCELLANEOUS GENITOURINARY AGENTS

Estradiol (Estrace)	\$\$ - \$\$\$\$\$\$\$\$
Estrogens, Conjugated (Premarin)	\$\$\$\$\$\$\$
Nitrofurantoin (Macrodantin)	\$\$ - \$\$\$
Sevelamer (Renagel)	\$\$\$ - \$\$\$

IMMUNOLOGICAL AGENTS

IMMUNE SERUMS

Hepatitis B Immune Globulin (HBIG)	\$\$\$\$\$\$\$
------------------------------------	----------------

BACTERIAL VACCINES

Pneumococcal Vaccine, Polyvalent (Pneumovax)	\$\$\$\$\$
--	------------

VIRAL VACCINES

Hepatitis A Vaccine (Vaqta)	\$\$\$\$\$\$
Hepatitis B Virus Vaccine, Recombinant (Recombivax HB, Engerix-B)	\$\$\$\$\$\$
Influenza Virus Vaccine (Fluzone, Fluviron)	\$\$
Measles, Mumps and Rubella Virus Vaccine, Live (MMR II)	\$\$\$\$\$\$
Poliovirus Vaccine, Inactivated (IPOL)	\$\$\$\$\$
Rubella Virus Vaccine Live (Meruvax II)	\$\$\$\$\$
Varicella Virus Vaccine, Live (Varivax)	\$\$\$\$\$\$\$

TOXOIDS

Diphtheria & Tetanus Toxoids Adsorbed (DT)	\$\$\$\$
Diphtheria & Tetanus Toxoids Adsorbed for Adult Use (Td)	\$\$\$

IN-VIVO DIAGNOSTIC BIOLOGICALS

Tuberculin, Purified Protein Derivative (P.P.D.)	\$\$ - \$\$\$\$\$\$
--	---------------------

INFECTIOUS DISEASE AGENTS

ANTIBIOTICS

Penicillins	
Amoxicillin (Amoxil, Polymox)	\$ - \$\$\$
Amoxicillin/Clavulanate (Augmentin)	\$\$\$ - \$\$\$\$\$
Ampicillin (Polycillin, Omnipen)	\$ - \$\$\$
Cloxacillin (Cloxapen, Tegopen) or Dicloxacillin (Dycill, Dynapen, Pathocil)	\$ - \$\$\$\$\$\$
Nafcillin (Unipen) or Oxacillin (Prostaphlin)	\$ - \$\$\$\$\$
Penicillin G Benzathine (Bicillin)	\$\$\$ - \$\$\$\$\$\$
Penicillin G Benzathine/Penicillin G Procaine (Bicillin C-R)	\$\$\$\$
Penicillin G Procaine (Wycillin)	\$\$ - \$\$\$\$\$\$
Penicillin G Sodium	\$\$ - \$\$\$
Penicillin V Potassium (Pen-Vee K, V-Cillin K)	\$ - \$\$
Ticarcillin (Ticar)	\$\$\$\$\$ - \$\$\$\$\$\$
Ticarcillin/Clavulanate (Timentin)	\$\$\$\$\$\$\$

Cephalosporins		
Cefazolin (Kefzol, Ancef)	\$\$\$\$ -\$\$\$\$\$	
Cefoperazone (Cefobid)	\$\$\$\$\$\$\$	
Ceftriaxone (Rocephin)	\$\$\$\$\$\$\$	
Cefuroxime Axetil (Ceftin) - Oral form only - RESERVE USE	\$\$\$ -\$\$\$\$\$	
Cephalexin (Keflex)	\$ - \$	
Macrolides		
Azithromycin (Zithromax) - RESERVE USE	\$\$\$\$\$\$\$	
Clarithromycin (Biaxin) - RESERVE USE	\$\$\$ - \$\$\$\$	
Erythromycin (Erythrocin)	\$ - \$	
Erythromycin Ethylsuccinate/Sulfisoxazole Suspension (Pediazole)	\$ - \$\$	
Tetracyclines		
Doxycycline (Vibramycin)	\$ - \$\$\$\$\$\$	
Tetracycline (Achromycin, Panmycin)	\$ - \$\$	
Quinolones		
Ciprofloxacin (Cipro)	\$\$\$\$ -\$\$\$\$\$\$\$	
Levofloxacin (Levaquin)	\$\$\$\$ -\$\$\$\$\$\$\$	
Aminoglycosides		
Gentamicin (Garamycin)	\$\$	
Neomycin (Mycifradin)	\$\$ - \$\$\$	
Tobramycin (Nebcin)	\$\$\$ - \$\$\$\$	
Miscellaneous Antibiotics		
Clindamycin (Cleocin)	\$ - \$\$\$\$\$\$\$\$	
Metronidazole (Flagyl)	\$ - \$\$\$\$\$\$\$\$	
Trimethoprim/Sulfamethoxazole (Co- Trimoxazole, Bactrim, Septra)	\$ - \$\$\$\$\$\$\$\$	
Vancomycin (Vancocin)	\$\$\$\$ -\$\$\$\$\$\$\$	

ANTIFUNGALS

Fluconazole (Diflucan)	\$\$\$\$ -\$\$\$\$\$
Griseofulvin (Fulvicin)	\$
Ketoconazole (Nizoral)	\$\$ - \$\$\$
Nystatin (Mycostatin)	\$\$ - \$\$

ANTIMALARIALS

Chloroquine (Aralen)	\$\$\$
----------------------	--------

ANTITUBERCULARS

Ethambutol (Myambutol)	\$\$ - \$\$\$\$\$
Ethionamide	\$\$\$
Isoniazid (INH)	\$
Pyrazinamide	\$ - \$\$\$
Rifampin (Rifadin)	\$\$ - \$\$\$\$\$\$\$\$\$
Rifampin/Isoniazid (Rifamate)	\$\$

ANTIVIRALS

Acyclovir (Zovirax)	\$\$ - \$\$\$\$\$
Amantadine (Symmetrel)	\$\$ - \$\$
Delavirdine (DLV, Rescriptor)	\$\$\$\$\$
Didanosine (ddl, Videx)	\$\$\$\$\$
Indinavir (Crixivan)	\$\$\$\$\$
Lamivudine (Epivir)	\$\$\$\$\$
Lamivudine/Zidovudine (Combivir)	\$\$\$\$\$\$
Nelfinavir (Viracept)	\$\$\$\$\$\$
Nevirapine (NVP, Viramune)	\$\$\$\$\$
Ritonavir (Norvir)	\$\$\$\$ - \$\$\$\$\$\$
Saquinavir (Invirase, Fortovase)	\$\$\$\$
Stavudine (d4T, Zerit)	\$\$\$\$
Zidovudine (AZT, Retrovir)	\$\$\$\$ - \$\$\$\$\$

ANTIHELMINTICS

Mebendazole (Vermox)	\$\$\$ - \$\$\$\$\$\$
Pyrantel (Antiminth)	\$ - \$\$\$\$\$
Thiabendazole (Mintezol)	\$\$\$\$

URINARY ANTI-INFECTIVES

Nitrofurantoin (Macrodantin)	\$\$ - \$\$\$\$
------------------------------	-----------------

MISCELLANEOUS ANTI-INFECTIVES

Pentamidine (Pentam) - RESERVE USE	\$\$\$\$\$\$
------------------------------------	--------------

INTRAVENOUS SOLUTIONS AND ADDITIVES

INTRAVENOUS SOLUTIONS

Amino Acid Injection (Aminosyn)	\$\$\$\$ -\$\$\$\$\$
Dextran (Centran, LMD, Macrodex, Rheomacrodex)	\$\$\$\$ -\$\$\$\$\$
Dextrose/Sodium Chloride Intravenous Solution	\$\$\$
Dextrose 5% in 0.2% Sodium Chloride	
Dextrose 5% in 0.45% Sodium Chloride	
Dextrose 5% in 0.9% Sodium Chloride	
Dextrose 5% in Water	\$ - \$\$
Dextrose 5% in Ringer's Lactate	\$
Dextrose 5% with Multiple Electrolytes (D5 E75, Baxter)	\$\$\$
Dextrose 5%/Sodium Chloride/Potassium Chloride Intravenous Solution	\$\$\$\$\$ -\$\$\$\$\$
Dextrose 5%/Sodium Chloride 0.2%/Potassium Chloride	
Dextrose 5%/Sodium Chloride 0.45%/Potassium Chloride	
Dextrose 5%/Sodium Chloride 0.9%/Potassium Chloride	
Dextrose 50% in Water	\$\$\$\$ -\$\$\$\$\$
Ringer's Lactate Solution (Hartmann's Solution)	\$\$
Sodium Chloride Intravenous Solution	\$\$\$\$ -\$\$\$\$\$
Sodium Chloride 0.2%	\$ - \$\$\$
Sodium Chloride 0.45%	
Sodium Chloride 0.9%	
Water for Injection	\$\$ - \$\$\$

ELECTROLYTE REPLACEMENT ADDITIVES

Calcium Gluconate	\$\$
Magnesium Sulfate	\$ - \$
Potassium Chloride	\$ - \$\$
Sodium Bicarbonate	\$ - \$\$
Sodium Chloride	\$ - \$\$
Sodium Lactate	\$\$\$\$
Zinc Sulfate	\$ - \$\$

NUTRITIONAL AGENTS

VITAMINS

Ascorbic Acid (Vitamin C)	\$ - \$
Cyanocobalamin (Vitamin B ₁₂)	\$ -\$\$\$\$\$
Folic Acid (Folvite)	\$ - \$\$\$
Leucovorin (Wellcovorin)	\$\$ - \$\$\$
Niacin/Nicotinamide (Nicobid)	\$\$
Phytonadione (Vitamin K ₁ , Mephyton, Konakion)	\$\$ - \$\$\$
Pyridoxine (Vitamin B ₆)	\$ - \$\$
Thiamine (Vitamin B ₁)	\$ - \$
Vitamin A (Aquasol A)	\$ -\$\$\$\$\$
Vitamin D (Ergocalciferol, Calciferol, Drisdol)	\$
Vitamin E (Aquasol E)	\$ -\$\$\$\$\$

MINERALS TRACE ELEMENTS AND ELECTROLYTES

Calcium Carbonate (Os-Cal, Titralac) - 40% elemental calcium	\$
Calcium Citrate (Citracal)	\$
Calcium Glubionate (Neo-Calglucon) - 6% elemental calcium	\$ - \$\$
Calcium Gluconate - 9% elemental calcium	\$\$
Ferrous Fumarate/Docusate Sodium (Ferro-Sequels) - 33% elemental iron	\$\$
Ferrous Sulfate (Feosol, Fer-In-Sol) -20% elemental iron	\$
Iron Dextran Complex (Imferon)	\$\$\$\$\$
Zinc Sulfate	\$ - \$\$

COMBINATION PRODUCTS

Calcium Carbonate/Vitamin D (Oscal + D)	\$
Multivitamin (Unicap, Hexavitamins)	\$ - \$
Multivitamin/Minerals	\$ - \$
Multivitamins, Pediatric (Poly-Vi-Sol)	\$\$
Multivitamins, Prenatal (Filibon)	\$
Vitamin B Complex/Vitamin C (Stresscaps, Allbee with C)	\$
Vitamin B Complex/Vitamin C/Zinc	\$

RESPIRATORY AGENTS

BRONCHODILATORS

Albuterol (Proventil, Ventolin)	\$ - \$\$\$
Aminophylline	\$\$ - \$\$
Metaproterenol (Alupent)	\$ -\$\$\$\$\$
Salmeterol (Serevent)	\$\$\$\$\$\$
Terbutaline (Brethine)	\$ -\$\$\$\$\$
Theophylline (Elixophyllin)	\$ - \$\$

DECONGESTANTS

Pseudoephedrine (Sudafed)	\$ - \$\$
---------------------------	-----------

STEROIDS

Bclomethasone (Vanceril, Beconase)	\$\$\$\$\$\$
Fluticasone (Flonase, Flovent)	\$\$\$\$\$\$
Mometasone (Nasonex)	\$\$\$\$\$\$
Triamcinolone (Azmacort, Nasacort)	\$\$\$\$\$\$

ANTITUSSIVES

Dextromethorphan	\$ - \$\$
------------------	-----------

EXPECTORANTS

Guaifenesin (Robitussin)	\$ - \$\$
Potassium Iodide (SSKI)	\$ - \$

COUGH AND COLD PREPARATIONS

Brompheniramine/Pseudoephedrine (Bromfed)	\$\$
Chlorpheniramine (Chlor-Trimeton, Teldrin)	\$ - \$\$
diphenhydRAMINE (Benadryl)	\$
Fexofenadine (Allegra)	\$\$
Fexofenadine/Pseudoephedrine (Allegra-D)	\$\$
Guaifenesin/Dextromethorphan (Robitussin DM)	\$ - \$\$
Guaifenesin/Pseudoephedrine (Entex PSE)	\$
Hydrocodone/Guaifenesin (Hycotuss, Kwelcof)	\$\$
Loratadine (Claritin)	\$\$
Triprolidine/Pseudoephedrine (Actifed)	\$

MISCELLANEOUS RESPIRATORY DRUGS

Acetylcysteine (Mucomyst)	\$\$\$\$ -\$\$\$\$\$
Cromolyn (Intal)	\$\$\$\$\$\$\$
Ipratropium (Atrovent Inhaler)	\$\$\$\$\$\$\$
Sodium Chloride	\$ - \$
Zafirlukast (Accolate)	\$\$

TOPICAL AGENTS

OPHTHALMICS

Agents for Glaucoma

Betaxolol (Betoptic S)	\$\$\$\$
Bimatoprost (Lumigan)	\$\$\$\$\$\$\$
Brimonidine (Alphagan)	\$\$\$\$\$\$\$
Latanoprost (Xalatan)	\$\$\$\$\$\$\$
Pilocarpine (Isopto Carpine)	\$ - \$\$\$\$\$\$
Timolol (Timoptic)	\$\$ - \$\$\$\$\$\$
Timolol/Dorzolamide (Cosopt)	\$\$\$\$\$\$\$
Travoprost (Travatan)	\$\$\$\$\$\$\$

Antibiotics

Bacitracin (Baciguent)	\$
Ciprofloxacin (Ciloxan)	\$\$\$\$\$ - \$\$\$\$\$\$
Erythromycin	\$\$
Gentamicin (Garamycin)	\$\$
Polymyxin B/Bacitracin (Polysporin)	\$\$\$\$\$
Polymyxin B/Trimethoprim (Polytrim)	\$\$\$
Sulfacetamide Sodium (Sulamyd)	\$ - \$\$\$\$\$\$
Tobramycin (Tobrex)	\$\$ - \$\$\$\$\$\$

Mydriatics

Atropine Sulfate (Isopto Atropine)	\$\$
Homatropine (Isopto Homatropine)	\$\$ - \$\$\$
Phenylephrine (Neo-Synephrine)	\$\$ - \$\$\$
Scopolamine (Isopto Hyoscine)	\$\$ - \$\$
Tropicamide (Mydriacyl)	\$\$ - \$\$\$

Lubricants

Ophthalmic Lubricant (HypoTears, HypoTears PF) [preservative-free, lanolin-free]	\$ - \$\$\$
--	-------------

Decongestant/Antiallergy		
Naphazoline (Naphcon, AK-Con)	\$\$	
Olopatadine (Patanol)	\$\$\$\$\$\$\$	
Tetrahydrozoline (Visine Allergy Relief, Visine Moisturizing)	\$\$\$	
Miscellaneous Ophthalmics		
Dexamethasone (Decadron) - RESERVE USE	\$\$	
Fluorescein Sodium	\$ - \$\$\$	
Proparacaine (Alcaine)	\$\$	
Sodium Chloride	\$\$ - \$\$\$\$	
Tobramycin/Dexamethasone (TobraDex)- RESERVE USE	\$\$\$\$\$\$ - \$\$\$\$\$\$	

OTICS

Acetic Acid (Acetasol, VoSol)	\$\$	
Acetic Acid/Aluminum Acetate (Domeboro Otic)	\$	
Acetic Acid/Hydrocortisone/Propylene Glycol/ Sodium Acetate/Benzethonium (VoSol HC)	\$\$	
Antipyrine/Benzocaine (Allergen, Auralgan)	\$ - \$\$	
Carbamide Peroxide/Glycerin/Propylene Glycol/ Sodium Stannate (Debrox)	\$	
Ciprofloxacin/Hydrocortisone (Cipro HC Otic)	\$\$\$\$\$\$	
Mineral Oil	\$\$	
Neomycin/Polymyxin B/Hydrocortisone (Cortisporin - ear drops only)	\$\$\$	

NASAL, MOUTH AND THROAT AGENTS

Benzocaine	\$\$ - \$\$\$
Carbamide Peroxide/Glycerin/Propylene Glycol/ Sodium Stannate (Gly-Oxide)	\$\$\$
Carboxymethylcellulose/Electrolytes (Saliva Substitute, Moi-Stir, Salivart, MouthKote, Salix)	\$ - \$\$\$
Cetylpyridinium (Cepacol)	\$ - \$\$\$
Chlorhexidine (Peridex)	\$\$
Clotrimazole (Mycelex, Fungoid)	\$\$
Doxycycline (Periostat)	\$\$\$
Nystatin (Mycostatin)	\$\$ - \$\$\$
Oxymetazoline (Afrin)	\$
Phenol (Chloraseptic)	\$\$
Phenylephrine (Neo-Synephrine)	\$\$
Sodium Fluoride	\$
Stannous Fluoride (OmniMed, PerioMed)	\$\$\$\$
Triamcinolone in Oral Adhesive Base (Kenalog in Orabase)	\$\$

DERMATOLOGICALS

Acne Agents

Adapalene (Differin)	\$\$\$\$\$\$\$
Benzoyl Peroxide	\$ - \$\$
Benzoyl Peroxide/Clindamycin (BenzaClin)	\$\$\$\$\$\$\$
Clindamycin (Cleocin T)	\$\$\$ - \$\$\$\$\$\$
Erythromycin/Benzoyl Peroxide (Benzamycin)	\$\$\$\$\$\$\$
Metronidazole (Noritate, MetroGel)	\$\$\$\$\$\$\$
Sulfur/Resorcinol (Sulforcin, Rezamide)	\$\$\$\$
Tretinoin Gel (Retin-A)	\$\$\$\$\$ - \$\$\$\$\$\$

Skin Cleansers

Abrasive Cleanser (Brasivol, Pernox)	\$\$\$\$
Non-Soap Cleanser (Cetaphil)	\$\$\$ - \$\$\$

Sunscreens

Cream/Lotion: contains a minimum SPF of 15 \$\$\$

Burn Agents

Bacitracin (Baciguent)	\$
Silver Sulfadiazine (Silvadene)	\$\$ - \$\$\$\$\$\$

Diaper Rash Agents

Diaper Rash Ointment (Desitin, Vitamin A&D Ointment, Diaperene)	\$ - \$\$
Diaper Rash Powder (Mexsana)	\$\$

Antiseborrheic Agents	
Selenium Sulfide (Selsun)	\$\$ - \$\$\$
Sulfacetamide Sodium (Sebizon)	\$\$\$\$\$\$\$
Antipsoriatics	
Calcipotriene (Dovonex)	\$\$\$\$\$\$\$
Methotrexate	\$\$\$\$
Selenium Sulfide (Selsun)	\$\$ - \$\$\$
Tazarotene (Tazorac, Avage)	\$\$\$\$\$\$\$
Anti-Histamine Agents	
Calamine/Zinc Oxide/Glycerin (Calamine Lotion)	\$
Calamine/Pramoxine (Caladryl)	\$\$\$
diphenhydrAMINE (Benadryl)	\$
Antiseptics & Germacides	
Benzalkonium Chloride (Zephran)	\$\$ - \$\$\$
Chlorhexidine (Hibiclens, Bactoshield)	\$\$ - \$\$\$
Hexachlorophene (pHisoHex)	\$\$\$ -\$\$\$\$\$
Povidone-Iodine (Betadine)	\$ -\$\$\$\$
Anti-Infectives	
Antibiotics	
Bacitracin (Baciguent)	\$
Bacitracin/Polymyxin B (Polysporin)	\$\$\$ - \$\$\$
Clindamycin (Cleocin T)	\$\$\$ -\$\$\$\$\$
Gentamicin (Garamycin)	\$\$\$
Metronidazole (Noritate, MetroGel)	\$\$\$\$\$\$\$
Mupirocin (Bactroban)	\$\$\$\$\$\$\$
Neomycin/Polymyxin B/Bacitracin (Triple Antibiotic Ointment)	\$
Polymyxin B/Neomycin (Neosporin)	\$\$ -\$\$\$\$\$
Antiviral	
Acyclovir (Zovirax)	\$\$\$\$\$\$\$

Antifungals	
Benzoic and Salicylic Acids (Whitfield's)	\$\$
Calcium Undecylenate (Caldesene)	\$\$
Clotrimazole (Lotrimin, Fungoid)	\$\$\$\$\$
Ketoconazole (Nizoral)	\$\$\$\$\$ -\$\$\$\$\$
Miconazole (Monistat)	\$\$\$ -\$\$\$\$
Nystatin (Mycostatin)	\$\$ -\$\$\$\$
Terbinafine (Lamisil)	\$\$\$\$\$
Tolnaftate (Tinactin)	\$\$ - \$\$\$
Zinc Undecylenate (Desenex)	\$\$\$ -\$\$\$\$
 Scabicides & Pediculicides	
Crotamiton (Eurax)	\$\$\$\$
Lindane (Gamma Benzene Hexachloride, Kwell)	\$\$\$\$
Permethrin 1% Liquid (NIX)	\$\$\$
Permethrin 5% Cream (Elimite)	\$\$\$\$
Pyrethrins/Piperonyl Butoxide (A-200, RID)	\$\$\$
 Corticosteroids	
Betamethasone Valerate (Valisone)	\$\$ - \$\$\$
Clobetasol (Temovate) - RESERVE USE	\$\$\$\$\$
Fluocinolone or Fluocinonide (Synalar, Lidex)	\$\$ - \$\$\$
Hydrocortisone Acetate (Lanacort, Corticaine)	\$ - \$\$\$
 Local Anesthetics	
Benzocaine (Lanacaine)	\$ - \$\$\$
Bupivacaine (Marcaine)	\$\$
Dibucaine (Nupercainal)	\$ - \$\$\$
Ethyl Chloride	\$\$\$\$
Lidocaine (Xylocaine)	\$\$ -\$\$\$\$\$
Pramoxine (Tronothane)	\$\$\$ -\$\$\$\$\$
 Emollients	
Emollient Lotion/Cream (Lubriderm, Allercreme, Keri Lotion, Cetaphil, Lac- Hydrin)	\$\$ - \$\$\$
Emollient Ointment (Lanolin)	\$ - \$\$\$
 Skin Protectants	
Benzoin, Compound Tincture	\$\$
Silver Sulfadiazine (Silvadene)	\$\$ -\$\$\$\$\$
Zinc Oxide	\$\$\$ -\$\$\$\$\$
 Ointment & Lotion Bases	
Petrolatum, White (Vaseline)	\$\$

Tar-Containing Agents		
Coal Tar (Ionil-T, Tegrin)		\$\$\$
Wet Dressings & Soaks		
Aluminum Acetate (Burow's Solution)		\$\$ - \$\$\$\$\$
Rubs and Liniments		
Methyl Salicylate (Ben-Gay)		\$
Keratolytics		
Podophyllum Resin		\$\$\$\$\$
Salicylic Acid		\$\$ - \$\$\$
Miscellaneous Dermatologicals		
Benzoic and Salicylic Acids (Whitfield's)		\$\$
Camphor-Phenol (Campho-Phenique)		\$\$
Collagenase (Santyl)		\$\$\$\$\$
Hydrogen Peroxide		\$
Silver Nitrate		\$
Trypsin/Balsam Peru/Castor Oil (Granulex)		\$\$\$\$
Zinc Oxide		\$\$\$ - \$\$\$\$\$\$

IRRIGATION SOLUTIONS

Sodium Chloride	\$\$\$
Water for Injection	\$\$ - \$\$\$

ALPHABETICAL INDEX

A-200	65, 105	Ambien	17, 77, 86
Ability	13, 27, 85	Amino Acid Injection.....	26, 98
Abrasive Cleanser	24, 104	Aminophylline	26, 100
Accolate	76, 101	Aminosyn.....	26, 98
Acetaminophen.....	24, 83	Amitriptyline	14, 26, 84
Acetaminophen/Codeine	24, 83	Amlodipine.....	26, 81
Acetaminophen/Hydrocodone	24, 83	Amobarbital	17, 18, 26, 86
Acetasol.....	24, 103	Amoxapine	14, 26, 85
acetaZOLAMIDE.....	24, 81	Amoxicillin	26, 95
Acetic Acid	24, 103	Amoxicillin/Clavulanate	27, 95
Acetic Acid/Aluminum Acetate.....	24, 103	Amoxil.....	26, 95
Acetic Acid/Hydrocortisone/Propylene Glycol/Sodium Acetate/Benzethonium	24, 103	Aphetamine Mixture	16, 27, 86
Acetylcysteine	25, 79, 101	Amphojel	25, 90
Achromycin	71, 95	Ampicillin	27, 95
ACTH	36, 90	Amytal.....	17, 18, 26, 86
Actifed	75, 79, 101	Anafranil	16, 35, 84
Activated Charcoal.....	25, 79, 93	Ancef	32, 95
Acyclovir.....	25, 97, 105	Android	54, 89
Adapalene	25, 104	Androlan	70, 89
Adapin.....	14, 40, 84	Antabuse	40, 79
Adderall.....	16, 27, 86	Antilirium.....	61, 79
Adderall XR.....	16, 27, 86	Antiminth	65, 97
Adenocard	25, 81	Antipyrine/Benzocaine	27, 103
Adenosine	25, 81	Antivert	52, 83, 93
Adrenalin.....	41, 82	Anusol	66, 92
Afrin	59, 103	Anusol-HC	66, 92
AK-Con	56, 102	Apresoline	47, 82
Akineton.....	30, 88	Aquasol A	76, 99
Albuterol.....	25, 100	Aquasol E	76, 99
Alcaine	64, 102	Aralen	33, 96
Aldactazide	69, 81	Aricept	19, 40, 88
Aldactone	69, 80	Aripiprazole.....	13, 27, 85
Aldomet.....	54, 82	Aristocort	74, 89
Alendronate	25, 90	Artane	74, 88
Allbee with C	76, 99	Asacol.....	53, 93
Allegra.....	43, 79, 101	Ascorbic Acid.....	27, 99
Allegra-D.....	43, 79, 101	Asendin	14, 26, 85
Allercreme	41, 106	Aspirin	27, 80, 83
Allergen.....	27, 103	Atarax	17, 47, 79, 83, 86
Allopurinol	25, 90	Atenolol.....	27, 81, 88
Alora	42, 88	Ativan.....	17, 52, 84, 86, 87
Alphagan.....	30, 101	Atomoxetine	27, 86
Alprazolam.....	17, 25, 84, 86	Atorvastatin	28, 82
Aludrox.....	26, 90	Atropine Sulfate.....	28, 102
Aluminum Acetate	25, 106	Atrovent	49, 101
Aluminum Hydroxide.....	25, 90	Attapulgite.....	28, 92
Aluminum Hydroxide/Magnesium Hydroxide.....	26, 90	Augmentin	27, 95
Aluminum Hydroxide/Magnesium Hydroxide/ Simethicone	26, 90	Auralgan	27, 103
Aluminum Hydroxide/Magnesium Trisilicate.....	26, 90	Avage	70, 104
Alupent.....	53, 100	Avandia.....	67, 78
Amantadine.....	26, 88, 97	Aventyl.....	14, 58, 84
		Azithromycin.....	18, 28, 95
		Azmacort	74, 100

AZT	77, 97	Calcipotriene	31, 104
Azulfidine	70, 93	Calcitonin-Salmon	31, 90
B.A.L.	39, 79	Calcium Carbonate.....	31, 90, 99
Bacid.....	49, 92	Calcium Carbonate/Vitamin D	31, 99
Baciguent.....	28, 102, 104, 105	Calcium Citrate.....	31, 99
Bacitracin.....	28, 102, 104, 105	Calcium Glubionate	31, 99
Bacitracin/Polymyxin B	28, 105	Calcium Gluconate	31, 98, 99
Baclofen.....	28, 87	Calcium Undecylenate	31, 105
Bactoshield	33, 105	Caldesene	31, 105
Bactrim.....	75, 96	Campho-Phenique	32, 107
Bactroban	56, 105	Camphor-Phenol	32, 107
Beclomethasone.....	28, 100	Capoten.....	32, 82
Beconase.....	28, 100	Captopril	32, 82
Benadryl.....	17, 39, 79, 83, 86, 101, 104	Carafate.....	69, 93
Benazepril.....	28, 82	Carbamazepine	16, 21, 32, 85, 87
Benemid.....	64, 90	Carbamide Peroxide/Glycerin/Propylene Glycol/Sodium	
Ben-Gay.....	54, 106	Stannate	32, 103
Bentyl.....	39, 90	Carbatrol.....	16, 21, 32
BenzaClin.....	29, 104	Carboxymethylcellulose/Electrolytes	32, 103
Benzalkonium Chloride.....	28, 105	Cardizem	39, 81
Benzamycin	42, 104	Cascara Aromatic.....	32, 91
Benzocaine	29, 103, 106	Cascara Sagrada	32, 91
Benzoic and Salicylic Acids	29, 105, 107	Catapres.....	16, 35, 82
Benzoin, Compound Tincture	29, 106	Cefazolin.....	32, 95
Benzoyl Peroxide	29, 104	Cefobid	32, 95
Benzoyl Peroxide/Clindamycin	29, 104	Cefoperazone	32, 95
Benztropine.....	29, 88	Ceftin	18, 32, 95
Betadine.....	63, 105	Ceftriaxone	32, 95
Betamethasone Valerate	29, 106	Cefuroxime Axetil	18, 32, 95
Betaxolol	30, 101	Celexa	14, 34, 84
Bethanechol.....	30, 93	Cellulose.....	33, 91
Betoptic S	30, 101	Cepacol	33, 103
Biaxin	18, 34, 95	Cephalexin	33, 95
Bicillin.....	60, 95	Cephulac	50, 92
Bicillin C-R	60, 95	Cerebyx	45, 87
Bicitra.....	68, 93	Cetaphil	41, 58, 104, 106
Bimatoprost.....	30, 101	Cetylpyridinium	33, 103
Biperiden.....	30, 88	Chloral Hydrate.....	17, 33, 86
Bisacodyl.....	30, 91	Chloraseptic	61, 103
Bismatrol.....	30, 92, 93	Chlordiazepoxide	17, 33, 84
Bismuth Subsalicylate.....	30, 92, 93	Chlorhexidine	33, 103, 105
Bonine.....	52, 83, 93	Chloroquine	33, 96
Brasivol	24, 104	Chlorpheniramine	33, 79, 101
Brethine	70, 100	chlorproMAZINE	13, 34, 85
Brimonidine	30, 101	chlorproPAMIDE.....	34, 78
Bromfed	30, 79, 101	Chlorthalidone	34, 80
Bromocriptine	30, 88	Chlor-Trimeton	33, 79, 101
Brompheniramine/Pseudoephedrine	30, 79, 101	Cholestyramine	34, 82
Bupivacaine	30, 106	Cilonan	34, 102
buPROPIOn	14, 31, 85	Cipro	34, 96
Burow's Solution	25, 106	Cipro HC Otic	34, 103
BuSpar	17, 31, 86	Ciprofloxacin.....	34, 96, 102
busPIRone	17, 31, 86	Ciprofloxacin/Hydrocortisone	34, 103
Caladryl.....	31, 104	Citalopram	14, 34, 84
Calamine Lotion	31, 104	CitraCal	31, 99
Calamine/Pramoxine	31, 104	Citrolith	63, 93
Calamine/Zinc Oxide/Glycerin	31, 104	Citrucel	54, 91
Calan	16, 76, 81, 88	Clarithromycin	18, 34, 95
Calciferol.....	41, 76, 99	Claritin	52, 79, 101

Cleocin.....	34, 96	Desipramine	14, 37, 84
Cleocin T.....	34, 104, 105	Desitin	36, 38, 104
Climara	42, 88	Desmopressin	37, 90
Clindamycin	34, 96, 104, 105	Desyrel	14, 17, 73, 85, 86
Clinoril.....	70, 83	Detrol.....	73, 93
Clobetasol.....	18, 35, 106	Detrol LA	73, 93
clomiPRAMINE	16, 35, 84	Dexamethasone	37, 89, 102
Clonazepam.....	17, 35, 84, 86, 87	Dexedrine	16, 37, 86
Clonidine.....	16, 35, 82	Dextran.....	37, 98
Clopidogrel.....	18, 35, 80	Dextroamphetamine	16, 37, 86
Clorazepate	17, 35, 84, 87	Dextromethorphan.....	37, 100
Clotrimazole.....	35, 94, 103, 105	Dextrose 5% in 0.2% Sodium Chloride	37, 98
Cloxacillin.....	35, 95	Dextrose 5% in 0.45% Sodium Chloride	37, 98
Clozapen.....	35, 95	Dextrose 5% in 0.9% Sodium Chloride	37, 98
Clozapine.....	13, 19, 35, 85	Dextrose 5% in Ringer's Lactate	38, 98
Clozaril.....	13, 19, 35, 85	Dextrose 5% in Water	38, 98
Coal Tar.....	36, 106	Dextrose 5% with Multiple Electrolytes.....	38, 98
Cod Liver Oil/Zinc Oxide/Talc.....	36, 38	Dextrose 5%/Sodium Chloride 0.2%/Potassium Chloride	38, 98
Cogentin	29, 88	Dextrose 5%/Sodium Chloride 0.45%/Potassium Chloride	38, 98
Colace.....	40, 92	Dextrose 5%/Sodium Chloride 0.9%/Potassium Chloride	38, 98
Colchicine	36, 90	Dextrose 5%/Sodium Chloride/Potassium Chloride Intravenous Solution	38, 98
Collagenase.....	36, 107	Dextrose 50% in Water	38, 78, 98
Co-Lyte	64, 92	Dextrose/Sodium Chloride Intravenous Solution....	37, 98
Combivir.....	50, 97	DiBeta.....	45, 78
Compazine.....	64, 83, 93	Diabinese	34, 78
Concerta	16, 54, 86	Diamox	24, 81
Corgard.....	56, 81, 88	Diaper Rash Powder	38, 104
Corticaine.....	47, 106	Diaperene.....	38, 77, 104
Corticotropin	36, 90	Diastat	38, 87
Cortisone	36, 89	Diazepam	17, 38, 84, 86, 87
Cortisporin	57, 103	Dibucaine	38, 106
Cosopt	72, 101	Dicloxacillin.....	39, 95
Co-Trimoxazole	75, 96	Dicyclomine	39, 90
Coumadin	76, 80	Didanosine	39, 97
Creon	59, 93	Differin	25, 104
Crixivan.....	48, 97	Diflucan	43, 96
Cromolyn	36, 101	Digoxin	39, 81
Crotamiton	36, 105	Dilantin.....	21, 61, 87
Cuprimine	60, 79	Diltiazem.....	39, 81
Cyanocobalamin	36, 99	Dimercaprol	39, 79
Cyproheptadine.....	36, 79	diphenhydrAMINE	17, 39, 79, 83, 86, 101, 104
Cytotec.....	55, 93	Diphtheria & Tetanus Toxoids Adsorbed	39, 94
d4T.....	69, 97	Diphtheria & Tetanus Toxoids Adsorbed for Adult Use	40, 94
D5 E75.....	38, 98	Disulfiram	40, 79
Dantrium	37, 87	Ditropan	59, 93
Dantrolene	37, 87	Ditropan XL	59, 93
DDAVP	37, 90	Divalproex	16, 21, 40, 85, 87, 88
ddl	39, 97	Divalproex ER	19, 40
Debrox	32, 103	DLV	37, 97
Decadron	37, 89, 102	Docusate Calcium	40, 92
Deferoxamine	37, 79	Docusate Sodium	40, 92
Delavirdine.....	37, 97	Docusate Sodium/Casanthrol.....	40, 92
Delta-Cortef	63, 89	Docusate Sodium/Sennosides	40, 92
Deltasone.....	63, 89	Dolophine	53, 83
Depakene	16, 21, 75, 85, 87, 88		
Depakote	16, 21, 40, 85, 88		
Depakote ER	19, 40, 88		
Desenex.....	77, 105		
Desferal	37, 79		

Domeboro Otic 24, 103
Donepezil 19, 40, 88
DOPamine 40, 82
Dovonex 31, 104
Doxepin 14, 40, 84
Doxinate 40, 92
Doxycycline 41, 95, 103
Drisdol 41, 76, 99
Dulcolax 30, 91
Duragesic 43, 83
Dyazide 74, 81
Dycill 39, 95
Dynapen 39, 95
Dyrenium 74, 80
Effexor 14, 75, 84
Elavil 14, 26, 84
Elimite 60, 105
Elixophyllin 71, 100
Emollient Lotion/Cream 41, 106
Emollient Ointment 41, 106
Enalapril 41, 82
Engerix-B 47, 94
Enoxaparin 41, 80
Entex PSE 46, 101
Epinephrine 41, 82
Epivir 50, 97
Epsom Salt 52, 91
Ergocalciferol 41, 76, 99
Erythrocin 41, 95
Erythromycin 41, 95, 102
Erythromycin Ethylsuccinate/Sulfisoxazole 42, 95
Erythromycin/Benzoyl Peroxide 42, 104
Escitalopram 14, 42, 84
Esidrix 47, 80
Eskalith 16, 51, 85
Estrace 42, 88, 94
Estraderm 42, 88
Estradiol 42, 88, 94
Estrogen/medroxyPROGESTERone 42, 89
Estrogens, Conjugated 42, 88, 94
Ethambutol 42, 96
Ethinyl Estradiol/Norethindrone 42, 89
Ethinyl Estradiol/Norgestrel 42, 89
Ethionamide 42, 96
Ethosuximide 21, 43, 87
Ethyl Chloride 43, 106
Eurax 36, 105
Euthroid 51, 89
Evista 65, 90
Exelon 19, 67, 88
Famotidine 43, 90
Felbamate 19, 43, 87
Felbatol 19, 43, 87
Felodipine 43, 81
Fentanyl 43, 83
Feosol 43, 80, 99
Fer-In-Sol 43, 80, 99
Ferro-Sequels 43, 80, 99
Ferrous Fumarate/Docusate Sodium 43, 80, 99

Ferrous Sulfate 43, 80, 99
Fexofenadine 43, 79, 101
Fexofenadine/Pseudoephedrine 43, 79, 101
Filibon 56, 99
Flagyl 55, 96
Flavoxate 43, 93
Fleet Phospho-Soda 69, 91
Fleet's Enema 69, 91
Flonase 44, 100
Florinef 44, 89
Flovent 44, 100
Fluconazole 43, 96
Fludrocortisone 44, 89
Fluocinolone 44, 106
Fluocinonide 44, 106
Fluorescein Sodium 44, 102
Fluoxetine 14, 44, 84
Fluphenazine 13, 44, 85
Fluticasone 44, 100
Fluvastatin 44, 82
Fluviron 48, 94
Fluvoxamine 14, 44, 84
Fluzone 48, 94
Folic Acid 45, 99
Folvite 45, 99
Fortovase 67, 97
Fosamax 25, 90
Fosphenytoin 45, 87
Fulvicin 46, 96
Fungoid 35, 103, 105
Furosemide 45, 80
Gabapentin 45, 87
Gabatril 72, 87
Galantamine 45, 88
Gamma Benzene Hexachloride 51, 105
Garamycin 45, 96, 102, 105
Gaviscon 26, 90
Gemfibrozil 45, 82
Gentamicin 45, 96, 102, 105
Gentran 37, 98
Geodon 13, 77, 85
gliPiZIDE 45, 78
Glucagon 45, 78, 79
Glucophage 53, 78
Glucophage XR 53, 78
Glucotrol 45, 78
glyBURIDE 45, 78
Glycerin 46, 92
Gly-Oxide 32, 103
GOLYTELY 64, 92
Granulex 75, 107
Griseofulvin 46, 96
Guaifenesin 46, 100
Guaifenesin/Dextromethorphan 46, 101
Guaifenesin/Pseudoephedrine 46, 101
Guanethidine 46, 82
Gyne-Lotrimin 35, 94
Habitrol 57, 79
Halcion 17, 74, 84, 86

Haldol.....	13, 46, 85	Isordil.....	49, 81
Haloperidol.....	13, 46, 85	Isosorbide Dinitrate	49, 81
Hartmann's Solution	66, 98	Isosorbide Mononitrate.....	49, 81
HBIG	46, 94	Kaolin-Pectin	49, 92
Heparin	46, 80	Kapectate	30, 92
Hepatitis A Vaccine.....	46, 94	Kayexalate.....	69, 82
Hepatitis B Immune Globulin	46, 94	Keflex	33, 95
Hepatitis B Virus Vaccine, Recombinant	47, 94	Kefzol.....	32, 95
Hexachlorophene.....	47, 105	Kenacort.....	74, 89
Hexavitamins	56, 99	Kenalog in Orabase.....	74, 103
Hibiclens	33, 105	Keppra.....	50, 87
Homatropine	47, 102	Keri Lotion	41, 106
Humalog	48, 78	Ketoconazole.....	49, 96, 105
Humalog Mix 75/25.....	48, 78	Ketorolac	49, 83
Hycotuss	47, 101	Klonopin.....	17, 35, 84, 86, 87
hydrALAZINE	47, 82	Konakion	61, 99
Hydrochlorothiazide	47, 80	Kwelcof.....	47, 101
Hydrocodone/Guaifenesin	47, 101	Kwell	51, 105
Hydrocortisone.....	47, 89, 106	Labetalol.....	49, 81
HydroDIURIL	47, 80	Lac-Hydrin	41, 106
Hydrogen Peroxide	47, 107	Lactinex	49, 92
hydrOXYzine.....	17, 47, 79, 83, 86	Lactobacillus Acidophilus	49, 92
Hygroton	34, 80	Lactulose	50, 92
HypoTears	59, 102	Lamictal.....	16, 50, 87
HypoTears PF.....	59, 102	Lamisil	70, 105
Ibuprofen.....	48, 83	Lamivudine	50, 97
Imdur.....	49, 81	Lamivudine/Zidovudine	50, 97
Imferon	49, 80, 99	Lamotrigine.....	16, 50, 87
Imipramine.....	14, 48, 84	Lanacaine.....	29, 106
Imitrex.....	70, 88	Lanacort	47, 106
Imodium.....	52, 92	Lanolin.....	41, 106
Inderal.....	16, 64, 81, 88	Lanoxin.....	39, 81
Indinavir	48, 97	Lansoprazole.....	50, 91
Influenza Virus Vaccine	48, 94	Lantus.....	48, 78
INH.....	49, 96	Larodopa	50, 88
Insulin, Combination (70/30).....	48, 78	Lasix	45, 80
Insulin, Glargin.....	48, 78	Latanoprost	50, 101
Insulin, Lente	48, 78	Lescol	44, 82
Insulin, Lispro.....	48, 78	Leucovorin.....	50, 79, 99
Insulin, Lispro/Insulin, Lispro Protamine	48, 78	Levaquin	51, 96
Insulin, NPH.....	48, 78	Levarterenol	50, 58, 82
Insulin, Regular.....	48, 78	Levetiracetam	50, 87
Insulin, Ultralente	48, 78	Levodopa.....	50, 88
Intal	36, 101	Levodopa/Carbidopa	50, 88
Intropin	40, 82	Levofoxacin	51, 96
Invirase	67, 97	Levonorgestrel/Ethynodiol	51, 89
Ionil-T	36, 106	Levophed.....	50, 58, 82
Ipecac Syrup.....	48, 79, 93	Levothyroxine	51, 89
IPOL.....	62, 94	Lexapro	14, 42, 84
Ipratropium.....	49, 101	Librium.....	17, 33, 84
Iron Dextran.....	49, 80, 99	Lidex	44, 106
Ismelin	46, 82	Lidocaine	51, 106
ISMO.....	49, 81	Lindane.....	51, 105
Isoniazid.....	49, 96	Lioresal.....	28, 87
Isoptin	16, 76, 81, 88	Liotrix	51, 89
Isopto Atropine.....	28, 102	Lipitor.....	28, 82
Isopto Carpine	61, 101	Lisinopril	51, 82
Isopto Homatropine	47, 102	Lithium	16, 51, 85
Isopto Hyoscine	67, 102	Lithobid	16, 51, 85

Lithonate	51, 85
LMD	37, 98
Loestrin	42, 89
Lo-Ovral	42, 89
Loperamide	52, 92
Lopid	45, 82
Lopressor	54, 81, 88
Loratadine	52, 79, 101
Lorazepam	17, 52, 84, 86, 87
Lortab	24, 83
Lotensin	28, 82
Lotrimin	35, 105
Lovenox	41, 80
Loxapine	13, 52, 85
Loxitane	13, 52, 85
Lubriderm	41, 106
Ludiomil	14, 52, 85
Lumigan	30, 101
Luminal	21, 61, 87
Luvox	14, 44, 84
Maalox	26, 90
Macrodantin	58, 94, 97
Macrodex	37, 98
Magnesium Citrate	52, 91
Magnesium Hydroxide	52, 91
Magnesium Sulfate	52, 91, 98
Maprotiline	14, 52, 85
Marcaine	30, 106
Maxzide	74, 81
Measles, Mumps and Rubella Virus Vaccine, Live	52, 94
Mebaral	53, 87
Mebendazole	52, 97
Meclizine	52, 83, 93
Medrol	54, 89
medroxyPROGESTERone	53, 88
Mellaril	13, 20, 71, 85
Mephobarbital	53, 87
Mephyton	61, 79, 80, 99
Meruvax II	67, 94
Mesalamine	53, 93
Mesoridazine	13, 19, 53, 85
Metamucil	65, 91
Metaproterenol	53, 100
Metformin	53, 78
Methadone	53, 83
Methimazole	53, 89
Methocarbamol	53, 87
Methotrexate	53, 79, 104
Methyl Salicylate	54, 106
Methylcellulose	54, 91
Methyldopa	54, 82
Methylphenidate	16, 54, 86
Methylprednisolone	54, 89
methylTESTOSTERone	54, 89
Meticorten	63, 89
Metoclopramide	54, 83, 91
Metoprolol	54, 81, 88
MetroGel	55, 104, 105
Metronidazole	55, 96, 104, 105
Mexsana	38, 104
Miacalcin	31, 90
Miconazole	55, 94, 105
Micronase	45, 78
Midazolam	55, 86
Milk of Magnesia	52, 91
Mineral Oil	55, 103
Minipress	63, 82
Mintezol	71, 97
MiraLax	62, 92
Mirtazapine	14, 17, 55, 85
Misoprostol	55, 93
MMR II	52, 94
Moban	13, 55, 85
Moi-Stir	32, 103
Molindone	13, 55, 85
Mometasone	55, 100
Monistat	55, 94, 105
Monoket	49, 81
Morphine	55, 83
Motrin	48, 83
MouthKote	32, 103
Mucomyst	25, 79, 101
Multivitamin	56, 99
Multivitamin, Prenatal	56, 99
Multivitamin/Minerals	56, 99
Multivitamins, Pediatric	56, 99
Mupirocin	56, 105
Myambutol	42, 96
Mycelex	35, 94, 103
Mycifradin	57, 96
Mycostatin	58, 96, 103, 105
Mydriacyl	75, 102
Mylanta	26, 90
Mylicon	68, 91
Mysoline	63, 87
Nabumetone	19, 56, 83
Nadolol	56, 81, 88
Nafcillin	56, 95
Naloxone	56, 79, 86
Naltrexone	56, 79, 86
Naphazoline	56, 102
Naphcon	56, 102
Naprosyn	57, 83, 88
Naproxen	57, 83, 88
Narcan	56, 79, 86
Nardil	14, 61, 84
Nasacort	74, 100
Nasonex	55, 100
Navane	13, 71, 85
Nebcin	72, 96
Nefazodone	14, 57, 85
Nelfinavir	57, 97
Neo-Calglucon	31, 99
Neomycin	57, 96
Neomycin/Polymyxin B/Bacitracin	57, 105
Neomycin/Polymyxin B/Hydrocortisone	57, 103
Neosporin	62, 105
Neo-Synephrine	61, 102, 103

Neurontin	45, 87	Paroxetine	14, 60, 84
Nevirapine.....	57, 97	Patanol	59, 102
Niacin/Nicotinamide.....	57, 82, 99	Pathocil.....	39, 95
Nicobid.....	57, 82, 99	Paxil.....	14, 60, 84
Nicoderm	57, 79	Pediazole.....	42, 95
Nicorette	57, 79	PEG.....	64, 92
Nicotine	57, 79	Penicillamine	60, 79
Nicotine Polacrilex	57, 79	Penicillin G Benzathine.....	60, 95
Nicotine Transdermal Patch	57, 79	Penicillin G Benzathine/Penicillin G Procaine	60, 95
Nicotrol.....	57, 79	Penicillin G Procaine	60, 95
NIFEEdipine.....	57, 81	Penicillin G Sodium	60, 95
Nitrofurantoin	58, 94, 97	Penicillin V Potassium	60, 95
Nitroglycerin.....	58, 81	Pentam	19, 60, 97
NIX	60, 105	Pentamidine	19, 60, 97
Nizoral.....	49, 96, 105	Pentasa	53, 93
Noctec.....	17, 33, 86	Pen-Vee K	60, 95
Nolvadex.....	70, 79	Pepcid	43, 90
Non-Soap Cleanser	58, 104	Pepto-Bismol	30, 92, 93
Norepinephrine	58, 82	Periactin	36, 79
Norgestimate/Ethinyl Estradiol.....	58, 89	Peri-Colace.....	40, 92
Noritate	55, 104, 105	Peridex	33, 103
Normodyne	49, 81	PerioMed	69, 103
Norpramin	14, 37, 84	Periostat	41, 103
Nortriptyline	14, 58, 84	Permethrin	60, 105
Norvasc.....	26, 81	Permitil	13, 44, 85
Norvir	66, 97	Pernox	24, 104
Nupercainal	38, 106	Perphenazine	13, 60, 85
NVP	57, 97	Pertofrane.....	37, 84
Nystatin	58, 96, 103, 105	Petrolatum, White.....	61, 106
OCL	64, 92	Phenazopyridine	61, 93
Olanzapine.....	13, 59, 85	Phenelzine	14, 61, 84
Olopatadine	59, 102	Phenergan	64, 79, 83
Omeprazole	59, 91	Phenobarbital	21, 61, 87
OmniMed	69, 103	Phenol	61, 103
Omnipen	27, 95	Phenylephrine	61, 102, 103
Ophthalmic Lubricant.....	59, 102	Phenytoin	21, 61, 87
Oracit	68, 93	pHisoHex	47, 105
Oreton	54, 89	Physostigmine	61, 79
Orinase	73, 78	Phytomedicine	61, 79, 80, 99
Ortho Tri-Cyclen	58, 89	Pilocarpine	61, 101
Ortho-Novum 777	42, 89	Plavix	18, 35, 80
Os-Cal.....	31, 90, 99	Plendil	43, 81
Oscal + D	31, 99	Pneumococcal Vaccine, Polyvalent	62, 94
Ovral	42, 89	Pneumovax	62, 94
Oxacillin	59, 95	Podophyllum Resin	62, 106
Oxazepam	17, 59, 84, 86	Poliovirus Vaccine, Inactivated	62, 94
Oxcarbazepine	16, 59, 87	Polycillin	27, 95
Oxybutynin	59, 93	Polycitra	63, 93
Oxycodone	59, 83	Polycitra K	63, 93
OxyContin	59, 83	Polycitra-LC	63, 93
Oxymetazoline	59, 103	Polyethylene Glycol	62, 92
P.P.D.	75, 95	Polymox	26, 95
Pamelor	14, 58, 84	Polymyxin B/Bacitracin	62, 102
Pancrease	59, 93	Polymyxin B/Neomycin	62, 105
Pancrelipase	59, 93	Polymyxin B/Trimethoprim	62, 102
Panmycin	71, 95	Polysporin	28, 62, 102, 105
Pantoprazole	60, 91	Polytrim	62, 102
Parlodol	30, 88	Poly-Vi-Sol	56, 99
Parnate	14, 73, 84	Potassium Chloride	62, 98

Potassium Citrate	62, 93	Renagel	67, 94
Potassium Citrate Combinations	63, 93	Repaglinide.....	66, 78
Potassium Iodide	63, 100	Rescriptor	37, 97
Povidone-Iodine	63, 105	Restoril	17, 70, 84, 86
Pramoxine.....	63, 92, 106	Retin-A	73, 104
Prandin	66, 78	Retrovir	77, 97
Prazosin.....	63, 82	ReVia.....	56, 79, 86
prednisoLONE	63, 89	Rezamid	70, 104
predniSONE.....	63, 89	Rheomacrodex	37, 98
Premarin	42, 88, 94	RID	65, 105
PremPro.....	42, 89	Rifadin	66, 96
Prevacid.....	50, 91	Rifamate	66, 96
Prilosec	59, 91	Rifampin	66, 96
Primidone.....	63, 87	Rifampin/Isoniazid	66, 96
Prinivil	51, 82	Ringer's Lactate Solution	66, 98
Pro-Banthine.....	64, 90	Risperdal	13, 66, 85
Probenecid.....	64, 90	Risperdal Consta.....	66, 85
Procainamide.....	64, 81	Risperidone	13, 66, 85
Procardia	57, 81	Ritalin.....	16, 54, 86
Prochlorperazine.....	64, 83, 93	Ritonavir	66, 97
Prolixin	13, 44, 85	Rivastigmine.....	19, 67, 88
Promethazine.....	64, 79, 83	Robaxin	53, 87
Pronestyl.....	64, 81	Robitussin.....	46, 100
Propantheline.....	64, 90	Robitussin DM	46, 101
Proparacaine	64, 102	Rocephin	32, 95
Propranolol	16, 64, 81, 88	Rosiglitazone	67, 78
Propylethylene Glycol Electrolyte Solution.....	64, 92	Rowasa	53, 93
Propylthiouracil	64, 89	Rubella Virus Vaccine Live.....	67, 94
Prostaphlin.....	59, 95	Salicylic Acid.....	67, 106
ProStep.....	57, 79	Saliva Substitute.....	32, 103
Protamine	64, 79, 80	Salivart.....	32, 103
Protonix.....	60, 91	Salix.....	32, 103
Protriptyline.....	14, 65, 84	Salmeterol	67, 100
Proventil.....	25, 100	Sani-Supp.....	46, 92
Provera	53, 88	Santyl.....	36, 107
Prozac.....	14, 44, 84	Saquinavir.....	67, 97
Pseudoephedrine.....	65, 100	Scopolamine.....	67, 102
Psyllium	65, 91	Sebizon.....	69, 104
Pyrantel.....	65, 97	Selenium Sulfide	67, 104
Pyrazinamide	65, 96	Selsun	67, 104
Pyrethrins/Piperonyl Butoxide	65, 105	Senna	67, 91
Pyridium	61, 93	Senokot	67, 91
Pyridoxine	65, 99	Septra	75, 96
Questran	34, 82	Serax	17, 59, 84, 86
Quetiapine	13, 65, 85	Serentil	13, 19, 53, 85
Quinidine Gluconate	65, 81	Serevent	67, 100
Quinidine Sulfate	65, 81	Seroquel	13, 65, 85
Raloxifene	65, 90	Sertraline	14, 67, 84
Ranitidine	66, 90	Serzone	14, 57, 85
Recombivax HB	47, 94	Sevelamer	67, 94
Rectal Hemorrhoidal Cream with Hydrocortisone ..	66, 92	Silvadene.....	68, 104, 106
Rectal Hemorrhoidal Ointment	66, 92	Silver Nitrate	68, 107
Rectal Hemorrhoidal Suppositories	66, 92	Silver Sulfadiazine	68, 104, 106
Rectal Hemorrhoidal Suppositories with Hydrocortisone.....	66, 92	Simethicone	68, 91
Reglan	54, 83, 91	Simvastatin	68, 82
Relafen	19, 56, 83	Sinemet	50, 88
Remeron.....	14, 17, 55, 85	Sinequan	14, 40, 84
Reminyl.....	45, 88	Sodium Bicarbonate	68, 93, 98
		Sodium Chloride.....	68, 93, 101, 102, 107

Sodium Chloride 0.2%	68, 98	Thyroid	72, 89
Sodium Chloride 0.45%	68, 98	Thyroid, Desiccated	72, 89
Sodium Chloride 0.9%	68, 98	Thyrolar	51, 89
Sodium Chloride Intravenous Solution	68, 98	Tiagabine	72, 87
Sodium Citrate/Citric Acid	68, 93	Ticar	72, 95
Sodium Fluoride	69, 103	Ticarcillin	72, 95
Sodium Lactate	69, 98	Ticarcillin/Clavulanate	72, 95
Sodium Phosphate/Biphosphate	69, 91	Tigan	74, 83, 93
Sodium Polystyrene Sulfonate	69, 82	Timentin	72, 95
Sonata	17, 76, 86	Timolol	72, 101
Sorbitol	69, 92	Timolol/Dorzolamide	72, 101
Sorbitrate	49, 81	Timoptic	72, 101
Spironolactone	69, 80	Tinactin	73, 105
Spironolactone/Hydrochlorothiazide	69, 81	Tioconazole	72, 94
SSKI	63, 100	Titralac	31, 90, 99
Stanous Fluoride	69, 103	Tizanidine	20, 72, 87
Stavudine	69, 97	TobraDex	72, 102
Stelazine	13, 74, 85	Tobramycin	72, 96, 102
Stimate	37, 90	Tobramycin/Dexamethasone	72, 102
Strattera	27, 86	Tobrex	72, 102
Stresscaps	76, 99	Tofranil	14, 48, 84
Sucralfate	69, 93	TOLBUTamide	73, 78
Sudafed	65, 100	Tolnaftate	73, 105
Sulamyd	69, 102	Tolterodine	73, 93
Sulfacetamide Sodium	69, 102, 104	Topamax	16, 73, 87
Sulfasalazine	70, 93	Topiramate	16, 73, 87
Sulforcin	70, 104	Toradol	49, 83
Sulfur/Resorcinol	70, 104	Tramadol	73, 83
Sulindac	70, 83	Tranxene	17, 35, 84, 87
Sumatriptan	70, 88	Tranxene SD	35, 87
Sunscreen/block	70, 104	Tranylcypromine	14, 73, 84
Surfak	40, 92	Travatan	73, 101
Surmontil	14, 75, 84	Travoprost	73, 101
Symmetrel	26, 88, 97	Trazodone	14, 17, 73, 85, 86
Synalar	44, 106	Tretinoin	73, 104
Synthroid	51, 89	Trexan	56, 79, 86
Tamoxifen	70, 79	Triamcinolone	74, 89, 100
Tapazole	53, 89	Triamcinolone in Oral Adhesive Base	74, 103
Tazarotene	70, 104	Triamterene	74, 80
Tazorac	70, 104	Triamterene/Hydrochlorothiazide	74, 81
Tegopen	35, 95	Triazolam	17, 74, 84, 86
Tegretol	16, 21, 32, 85, 87	Trifluoperazine	13, 74, 85
Tegretol XR	16, 21, 32	Trihexyphenidyl	74, 88
Tegrin	36, 106	Trilafon	13, 60, 85
Teldrin	33, 79, 101	Trileptal	16, 59, 87
Temazepam	17, 70, 84, 86	Tri-Levlen	51, 89
Temovate	18, 35, 106	Trimethobenzamide	74, 83, 93
Tenormin	27, 81, 88	Trimethoprim/Sulfamethoxazole	75, 96
Terbinafine	70, 105	Trimipramine	14, 75, 84
Terbutaline	70, 100	Triphasil	51, 89
Testosterone	70, 89	Triple Antibiotic Ointment	57, 105
Tetracycline	71, 95	Triprolidine/Pseudoephedrine	75, 79, 101
Tetrahydrozoline	71, 102	Tronothane	63, 92, 106
Theophylline	71, 100	Tropicamide	75, 102
Thiabendazole	71, 97	Trypsin/Balsam Peru/Castor Oil	75, 107
Thiamine	71, 99	Tuberculin, Purified Protein Derivative	75, 95
Thioridazine	13, 20, 71, 85	Tylenol	24, 83
Thiothixene	13, 71, 85	Ultram	73, 83
Thorazine	13, 34, 85	Unicap	56, 99

Unifiber	33, 91	Vivactil	14, 65, 84
Unipen	56, 95	Vivelle	42, 88
Urecholine	30, 93	VoSol	24, 103
Urispas.....	43, 93	VoSol HC.....	24, 103
Urocit K.....	62, 93	Warfarin.....	76, 80
Vagistat-1.....	72, 94	Water for Injection.....	76, 98, 107
Valisone	29, 106	Water for Irrigation	76, 93
Valium.....	17, 38, 84, 86, 87	Wellbutrin	14, 31, 85
Valproate	16, 21, 75, 85, 87, 88	Wellbutrin SR	14, 31, 85
Valproic Acid.....	16, 21, 75	Wellbutrin XL.....	14, 31, 85
Vanceril.....	28, 100	Wellcovorin.....	50, 79, 99
Vancocin	75, 96	Whitfield's	29, 105, 107
Vancomycin	75, 96	Wyanoids.....	66, 92
Vaqta	46, 94	Wycillin	60, 95
Varicella Virus Vaccine, Live	75, 94	Xalatan	50, 101
Varivax.....	75, 94	Xanax	17, 25, 84, 86
Vaseline	61, 106	Xylocaine	51, 106
Vasotec.....	41, 82	Zafirlukast.....	76, 101
V-Cillin K	60, 95	Zaleplon.....	17, 76, 86
Venlafaxine	14, 75, 84	Zanaflex.....	20, 72, 87
Ventolin	25, 100	Zantac	66, 90
Verapamil.....	16, 76, 81, 88	Zarontin	21, 43, 87
Vermox	52, 97	Zephiran	28, 105
Versed	55, 86	Zerit	69, 97
Vibramycin	41, 95	Zestril.....	51, 82
Vicodin	24, 83	Zidovudine	77, 97
Videx.....	39, 97	Zinc Oxide	77, 106, 107
Viracept.....	57, 97	Zinc Oxide/Petrolatum/Imidazolidinyl Urea	38, 77
Viramune	57, 97	Zinc Sulfate	77, 98, 99
Visine Allergy Relief.....	71, 102	Zinc Undecylenate.....	77, 105
Visine Moisturizing.....	71, 102	Ziprasidone.....	13, 77, 85
Vitamin A	76, 99	Zithromax	18, 28, 95
Vitamin A&D Ointment.....	38, 76, 104	Zocor	68, 82
Vitamin B Complex/Vitamin C	76, 99	Zoloft	14, 67, 84
Vitamin B Complex/Vitamin C/Zinc.....	76, 99	Zolpidem.....	17, 77, 86
Vitamin B ₁	71, 99	Zonegran	77, 87
Vitamin B ₁₂	36, 99	Zonisamide	77, 87
Vitamin B ₆	65, 99	Zovirax.....	25, 97, 105
Vitamin C	27, 99	Zydis	59, 85
Vitamin D	76, 99	Zyloprim	25, 90
Vitamin E	76, 99	Zyprexa.....	13, 59, 85
Vitamin K ₁	61, 79, 80, 99		

APPENDIX 1: NEW DRUG APPLICATION FORM

415 — C
EXHIBIT A

TEXAS DEPARTMENT OF MENTAL HEALTH AND MENTAL RETARDATION

NEW DRUG APPLICATION

(for inclusion in the *TDMHMR Drug Formulary*)

** (THE NEW DRUG APPLICATION PROCESS IS DESCRIBED ON THE BACK OF THIS FORM.) **

Date: _____

Name of practitioner submitting the application: _____

Name of entity with which the practitioner is associated by employment or contract (i.e., state hospital, state school, state center, or local authority (state-operated community services (SOCS) or community MHMR center)): _____

Information regarding new drug:

Therapeutic Classification	
Generic Name	
Trade Name(s)	
Manufacturer(s)	
Dosage Form(s)	

Explain the pharmacological action or use of this drug:

Explain the advantages of this drug over those listed in the formulary:

State which drugs this new drug would replace or supplement:

application is approved _____ signature of chairman of facility pharmacy and therapeutics committee
OR
 application is appropriate and complete _____ signature of clinical/medical director or designee

Section 415.108 of TDMHMR rules governing the use and maintenance of the *TDMHMR Drug Formulary* (25 TAC, Chapter 415, Subchapter C) describes the procedures for applying to have a drug added to the formulary, which are:

§415.108. Applying to Have a Drug Added to the Formulary.

- (a) Any member of the Executive Formulary Committee, any service system component* practitioner, or any contract practitioner may apply to have a drug added to the *TDMHMR Drug Formulary* by completing the New Drug Application Form DF-1, referenced as Exhibit A in §415.112 of this title (relating to Exhibit) and including:
 - (1) published articles in biomedical literature that substantiate the efficacy and safety of the proposed drug;
 - (2) information on the advantages of the proposed drug compared with similar formulary drugs;
 - (3) a list of formulary drugs that the proposed drug would replace or supplement; and
 - (4) cost effectiveness data.
- (b) Submitting the application.
 - (1) If the person submitting the application is a facility** practitioner or a facility contract practitioner, then that practitioner submits the application to the facility's pharmacy and therapeutics committee for approval. If the committee approves the application, then it forwards the application to the Executive Formulary Committee.
 - (2) If the person submitting the application is a non-facility service system component practitioner or a non-facility service system component contract practitioner, then that practitioner submits the application to the component's clinical/medical director or designee who determines if the application is appropriate and complete, and if so, forwards the application to the Executive Formulary Committee.
 - (3) If the person completing the application is a member of the Executive Formulary Committee, then that person submits the application directly to Executive Formulary Committee.
- (c) The Executive Formulary Committee considers the drug application and recommends:
 - (1) approving the proposed drug's inclusion and, if appropriate, approving audit criteria and recommending dosage guidelines;
 - (2) denying the proposed drug's inclusion;
 - (3) approving the proposed drug on a trial basis for a specified period of time;
 - (4) approving the proposed drug as a reserve drug, with guidelines; or
 - (5) postponing the decision until a later meeting.

* The term "service system component" means a state hospital, state school, state center, or local authority (state-operated community services (SOCS) or community MHMR center).

** The term "facility" means a state hospital, state school, or state center.

APPENDIX 2: NON-FORMULARY DRUG JUSTIFICATION FORM

TEXAS DEPARTMENT OF MENTAL HEALTH
AND MENTAL RETARDATION

Non-Formulary Drug Justification

Facility/Component _____

Drug Name

(Generic) (Trade) (Strength) (AHFS Therapeutic Class)

The purpose of the *Formulary* is to ensure that the treatments available to TDMHMR consumers are consistent with need, effectiveness, risk, and cost. Exceptions to the *Formulary* are limited to the three categories listed below.

This request is for:

- an individual patient treatment course. Is this chronic or acute therapy?
- more than one patient/course of therapy. It is estimated that _____ patients will be treated with this drug. Is this chronic or acute therapy?

Reason for request:

- An illness for which no *Formulary* drug is as safe or effective.
- A trial supply in anticipation of application to *Formulary*.
- To prevent interruption of course of therapy established prior to admission.

Quantity Ordered	Cost/Unit	Total Purchase Cost	*Estimate Course Chronic Cost Per Month	*Estimate Course Acute Cost

- Alternative drugs are available on the *TXMHMR Formulary*.

Attending Physician (signature)

Date

Was the drug ordered yes or no?

AGREE

DISAGREE

Pharmacy Director or designee (signature)

Date

APPROVED

DISAPPROVED

EMERGENCY

Facility clinical/medical director's approval must be obtained within three working days.

Facility clinical/medical director or designee (signature)

Date

APPENDIX 3: ADVERSE DRUG REACTION REPORTING (MEDWATCH) FORM



THE FDA MEDICAL PRODUCTS REPORTING PROGRAM

For VOLUNTARY reporting
by health professionals of adverse
events and product problems

Form Approved: OMB No. 0915-0281 Expires: 11/30/00
See OMB statement on reverse

FDA Use Only
Triage unit sequence #

Page _____ of _____

A. Patient information

1. Patient identifier In confidence	2. Age at time of event: or Date of birth:	3. Sex <input type="checkbox"/> female <input type="checkbox"/> male	4. Weight lbs or kgs
--	--	--	-------------------------------

B. Adverse event or product problem

1. Adverse event and/or Product problem (e.g., defects/malfunctions)

2. Outcomes attributed to adverse event (check all that apply)
- | | |
|---|---|
| <input type="checkbox"/> death | <input type="checkbox"/> disability |
| <input type="checkbox"/> life-threatening | <input type="checkbox"/> congenital anomaly |
| <input type="checkbox"/> hospitalization – initial or prolonged | <input type="checkbox"/> required intervention to prevent permanent impairment/damage |
| | <input type="checkbox"/> other: |

3. Date of event
(month/yr)

4. Date of this report
(month/yr)

5. Describe event or problem

6. Relevant tests/laboratory data, including dates

7. Other relevant history, including preexisting medical conditions (e.g., allergies, race, pregnancy, smoking and alcohol use, hepatic/renal dysfunction, etc.)

FDA Mail to: MEDWATCH or FAX to:
5600 Fishers Lane 1-800-FDA-0178
Rockville, MD 20852-9787

FDA Form 3500

Submission of a report does not constitute an admission that medical personnel or the product caused or contributed to the event.

C. Suspect medication(s)

1. Name (give labeled strength & mfr/labeler, if known)

#1

#2

2. Dose, frequency & route used

#1

3. Therapy dates (if unknown, give duration)
from/to (or best estimate)

#1

#2

4. Diagnosis for use (indication)

#1

5. Event abated after use stopped or dose reduced

#1 yes no doesn't apply

#2 yes no doesn't apply

#2

6. Lot # (if known)

#1

7. Exp. date (if known)

#1

#2

9. NDC # (for product problems only)

—

—

10. Concomitant medical products and therapy dates (exclude treatment of event)

D. Suspect medical device

1. Brand name

2. Type of device

3. Manufacturer name & address

4. Operator of device

health professional
 lay user/patient
 other:

6. model #

5. Expiration date
(month/yr)

catalog #

7. If implanted, give date
(month/yr)

serial #

8. If explanted, give date
(month/yr)

lot #

other #

9. Device available for evaluation? (Do not send to FDA)

yes no returned to manufacturer on (month/yr)

10. Concomitant medical products and therapy dates (exclude treatment of event)

E. Reporter (see confidentiality section on back)

1. Name & address

phone #

2. Health professional?

yes no

3. Occupation

4. Also reported to

manufacturer
 user facility
 distributor

5. If you do NOT want your identity disclosed to the manufacturer, place an "X" in this box.

ADVICE ABOUT VOLUNTARY REPORTING

Report experiences with:

- medications (drugs or biologics)
- medical devices (including in-vitro diagnostics)
- special nutritional products (dietary supplements, medical foods, infant formulas)
- other products regulated by FDA

Report SERIOUS adverse events. An event is serious when the patient outcome is:

- death
- life-threatening (real risk of dying)
- hospitalization (initial or prolonged)
- disability (significant, persistent or permanent)
- congenital anomaly
- required intervention to prevent permanent impairment or damage

Report even if:

- you're not certain the product caused the event
- you don't have all the details

Report product problems – quality, performance or safety concerns such as:

- suspected contamination
- questionable stability
- defective components
- poor packaging or labeling
- therapeutic failures

How to report:

- just fill in the sections that apply to your report
- use section C for all products except medical devices
- attach additional blank pages if needed
- use a separate form for each patient
- report either to FDA or the manufacturer (or both)

Important numbers:

- 1-800-FDA-0178 to FAX report
- 1-800-FDA-7737 to report by modem
- 1-800-FDA-1088 to report by phone or for more information
- 1-800-822-7967 for a VAERS form for vaccines

If your report involves a serious adverse event with a device and it occurred in a facility outside a doctor's office, that facility may be legally required to report to FDA and/or the manufacturer. Please notify the person in that facility who would handle such reporting.

Confidentiality: The patient's identity is held in strict confidence by FDA and protected to the fullest extent of the law. The reporter's identity, including the identity of a self-reporter, may be shared with the manufacturer unless requested otherwise. However, FDA will not disclose the reporter's identity in response to a request from the public, pursuant to the Freedom of Information Act.

The public reporting burden for this collection of information has been estimated to average 30 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to:

DRA Reports Clearance Office
Paperwork Reduction Project, (0910-0221)
Hubert H. Humphrey Building, Room 531-H
200 Independence Avenue, S.W.
Washington, DC 20201

"An agency may not conduct or sponsor,
and a person is not required to respond to,
a collection of information unless it displays
a currently valid OMB control number."

Please DO NOT
RETURN this form
to this address.

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Public Health Service • Food and Drug Administration

FDA Form 3500-back

Please Use Address Provided Below – Just Fold In Thirds, Tape and Mail

**Department of
Health and Human Services**
Public Health Service
Food and Drug Administration
Rockville, MD 20857

Official Business
Penalty for Private Use \$300

BUSINESS REPLY MAIL
FIRST CLASS MAIL PERMIT NO. 946 ROCKVILLE, MD

POSTAGE WILL BE PAID BY FOOD AND DRUG ADMINISTRATION

MEDWATCH
The FDA Medical Products Reporting Program
Food and Drug Administration
5600 Fishers Lane
Rockville, MD 20852-9787

Medication Audit Criteria and Guidelines

-- INDEX --

The audit criteria and guidelines are developed for use in the treatment of psychiatric conditions and not medical conditions.

Criteria and Drug Audit Form Number	Description
CURRENT #	
Appendix A	Purpose of Laboratory Monitoring
Table A	Cytochrome P450 Drug Metabolism/Inhibition
1.	CARBAMAZEPINE (TEGRETOL®)
2.	LAMOTRIGINE (LAMICTAL®)
3.	LITHIUM (ESKALITH®, LITHOBID®, ESKALITH CR®, etc.)
4.	OXCARBAZEPINE (TRILEPTAL®)
5.	TOPIRAMATE (TOPAMAX®)
6.	VALPROIC ACID (DEPAKENE®), DIVALPROEX SODIUM (DEPAKOTE®)
7.	VERAPAMIL (CALAN®, ISOPTIN®)
8.	BENZODIAZEPINES alprazolam (Xanax®), chlordiazepoxide (Librium®), clorazepate (Tranxene®), diazepam (Valium®), lorazepam (Ativan®), Oxazepam (Serax®), temazepam (Restoril®), triazolam (Halcion®), Clonazepam (Klonopin®)
9.	BUSPIRONE (BUSPAR®)
10.	AMOXAPINE (ASENDIN®)
11.	BUPROPION (WELLBUTRIN® and WELLBUTRIN® SR)
12.	MIRTAZAPINE (REMERON®)
13.	MONOAMINE OXIDASE INHIBITORS phenelzine (Nardil®), tranylcypromine (Parnate®)
14.	NEFAZODONE (SERZONE®)
15.	SSRIs: CITALORPAM (CELEX®), FLUOXETINE (PROZAC®), SERTRALINE (ZOLOFT®), PAROXETINE (PAXIL®), FLUVOXAMINE (LUVOX®)
16.	TRAZODONE (DESYREL®)
17.	TRICYCLIC ANTIDEPRESSANTS amitriptyline (Elavil®), desipramine (Norpramin®, Pertofrane®), doxepin (Sinequan®), imipramine (Tofranil®), maprotiline (Ludiomil®), nortriptyline (Pamelor®, Aventyl®), protriptyline (Vivactil®), trimipramine (Surmontil®)
18.	VENLAFAKINE (EFFEXOR® and EFFEXOR® ER)
19.	ANTIPSYCHOTICS chlorpromazine (Thorazine®), fluphenazine (Prolixin®), haloperidol (Haldol®), loxapine (Loxitane®), molindone (Molan®), perphenazine (Trilafon®), thiothixene (Navane®), trifluoperazine (Stelazine®)
20.	ANTIPSYCHOTICS mesoridazine (Serentil®)thioridazine (Mellaril®)
21.	CLOZAPINE (CLOZARIL®)
22.	DECANOATES fluphenazine decanoate (Prolixin® Decanoate), haloperidol decanoate (Haldol® Decanoate)

Criteria and Drug Audit Form Number CURRENT #	Description
23.	RISPERIDONE (RISPERDAL®), OLANZAPINE (ZYPREXA®), QUETIAPINE (SEROQUEL®), ZIPRASIDONE (GEODON®)
24.	SEDATING ANTIHISTAMINES diphenhydramine HCL (Benadryl®), hydroxyzine HCL (Atarax®)
25.	ZALEPLON (SONATA®)
26.	ZOLPIDEM (AMBIEN®)
27.	BETA-BLOCKERS propranolol (Inderal®), atenolol (Tenormin®), metoprolol (Lopressor®), nadolol (Corgard®)
28.	CLOMIPRAMINE (ANAFRANIL®)
29.	GABAPENTIN (NEURONTIN®)
30.	NALTREXONE (REVIA®)
31.	STIMULANTS methylphenidate (Ritalin®, Concerta™), dextroamphetamine (Dexedrine®), dextroamphetamine/amphetamine mixture (Adderall®)

Medication Audit Criteria and Guidelines

Purpose of Laboratory Monitoring

This document was developed based on the premise that the laboratory tests needed for prescribing psychotropic medications are apart from the laboratory tests obtained for the evaluation of the patient's general health status. The required laboratory tests listed are specific for risk factors associated with that particular psychotropic medication. The required psychotropic medication laboratory screening does not substitute for a good history and physical and subsequent healthcare screening needed for the provision of good general medical care for the person who has become a psychiatric patient.

The specific laboratory tests required for the use of psychotropic medication can be obtained from other treatment settings provided:

- The laboratory tests were obtained within 90 days prior to initiation of treatment.
- The actual values of the tests are documented in the chart. Other documentation shall include the date the lab work was obtained and the name of the laboratory.
- There are no intervening illnesses within those 90 days which would necessitate repeating the lab work.

The laboratory tests listed in this document are minimum requirements. The clinician is encouraged to obtain any necessary lab work which he/she feels is clinically justified.

Table A**Cytochrome P450 Drug Metabolism/Inhibition**Adapted from *Bazire and Benefield's Psychotropic Drug Directory 2001, Alden Press, Oxford, UK***CYP1A2**

Substrates	Significant enzyme inducers	Significant enzyme inhibitors
Caffeine	Caffeine	Cimetidine
Clozapine (most)	Charcoal broiled foods	Ciprofoxacin
Dizaepam	Cigarette smoke	Clarithromycin
Fluvoxamine (partly)	Omeprazole	Erythromycin
Haloperidol	Phenobarbital (weak)	Fluvoxamine (potent – other SSRIs very weak)
Mirtazapine	Phenytoin (weak)	Isoniazid
Olanzapine	Rifampin	Ketoconazole
Ondansetron	Ritonavir	Mirtazapine (very weak)
Pimozide		Nefazodone (very weak)
Propranolol?		Norfloxacin
Tacrine		Omeprazole
Tamoxifen		
Theophylline		
Tricyclic antidepressants		
Verapamil		
Wargfarin		
Ziprasidone		

CYP2C family

Substrates	Significant enzyme inducers	Significant enzyme inhibitors
Amitriptyline	Carbamazepine (weak)	Fluoxetine?
Clomipramine	Phenobarbital (weak)	Fluvoxamine?
Imipramine	Phenytoin (weak)	Sertraline?
Diazepam	Rifampin (weak)	
Omeprazole		
NSAIDs (some)		
Phenytoin		
Tolbutamide		
Warfarin		

CYP2C9/19

Substrates	Significant enzyme inducers	Significant enzyme inhibitors
Barbiturates (19)	Rifampin (9)	Amiodarone (9)
Bupropion (9)		Carbamazepine (9)
Citalopram (19)		Chloramphenical (9)
Diazepam (19)		Cimetadine (9)
Fluoxetine (9)		Fluconazole
Mephenytoin (19)		Fluoxetine (9-weak, 19-moderate)
NSAIDs (9)		Fluvoxamine (9/19)
Omeprazole (19)		Ketoconazole
Propranolol (partly 19)		Omeprazole (9/19)
Tricyclic antidepressants		Oxcarbazepine (19)
Tolbutamide (19)		Sertraline (9 – moderate)
Topiramate (19)		Topiramate
Warfarin (9/19)		Venlafaxine (weak)

Cytochrome P450 Drug Metabolism/Inhibition (*continued*)

Adapted from *Bazire and Benefield's Psychotropic Drug Directory 2001, Alden Press, Oxford, UK*

CYP2D6

Substrates	Significant enzyme inducers	Significant enzyme inhibitors
Amphetamines	Carbamazepine (weak)	Amiodarone
Antiarrhythmics Type 1C	Phenobarital (weak)	Chloroquine
Beta Blockers	Phenytoin (weak)	Chlorpromazine
Chlorpromazine	Rifampin (weak)	Cimetidine
Citalopram (minor)	Ritonavir (weak)	Citalopram (very weak)
Clozapine (minor)		Dextromethorphan
Codeine (to morphine)		Ditiazem (weak)
Desipramine (weak)		Diphenhydramine
Dextromethorphan		Fenfluramine ?)
Donepezil		Flecainide
Fluoxetine (partly)		Fluoxetine (strong)
Fluphenazine		Fluphenazine
Fluvoxamine (very weakly)		Fluvoxamine (very weak)
Haloperidol		Haloperidol
Hydrocodone		Methadone
Loratadine		Metoclopramide
Methadone		Metoprolol
Mirtazapine		Mirtazapine (very weak)
Nefazodone		Nefazodone (very weak)
Nicotine (partly)		Nicardipine
Olanzapine (partly)		Norfluoxetine (strong)
Oxycodone		Paroxetine (strong-dose related)
Paroxetine		Perphenazine
Perphenazine		Pindolol
Risperidone		Propranolol
Thioridazine		Quinidine (strong)
Timolol		Quinine
Trazodone		Sertraline (weak – doses <150mg/day)
Secondary and tertiary TCAs		Thioridazine
Venlafaxine (partly)		Timolol
		Tricyclics (all – strong)
		Venlafaxine (very weak)
		Yohimbine

Cytochrome P450 Drug Metabolism/Inhibition (*continued*)

Adapted from *Bazire and Benfield's Psychotropic Drug Directory 2001, Alden Press, Oxford, UK*

CYP3A4 (THE MOST IMPORTANT P450 ENZYME)

Substrates	Significant enzyme inducers	Significant enzyme inhibitors
Amiodarone	Barbiturates (all)	Acetazolamide
Benzodiazepines (not lorazepam)	Carbamazepine	Amiodarone
Buspirone	Cortisol	Cannabinoids
Calcium Channel Blockers	Dexamethasone	Cimetidine (moderate)
Cannabinoids	Ethosuximide	Citalopram (weak)
Carbamazepine	Oxcarbazepine	Clarithromycin
Cisapride (restricted use)	Phenytoin	Clotrimazole
Citalopram (major)	Prednisone	Diltiazem (weak)
Clindamycin	Rifampin	Fluconazole (strong)
Clozapine (partly)	Troglitazone	Fluoxetine (weak)
Cocaine		Fluvoxamine (moderate)
Codeine		Grapefruit juice (weaker)
Cortisol		Indinavir (moderate)
Cyclosporin		Itraconazole (strong)
Dapsone		Ketoconazole (strong)
Dexamethasone		Macrolides (erythromycin, clarithromycin)
Disopyramine		Metronidazole
Donepezil		Miconazole (strong)
Doxorubicin		Mirtazapine (very weak)
Estradiol		Nefazodone (strong)
Ethosuximide		Nelfinavir)
Fentanyl		Norfluoxetine (moderate)
Fluoxetine		Omeprazole (weak)
Flyburide		Paroxetine (weak)
Indinavir		Quinine
Itraconazole		Ritonavir (moderate)
Ketoconazole		Sertraline (minor/moderate – dose dependant)
Lidocaine		Trazodone
Lortadine		Tricyclics (moderate)
Lovastatin		Troleandomycin (strong)
Macrolides (not azithromycin)		Venlafaxine (weak)
Miconazole		Verapamil
Mirtazapine (partly)		
Nefazodone		
Omeprazole		
Ondansetron		
Pimozide (mostly)		
Pravastatin		
Prednisone		
Progesterone		
Quetiapine		
Quinidine		
Rifampin		
Risperidone		
Ritonavir		
Saquinavir		
Sertraline		
Testosterone		
Tamoxifen		
Tertiary Tricyclics		
Valproic acid		
Venlafaxine		
Vinblastine		
Vincristine		
Warfarin		
Zaleplon (secondary route)		
Ziprasidone (major)		
Zolpidem (mainly)		

CARBAMAZEPINE (TEGRETOL®)

INDICATIONS

- 1) Cyclic mood disorders
- 2) Aggressive behavior secondary to a psychiatric disorder
- 3) Chronic Pain
- 4) Acute mania

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction or similarly severe significant hypersensitivity to carbamazepine or tricyclic antidepressants

Relative:

- 1) History of blood dyscrasias
- 2) Myoclonic seizure, atonic seizures
- 3) AV heart block
- 4) History of bone marrow -suppression
- 5) Pregnancy/nursing mothers
- 6) Concomitant use of clozapine (Clozaril®)

Precautions

- 1) Diabetes Mellitus
- 2) SIADH
- 3) Glaucoma or urinary retention
- 4) Concomitant use of monoamine oxidase (MAO) inhibitors
- 5) Coronary artery disease
- 6) Hyponatremia, dilutional
- 7) Renal function impairment

Pregnancy and Breast-Feeding

See relative contraindications. FDA Pregnancy Category C.

Drug Interactions of Major Significance

- 1) Anticoagulants, coumadin
- 2) Anticonvulsants, hydantoin, succinimide, primidone, felbamate
- 3) Barbiturates
- 4) Benzodiazepines metabolized via hepatic microsomal enzymes, especially clonazepam
- 5) Valproic acid
- 6) Antidepressants, tricyclic
- 7) Cimetidine
- 8) Clarithromycin, erythromycin or troleandomycin
- 9) Contraceptives, estrogen-containing, oral
- 10) Estrogens, including estramustine
- 11) Quinidine
- 12) Corticosteroids
- 13) Calcium channel blockers (especially diltiazem and verapamil)
- 14) Isoniazid
- 15) Monoamine oxidase (MAO) inhibitors, including furazolidone and procarbazine
- 16) Propoxyphene
- 17) Antipsychotics (especially haloperidol and risperidone)
- 18) Nefazodone

SEE TABLE A: Cytochrome P450 Drug Metabolism/Inhibition

CARBAMAZEPINE (TEGRETOL®) - continued

PRECAUTIONS TO CONSIDER (continued)

Age-Specific considerations

None

Side Effects Which Require Medical Attention

- 1) Blurred or double vision
- 2) Rash
- 3) Sore throat or fever
- 4) Worsening confusion or disorientation
- 5) Nausea, vomiting diarrhea or abdominal discomfort
- 6) Drowsiness, lethargy
- 7) Headache
- 8) Bone or joint pain

PATIENT MONITORING

Patient Monitoring Parameters

- 1) CBC with platelets – baseline and 1 to 2 weeks after each dose increase and as clinically indicated
- 2) Hepatic function panel and electrolytes; baseline and as clinically indicated.
- 3) Pregnancy Test – as clinically indicated
- 4) Carbamazepine Levels – 3-4 weeks after dose adjustment, then as clinically indicated

Usual therapeutic levels 4-12 mcg/ml

Therapeutic ranges for the lab used should be listed on the report.

Dosing

Take with food to avoid stomach upset

See TDMHMR Formulary for dosage guidelines.

LAMOTRIGINE (LAMICTAL®)

INDICATIONS

- 1) Cyclic mood disorders

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction and similarly severe significant hypersensitivity to medication prescribed

Relative:

- 1) Pregnancy/nursing mothers
- 2) Age less than 16 years of age

Precautions

- 1) Combined use with Valproic acid
- 2) Photosensitivity
- 3) Renal or hepatic impairment

Pregnancy and Breast-Feeding

See relative contraindications. FDA Pregnancy Category C.

Drug Interactions of Major Significance

- 1) Valproic acid
- 2) Acetaminophen
- 3) Carbamazepine, phenytoin, phenobarbital

Age-Specific Considerations

- 1) Safety and efficacy in children<16 has not been established

Side Effects Which Require Medical Attention

- 1) Rash
- 2) Headache
- 3) Diplopia
- 4) Rhinitis
- 5) Nausea, vomiting, diarrhea or abdominal discomfort
- 6) Drowsiness, lethargy, dizziness

PATIENT MONITORING

Patient Monitoring Parameters

- 1) Renal Function Test - baseline and as clinically indicated
- 2) Hepatic Function Test - baseline, yearly and as clinically indicated
- 3) Pregnancy Test - as clinically indicated

Dosing

See TDMHMR Drug Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

LITHIUM (ESKALITH®, LITHOBID®, ESKALITH CR®, etc.)

INDICATIONS

- 1) Cyclic mood disorders
- 2) Augmentation of antidepressant therapy
- 3) Aggressive behavior secondary to a psychiatric disorder
- 4) Acute mania

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction or similarly severe significant hypersensitivity to the medication prescribed

Relative:

- | | |
|-----------------------------|---------------------------------|
| 1) Cardiovascular disease | 5) Pregnancy/nursing mothers |
| 2) Severe dehydration | 6) Renal insufficiency |
| 3) Goiter or hypothyroidism | 7) Hyperparathyroidism |
| 4) Psoriasis | 8) Concomitant use of diuretics |

Precautions

- | | |
|------------------------------------|----------------------|
| 1) Diagnosis of a seizure disorder | 6) Urinary retention |
| 2) Parkinson's disease | 7) Thyroid disorders |
| 3) Dehydration | 8) Psoriasis |
| 4) Severe infections | 9) Severe acne |
| 5) Dementia, brain injuries | |

Pregnancy and Breast-Feeding

See relative contraindications. FDA Pregnancy Category D.

Drug Interactions of Major Significance

- 1) Concomitant use of diuretics
- 2) Concomitant use of non-steroidal anti-inflammatory drugs
- 3) Iodine containing substances
- 4) Antipsychotics

Age-Specific Considerations

Monitoring of skeletal development in children if chronic lithium therapy is advised. Geriatric patients usually require lower doses and more frequent monitoring.

PRECAUTIONS TO CONSIDER

Side Effects Which Require Medical Attention

- 1) Weight gain
- 2) Edema
- 3) Hypothyroidism
- 4) Slurred speech
- 5) Drowsiness, lethargy
- 6) Nausea, vomiting, diarrhea
- 7) Ataxia
- 8) Trembling
- 9) Polydipsia
- 10) Polyuria
- 11) Headache

PATIENT MONITORING

Patient Monitoring Parameters

- 1) EKG (mandatory for everyone – baseline, yearly and as clinically indicated)
- 2) CBC – baseline, yearly and as clinically indicated
- 3) Thyroid studies – baseline; then TSH every 6 months and as clinically indicated
- 4) BUN, creatinine, glucose and electrolytes; baseline and as clinically indicated.
- 5) UA - baseline and as clinically indicated
- 6) Pregnancy Test - as clinically indicated
- 7) Lithium Levels – one week after initiation or dosage change and as clinically indicated
- 8) Calcium and phosphate, in children under 12, prior to initiation and as clinically indicated

Usual therapeutic Level: 0.6-1.5 meq/L (12 hour post dose)

Therapeutic ranges for the lab used should be listed on the report.

Dosing

Take with food to avoid stomach upset

See TDMHMR Drug Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

OXCARBAZEPINE (TRILEPTAL®)

INDICATIONS

- 1) Cyclic mood disorders

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction or similarly severe significant hypersensitivity to carbamazepine or oxcarbazepine

Relative:

- 1) Hyponatremia

Precautions

- 1) Renal insufficiency

Pregnancy and Breast-Feeding

See relative contraindications. FDA Pregnancy Category C.

Drug Interactions of Major Significance

- 1) Oral contraceptives
- 2) Phenytoin

Age-Specific considerations

None

Side Effects Which Require Medical Attention

- 1) Psychomotor retardation
- 2) Concentration difficulties
- 3) Somnolence and fatigue
- 4) Ataxia, gait disturbances
- 5) Nausea, vomiting
- 6) Signs and symptoms of hyponatremia

PATIENT MONITORING

Patient Monitoring Parameters

- 1) Electrolytes – baseline and as clinically indicated
- 2) Pregnancy Test – as clinically indicated

Dosing

See TDMHMR Formulary for dosage guidelines.

TOPIRAMATE (TOPAMAX®)

INDICATIONS

- 1) Cyclic mood disorders

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction and similarly severe significant hypersensitivity to medication prescribed

Relative:

- 1) Pregnancy/nursing mothers

Precautions

- 1) Kidney stones
- 2) Parasthesias
- 3) Renal or hepatic impairment

Pregnancy and Breast-Feeding

See relative contraindications. FDA Pregnancy Category C.

Drug Interactions of Major Significance

- 1) oral contraceptives
- 2) carbonic anhydrase inhibitors
- 3) valproic acid, carbamazepine, phenytoin

Age-Specific Considerations

- 1) Safety and efficacy in children have not been established

Side Effects Which May Require Medical Attention

- | | |
|-------------------------------|-------------------------------------|
| 1) Psychomotor slowing | 6) Nervousness |
| 2) Headache | 7) Drowsiness, lethargy, somnolence |
| 3) Diplopia | 8) Depression |
| 4) Dizziness | 9) Weight loss |
| 5) Nausea, vomiting, anorexia | |

PATIENT MONITORING

Patient Monitoring Parameters

- 1) Renal Function Test - baseline, and as clinically indicated
- 2) Hepatic Function Test - baseline, and as clinically indicated
- 3) Pregnancy Test - as clinically indicated

Dosing

See TDMHMR Drug Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

VALPROIC ACID (DEPAKENE®), DIVALPROEX SODIUM (DEPAKOTE®)

INDICATIONS

- 1) Cyclic mood disorders
- 2) Aggressive behavior secondary to a psychiatric disorder
- 3) Chronic Pain
- 4) Acute mania

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction or similarly severe significant hypersensitivity to the medication prescribed

Relative

- 1) Hepatic disease/impairment
- 2) Blood dyscrasias, clotting disorders or concomitant drugs that alter clotting function (aspirin, non-steroidal anti-inflammatory drugs, warfarin, heparin, low molecular weight heparins, clopidogrel etc.)
- 3) Pregnancy/nursing mothers

Precautions

- 1) Hypoalbuminemia
- 2) Renal impairment

Pregnancy and Breast-Feeding

See relative contraindications. FDA Pregnancy Category D.

Drug Interactions of Major Significance

- | | |
|--|--|
| 1) Concomitant CNS depressants | 7) Phenytoin |
| 2) Anticoagulants | 8) Lamotrigine |
| 3) Carbamazepine | 9) Non-steroidal anti-inflammatory drugs |
| 4) Felbamate | 10) Aspirin |
| 5) Concomitant hepatotoxic medications | 11) Phenobarbital |
| 6) Mefloquine | 12) Diazepam |

Age-specific Considerations

Age younger than 10 years old due to high risk of hepatic toxicity

Geriatric patients have increased amounts of free drug (use lower total plasma concentration or get free VPA plasma concentration)

Side Effects Which Require Medical Attention

- 1) Worsening confusion or disorientation
- 2) Nausea, vomiting, diarrhea, abdominal discomfort or anorexia
- 3) Bruising or bleeding
- 4) Clinically significant weight gain
- 5) Tremors

VALPROIC ACID (DEPAKENE®), DIVALPROEX SODIUM (DEPAKOTE®) - continued

PATIENT MONITORING

Patient Monitoring Parameters

- 1) CBC - with differential and platelet count - baseline then one (1) to two (2) weeks after each dosage increase, and as clinically indicated
- 2) Hepatic function panel- baseline and as clinically indicated
- 3) Pregnancy Test – baseline and as clinically indicated
- 4) Valproic acid level – 1-2 weeks after initiation and dosage change, then as clinically indicated.
- 5) Serum creatinine and BUN at baseline and as clinically indicated

Usual therapeutic levels 50-150 mcg/ml

Therapeutic ranges for the lab used should be listed on the report.

Dosing

Take with food to avoid stomach upset

See TDMHMR Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

VERAPAMIL (CALAN® , ISOPTIN®)

INDICATIONS

- | |
|--------------------------|
| 1) Cyclic mood disorders |
|--------------------------|

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction and similarly severe significant hypersensitivity to medication prescribed
- 2) Severe left ventricular dysfunction
- 3) Hypotension (systolic pressure < 90 mm Hg) or cardiogenic shock
- 4) Sick sinus syndrome (except in patients with a functioning artificial ventricular pacemaker)
- 5) Second or third AV block (except in patients with a functioning artificial ventricular pacemaker)
- 6) Patients with atrial flutter or atrial fibrillation and an accessory bypass tract (Wolff-Parkinson-White)

Relative:

- 1) Hepatic function impairment
- 2) Renal impairment
- 3) Cardiac conduction disturbances not outlined in absolute contraindications

Precautions

Pregnancy/nursing mothers

Pregnancy and Breast-Feeding

See precautions. FDA Pregnancy Category C.

Drug Interactions of Major Significance

- 1) Carbamazepine
- 2) Lithium

SEE TABLE A: Cytochrome P450 Drug Metabolism/Inhibition

Age-Specific Considerations

Safety and efficacy not established for those under 18 years old.

Side Effects Which Require Medical Attention

- | | |
|-----------------|-----------------------------------|
| 1) Hypotension | 4) Headache, chronic or recurrent |
| 2) Constipation | 5) Dizziness or lightheadedness |
| 3) Nausea | 6) Bradycardia |

PATIENT MONITORING

Patient Monitoring Parameters

- 1) Pregnancy Test - as clinically indicated
- 2) Renal and Liver Function Test - baseline and as clinically indicated
- 3) Vital signs with initial dosing and with any dosage change.
- 4) EKG within one year prior to initiation of drug.

Dosing

See TDMHMR Drug Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

BENZODIAZEPINES

alprazolam (Xanax®), chlordiazepoxide (Librium®), clorazepate (Tranxene®), diazepam (Valium®), lorazepam (Ativan®), Oxazepam (Serax®), temazepam (Restoril®), triazolam (Halcion®), Clonazepam (Klonopin®)

INDICATIONS

- | | |
|---|---|
| 1) Anxiety disorders | 6) Akathisia |
| 2) Panic disorder | 7) Acute intervention for agitation/violent behavior |
| 3) Anxiety associated with depression | 8) Bipolar disorder, mania
- adjunctive or second line therapy |
| 4) Short term use for the treatment of insomnia | 9) Alcohol/substance abuse withdrawal |
| 5) Sedative hypnotic withdrawal | |

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction or similarly severe significant hypersensitivity to the medication prescribed

Relative:

- 1) Pregnancy/nursing mothers
- 2) Myasthenia gravis
- 3) Severe COPD

Precautions

- | | |
|--|--|
| 1) Hepatic impairment | 6) Discontinuation or rapid dose reduction |
| 2) Porphyria | 7) Attention Deficit Hyperactivity Disorder (ADHD) |
| 3) History of alcohol and drug abuse | 8) Dementias/delirium |
| 4) Sleep apnea | |
| 5) Sedative hypnotic intoxication/dependence | |

Pregnancy and Breast-Feeding

See relative contraindications. Most benzodiazepines are FDA Pregnancy Category D or X.

Drug Interactions of Major Significance

- 1) Alcohol or CNS depressants
- 2) Clozapine (excessive sedation or respiratory depression)
- 3) Other drugs with respiratory depression

SEE TABLE A: Cytochrome P450 Drug Metabolism/Inhibition

Age-Specific Considerations

- 1) Lower doses should be used in children and elderly
- 2) Avoid long half-life drugs in elderly (causes falls)
- 3) May cause excitability in children, elderly and persons with developmental disabilities

Side Effects Which Require Medical Attention

- 1) Worsening agitation, disinhibition or aggression
- 2) Obtundation
- 3) Ataxia
- 4) Redness, swelling or pain at injection site

BENZODIAZEPINES - continued

alprazolam (Xanax®), chlordiazepoxide (Librium®), clorazepate (Tranxene®), diazepam (Valium®), lorazepam (Ativan®), Oxazepam (Serax®), temazepam (Restoril®), triazolam (Halcion®), Clonazepam (Klonopin®)

PATIENT MONITORING

Patient Monitoring Parameters

- 1) Pregnancy Test - as clinically indicated

Dosing

See TDMHMR Drug Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

BUSPIRONE (BUPPAR®)

INDICATIONS

- | | |
|------------------------|--|
| 1) Anxiety Disorder | 3) Self injurious behavior |
| 2) Aggressive behavior | 4) Augmentation for resistant depression |

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction or similarly severe significant hypersensitivity to the medication prescribed

Relative:

- 1) Hepatic function impairment
- 2) Renal impairment

Precautions

- 1) Pregnancy/nursing mothers
- 2) Doses > 45 mg/day in patients with developmental disabilities

Pregnancy and Breast-Feeding

See precautions. FDA Pregnancy Category B.

Drug Interactions of Major Significance

- 1) Concomitant monoamine oxidase (MAO) inhibitors
- 2) Erythromycin
- 3) Itraconazole
- 4) Nefazodone

Age-Specific Considerations

Safety and efficacy not established for those under 18 years old.

Side Effects Which Require Medical Attention

- 1) Worsening confusion or agitation
- 2) Restlessness (akathisia)
- 3) Nausea
- 4) Headache, chronic or recurrent
- 5) Dizziness or lightheadedness

PATIENT MONITORING

Patient Monitoring Parameters

- 1) Pregnancy Test - as clinically indicated

Dosing

See TDMHMR Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

AMOXAPINE (ASENDIN®)

INDICATIONS

- 1) Depression with psychotic features

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction or similarly severe significant hypersensitivity to the medication prescribed
- 2) Recovery phase of myocardial infarction

Relative:

- 1) Pregnancy/nursing mothers
- 2) History of neuroleptic malignant syndrome

Precautions

Alcohol intoxication, bipolar disorder in the absence of a mood stabilizer, recent or current blood dyscrasias, cardiovascular disorders including arrhythmia, heart block and failure, patients at risk for paralytic ileus, glaucoma, hepatic function impairment, hyperthyroidism, prostatic hypertrophy, renal failure, diagnosis of a seizure disorder, urinary retention, Parkinson's disease, tardive dyskinesia or history of EPS.

Pregnancy and Breast-Feeding

See relative contraindications. FDA Pregnancy Category C.

Drug Interactions of Major Significance

None

Age-Specific Considerations

- 1) Not indicated in children under age 16

Side Effects Which Require Medical Attention

- 1) Anticholinergic effects
- 2) Sexual function impairment
- 3) Seizures
- 4) Dizziness, lightheadedness or fainting
- 5) EPS
- 6) Akathisia
- 7) Tardive dyskinesia
- 7) Symptoms of prolactin elevation (galactorrhea, amenorrhea, gynecomastia)
- 8) Signs and symptoms of neuroleptic malignant syndrome

PATIENT MONITORING

Patient Monitoring Parameters

- 1) EKG – baseline and as clinically indicated.
- 2) Pregnancy test - as clinically indicated.
- 3) Screening for abnormal involuntary movements using a standardized test-prior to initiation, six months, annually and as clinically indicated.

Dosing

See TDMHMR Drug Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

BUPROPION (WELLBUTRIN® and WELLBUTRIN® SR)

INDICATIONS

- 1) Depressive Disorders
- 2) Attention deficit hyperactivity disorder
- 3) Nicotine Dependence

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction or similarly severe significant hypersensitivity to the medication prescribed
- 2) Anorexia nervosa and bulimia
- 3) Diagnosis of a seizure disorder

Relative:

None

Precautions

Bipolar disorder in the absence of a mood stabilizer, hepatic function impairment, renal failure, CNS tumor, head trauma or history of seizures, pregnancy/nursing mothers.

Pregnancy and Breast-Feeding

See precautions. FDA Pregnancy Category B.

Drug Interactions of Major Significance

- 1) Concomitant monoamine oxidase inhibitors (furazolidone, procarbazine, selegiline, tranylcypromine, phenelzine, isoniazid)
- 2) Concomitant use of CNS depressants
- 3) Cimetidine
- 4) Concomitant use of medications with anticholinergic effects
- 5) Noradrenergic anti-hypertensive agents (clonidine, guanabenz, guanadrel, guanethidine)
- 6) Concomitant use of drugs that lower seizure threshold

Age-Specific Considerations

None

Side Effects Which Require Medical Attention

- 1) Delirium or cognitive dysfunction
- 2) Seizure
- 3) Headache, severe
- 4) Restlessness

PATIENT MONITORING

Patient Monitoring Parameters

- 1) Pregnancy test - as clinically indicated.

Dosing

See TDMHMR Drug Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

MIRTAZAPINE (REMERON®)

INDICATIONS

- 1) Depressive Disorders
- 2) Insomnia

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction and similarly severe significant hypersensitivity to medication prescribed

Relative:

- 1) Pregnancy/nursing mothers

Precautions

Bipolar disorder in the absence of mood stabilizer, recovery phase of myocardial infarction, hepatic function impairment, renal failure, diagnosis of a seizure disorder

Pregnancy and Breast-Feeding

See relative contraindications. FDA Pregnancy Category C.

Drug Interactions of Major Significance

Concomitant monoamine oxidase inhibitors (or within 14 days of an MAOI)

Age-Specific Considerations

No data available in individuals under the age of 18

Side Effects Which Require Medical Attention

- 1) Agranulocytosis (or signs of infection: sore throat, fever, etc.)
- 2) Elevated cholesterol
- 3) Dizziness, unsteadiness, lightheadedness or fainting
- 4) Increased weight gain/appetite
- 5) Elevation in liver enzymes (ALT)
- 6) Orthostatic hypotension

PATIENT MONITORING

Patient Monitoring Parameters

- 1) Pregnancy Test - as clinically indicated

Dosing

See TDMHMR Drug Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

MONOAMINE OXIDASE INHIBITORS

phenelzine (Nardil®), tranylcypromine (Parnate®)

INDICATIONS

- 1) Depressive Disorders
- 2) Panic Disorders
- 3) Anxiety Disorders

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction or similarly severe significant hypersensitivity to the medication prescribed
- 2) Pheochromocytoma
- 3) Congestive heart failure
- 4) Concomitant use of another monoamine oxidase inhibitor
- 5) Concomitant use with meperidine
- 6) Concomitant use with SSRI's, buspirone or venlafaxine
- 7) Concomitant use of pressor amines (e.g. ephedrine, phenylpropanolamine, pseudoephedrine)
- 8) Stimulants

Relative:

- 1) Impaired renal function
- 2) Severe hepatic disease
- 3) Pregnancy/nursing mothers
- 4) Hyperthyroidism
- 5) Concomitant use of tricyclic antidepressant, methyldopa, dopamine, levodopa, selegiline, dextromethorphan

Precautions

Bipolar disorder in the absence of a mood stabilizer, hepatic function impairment, renal function impairment, hypertension or history of hypertension, diagnosis of a seizure disorder or history of seizures, recent cardiac disease including myocardial infarction, concomitant use of antihypertensives.

Pregnancy and Breast-Feeding

See relative contraindications. Phenelzine and tranylcypromine are FDA Pregnancy Category C.

Age-Specific Considerations

No data available in individuals under the age of 18.

Side Effects Which Require Medical Attention

- 1) Headache
- 2) Sexual dysfunction
- 3) Blood pressure alteration, especially hypertension
- 4) Delirium
- 4) Dizziness, lightheadedness or fainting (orthostatic hypotension)
- 5) Clinically significant weight gain

MONOAMINE OXIDASE INHIBITORS - continued
phenelzine (Nardil®), tranylcypromine (Parnate®)

PRECAUTIONS TO CONSIDER (continued)

Drug Interactions of Major Significance

See absolute and relative contraindications

FOODS CONTAINING TYRAMINE

****High Amounts of Tyramine***

Smoked, aged or pickled meat or fish
Sauerkraut
Aged Cheeses such as Swiss and Cheddar
Yeast extracts
Fava beans

*****Moderate Amounts of Tyramine***

Beer
Avocados
Meat extracts
Red wines such as Chianti

******Low Amounts of Tyramine***

Caffeine-containing beverages
Distilled spirits
Chocolate
Soy sauce
Cottage and cream cheeses
Yogurt and sour cream

*May not consume

**May consume in moderation

***May consume

Adapted from Shulman KI, et al. Dietary restriction, tyramine, and the use of monoamine oxidase inhibitors. J Clin Psychopharmacol. 1989; 9: 397.

PATIENT MONITORING

Patient Monitoring Parameters

- 1) Blood chemistries with emphasis on hepatic and renal functions; baseline, yearly and as clinically indicated during prolonged or high dose therapy.
- 2) Pregnancy test – as clinically indicated.
- 3) Blood pressure at baseline and during dosage adjustments and as clinically indicated. Therapeutic ranges for the lab used should be listed on the report.

Dosing

See TDMHMR Drug Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

NEFAZODONE (SERZONE®)

INDICATIONS

- 1) Depressive Disorders

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction or similarly severe significant hypersensitivity to the medication prescribed or trazodone.
- 2) Patients who were withdrawn from nefazodone because of evidence of liver injury.

Relative:

- 1) Pregnancy/nursing mothers

Precautions

- 1) Bipolar disorder in the absence of a mood stabilizer, recovery phase of myocardial infarction, hepatic function impairment, renal failure, diagnosis of a seizure disorder.
- 2) Active liver disease, elevated baseline serum transaminases.

Pregnancy and Breast-Feeding

See relative contraindications. FDA Pregnancy Category C.

Drug Interactions of Major Significance

- 1) Concomitant monoamine oxidase inhibitors (or within 14 days of an MAOI)
- 2) Alprazolam, triazolam
- 3) Carbamazepine
- 4) Pimozide

SEE TABLE A: Cytochrome P450 Drug Metabolism/Inhibition

Age-Specific Considerations

No data available in individuals under the age of 18.

Side Effects Which Require Medical Attention

- | | |
|--|--|
| 1) Priapism | 5) Tinnitus |
| 2) Dizziness, unsteadiness,
lightheadedness or fainting | 6) Bradycardia less than 50 beats/min. |
| 3) Visual changes | 7) Elevated liver function tests |
| 4) Rash | 8) Dark urine, anorexia, malaise and GI symptoms |

PATIENT MONITORING

Patient Monitoring Parameters

- 1) Pregnancy test – as clinically indicated.
- 2) ALT and AST – baseline, at 1, 2, 4, 6 and 12 months, then annually and as clinically indicated. Stop drug if AST or ALT levels are 3 times (or greater) the upper limit of normal.

Dosing

See TDMHMR Drug Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

SSRIs: CITALOPRAM (CELEX®), FLUOXETINE (PROZAC®), SERTRALINE (ZOLOFT®), PAROXETINE (PAXIL®), FLUVOXAMINE (LUVOX®)

INDICATIONS

- | | |
|----------------------------------|-------------------------------|
| 1) Depressive Disorders | 5) Self injurious behavior |
| 2) Obsessive-compulsive disorder | 6) Late Luteal Phase Disorder |
| 3) Panic disorder | 7) Anxiety Disorders |
| 4) Eating disorders | 8) Social Phobia |

PRECAUTIONS TO CONSIDER

Contraindications

Absolute

- 1) History of anaphylactic reaction or similarly severe significant hypersensitivity to the medication prescribed
- 2) Concurrent administration of MAOI (or within 14 days of receiving citalopram, sertraline, paroxetine or fluvoxamine; or within 35 days of receiving fluoxetine)

Relative

- | | |
|---------------------------------------|--|
| 1) Severe hepatic function impairment | 3) Seizure disorder or history of seizure disorder |
| 2) Severe renal function impairment | |

Precautions

Bipolar disorder in the absence of a mood stabilizer, hepatic function impairment, renal failure, diagnosis of a seizure disorder, diabetes mellitus, pregnancy/nursing mothers.

Pregnancy and Breast-Feeding

See precautions. FDA Pregnancy Category B

Drug Interactions of Major Significance

- | | |
|---------------------------------------|-------------------------------------|
| 1) See contraindications | 5) Valproic acid, divalproex sodium |
| 2) Alcohol | 6) Tolbutamide |
| 3) Concomitant use of CNS depressants | 7) Carbamazepine |
| 4) Phenytoin | |

SEE TABLE A: Cytochrome P450 Drug Metabolism/Inhibition

Age-Specific Considerations

None

Side Effects Which Require Medical Attention

- | | |
|-------------------------|-------------------------------|
| 1) Chills or fever | 6) Sexual function impairment |
| 2) Joint or muscle pain | 7) Severe GI distress |
| 3) Skin rash | 8) Akathisia |
| 4) Hives or itching | 9) Hyponatremia |
| 5) Trouble breathing | 10) Hypothyroidism |

PATIENT MONITORING

Patient Monitoring Parameters

- 1) Pregnancy test - as clinically indicated.

Dosing

See TDMHMR Drug Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

TRAZODONE (DESYREL®)

INDICATIONS

- 1) Depressive Disorders
- 2) Insomnia

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction or similarly severe significant hypersensitivity to the medication prescribed or nefazodone
- 2) Recovery phase of myocardial infarction

Relative:

- 1) Pregnancy/nursing mothers

Precautions

Alcohol intoxication, bipolar disorder in the absence of a mood stabilizer, cardiovascular disorders including arrhythmia, heart block and failure, hepatic function impairment, renal failure, diagnosis of a seizure disorder, hypotension and history of priapism.

Pregnancy and Breast-Feeding

See relative contraindications. FDA Pregnancy Category C.

Drug Interactions of Major Significance

None

Age-Specific Considerations

Has not been studied in persons younger than 18.

Side Effects Which Require Medical Attention

- 1) Priapism
- 2) Dizziness, lightheadedness or fainting

PATIENT MONITORING

Patient Monitoring Parameters

- 1) EKG - as clinically.
- 2) Pregnancy test - as clinically indicated.

Dosing

See TDMHMR Drug Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

TRICYCLIC ANTIDEPRESSANTS

amitriptyline (Elavil®), desipramine (Norpramin®, Pertofrane®), doxepin (Sinequan®), imipramine (Tofranil®), maprotiline (Ludiomil®), nortriptyline (Pamelor®, Aventyl®), protriptyline (Vivactil®), trimipramine (Surmontil®)

INDICATIONS

- | | |
|---|----------------------------------|
| 1) Depressive Disorders | 6) Anxiety disorders |
| 2) Panic disorder | 7) Chronic Pain |
| 3) Bulimia nervosa | 8) Insomnia |
| 4) Attention deficit hyperactivity disorder | 9) Obsessive-Compulsive Disorder |
| 5) Functional enuresis | |

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction or similarly severe significant hypersensitivity to the medication prescribed
- 2) Recovery phase of myocardial infarction
- 3) Pheochromocytoma

Relative:

- 1) Pregnancy/nursing mothers

Precautions

Alcohol intoxication, bipolar disorder in the absence of a mood stabilizer, recent or current blood dyscrasias, cardiovascular disorders including arrhythmia, diseases states where increased anticholinergic activity may complicate disease course (narrow-angle glaucoma, benign prostatic hypertrophy), heart block and failure, lower respiratory tract symptoms (asthma), , hepatic function impairment, hyperthyroidism, renal failure, diagnosis of a seizure disorder.

Pregnancy and Breast-Feeding

See relative contraindications. Most tricyclic antidepressants are FDA Pregnancy Category C or D.

Drug Interactions of Major Significance

- 1) Concomitant monoamine oxidase inhibitors (furazolidone, procarbazine, selegiline, tranylcypromine, phenelzine, isoniazid)
- 2) Concomitant use of CNS depressants
- 3) Cimetidine
- 4) Concomitant use of medications with anticholinergic effects
- 5) Noradrenergic anti-hypertensive agents (clonidine, guanabenz, guanadrel, guanethidine)
- 6) SSRI

SEE TABLE A: Cytochrome P450 Drug Metabolism/Inhibition

Age-Specific Considerations

Most agents are not recommended for use in children; if used, conservative dosing, EKG prior to dosage increase and plasma concentration monitoring are advised.

TRICYCLIC ANTIDEPRESSANTS

- continued

amitriptyline (Elavil®), desipramine (Norpramin®, Pertofrane®), doxepin (Sinequan®), imipramine (Tofranil®), maprotiline (Ludiomil®), nortriptyline (Pamelor®, Aventyl®), protriptyline (Vivactil®), trimipramine (Surmontil®)

PRECAUTIONS TO CONSIDER (continued)Side Effects Which Require Medical Attention

- 1) Anticholinergic effects
- 2) Sexual function impairment
- 3) Seizures
- 4) Dizziness, lightheadedness or fainting (orthostatic hypotension)
- 5) Tachycardia greater than 100 beats/min
- 6) Jaundice
- 7) QTc >500 msec

PATIENT MONITORINGPatient Monitoring Parameters

- 1) EKG – baseline and as clinically indicated.
- 2) Pregnancy test - as clinically indicated.
- 3) Blood levels as clinically indicated. Therapeutic ranges for the lab used should be listed on the report. See Antidepressant Table in current TDMHMR Drug Formulary Book.

Dosing

See TDMHMR Drug Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

VENLAFAXINE (EFFEXOR® and EFFEXOR® ER)

INDICATIONS

- | | |
|---|----------------------|
| 1) Depressive Disorders | 3) Anxiety disorders |
| 2) Attention deficit hyperactivity disorder | 4) Chronic Pain |

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction or similarly severe significant hypersensitivity to the medication prescribed
- 2) Pheochromocytoma
- 3) Concomitant use of monoamine oxidase inhibitor

Relative:

- | | |
|------------------------------|--|
| 1) Impaired renal function | 4) Hyperthyroidism |
| 2) Severe hepatic disease | 5) Hypertension or history of hypertension |
| 3) Pregnancy/nursing mothers | |

Precautions

Bipolar disorder in the absence of a mood stabilizer, hepatic function impairment, renal function impairment, diagnosis of a seizure disorder or history of seizures.

Pregnancy and Breast-Feeding

See relative contraindications. FDA Pregnancy Category C.

Drug Interactions of Major Significance

- 1) Concurrent administration of MAOI, or within 14 days of MAOI
- 2) SSRIs or other serotonergic drugs

SEE TABLE A: Cytochrome P450 Drug Metabolism/Inhibition

Age-specific Considerations

No data available in individuals under the age of 18.

Side Effects Which Require Medical Attention

- | | |
|--|------------------------|
| 1) Headache | 4) Delirium |
| 2) Sexual dysfunction | 5) Visual disturbances |
| 3) Blood pressure alteration,
especially hypertension | |

PATIENT MONITORING

Patient Monitoring Parameters

- 1) Pregnancy test – as clinically indicated.
- 2) Blood pressure during dosage titration and as clinically necessary.

Dosing

See TDMHMR Drug Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

ANTIPSYCHOTICS

chlorpromazine (Thorazine®), fluphenazine (Prolixin®), haloperidol (Haldol®), loxapine (Loxitane®), molindone (Molan®), perphenazine (Trilafon®), thiothixene (Navane®), trifluoperazine (Stelazine®)

INDICATIONS

- 1) Disorders with psychotic symptoms (schizophrenia, schizoaffective disorder, manic disorders, depression with psychotic features, drug-induced psychosis, psychosis associated with other organic conditions)
- 2) Tourette's disorder (haloperidol only)
- 3) Personality disorders – schizotypal, paranoid and borderline
- 4) Acute and/or short term use for management of aggressive or violent behavior
- 5) Stereotypies

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction and similarly severe significant hypersensitivity to medication prescribed or structurally related medication
- 2) Severe CNS depression

Relative:

- 1) Pregnancy/nursing mothers
- 2) History of drug-induced agranulocytosis or leukopenia
- 3) Breast cancer
- 4) History of neuroleptic malignant syndrome
- 5) Narrow angle glaucoma (for chlorpromazine)
- 6) Impaired hepatic function
- 7) Prostatic hypertrophy (for chlorpromazine)
- 8) Parkinson's disease
- 9) Severe cardiovascular diseases, including certain conduction disturbances

Precautions

Alcoholism (active), recent or current blood dyscrasias, hepatic function impairment, angina, hypotension, congestive heart failure, arrhythmias, breast cancer, glaucoma, history of neuroleptic malignant syndrome, benign prostatic hypertrophy, Parkinson's disease, poorly controlled seizure disorder, urinary retention, patients at risk for paralytic ileus.

Pregnancy and Breast-Feeding

See relative contraindications. Most antipsychotics are FDA Pregnancy Category C.

ANTIPSYCHOTICS - continued

chlorpromazine (Thorazine®), fluphenazine (Prolixin®), haloperidol (Haldol®), loxapine (Loxitane®), molindone (Molan®), perphenazine (Trilafon®), thiothixene (Navane®), trifluoperazine (Stelazine®)

PRECAUTIONS TO CONSIDER (continued)

Drug Interactions of Major Significance

- 1) Concomitant use of CNS depressants
- 2) Antithyroid agents
- 3) Concomitant use of agents that cause EPS (including droperidol metoclopramide, amoxapine, metyrosine, pimozide, reserpine)
- 4) Concomitant use of hypotension producing agents
- 5) Levodopa
- 6) Concomitant anticholinergic drugs (for chlorpromazine)

SEE TABLE A: Cytochrome P450 Drug Metabolism/Inhibition

Age-Specific Considerations

- 1) Conservative dosing and careful monitoring are advised in children and the elderly

Side Effects Which Require Medical Attention

- 1) Anticholinergic effects (for chlorpromazine)
- 2) Acute extrapyramidal side effects (dystonia, pseudo-Parkinsonism)
- 3) Akathisia
- 4) Hypotension
- 5) Rashes, photosensitivity and altered pigmentation
- 6) Tardive dyskinesia or other late-onset EPS
- 7) Visual changes
- 8) Early symptoms of agranulocytosis effects (fever, sore throat, weakness)
- 9) Fluctuating vital signs
- 10) Altered consciousness
- 11) Galactorrhea
- 12) Amenorrhea
- 13) Gynecomastia
- 14) Poikilothermia
- 15) Signs and symptoms of neuroleptic malignant syndrome

PATIENT MONITORING

Patient Monitoring Parameters

- 1) Pregnancy Test - as clinically indicated
- 2) Screening for abnormal involuntary movements using a standardized test - prior to initiation, six months, annually and as clinically indicated
- 3) Eye exams (prolonged or high-dose therapy with chlorpromazine dose of 2,000mg/day) yearly or as indicated
- 4) EKG - as clinically indicated.

Dosing

See TDMHMR Drug Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

ANTIPSYCHOTICS

mesoridazine (Serentil®), thioridazine (Mellaril®)

INDICATIONS

- 1) Schizophrenia, refractory (failed other classes of antipsychotics)

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction and similarly severe significant hypersensitivity to medication prescribed or structurally related medication
- 2) Severe CNS depression
- 3) QTc > 450 msec
- 4) Hypomagnesemia
- 5) Hypokalemia
- 6) Retinitis Pigmentosa
- 7) Concomitant use of other drugs known to prolong QTc interval

Relative:

- 1) Pregnancy/nursing mothers
- 2) History of drug induced agranulocytosis or leukopenia
- 3) Breast cancer
- 4) History of neuroleptic malignant syndrome
- 5) Narrow angle glaucoma
- 6) Impaired hepatic function
- 7) Prostatic hypertrophy
- 8) Parkinson's disease
- 9) Severe cardiovascular diseases, including certain conduction disturbances

Precautions

Alcoholism (active), recent or current blood dyscrasias, hepatic function impairment, angina, hypotension, congestive heart failure, arrhythmias, breast cancer, glaucoma, history of neuroleptic malignant syndrome, prostatic hypertrophy, Parkinson's disease, poorly controlled seizure disorder, urinary retention, patients at risk for paralytic ileus.

Pregnancy and Breast-Feeding

See relative contraindications. Most antipsychotics are FDA Pregnancy Category C.

ANTIPSYCHOTICS - continued

mesoridazine (Serentil®), thioridazine (Mellaril®)

PRECAUTIONS TO CONSIDER (continued)

Drug Interactions of Major Significance

- 1) Concomitant use of CNS depressants
- 2) Antithyroid agents
- 3) Concomitant use of agents that cause EPS (including droperidol, metoclopramide, amoxapine, metyrosine, pimozide, reserpine)
- 4) Concomitant use of hypotension producing agents
- 5) Levodopa
- 6) Concomitant anticholinergic drugs

SEE TABLE A: Cytochrome P450 Drug Metabolism/Inhibition

Age-Specific Considerations

- 1) Conservative dosing and careful monitoring are advised in children and the elderly

Side Effects Which Require Medical Attention

- 1) Anticholinergic effects
- 2) Acute extrapyramidal side effects (akathisia, dystonia, pseudo-Parkinsonism)
- 3) Hypotension
- 4) Rashes, photosensitivity and altered pigmentation
- 5) Tardive dyskinesia or other late-onset EPS
- 6) Visual changes
- 7) Early symptoms of agranulocytosis effects (fever, sore throat, weakness)
- 8) Fluctuating vital signs
- 9) Altered consciousness
- 10) Galactorrhea
- 11) Amenorrhea
- 12) Gynecomastia

PATIENT MONITORING

Patient Monitoring Parameters

- 1) Pregnancy Test - as clinically indicated
- 2) Screening for abnormal involuntary movements using a standardized test - prior to initiation, six months, annually and as clinically indicated
- 3) Eye exams (prolonged or high-dose therapy with thioridazine dose if 800 mg/day) yearly or as indicated
- 4) EKG - baseline; 7-14 days after dosage change; 7-14 days after other medication changes that could significantly alter the cardiac effects of mesoridazine and thioridazine; every six months, and as clinically indicated
- 5) Serum potassium level – baseline, every six months, and as clinically indicated
- 6) Serum magnesium level – baseline and as clinically indicated (especially if potassium level is low)

Dosing

See TDMHMR Drug Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

CLOZAPINE (CLOZARIL®)

INDICATIONS

- 1) For use in patients with refractory schizophrenia or schizoaffective disorder, defined as failure on two antipsychotics from two different chemical families given for sufficient time (6-12 weeks) at a sufficient dose (1000 mg/day of chlorpromazine equivalents).
- 2) For use in schizophrenic or schizoaffective patients who cannot tolerate other antipsychotics.
- 3) Psychosis associated with other organic conditions, (who have failed two antipsychotics, or who cannot tolerate other antipsychotics)
- 4) Manic disorders with psychosis (in patients who have failed two antipsychotics)

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction or similarly severe significant hypersensitivity to the medication prescribed
- 2) Myeloproliferative disorders
- 3) History of clozapine-induced agranulocytosis

Relative:

- 1) History of drug induced agranulocytosis or leukopenia
- 2) Breast cancer
- 3) History of neuroleptic malignant syndrome
- 4) Narrow angle glaucoma
- 5) Impaired hepatic function
- 6) Prostatic hypertrophy
- 7) Parkinson's disease
- 8) Severe cardiovascular diseases
- 9) Concomitant use of agents that may cause bone marrow suppression, including carbamazepine (Tegretol®)

Precautions

Alcoholism (active), recent or current blood dyscrasias, diabetes mellitus, hepatic function impairment, angina, hypotension, congestive heart failure, arrhythmias, breast cancer, glaucoma, history of neuroleptic malignant syndrome, prostatic hypertrophy, obesity, Parkinson's disease, diagnosis of a seizure disorder, urinary retention, patients at risk for paralytic ileus, pregnancy/nursing mothers.

Pregnancy and Breast-Feeding

See precautions. FDA Pregnancy Category B.

CLOZAPINE (CLOZARIL®) - continued

PRECAUTIONS TO CONSIDER

Drug Interactions of Major Significance

- 1) Concomitant use of CNS depressants
- 2) Antithyroid agents
- 3) Concomitant use of agents that cause EPS (including droperidol, metoclopramide, amoxapine, metyrosine, pimozide, reserpine)
- 4) Concomitant use of hypotension producing agents
- 5) Levodopa

SEE TABLE A: Cytochrome P450 Drug Metabolism/Inhibition

Age-Specific Considerations

Safety and efficiency have not been established in children under the age of 16. Geriatric patients may be more susceptible to orthostatic and anticholinergic effects.

Side Effects Which Require Medical Attention

- 1) Anticholinergic effects
- 2) Hypotension
- 3) Rashes, photosensitivity and altered pigmentation
- 4) Tardive dyskinesia or other late-onset EPS
- 5) Visual changes
- 6) Early symptoms of agranulocytosis (fever, sore throat, weakness)
- 7) Fluctuating vital signs
- 8) Altered consciousness
- 9) Fever
- 10) Drooling
- 11) Hyperglycemia
- 12) Clinically significant weight gain
- 13) Seizure

PATIENT MONITORING

Patient Monitoring Parameters

- 1) CBC as indicated by guidelines established by the manufacturer.
- 2) Pregnancy Test – as clinically indicated
- 3) Screening for abnormal involuntary movements using a standardized test - prior to initiation, annually and as clinically indicated.
- 4) EKG and CPK – as clinically indicated
- 5) Fasting glucose monitoring (finger stick or serum) every three months
- 6) HgbA_{1c} as clinically indicated

Dosing

See TDMHMR Drug Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

DECANOATES

fluphenazine decanoate (Prolixin® Decanoate), haloperidol decanoate (Haldol® Decanoate)

INDICATIONS

- 1) Chronic psychotic disorder requiring prolonged parenteral treatment

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction and similarly severe significant hypersensitivity to medication prescribed or structurally related medication
- 2) Severe CNS depression

Relative:

- | | |
|--|-----------------------------------|
| 1) Pregnancy/nursing mothers | 5) Impaired hepatic function |
| 2) History of drug induced agranulocytosis or leukopenia | 6) Parkinson's disease |
| 3) Breast cancer | 7) Severe cardiovascular diseases |
| 4) History of neuroleptic malignant syndrome | |

Precautions

Alcoholism (active), recent or current blood dyscrasias, hepatic function impairment, angina, hypotension, congestive heart failure, arrhythmias, breast cancer, , history of neuroleptic malignant syndrome, Parkinson's disease, poorly controlled seizure disorder.

Pregnancy and Breast-Feeding

See relative contraindications. FDA Pregnancy Category C.

Drug Interactions of Major Significance

- 1) Concomitant use of CNS depressants
- 2) Antithyroid agents
- 3) Concomitant use of agents that cause EPS (including droperidol, metoclopramide, amoxapine, metyrosine, pimozide, reserpine)
- 4) Levodopa

DECANOATES - continued

fluphenazine decanoate (Prolixin® Decanoate), haloperidol decanoate (Haldol® Decanoate)

PRECAUTIONS TO CONSIDER

Age-Specific Considerations

Conservative dosing and careful monitoring are advised in children and the elderly

Side Effects Which Require Medical Attention

- 1) Extrapyramidal side effects
- 2) Akathisia
- 3) Rashes
- 4) Tardive dyskinesia or other late-onset EPS
- 5) Early symptoms of agranulocytosis (fever, sore throat, weakness)
- 6) Fluctuating vital signs
- 7) Altered consciousness
- 8) Galactorrhea
- 9) Amenorrhea
- 10) Gynecomastia
- 11) Signs and symptoms of neuroleptic malignant syndrome

PATIENT MONITORING

Patient Monitoring Parameters

- 1) Pregnancy Test – as clinically indicated.
- 2) Screening for abnormal involuntary movements using a standardized test - prior to initiation, six months, annually and as clinically indicated.
- 3) Blood levels as clinically indicated. Therapeutic ranges for the lab used should be listed on the report.
- 4) EKG - as clinically indicated.

Dosing

See TDMHMR Drug Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

RISPERIDONE (RISPERDAL®), OLANZAPINE (ZYPREXA®), QUETIAPINE (SEROQUEL®), ZIPRASIDONE (GEODON®)

INDICATIONS

- 1) Disorders with psychotic symptoms (schizophrenia, schizoaffective disorder, manic disorders, depression with psychotic features, drug-induced psychosis, psychosis associated with other medical conditions)
- 2) Severe aggression secondary to a psychiatric disorder
- 3) Self Injurious Behavior secondary to a psychiatric disorder

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction and similarly severe significant hypersensitivity to medication prescribed
- 2) Severe CNS depression
- 3) QTc > 500 msec
- 4) For ziprasidone - Recent myocardial infarction, uncompensated congestive heart failure or when other drugs are being used that also prolong the QT interval such as (not complete list) quinidine, dofetilide, pimozide, sotalol, thioridazine, moxifloxacin, and sparfloxacin

Relative:

- | | |
|--|-----------------------------------|
| 1) Pregnancy/nursing mothers | 5) Impaired hepatic function |
| 2) History of drug induced agranulocytosis or leukopenia | 6) Parkinson's disease |
| 3) Breast cancer | 7) Severe cardiovascular diseases |
| 4) History of neuroleptic malignant syndrome | |

Precautions

Alcoholism (active), cataracts (quetiapine) recent or current blood dyscrasias, diabetes mellitus (olanzapine), hepatic function impairment, , angina, hypotension, congestive heart failure, arrhythmias, breast cancer, history of neuroleptic malignant syndrome, obesity, Parkinson's disease, poorly controlled seizure disorder.

Pregnancy and Breast-Feeding

See relative contraindications. FDA Pregnancy Category C.

Drug Interactions of Major Significance

- 1) Concomitant use of CNS depressants
- 2) Concomitant use of agents that cause EPS (including droperidol, metoclopramide, amoxapine, metyrosine, pimozide, reserpine)
- 3) Concomitant use of hypotension producing agents
- 4) Levodopa
- 5) Antithyroid agents
- 6) Drugs that prolong the QT interval

SEE TABLE A: Cytochrome P450 Drug Metabolism/Inhibition

RISPERIDONE (RISPERDAL®), OLANZAPINE (ZYPREXA®), QUETIAPINE (SEROQUEL®), ZIPRASIDONE (GEODON®) - continued

PRECAUTIONS TO CONSIDER - *continued*

Age-Specific Considerations

Safety and efficacy have not been established in children under the age of 18. Conservative dosing is advised in the elderly.

Side Effects Which Require Medical Attention

- 1) Anticholinergic effects
- 2) Acute extrapyramidal side effects (dystonia, pseudo-Parkinsonism)
- 3) Akathisia
- 3) Hypotension
- 4) Rashes, photosensitivity and altered pigmentation
- 5) Tardive dyskinesia or other late-onset EPS
- 6) Visual changes
- 7) Early symptoms of agranulocytosis (fever, sore throat, weakness)
- 8) Fluctuating vital signs
- 9) Altered consciousness
- 10) Galactorrhea (risperidone)
- 11) Amenorrhea (risperidone)
- 12) Gynecomastia (risperidone)
- 13) Hyperglycemia (olanzapine)
- 14) Weight gain
- 15) Cataracts (quetiapine)
- 16) QTc > 500 msec

PATIENT MONITORING

Patient Monitoring Parameters

- 1) Pregnancy Test - as clinically indicated
- 2) Screening for abnormal involuntary movements using a standardized test - prior to initiation, annually and as clinically indicated.
- 3) Quetiapine (Seroquel®) – Eye exam (direct ophthalmoscope or slit lamp) at initiation or shortly thereafter within 4 to 6 weeks and every six (6) months while on drug. This is completed for the monitoring of cataracts.
- 4) EKG – as clinically indicated
- 5) Glucose monitoring every three months (olanzapine)
- 6) HgbA_{1c} as clinically indicated (olanzapine)
- 7) Patients at risk for hypokalemia (and/or hypomagnesemia) should have baseline serum potassium and magnesium before starting ziprasidone

Dosing

See TDMHMR Drug Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

SEDATING ANTIHISTAMINES

diphenhydramine HCL (Benadryl®), hydroxyzine HCL (Atarax®)

INDICATIONS

- | | |
|-----------------------------|-------------------------------|
| 1) Anxiety | 3) Parkinsonism and other EPS |
| 2) Aggression and agitation | 4) Insomnia |

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) Delirium
- 2) Anticholinergic intoxication

Relative:

- 1) Nursing mothers
- 2) Renal impairment
- 3) Hepatic impairment
- 4) Elderly, debilitated patients
- 5) Lower respiratory tract symptoms (asthma)
- 6) Diseases states where increased anticholinergic activity may complicate disease course (narrow-angle glaucoma, benign prostatic hypertrophy)

Precautions

- 1) Photosensitivity
- 2) Respiratory impairment

Pregnancy and Breast-Feeding

FDA Pregnancy Category B (diphenhydramine), Category C (hydroxyzine)

Drug Interactions of Major Significance

- 1) Concomitant use of monoamine oxidase (MAO) inhibitors

Age-Specific Considerations

- 1) May cause excitability in children, elderly and persons with developmental disabilities

Side Effects Which Require Medical Attention

- 1) Worsening confusion or agitation
- 2) Somnolence
- 3) Nausea
- 4) Headache
- 5) Dizziness or lightheadedness

PATIENT MONITORING

Patient Monitoring Parameters

- 1) Pregnancy Test - as clinically indicated

Dosing

See TDMHMR Drug Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

ZALEPLON (SONATA®)

INDICATIONS

- 1) Short term treatment of insomnia

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction or similarly significant hypersensitivity to the medication prescribed

Relative:

None

Precautions

- 1) Pregnancy/nursing mothers
- 2) Impaired hepatic function
- 3) History of alcohol or drug abuse
- 4) Sleep Apnea

Pregnancy and Breast-Feeding

See precautions. FDA Pregnancy Category C.

Drug interactions of major significance

- 1) Alcohol or CNS Depressants
- 2) Cimetidine

Age-Specific considerations

- 1) Safety and efficacy not established for those under 18 years old.

Side Effects Which Require Medical Attention

- 1) Ataxia
- 2) Confusion or disorientation
- 3) Rash
- 4) Falling or dizziness
- 5) Worsening agitation, disinhibition or aggression

PATIENT MONITORING

Patient Monitoring Parameters

- 1) Pregnancy Test - as clinically indicated

Dosing

See TDMHMR Drug Formulary for dosage guidelines.

See medication rule for exceptions (documentation required).

ZOLPIDEM (AMBIEN®)

INDICATIONS

- 1) Short term treatment of insomnia

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction or similarly significant hypersensitivity to the medication prescribed

Relative:

None

Precautions

- 1) Pregnancy/nursing mothers
- 2) Impaired hepatic function
- 3) History of alcohol or drug abuse
- 4) Sleep Apnea

Pregnancy and Breast-Feeding

See precautions. FDA Pregnancy Category B.

Drug interactions of major significance

- 1) Alcohol or CNS Depressants

Age-Specific considerations

- 1) Safety and efficacy not established for those under 18 years old.

Side Effects Which Require Medical Attention

- 1) Ataxia
- 2) Confusion or disorientation
- 3) Rash
- 4) Falling or dizziness
- 5) Worsening agitation, disinhibition or aggression

PATIENT MONITORING

Patient Monitoring Parameters

- 1) Pregnancy Test - as clinically indicated

Dosing

See TDMHMR Drug Formulary for dosage guidelines.

See medication rule for exceptions (documentation required).

BETA-BLOCKERS

propranolol (Inderal®), atenolol (Tenormin®), metoprolol (Lopressor®), nadolol (Corgard®)

INDICATIONS

- 1) Aggressive behavior
- 2) Performance anxiety
- 3) Lithium- or valproic acid- induced tremors
- 4) Akathisia

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) Second and third degree heart block
- 2) Cardiogenic shock
- 3) Sinus bradycardia
- 4) History of anaphylactic reaction or similarly severe significant hypersensitivity to the medication prescribed

Relative:

- 1) Impaired pulmonary function (COPD, etc.)
- 2) Pregnancy/nursing mothers
- 3) Diabetes mellitus
- 4) Asthma

Precautions

- 1) Hyperthyroidism
- 2) Peripheral vascular disease
- 3) Diabetes mellitus
- 4) Hepatic function impairment
- 5) Myasthenia Gravis
- 6) Psoriasis
- 7) Renal function impairment
- 8) Hyperlipidemia
- 9) Discontinuation or rapid dose reduction
- 10) Congestive heart failure

Pregnancy and Breast-Feeding

See relative contraindications. Most beta-blockers are Pregnancy Category B, C, or D.

Drug Interactions of Major Significance

- 1) Allergy extracts
- 2) Antidiabetic agents
- 3) Antihypertensives
- 4) Sympathomimetics
- 5) Xanthenes (caffeine, theophylline)

SEE TABLE A: Cytochrome P450 Drug Metabolism/Inhibition

BETA-BLOCKERS- continued

propranolol (Inderal®), atenolol (Tenormin®), metoprolol (Lopressor®), nadolol (Corgard®)

PRECAUTIONS TO CONSIDER (continued)

Age-Specific Considerations

None

Side Effects Which Require Medical Attention

- 1) Dizziness
- 2) Difficulty breathing
- 3) Edema or swelling
- 4) Cold hands or feet
- 5) Tiredness or weakness
- 6) Nightmares
- 7) Confusion or disorientation
- 8) Bradycardia or hypotension

PATIENT MONITORING

Patient Monitoring Parameters

- 1) EKG (Age 45 or over) - baseline and as clinically indicated
- 2) Pregnancy Test - as clinically indicated
- 3) Pulse rate, blood pressure – baseline, prior to each dosage increase, quarterly, and as clinically indicated

Dosing

See TDMHMR Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

CLOMIPRAMINE (ANAFRANIL®)

INDICATIONS

- 1) Obsessive - compulsive disorder

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction or similarly severe significant hypersensitivity to the medication prescribed
- 2) Recovery phase of myocardial infarction
- 3) Pheochromocytoma

Relative:

- 1) Pregnancy/nursing mothers

Precautions

Alcohol intoxication, bipolar disorder in the absence of a mood stabilizer, recent or current blood dyscrasias, cardiovascular disorders including arrhythmia, diseases states where increased anticholinergic activity may complicate disease course (narrow-angle glaucoma, benign prostatic hypertrophy), heart block and failure, hepatic function impairment, hyperthyroidism, renal failure, diagnosis of a seizure disorder

Pregnancy and Breast-Feeding

See relative contraindications. FDA Pregnancy Category D.

Drug Interactions of Major Significance

- 1) Concomitant monoamine oxidase inhibitors (furazolidone, procarbazine, selegiline, tranylcypromine, phenelzine, isoniazid)
- 2) Concomitant use of CNS depressants
- 3) Cimetidine
- 4) Concomitant use of medications with anticholinergic effects
- 5) Noradrenergic anti-hypertensive agents (clonidine, guanabenz, guanadrel, guanethidine)
- 6) SSRI

SEE TABLE A: Cytochrome P450 Drug Metabolism/Inhibition

Age-Specific Considerations

Not recommended for use in children under age 10, conservative dosing is advised. EKG prior to dosage increase.

Side Effects Which Require Medical Attention

- 1) Anticholinergic effects
- 2) Sexual function impairment
- 3) Seizures
- 4) Dizziness, lightheadedness or fainting (orthostatic hypotension)
- 5) Tachycardia greater than 100 beats/min.
- 8) Jaundice
- 9) QTc >500 msec

CLOMIPRAMINE (ANAFRANIL®) - continued

PATIENT MONITORING

Patient Monitoring Parameters

- 1) EKG – baseline and as clinically indicated.
- 2) Pregnancy test – as clinically indicated.

Dosing

See TDMHMR Drug Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

GABAPENTIN (NEURONTIN®)

INDICATIONS

- 1) Chronic Pain Disorders

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction and similarly severe significant hypersensitivity to medication prescribed

Relative:

- 1) Renal Failure
- 2) Pregnancy/nursing mothers

Precautions

- 1) Compromised renal function

Pregnancy and Breast-Feeding

See relative contraindications. FDA Pregnancy Category C.

Drug Interactions of Major Significance

- 1) Antacids
- 2) Cimetidine
- 3) Oral contraceptives

Age-Specific Considerations

Safety and efficacy in children <12 has not been established

Side Effects Which Require Medical Attention

- 1) Blurred or double vision
- 2) Clinically significant weight gain
- 3) Rhinitis
- 4) Tremor
- 5) Nausea, vomiting diarrhea or abdominal discomfort
- 6) Drowsiness, lethargy
- 7) Nystagmus

PATIENT MONITORING

Patient Monitoring Parameters

- 1) Renal Function Test - baseline and as clinically indicated

Dosing

See TDMHMR Drug Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

NALTREXONE (REVIA®)

INDICATIONS

- | | |
|-----------------------|----------------------------|
| 1) Alcoholism | 3) Self injurious behavior |
| 2) Narcotic addiction | 4) Eating disorders |

PRECAUTIONS TO CONSIDER

Contraindications

Absolute:

- 1) History of anaphylactic reaction and similarly severe significant hypersensitivity to medication prescribed
- 2) Acute hepatitis or liver failure

Relative:

- 1) Hepatic function impairment
- 2) Renal impairment

Precautions

- 1) Pregnancy/nursing mothers

Pregnancy and Breast-Feeding

See precautions. FDA Pregnancy Category C.

Drug Interactions of Major Significance

Concomitant opioid-containing products

Age-Specific Considerations

Safety and efficacy not established for those under 18 years old.

Side Effects Which Require Medical Attention

- 1) Hepatotoxicity
- 2) Opioid withdrawal
- 3) Nausea
- 4) Headache, chronic or recurrent
- 5) Dizziness or lightheadedness

PATIENT MONITORING

Patient Monitoring Parameters

- 1) Pregnancy Test - as clinically indicated
- 2) Hepatic Function Panel - baseline and as clinically indicated

Dosing

See TDMHMR Drug Formulary for dosage guidelines.

Exceptions to maximum dosage must be justified as per medication rule.

STIMULANTS

methylphenidate (Ritalin®, Concerta™), dextroamphetamine (Dexedrine®), dextroamphetamine/amphetamine mixture (Adderall®)

INDICATIONS

- 1) Attention deficit disorder, with or without hyperactivity
- 2) Narcolepsy (methylphenidate; dextroamphetamine; dextroamphetamine/amphetamine mixture)
- 3) Severe treatment resistant depression or depression in medically compromised patients

PRECAUTIONS TO CONSIDER

Contraindications

Absolute

- 1) History of anaphylactic reaction or similarly severe significant hypersensitivity to the medication prescribed
- 2) Severe depression in children (methylphenidate)

Relative

- 1) Tourette's syndrome or other motor or vocal tics
- 2) Pre-existing psychosis
- 3) Hypertension
- 4) Cardiovascular disease (dextroamphetamine, dextroamphetamine/amphetamine mixture)
- 5) Glaucoma (dextroamphetamine, methylphenidate, dextroamphetamine/amphetamine mixture)
- 6) History of drug abuse/dependence
- 7) Hyperthyroidism
- 8) Pregnant or nursing mothers

Precautions

- 1) Family history of tics
- 2) Epilepsy or other seizure history

Pregnancy and Breast-Feeding

See relative contraindications. Stimulant either are FDA Pregnancy Category C or unknown.

Drug Interactions of Major Significance

- 1) Monoamine oxidase (MAO) inhibitors
- 2) Stimulants/Sympathomimetics
- 3) Beta-Blockers (amphetamine)
- 4) Antidepressants (amphetamine)
- 5) Digitalis glycosides (amphetamine)
- 6) Meperidine (amphetamine)
- 7) Thyroid hormones (amphetamine)

STIMULANTS

- continued

methylphenidate (Ritalin®, Concerta™), dextroamphetamine (Dexedrine®), dextroamphetamine/amphetamine mixture (Adderall®)

PRECAUTIONS TO CONSIDER (continued)Age-Specific Considerations

None

Side Effects Which Require Medical Attention

- 1) Hypertension
- 2) Tachycardia
- 3) Weight loss
- 4) Abnormal motor movements or tics
- 5) Psychosis
- 6) Hyperthermia
- 7) Irritability or nervousness
- 8) Insomnia
- 9) Chest pain

PATIENT MONITORINGPatient Monitoring Parameters

- 1) Height and weight in children (baseline and as clinically indicated)

Dosing

See TDMHMR Drug Formulary for dosage guidelines

Exceptions to maximum dosage must be justified as per medication rule.