

Metadata for Nearshore Fish Atlas of Alaska

Definitions of field names

Hauls: number of beach seine hauls

Locales: number of locales

Measured: number of fish measured for length; fork length or total length depending on species

Sites: number of sampling sites

Species: number of species captured; see caveats concerning unidentified fish

% **Occurrence**: frequency of occurrence (i.e., number of seine hauls where a species was captured/total number of seine hauls)

Average Length: fork length = average length (millimeters) of fish from tip of nose to fork of tail; total length = average length (millimeters) of fish from tip of nose to end of tail

Bedrock: steep rock cliffs in exposed locations outside of bays; kelps *Laminaria saccharina* and *Alaria marginata* often attached to bedrock faces

Catch: number of fish captured

Dates Sampled: mo/dy/year

Eelgrass: *Zostera marina*; occupies lower intertidal and shallow subtidal zones in soft substrates

Habitat: bedrock, eelgrass, kelp, sand-gravel

Kelp: mostly understory kelps growing as dense, low-lying mats on rocky substrates; dominant kelps *Laminaria saccharina*, *Cymathere triplicata*, and *Alaria* spp.

Latitude: decimal degrees, WGS84 datum, northern hemisphere

Locales: general sampling area

Locale Photo: photo(s) of general sampling area

Longitude: decimal degrees, WGS84 datum, western hemisphere

Mean Catch/Haul: mean number of fish captured per beach seine haul (i.e., catch/total number of seine hauls)

PSS: practical salinity scale

Sand-Gravel: mostly sand or gravel beaches with no rooted vegetation; some algae may be present

Site: specific sampling site within locale

Site Photo: photo(s) of site

SST: sea surface temperature in degrees celsius

Species: common name of species

Caveats concerning unidentified fish

Larval and juvenile fish smaller than 50 mm length were sometimes difficult to identify to species in the field. Unidentifiable fish were classified to family groups and are referred to as “juvenile family group name” (e.g., juvenile smelt). Some larval fish, however, were unidentifiable to family groups; these fish are referred to as “larvae”. “Larvae” are accounted for in **Catch** totals, but are not accounted for in # **Species** unless no other identifiable fish were captured. Fish identified to family group are accounted for in **Catch** totals, but are not accounted for in # **Species** unless no species from the same family group were captured. For example, unidentified juvenile rockfish are counted as a species if no other identifiable rockfish species were captured.

Beach seine description

Most sampling was done with a 37-m variable-mesh beach seine that tapered from 5-m wide at the center to 1-m wide at the ends. Outer panels were each 10 m of 32-mm mesh, intermediate panels were each 4 m of 6-mm mesh, and the bunt was 9 m of 3.2 mm mesh. The seine had a lead line and float line so that the bottom contacted the substrate and the top floated. The seine was set as a “round haul” by holding one end of the seine on the beach and backing around in a skiff in a semi-circle with the other end of the seine.