

Appendix A

Glossary

The following references were among those used in developing the glossary (see full reference citations in Appendix B):

- USDA Forest Service, et al. 1994a
- Dunster and Dunster 1996

abiotic	The non-living components of the planet not currently part of living organisms, such as soils, rocks, water, air, light, and nutrients.
active channel width	The average width of the stream channel at the normal high water level. The normal high water level is the stage reached during average annual high flow. This high water level mark often corresponds with the edge of streamside terraces; a change in vegetation, soil, or litter characteristics; or the uppermost scour limit (bankfull stage) of a channel.
active nest tree	A tree or snag in which a nest is tended during the breeding season by a pair of northern spotted owls.
activity center	The nest tree, or the location best describing the focal point of the activity of a northern spotted owl or pair of northern spotted owls when the nest location is not known.
adaptive kernel	A method for determining home range. In the adaptive kernel method, local adjustments are applied to the width of individual kernels. Observations in areas of high density get less smoothing (tighter fit), and observations in areas of low density get more smoothing (looser fit).
adaptive management	Adaptive management is a system of making, implementing, and evaluating decisions, which recognizes that ecosystems and society are always changing. It is a systematic and rigorous approach to learning from actions, improving management, and accommodating change.
advanced structure stand	Stands with advanced structure are more developed than intermediate structure stands in the understory reinitiation stage. Tree crowns show significant layering from the tallest trees to the forest floor. Advanced structure stands that are highly diverse may develop structural characteristics typically linked with older forests or old growth.
aggregate	Sand and pebbles added to cement to make concrete, or that are used in road construction.
alluvial	Soil, debris, and other materials that have been deposited by currents of water.
ambient	Surrounding.
anadromous fish	Species of fish (e.g., salmon) that hatch and rear for a portion of their life history in fresh water rivers and streams, then mature in the ocean, and then migrate back into freshwater rivers and streams to spawn.

aquatic	In or on the water; aquatic habitats are in streams or other bodies of water, as contrasted with riparian habitats, which are near water.
aquatic zone	The area that includes the stream channel(s) and associated aquatic habitat features. This zone includes beaver ponds, stream-associated wetlands, side channels, and the channel migration zone.
aquifer	A sand, gravel, or rock formation that is capable of storing or transporting water below the surface of the ground.
archaeological and historical resources	Districts, sites, buildings, structures, and artifacts that possess material evidence of human life and culture of the prehistoric and historic past.
archaeological object	An object that is at least 75 years old; is part of the physical record of an indigenous or other culture found in the state or waters of the state; and is material remains of past human life or activity that are of archaeological significance, including, but not limited to, monuments, symbols, tools, facilities, technological by-products, and dietary by-products (Oregon Revised Statutes [ORS] 358.905).
archaeological site	<p>A geographic locality in Oregon, including but not limited to, submerged and submersible lands and the bed of the sea within the state’s jurisdiction, that contains archaeological objects and the contextual associations of the archaeological objects with each other, or with biotic or geological remains or deposits (ORS 358.905). Specific types of sites, as defined in Oregon law, are:</p> <p>pre-historic archaeological site—Created and/or used by humans indigenous to the area before Euro-American inhabitation.</p> <p>historic archaeological site—Created and/or used by humans since the time of Euro-American inhabitation; usually belowground and/or aboveground diminishing remains.</p> <p>historic site—Created and/or used by humans since the time of Euro-American inhabitation; usually aboveground structurally intact remains.</p> <p>site of archaeological significance—Any archaeological site on, or eligible for inclusion on, the National Register of Historic Places as determined in writing by the State Historic Preservation Officer, or any archaeological site that has been determined significant in writing by an Indian tribe (ORS 358.905).</p>

average annual high flow period	High flows generally occur between November and March, with some variability in timing year to year. Average high flows are typically represented with a 2.5-year return interval.
average high water level	The stage reached during the average annual high flow period. This level often corresponds with the edge of streamside terraces, marked changes in vegetation, changes in soil or litter characteristics, or the bankfull stage of a channel.
basal area	The area of the cross-section of a tree stem near the base, generally at breast height (4.5 feet above ground) and including the bark. The basal area per acre is the total basal area of all trees on that acre.
biodiversity <i>or</i> biological diversity	The genetic variation and the variety of microbial, plant, and animal life.
biotic	Any living aspect of the planet.
board foot	The amount of wood equivalent to a piece of wood one foot wide by one foot high by one inch thick.
Board of Forestry	The Oregon Board of Forestry (BOF) is a seven-member board appointed by the Governor and confirmed by the state Senate. At least one member must reside in each of the state's three administrative regions (east, south, and northwest). No more than three members may receive any significant portion of their income from the forest products industry. The BOF supervises all matters of forest policy within Oregon; appoints the State Forester; adopts rules regulating forest practices; and provides general supervision of the State Forester's management of the Oregon Department of Forestry (ODF).
Board of Forestry Lands	Board of Forestry Lands were acquired by the BOF under ORS 530.010 to 530.040. Most were transferred from counties to the BOF in exchange for a portion of future revenue from the lands. Some lands were acquired by direct purchase.
bog	A wetland that is characterized by the formation of peat soils and that supports specialized plant communities. A bog is a hydrologically closed system without flowing water. It is usually saturated, relatively acidic, and dominated by ground mosses, especially sphagnum. Bogs are distinguished from other wetlands by the dominance of mosses and the presence of extensive peat deposits.
burial	Any natural or prepared physical location, whether originally below, on, or above the surface of the earth, into which, as a part of a death rite or death ceremony of a culture, human remains were deposited (ORS 358.905).

buffer habitat	Stands surrounding occupied stands that do not have the characteristics of suitable habitat, but that buffer the occupied stand from wind and other environmental factors as well as from other potential deleterious effects of edge, such as increased predation.
candidate species	Species being considered by the Secretary of the Interior for listing as an endangered or a threatened species, but not yet the subject of a proposed rule.
channel migration zone	An area adjacent to an unconfined stream channel where channel migration is likely to occur during high-flow events. The presence of side channels or oxbows, stream-associated wetlands, and low terraces are indicators of these zones. The extent of these areas will be determined through site inspections using professional judgment.
chlorosis	Yellowing of normally green plant tissue due to destruction or limited production of chlorophyll; often a symptom of mineral deficiencies, disease (such as Swiss needle cast), feeding by sucking insects, root or stem girdling, or serious light deficiencies.
Clean Air Act	Federal law passed in 1970, and amended several times since. The authority to implement the act is delegated to the states. The Clean Air Act is implemented, in part, through a permit system.
clearcut	Traditionally, a silvicultural system in which the entire stand of trees is cleared from an area at one time. Clearcutting and planting (if needed) results in the establishment of a new even-aged stand of trees. In the Elliott State Forest, a modified clearcutting system is used, in which live trees, snags, and downed wood remain on the unit after harvest.
coarse filter – fine filter	For the Elliott State Forest, an operational approach to managing for biological diversity is the “coarse filter – fine filter” concept proposed by. The coarse-filter component is based on the premise that maintaining a range of seral stages, stand structures, and sizes, across a variety of ecosystems and landscapes, will meet the needs of most organisms. Fine-filter management superimposes specific management actions for individual species or habitats that require special consideration, such as species with unique or limited distributions.
co-dominant	Trees with crowns that form a general level of crown stratum and are not physically restricted from above, but are more or less crowded by other trees from the sides.

cohort	A group of trees regenerating after a single disturbance. The age range within a cohort may be as narrow as one year or as wide as several decades, depending on how long trees continue invading after a disturbance.
colluvial	Soil, debris, and other materials that have been moved downslope by gravity and biological activity.
Common School Forest Lands	A subset of the Common School Trust Lands that have been listed by the State Land Board for the primary use of timber production.
Common School Fund	A permanent fund or account managed to provide revenues to the common schools. The State Land Board (Governor, Secretary of State, and Treasurer) is the trustee of the Common School Fund (CSF).
Common School Trust Lands	State lands owned by the State Land Board. In Oregon, the lands originally granted by Congress under the Oregon Admission Act included approximately six percent of the new state's land for the use of schools. The primary goal in managing Common School Trust Lands is the generation of the greatest amount of income for the CSF over the long-term, consistent with sound techniques of land management.
commonly used road	A road that receives frequent traffic during the marbled murrelet breeding season, including but not limited to, a mainline road and roads connecting mainline roads. Roads not commonly used may include, but are not limited to, spur roads and blocked or decommissioned roads.
composition	The different species of plants and animals that live in an ecosystem. The dynamic attributes of a forest ecosystem are composition, function, and structure. Composition is the proportion of various species. Function is the processes taking place in the system. Structure includes kinds and distribution of stand components such as trees, snags and logs of various sizes and shapes.
concept	An abstract or generic idea generalized from particular instances.
confirmed occupancy	Occupied behaviors observed on more than one visit.

conifer stand	These stands occupy most of the Elliott State Forest. The ODF classifies as conifer stands those in which conifer species compose 30 percent or more of the tree canopy. Although conifers are the principal species with economic value in these stands, the stands may also include substantial amounts of other vegetation types such as hardwoods, brush, grass, and ferns, which contribute to a diverse forest ecosystem. These types are either intermixed with the conifers or are in clumps too small to map and inventory separately.
connectivity	A measure of how well different areas (patches) of a landscape are connected by linkages, such as habitat patches or corridors. At a landscape level, the connectivity of ecosystem functions and processes is of equal importance to the connectivity of habitats.
conservation area	Designated land where conservation strategies are applied for the purpose of attaining specific conservation objectives; this may include cultural or biological aspects. In the Elliott State Forest, conservation areas include habitats utilized by northern spotted owls and marbled murrelets, riparian management areas, rare or unique habitats, and areas requiring special protection for visual or other resource values. Management within conservation areas is aimed at maintaining desired conditions.
core area	An area of contiguous suitable habitat surrounding a nest site or activity center.
core use area	Areas of concentrated use within the home range identified by calculating an average observation density of all locations for an individual northern spotted owl and determining the contour where the observation density is greater than average. This contour does not have a connotation of statistical significance, but it delimits an area of concentrated use. The advantage to this approach is that it avoids arbitrary selection of contours, and each core area is based only on the density of locations for that particular northern spotted owl.
corridor	Areas of habitat that connect separate but similar habitat patches, within the landscape mosaic. For example, an area of mature timber, such as a riparian buffer, may connect larger patches of mature timber.

critical habitat	The specific areas within the general geographic area occupied by a federally listed species in which physical and biological features occur that the U.S. Fish and Wildlife Service (USFWS) has determined to be essential to the conservation of the species. Critical habitat is designated by USFWS pursuant to the federal Endangered Species Act (ESA). Not all of the area encompassed by critical habitat contains the necessary habitat characteristics to support a particular species.
culmination of mean annual increment	Mean annual increment is the total increment of growth of a stand divided by the age of the stand. The culmination age is the age at which the mean annual increment reaches its maximum. If maximization of wood volume is the objective for the stand, this age is generally used as the rotation age. Periodic thinning enhances growth and extends the culmination age.
danger tree	A standing tree, alive or dead, that presents a hazard to personnel due to deterioration or physical damage to the root system, trunk (stem), or limbs, and the degree and direction of lean.
debris flow	A rapidly moving mass of rock fragments, soil, and mud, with more than half of the particles being larger than grains of sand.
debris torrent	Rapid movement of a large quantity of materials, including wood and sediment, down a stream channel. This generally occurs in smaller streams during storms or floods, and scours the stream bed.
decadence	Process of decay, or condition of being in a decayed state, particularly as related to trees or stands of trees. Typified by the presence of pathogens causing various forms of rot, and often used to refer to the presence of snags and downed wood. A process influential in multiple aspects of ecosystem development from providing cavities for wildlife, to creating gaps in the canopy, to altering forest floor climate and structure.
demographic study	A study of population dynamics; the quantitative analysis of population structure and trends in size, growth rate, and distribution.
density	The number or size of a population (trees, species, etc.) in relation to a unit of space. In silviculture, stand density is measured as the amount of tree biomass per unit area of land. This can be measured as the number of trees, basal area, wood volume, or foliage cover. Also see “stand density” and “stand density index.”
desired future condition	An explicit description of the physical and biological characteristics of the Elliott State Forest in the future, as described in the forest vision.

detection	Sighting or hearing of one or more birds acting in a similar manner, i.e., a single bird or flock.
diameter breast height	The diameter of a tree, measured 4.5 feet above the ground on the uphill side of the tree.
dissected	A landscape that has been cut into hills and valleys by the process of erosion.
disturbance	A force that causes significant change in an ecosystem's structure and/or composition. Disturbance can be caused by natural events such as fire, flood, wind, earthquake, and insect or disease outbreak, or by human activities. The disruption of marbled murrelet reproductive activities.
dominance	Trees with crowns that extend above the general level of crown cover of other trees of the same stratum and are not physically restricted from above, although possibly somewhat crowded by other trees on the sides.
downed wood	Fallen trees or pieces of trees on the forest floor or in the stream channel that provide many important functions such as mineral cycling, nutrient mobilization, maintenance of site productivity, natural forest regeneration (nurse logs), substrates for mycorrhizal formation, and diverse habitats for fish and wildlife species.
early structure stand	Following a disturbance, an early structure stand develops through the stand initiation process. In the early years of this stage, the site is occupied primarily by tree seedlings or saplings, herbs, grass, or shrubs. In later years, increasing crown closure shades the ground, and herbs, shrubs, and grasses begin to die out or lose vigor. At this point, the stand transitions from an early stand initiation stage to an intermediate stem exclusion stage, leading to an intermediate structure stand.
earthflow	Movement of material, both sediment and vegetation, down a slope. Earthflows are typically large, but move only a few centimeters each year. (See also "landslide.")
ecosystem	A complex system comprising populations of organisms considered together with their physical environment and the interacting processes between them (e.g., marsh, watershed, lake ecosystem). Ecosystems do not have boundaries fixed in time or space because the form and function of ecosystems change at various rates, depending on prevailing environmental factors.
ecosystem functions	The many and varied biotic and abiotic processes that make an ecosystem functional, changing, and interactive (e.g., biogeochemical processes, nutrient cycling, decomposition, regeneration, and succession).

ecosystem management	A management practice and philosophy aimed at selecting, maintaining, and/or enhancing the ecological integrity of an ecosystem to ensure continued ecosystem health while providing resources, products, or non-consumptive values for humans. The actions taken reflect the management goals and range from protection from human influence through to an increasing intensity of interventions to serve human needs.
edge	The point where two different plant communities (different vegetation types, successional stages, or conditions) meet. Edges may be created by a soil or topographical feature of the site, or where short-term effects are created by natural or human-caused disturbances.
endangered species	As defined by the ESA: Any species (including subspecies or qualifying population) that is in danger of extinction throughout all or a significant portion of its range.
ephemeral stream	Ephemeral streams occur in direct response to precipitation, running only during or shortly after periods of heavy rainfall or rapid snowmelt.
effectiveness monitoring	Used to determine if the design and execution of the prescribed management practices are achieving the Habitat Conservation Plan (HCP) conservation objectives. Every management decision is intended to achieve a given set of future conditions. Effectiveness monitoring can be used to compare existing conditions to both past and the desired future conditions to describe the overall progress or success of the management activities.
evolutionarily significant unit	An evolutionarily significant unit (ESU) is a group of stocks or populations that: 1) are substantially reproductively isolated from other population units of the same species; and 2) represent an important component in the evolutionary legacy of the species. This term is used by the National Marine Fisheries Service (NMFS) as guidance for determining what constitutes a “distinct population segment” for the purposes of listing Pacific salmon species under the ESA. For example, the “Oregon Coast chinook ESU” is a delineation that encompasses all populations of chinook salmon from the Necanicum River on the northern Oregon coast, to Cape Blanco on the south coast.
extensive management	Extensive forest management is a term used for protection of the forest from fire and insects, and the reliance on natural regeneration for provision of the next forest.
federally listed species	Species, including subspecies and distinct vertebrate populations, of fish, wildlife, or plants, listed at 50 CFR 17.11 and 17.12 as either endangered or threatened.

fixed kernel	A method of determining home range. In the fixed kernel method, a single smoothing width is used on all the observations in the sample. The fixed kernel generally produces estimates of home-range size and contours with lower bias than the adaptive kernel in simulation studies.
Forest Land Management Classification System	Under OAR 629-035-055, state forest lands are classified according to the management that will be applied. The classification describes the management emphasis for the land as determined by Forest Management Plans and any applicable HCP. State forest lands are classified as General Stewardship, Focused Stewardship, or Special Stewardship. Focused and Special Stewardship classifications are used when a particular forest resource may need a more focused approach or priority in management compared to other resources.
formation	A group of strata, or layers, of the same sort of rock or mineral, or rock having common characteristics.
fractal	Irregular shapes and surfaces that cannot be represented by classical geometry. Fractal dimension is an index of the complexity of spatial patterns.
fragmentation	The relationship of the landscape matrix to other types of patches; as fragmentation increases, the matrix becomes geometrically more complex. Maximum landscape fragmentation occurs when no dominant patch exists. Fragmentation is also defined as the spatial arrangement of successional stages across the landscape as the result of disturbance, and is often used to refer specifically to the process of reducing the size and connectivity of late successional or old growth forests.
fry	For salmonids, young fish that have just emerged from the gravel and are actively feeding.
function	An activity or process that occurs in an ecosystem; some typical functions are plant growth, animal reproduction, and decay of dead plants.
geographic information system	A system for management analysis and display of geographic knowledge that is represented using a series of information sets such as maps and globes, geographic data sets, processing and workflow models, data models, and meta data.
geotechnical	The study of soil stability in relation to engineering.
geothermal	Of or relating to the internal heat of the earth.

goals	A concise, broad statement of an organization’s end or process that programs are designed to achieve. A goal is normally expressed as a broad, general statement of purpose, is usually not quantifiable, and is timeless in that it usually has no specific date by which it is to be completed.
groundwater	The subsurface water supply (below the water table) that saturates the pores and fractures of sand, gravel, and rock formations.
guidelines	A set of recommended or suggested methods or actions that should be followed in most circumstances to assist administrative and planning decisions, and their implementation in the field. They are provided as a broad framework of recommended actions to be taken, and thus provide some flexibility for decision-making.
guiding principles	The overall rules, goals, and responsibilities that guide the planning process for the Elliott State Forest.
Habitat Conservation Plan	A comprehensive planning document that is a mandatory component of an Incidental Take Permit (ITP) application pursuant to section 10(a)(2)(A) of the ESA.
harass	“... an intentional or negligent act or omission which creates the likelihood of injury to wildlife by annoying it to such an extent as to significantly disrupt normal behavioral patterns which include, but are not limited to, breeding, feeding, and sheltering.” (50 CFR 17.3).
Harvest Unit	Delineated forest parcels that reflect potential logical harvest operation areas considering topography and access. A unit for clearcut and thinning choices.
hardwood stand	These stands are found on a minority of Elliott State Forest lands. The ODF classifies as hardwood stands those in which hardwood species comprise more than 70 percent of the tree canopy.
harm	An act “which actually kills or injures” listed wildlife; may include “significant habitat modification or degradation where it actually kills or injures wildlife by significantly impairing essential behavioral patterns, including breeding, feeding, or sheltering ...” (50 CFR 17.3).
headwall	The steep slope or rocky cliffs at the head of a valley.
high water line	The stage reached during the average annual high flow period. This level often corresponds with the edge of streamside terraces, marked changes in vegetation, or changes in soil or litter characteristics.

historic artifacts	Three-dimensional objects including furnishings, art objects, and items of personal property that have historic significance. “Historic artifacts” do not include paper, electronic media, or other media that are classified as public records (ORS 358.635).
historic property	Real property that is currently listed in the National Register of Historic Places, established and maintained under the National Historic Preservation Act of 1966, or approved for listing on an Oregon Register of Historic Places.
home range	The area within which an animal conducts its activities during a defined period of time (generally determined through radio-telemetry monitoring).
hydrology	The study of the properties, distribution, and effects of water on the landscape, under the surface, in the rocks, and in the atmosphere.
implementation monitoring	Used to determine if objectives, standards, and management practices specified in the HCP conservation strategies are being accomplished. Implementation monitoring is used to determine whether specified actions or criteria are being met.
Implementation Plan	An ODF plan that describes in more detail than the long-range Forest Management Plan how the management strategies will be applied. These plans are designed to describe forest management activities for a ten-year period, and are revised at least every ten years.
incidental take	Take of any federally listed wildlife species that is incidental to, but not the purpose of, otherwise lawful activities.
Incidental Take Permit	An Incidental Take Permit (ITP) is a federal exemption to take prohibition of Section 9 of the ESA; the ITP is issued by the USFWS pursuant to Section 10(a)(1)(B) of the ESA. An ITP is also referred to as a Section 10 Permit or Section 10(a)(1)(B) Permit.
inner RMA zone	The next area away from the stream, adjacent to the stream bank zone.
integrated pest management	A systematic approach that uses a variety of techniques to reduce pest damage or unwanted vegetation to economically and socially tolerable levels. Integrated pest management techniques may include the use of natural predators and parasites, genetically resistant hosts, environmental modifications, and, when necessary and appropriate, chemical pesticides or herbicides.

integrated resource management	The management of two or more resources in the same general area and period of time (e.g., water, soil, timber, grazing, fish, wildlife, and forests). For the Elliott State Forest, integrated resource management means that the design and application of management practices must consider the effects and benefits of all of the forest resources in such a way that those effects and benefits lead to achieving the goals in the Forest Management Plan over time and across the landscape.
intensive management	<p>Intensive forest management: A management concept promoting basic forest management in combination with juvenile-stand improvement and/or the use of artificial regeneration to ensure reasonably uniform stand establishment and stocking.</p> <p>Intensive silviculture: Any silvicultural practices designed to accelerate stand development and improve the stand value and final yields in stands that are well established.</p>
interior habitat area	The portion of the older forest patch that remains effective when the negative effects of high contrast edge are removed.
intermediate structure stand	As early structure stands develop and transition into the stem exclusion stage, trees fully occupy the site and form a single, main canopy layer. The stem exclusion process begins when new trees, shrubs, and herbs no longer appear and existing ones begin to die, due to competition for light, nutrients, and moisture. Later, as more of the trees die, the understory reinitiation process begins, when enough light and nutrients become available so that herbs, shrubs, and young trees again appear in the understory.
intermittent stream	A stream with surface flow only part of the year. In the Forest Practices Act, defined as a stream that normally does not have summer surface flow after July 15.
jacks	Sexually mature male salmon that reached maturity earlier than usual for their species.
landscape	Many sets of stands that cover an area ranging from many hundreds to tens of thousands of acres.
landslide	The dislodging and fall of a mass of earth and rock. There are many types of landslides, including debris slides, earthflows, rock block slides, slumps, slump blocks, and slump earthflows. The different types of landslides vary tremendously in how they occur, how far they move, what type of materials move, etc.

late successional habitat	A forest stand whose typical characteristics are a multi-layered, multi-species canopy dominated by large overstory trees; numerous large snags; and abundant large woody debris (such as fallen trees) on the ground. Other characteristics such as canopy closure may vary by the forest zone (lodgepole, ponderosa, mixed conifer, etc.).
leave area	An area of standing timber retained among areas of logging activity to satisfy management objectives, such as seed source, wildlife habitat, or landscape management constraints.
legacy structures	Structural components within a forest stand that are retained during harvest operations, and that provide habitat diversity in the future stand. Examples of legacy structure include live trees, snags, and downed wood.
lieu lands	“Lieu lands” were offered by the federal government to the state to compensate for original land grants that had conflicting claims. The Elliott State Forest includes approximately 7,700 acres of lieu lands.
likely nesting habitat	Occupied marbled murrelet habitat that is considered to be the most likely location for nesting sites, based on information from surveys, aerial photos, stand information, and the judgment of biologists or others familiar with the area. Stand type breaks or topography may be used to delineate the boundaries of likely nesting habitat.
lithic scatter	A location where prehistoric stone tools were made, usually from obsidian. The tools and weapons were used locally or traded.
loading	The quantity of a substance entering a body of water.
management basin	An area used for forest planning. Management basins are designated and displayed in the district Implementation Plan, and are shown in Figure 5-1 in Chapter 5.. Their boundaries are based primarily on drainage and topographic patterns within the major drainage basins and watersheds. Basin boundaries are generally consistent with 5th and 6th field Hydrologic Unit Codes as developed by the U.S. Geological Survey.
management prescription	The management practices and intensity selected and scheduled for application on a specific area to attain predefined goals and objectives.
marbled murrelet management area	The area designated for the protection of marbled murrelets, according to ODF policy. (The acronym “MMMA” is sometimes pronounced “trima.”)

mature forest condition	Desired mature forest condition consists of a stand dominated by large conifer trees, or where hardwood-dominated conditions are expected to be the natural plant community, a mature hardwood/shrub community. For conifer stands, this equates to a basal area of 220 square feet or more per acre, inclusive of all conifers over 11 inches diameter breast height. At a mature age (80 to 100 years or greater), this equals 40 to 45 conifer trees 32 inches in diameter breast height per acre.
minor tree species	For a given stand, tree species that occur as a relatively small component of the stand, such as western redcedar or alder in a stand consisting mostly of Douglas-fir trees.
monitoring	The measurement of environmental characteristics and conditions over an extended period of time to determine status or trends in some aspect of environmental quality. implementation monitoring —Asks the question, “Did we do what we said we would do?” effectiveness monitoring —Asks the question, “Are the management practices producing the desired results?” validation monitoring —Asks the question, “Are the planning assumptions valid, or are there better ways to meet planning goals and objectives?”
morphology	Form and structure.
National Environmental Policy Act	The National Environmental Policy Act (NEPA) was signed into law in 1969. NEPA requires all federal agencies to consider and analyze all significant environmental impacts of any action proposed by those agencies; to inform and involve the public in the agency’s decision-making process; and to consider the environmental impacts in the agency’s decision-making process.
native	Indigenous to Oregon and not introduced.
natural ecosystem	An ecosystem that is minimally influenced by humans and that is, in the larger sense, diverse, resilient, and sustainable.
near-stream riparian	Areas directly adjacent to the stream. Large wood is delivered by the tree falling directly into the stream from the adjacent streambank or hillslope.
nest stand	A stand with an active nest or a recent nest site as determined from a fecal ring or eggshell fragment, or discovery of a chick or eggshell fragment on the forest floor.
nonpoint source	Entry of a pollutant into a body of water from widespread or diffuse sources, with no identifiable point of entry. The source is not a distinct, identifiable source such as a discharge pipe. Erosion is one example of a nonpoint source.

non-salmonid fish	Any fish species outside the family <i>Salmonidae</i> . A salmonid may be resident or anadromous; examples are Pacific lamprey and sculpins.
non-silviculturally capable	Areas that are rocky, swampy, covered by water, or for various other reasons have little to no commercial value for timber production. The Elliott State Forest has a few parcels of rocky or swampy lands scattered throughout the forest. Most are less than 5 acres, although a few are as large as 20 acres.
northern spotted owl circle	An area defined by the provincial radius circle around a northern spotted owl activity center.
northern spotted owl site	A territory occupied by northern spotted owls.
not commonly used road	Roads not commonly used may include, but are not limited to, spur roads and blocked or decommissioned roads.
nutrient cycling	Circulation or exchange of elements, such as nitrogen and carbon dioxide, between living and nonliving portions of the environment.
objective	A clear and specific statement of results to be achieved within a stated time period. An objective is measurable and implies precise time-phased steps to be taken and resources to be used, which, together, represent the basis for defining and controlling the work to be done.
occupied stand	A stand of potential habitat where marbled murrelets have been observed exhibiting behaviors that have been observed in stands with evidence of nesting, such as subcanopy behaviors or circling.
occupied habitat	Suitable habitat that has been surveyed and determined to be occupied by marbled murrelets.
occupied sites	Sites determined to be occupied by marbled murrelets based on the observation of subcanopy behaviors during protocol surveys, or the observation of nest trees, eggshell fragments, or other evidence of marbled murrelet reproductive activities.
old growth	A forest stand whose typical characteristics are a patchy, multi-layered, multi-species canopy dominated by large overstory trees, some with broken tops and decaying wood; numerous large snags; and abundant large woody debris (such as fallen trees) on the ground. In western Oregon, old-growth characteristics begin to appear in unmanaged forests at 175 to 250 years of age. (See “late successional habitat.”)

open road/active use	This category includes any road open for travel with a motorized vehicle. It includes permanent roads and also temporary roads that are currently in use or will be used in the near future. These roads are usually available for use at any time of the year. Use may be continuous or intermittent. Roads in this category require active maintenance and have a full maintenance obligation under the Forest Practices Act.
orographic	A process in which air masses are lifted up by mountains or similar obstructions, leading to higher amounts of precipitation on the windward side of the mountain.
outer RMA zone	The portion of the riparian management area farthest away from the stream.
Ownership, Site, Cover, Use, and Recommendations	The old inventory system developed by the ODF, that includes 1:12,000 scale maps and overlays, data files by type and various sorts, and data summaries. The system is now being replaced by the Stand Level Inventory.
owl circle	The area defined for the purpose of identifying the home range of a northern spotted owl pair or resident single northern spotted owl; the circle size varies by physiographic province. In the Oregon Coast Range, the radius of an owl circle is 1.5 miles, encompassing the area of 4,766 acres. Guidelines established by the USFWS (later rescinded) required protecting 70 acres of northern spotted owl habitat immediately around a northern spotted owl activity center, 500 acres within 0.7 miles, and 1,906 acres within 1.5 miles.
owl site	A territory occupied by northern spotted owls.
particulate	Small particles in smoke produced by burning wood and other forest debris. Two kinds of particulate are controlled under federal and/or state requirements: total suspended particulates and PM ₁₀ (particulate matter less than 10 microns in diameter).
patch	The landscape patch is an environmental unit between which “quality” differs, such as a habitat patch.
perennial stream	A stream with year-round surface flow. In the Oregon Forest Practices Act, a perennial stream is defined as a stream that normally has summer surface flow after July 15.
platform	A relatively flat surface at least 5 inches in diameter and at least 50 feet high in the live crown of a coniferous tree. A platform includes the limb and any deformities of, or epiphyte cover growing on, the limb. For instance, a four-inch-diameter limb with moss cover that increases the overall diameter to five inches is a platform.

platform tree	Any tree having a single platform capable of hosting a nest for a marbled murrelet.
point source	The release of a pollutant from a pipe or other distinct, identifiable point, directly into a body of water or into a water course leading to a body of water.
policy	A definite, stated method or course of action adopted and pursued by an entity that guides and determines present and future decisions and actions. A policy establishes a commitment by which an entity is held accountable.
pollutant	A substance of such character and existing in such quantities as to degrade an environmental resource (i.e., water, air, or soil) by impairing its usefulness (including its ability to support living organisms).
population	The organisms that constitute a particular group of a species, or that live in a particular habitat or area. A group of fish (e.g., Nehalem River fall chinook salmon) that spawn in a particular area at a particular time, and that do not interbreed to any substantial degree with any other group spawning in a different area, or in the same area at a different time are considered a population (OAR, Division 7, 635-07-501(38)).
potential murrelet habitat	Potential murrelet habitat is defined as: 1) mature (with or without an old-growth component) and old-growth coniferous forests; and 2) younger coniferous forests that have deformations or structures suitable for nesting. Potential habitat can be as far as 50 miles from the ocean.
potential habitat	Stands with the characteristics of occupied marbled murrelet habitat, but that have not been surveyed for the presence of this species.
prescribed burning	Controlled fire burning under specified conditions to accomplish planned objectives; also called slash burning, as a frequent objective is to reduce the amount of slash left after logging. Objectives may include site preparation for planting and reduction of fire hazards or pest problems.
presence	A stand of potential habitat where marbled murrelets have been detected at the stand, but subcanopy behaviors have not been documented.
properly functioning aquatic systems	The range of diverse aquatic and riparian conditions over time and space that emulate the habitat conditions that resulted from natural disturbance regimes under which native species evolved. There is no one condition that is properly functioning.

recruitment nesting habitat	Stands that do not exhibit the characteristics of occupied habitat, but that could be managed to develop such characteristics in the future.
redd	Location selected by a female salmon or trout for laying eggs; female digs a “nest” in the stream gravels with her tail.
Reforestation Organization Operations Tracking System	The Reforestation Organization Operations Tracking System (ROOTS) is the next step in the development of the State Forests Program Integrated Information System. ROOTS contains the following three main functions: 1) Stand Level Inventory (SLI): Tools for importing, updating, analyzing, viewing, and reporting on SLI information. 2) Silvicultural Treatment Records: Tools for managing, Analyzing, viewing, and reporting on information related to forest management activities such as harvesting, site preparation, planting, animal damage protection, vegetation management, interplanting, fertilization, and pruning. 3) Planning Units: Tools for making and recording a plan for future activities needed to achieve the desired future condition (DFC) for a specific geographic location.
refugia	Locations and habitats that support population of organisms that are limited to small fragments of their previous geographic range, and areas that remain unchanged while surrounding areas change markedly (the areas serve as a refuge for those species requiring specific habitats). The changes could be short term, such as wildfires or human activity, or much longer term, such as periods of glaciation.
regeneration	Regeneration refers to the process of renewal of a forest or stand of trees, or to the young trees in a stand.
regeneration harvest	The removal of trees to make regeneration possible or to assist in the development of the established regeneration (young trees). The most common type of regeneration harvest in the Elliott State Forest is a modified clearcut, leaving specified amounts of live trees, snags, and downed wood.
reserve	An area of land set aside to maintain it in a desired condition, e.g., as functional habitat for wildlife.
resident fish	Fish species that complete their entire life cycle in freshwater; non-anadromous fish. One example is a resident population of cutthroat trout.
residual live trees	Live trees that are retained to provide short-term habitat needs of wildlife species, to serve as a source of future snags and downed wood, and to provide legacy trees in future stands. This term also refers to live trees present in a stand that are legacies of a previous cohort of trees.

riparian area	Three-dimensional zone of direct influence and/or interaction between terrestrial and aquatic ecosystems. The boundaries of the riparian area extend outward from the stream bed or lakeshore.
riparian management area	A riparian management area (RMA) is a protected area with site-specific boundaries established by the ODF; the width varies according to the stream classification or special protection needs. The purpose of the RMA is to protect the stream, aquatic resources, and riparian area. Aquatic resources include water quality, water temperature, fish, stream structure, and other resources.
rock block slide	Type of landslide in which the weakness and initial breaking is in the underlying rock, not the soil. (See also “landslide.”)
rotation	Also called tree age rotation. The time needed from regeneration of a crop of trees through to harvestable timber, or the time period to reach other stand criteria (e.g., complex habitat function).
salmonid	Fish species belonging to the family <i>Salmonidae</i> ; includes trout, salmon, and whitefish species.
salvage	Salvage cutting is the utilization of standing or down trees that are dead, dying, or deteriorating, for whatever reason, before the timber values are lost.
seasonal stream	A stream with surface flow only part of the year. In the Forest Practices Act, defined as a stream that normally does not have summer surface flow after July 15.
seral stages	Developmental stages that succeed each other as an ecosystem changes over time; specifically, the stages of ecological succession as a forest develops.
silviculture	The art, science, and practice of controlling the establishment, composition, health, quality, and growth of the vegetation of forest stands. Silviculture involves the manipulation, at the stand and landscape levels, of forest and woodland vegetation, and the control or production of stand structures such as snags and down logs to meet the needs and values of society and landowners.

site class

Site class is a measure of an area’s relative capacity for producing timber or other vegetation. It is an index of the rate of tree height growth, with lower values indicating faster growing trees. The site index is expressed as the height of the tallest trees in a stand at an index age. In this document, an age of 50 years is used. The five site classes are defined below. Most of the Elliott State Forest is site class II or III.

- Site class I135 feet and up
- Site class II115 to 134 feet
- Site class III95 to 114 feet
- Site class IV75 to 94 feet
- Site class V below 75 feet

site index

A measure of forest productivity, expressed as the height of the tallest trees in a stand at an index age. In this document, an age of 50 years is used. (See “site class.”)

site status

The occupancy status of a surveyed area, as defined by the survey protocol.

Pair Status—Established by any of the following:

- A male and female are heard and/or observed (either initially or through their movement) in proximity (less than one-quarter mile apart) to each other on the same visit; or
- A male takes a mouse to a female; or
- A female is detected (seen) on a nest; or
- One or both adults are observed with young.

Resident Single Status—Established by:

- The presence or response of a single northern spotted owl within the same general area on three or more occasions within a breeding season; or
- Multiple responses over several years (e.g., two responses in year 1 and one response in year 2, from the same general area).

The presence or response of two birds of the opposite sex where pair status cannot be determined and where at least one member meets the other resident single requirements. (Note: This is considered “two birds, pair status unknown” in the survey protocol. This is lumped with the resident single category because management options are the same as for resident singles.)

slope stability

The degree to which a slope resists the downward pull of gravity. The more resistant, the more stable.

slump	A type of landslide; involves a failure in the soil, tends to be spoon-shaped, and the base often oozes out. (See also “landslide.”)
slump blocks, slump earthflows	Types of landslides. (See “landslide,” “slump,” and “earthflow.”)
smolts	Juvenile salmon that are leaving freshwater and migrating to the ocean.
snag	A standing dead tree.
spatial forest modeling	Spatial forest modeling is the assignment of harvest activities to specific forest parcels, thereby controlling the size and juxtaposition of treatment areas. Examples of spatial control include the size of regeneration harvests, the shape and size of older forest patches, establishing and maintaining connectivity, scheduling of transportation, and coordination of upslope and riparian activities. Spatial forest modeling is contrasted with strata-based forest modeling where parcels with common characteristics are merged together into strata with harvest activities assigned to percentages of the strata. However, with strata-based modeling, it is not known which parcels in the strata are scheduled, and it is not possible to control the size and juxtaposition of treatments.
species	When referring to the federal ESA, species also means: “...any subspecies of fish or wildlife or plants, and any distinct population segment of any species of vertebrate fish or wildlife which interbreeds when mature” [Section 3(15) of the ESA].
stand	A group of trees that forms contiguous potential marbled murrelet habitat with no gaps wider than 100 meters.
stand density	In silviculture, stand density is measured as the amount of tree biomass per unit area of land. This can be measured as the number of trees, basal area, wood volume, or foliage cover.
stand density index	A relative measure of stand density; converting a stand’s current density into a density at a reference size. It is usually expressed in the equivalent number of trees that are 10 inches in diameter, e.g., 65 trees per acre that average 26 inches in diameter have the same stand density index as 300 trees per acre that average 10 inches in diameter.
stand initiation	This stand development process begins when a disturbance such as timber harvest, fire, or wind has killed or removed most or all of the larger trees, or when brush fields are cleared for planting.

Stand Level Inventory	The ODF’s Stand Level Inventory acquires and updates state forest vegetation information at the specific site level (forest stand). This information is used for tactical and operational decision-making. The Stand Level Inventory includes vegetation sampling protocols, forest stand data arranged in a database, computer programs for managing and using the information, and documentation of inventory elements.
stand structure	For the purposes of this HCP, a series of three stand structures have been defined depicting the typical progression of stand development following a natural or human-caused disturbance. The stand initiation process is represented by the early stand structure . The stem exclusion and early understory reinitiation processes are represented by the intermediate structure . Structural complexity and larger tree size inherent to the advanced understory reinitiation process are characteristic of the advanced stand structure . Old growth stands are included in the advanced stand structure.
standard	A working principle that establishes the measure of performance extent, values, quantity, or quality for a given activity or item.
State Agency Coordination Program	Required under law for each state agency, to establish procedures to assure compliance with statewide land use goals and acknowledged city and county comprehensive plans and land use regulations.
state forests	In this HCP, “state forests” or “state forest lands” refers to Common School Forest Lands owned by the State Land Board, and to lands owned by the BOF, and managed by the ODF.
State Historic Preservation Office	Oregon’s State Historic Preservation Office was created in 1966 by federal statute. It administers the Statewide Plan for Historic Preservation and submits Oregon’s nominations for the National Register of Historic Places.
State Land Board	The Oregon State Land Board is composed of the Governor, Secretary of State, and State Treasurer. It was established under the Oregon Constitution to manage Common School Trust Lands and serve as trustee of the CSF.
Statewide Planning Goals	Statewide Planning Goals are adopted by the Land Conservation and Development Commission to set standards for local land use planning. They have the force of law.

steep, unique, or visual lands

- Areas almost exclusively associated with the steep, rocky slopes on either side of major rivers or streams, including the Umpqua River, Mill Creek, and the West Fork Millicoma River. These protected corridors vary from 1,000 to 4,000 feet in width. Slopes affected by public safety considerations fall within this category.
- Areas classified as non-silviculturally capable because they are rocky, swampy, or covered by water, or for various other reasons have little to no commercial value for timber production. Currently, the Elliott State Forest has a few parcels of rocky or swampy lands scattered throughout the forest. Most parcels are less than 5 acres, though a few are as large as 20 acres.
- Areas where scenic values are the primary values to be maintained, including areas buffering recreational areas, highway corridors, river corridors, lakeshores, and other scenic attractions.

stem exclusion process

The stem exclusion process begins when new trees, shrubs, and herbs no longer appear and existing ones begin to die, due to competition for light, nutrients, and moisture.

stock

For the purposes of fisheries management, a stock is an aggregation of fish populations that typically share common characteristics such as life histories, migration patterns, or habitats (OAR, Division 7, 635-07-501(51)). For example, “north-mid coast fall chinook salmon” can be defined as a stock. This stock includes a number of fall chinook “populations” from basins in this area such as the Siuslaw, Yaquina, and Tillamook Bay watersheds.

stocking

A measure of the adequacy of tree cover on an area. Unless otherwise specified, stocking includes trees of all ages.

strategy

A carefully considered plan or method, more encompassing and larger scale than tactics, for achieving an objective.

stream

To qualify as a stream, a water course must have a distinct channel that carries flowing surface water during some portion of the year, including associated beaver ponds, oxbows, side channels, and stream-associated wetlands if these features are connected to the stream by surface flow during any portion of the year. Ephemeral overland flow is not a stream, as this type of flow does not have a defined channel.

stream-associated wetland

A wetland that is immediately adjacent to a stream. This includes wetlands that are adjacent to beaver ponds, side channels, or oxbows that are hydrologically connected to the stream channel by surface flow at any time of the year.

stream bank zone	The land nearest to the stream, including the stream banks. Most riparian functions are supported to some extent by vegetation in this zone, which provides aquatic shade, delivers downed wood and organic inputs (leaves and tree litter) to the stream and riparian area, stabilizes the stream bank, contributes to floodplain functions, and influences sediment routing processes
stream classification	<p>Under the ODF’s Forest Practices Act, streams are classified in two categories based on their beneficial use.</p> <p style="padding-left: 40px;">Type F — Fish-bearing stream</p> <p style="padding-left: 40px;">Type N — Not a fish-bearing stream</p> <p>Streams are also classified by size and amount of flow. Large streams have an average annual flow greater than ten cubic feet per second (cfs); medium streams have an average annual flow of two to ten cfs; and small streams have an average annual flow less than two cfs.</p> <p>perennial streams—Year-round surface flow. In the Forest Practices Act, defined as a stream that normally has summer surface flow after July 15.</p> <p>intermittent streams—Surface flow only part of the year. In the Forest Practices Act, defined as a stream that normally does not have summer surface flow after July 15. Ephemeral streams may run only during or shortly after periods of heavy rainfall or rapid snowmelt.</p>
stream reach	A section of stream that is geomorphically distinct, and that can be delineated from other adjacent sections based on channel gradient, form, or other physical parameters.
structure	The physical parts of an ecosystem that can be seen and touched; typical structures in a forest are tree sizes, standing dead trees (snags), and fallen dead trees.
stumpage	The price charged for the right to harvest timber from publicly or privately owned forest land.
sub-canopy behavior	Behaviors occurring at or below the forest canopy, and strongly indicating that the site has some importance for breeding, including flying through the canopy, circling below canopy, and landing.
succession	A series of changes by which one group of organisms succeeds another group; a series of developmental stages in a plant community.
suitable habitat	Stands with the characteristics suitable for marbled murrelet nesting (including occupied habitat and potential habitat).

suppressed	Trees with crowns entirely below the general level of dominant and codominant trees and are physically restricted from immediately above.
sustainability	<p>Sustainability is the ability of an ecosystem to maintain ecological processes and functions, biological diversity, and productivity over time.</p> <p>Sustainable forest management describes forest management regimes that maintain the productive and renewal capacities, as well as the genetic, species, and ecological diversity of forest ecosystems.</p>
Sustainable Economic and Social Benefit	Key strategies in the Elliott State Forest Management Plan. These strategies focus on legal mandates and trust obligations, predictable and dependable products and revenues, and social benefit through forest management.
Sustainable Forest Ecosystem Management	Key strategies in the Elliott State Forest Management Plan. A management approach that is based on the synthesis of knowledge from various disciplines, including forestry, fisheries, wildlife, and hydrology. It is an approach to forest management that seeks to achieve a broad range of resource goals and provide a balance of social, economic, and environmental benefits from the forest over time. In addition, Sustainable Forest Ecosystem Management is a landscape approach to the management of forested ecosystems utilizing silvicultural tools emulating natural disturbances to provide forest products, maintain forest health, and retain a high level of social value.
sustained yield	<p>(1) An ideal forest management objective at which point the volume of wood removed is equal to growth within the total forest. Sustained-yield management implies continuous production planned to achieve at the earliest practical time a balance between increment to the mature forest and its cutting.</p> <p>(2) Achieving and maintaining in perpetuity a nearly equal annual or regular periodic output of the various renewable resources, without impairment of the productivity of the land.</p>
take	“... to harass, harm, pursue, hunt, shoot, wound, kill, trap, capture, or collect, or to attempt to engage in any such conduct” with regard to federally listed endangered species of wildlife (Section 3(18) of the ESA). Federal regulations provide the same taking prohibitions for threatened wildlife species (50 CFR 17.31(a)).
tectonic	Resulting from changes in the earth’s crust.

telemetry	The process of remotely monitoring an animal and its movements by radio transmissions from a device attached to the animal.
territory	The area that an animal defends, usually during breeding season, against intruders of its own species.
threatened and endangered species	Federal and state agencies make formal classifications of wildlife species, according to standards set by federal and state ESAs. The various classifications are defined below. Federal designations are made by the UUSFWS or NMFS. State of Oregon designations are made by the Oregon Department of Fish and Wildlife (ODFW).

Federal Classifications

candidate species—Those species for which the USFWS or NMFS has sufficient information on hand to support proposals to list as threatened or endangered.

endangered species—A species determined to be in danger of extinction throughout all or a significant portion of its range.

federally listed species—Species, including subspecies and distinct vertebrate populations, of fish, wildlife, or plants listed at 50 CFR 17.11 and 17.12 as either endangered or threatened.

proposed threatened or endangered species—Species proposed by the USFWS or NMFS for listing as threatened or endangered; not a final designation.

threatened species—Species likely to become endangered throughout all or a significant portion of their range within the foreseeable future.

State Classifications

endangered species—Any native wildlife species determined by the State Fish and Wildlife Commission to be in danger of extinction throughout any significant portion of its range within Oregon, or any native wildlife species listed as endangered by the federal ESA.

sensitive species—A watchlist, developed by the ODFW, of wildlife species that are likely to become threatened or endangered throughout all or a significant portion of their range in Oregon. Subdivided into four categories: critical, vulnerable, peripheral, and undetermined status.

threatened species—Any native wildlife species that the State Fish and Wildlife Commission determines is likely to become endangered within the foreseeable future throughout any significant portion of its range within Oregon.

threshold phenomenon	Pattern or trend in population growth (climate, etc.) that exhibits relatively long periods of slow change followed by sudden increase or decrease in response to a gradually changing environment.
triad approach	An approach in which three land use types are distinguished that can coexist without compromising the goal of sustaining biological diversity. The land use types are: 1) intensive commodity production areas; 2) areas with little or no resource use by people except low-intensity recreation; and 3) areas in which modest resource use is allowed while ecological values are protected (maintenance of diversity and ecosystem function takes precedence over commodity production).
unclassified stand	These stands are currently under contract for harvesting, or have already been harvested and will be planted soon.
understory reinitiation	The understory reinitiation process begins when enough light and nutrients become available to allow forest floor herbs, shrubs, and tree regeneration to again appear in the understory. The amount of brush and herbaceous species is minimal at the beginning, but increases to a substantial part of the stand by the end of the stage.
unsaturated zone	The layer of soil or rock between the aquifer and the surface of the ground. In this layer, some water is suspended in the spaces between soil or rocks, but the zone is not completely saturated.
upslope	Zero-order channels (zero-order channels are small unbranched draws), hollows, or hillslopes. Areas outside of the riparian area. Large wood is delivered by a landslide or landslide-debris flow combination that moves the wood into the stream channel from these areas.
upstream riparian	Near-stream riparian sources that are upstream of the reach of concern. High water and/or a debris flow transport large wood to its current location after initially falling into the stream from the riparian area.
validation monitoring	Used to determine whether data and assumptions for predicting outcomes and effects are correct. Validation monitoring seeks to verify the assumed linkages between cause and effect. Validation monitoring is long term and will be accomplished through formal research and effectiveness monitoring projects.
watershed	In general, a watershed is defined as an area within which all water that falls as rain or snow drains to the same stream or river. Watersheds can vary greatly in size, from that of a small stream to a larger waterbody such as the Umpqua River.

watershed analysis	A process in which data are evaluated and interpreted in order to understand causal linkages between watershed-scale processes. This process informs the design and execution of management plans and activities.
water table	The top of the groundwater. The water table is generally subsurface; marshes and lakes form where the water table meets the land surface.
wetland	As defined in Oregon’s Forest Practice Rules OAR 629-24-101 (77), wetlands are “those areas that are inundated or saturated by surface or ground water at a frequency and duration sufficient to support, and that under normal circumstances do support, a prevalence of vegetation typically adapted for life in saturated soil conditions.”
windthrow	Trees felled by high winds.

Appendix B

References

This appendix lists the books, reports, and other publications referred to in the plan. Listings are alphabetical. The following format is used.

Author's name in bold. Year published. Title of publication. Publisher, publisher's location.

The following abbreviations are used in this appendix. Standard two-letter postal abbreviations are used for the names of states.

BLM	Bureau of Land Management
DEQ	Oregon Department of Environmental Quality
GTR	General Technical Report
ODF	Oregon Department of Forestry
ODFW	Oregon Department of Fish and Wildlife
ONHP	Oregon Natural Heritage Program
PNW	Pacific Northwest Research Station (part of USDA Forest Service)
USDA	U.S. Department of Agriculture
USDI	U.S. Department of the Interior

References begin on the following page.

Agee, J.K. 1993. Fire Ecology of Pacific Northwest Forests. Covelo, CA Island Press, Washington, D.C. p. 5-6, 5-27.

Allbritten, M. 2002. A report on wildlife surveys conducted on the Elliott State Forest, 2000-20-01-2002. Unpublished report to Oregon Department of Forestry prepared by Marnie Allbritten, Oregon Department of Fish and Wildlife, Umpqua Watershed District. October 28, 2002. 13 p. plus appendices.

Allen and Dent. 2001. Shade conditions over forested streams in the blue mountain and coast range georegions of Oregon. ODF Technical Report #13. Oregon Department of Forestry, Salem, Oregon.

Altman, B. 2003. Olive-sided flycatcher. Pp. 374-376 *in* Birds of Oregon: A General Reference. D.B. Marshall, M.G. Hunter, and A.L. Contreras, Eds. Oregon State University Press, Corvallis, OR.

Altman, B. and R. Sallabanks. 2000. Olive-sided flycatcher (*Contopus cooperi*). *In* The Birds of North America, No. 502 (A. Poole and F. Gills, eds). The Birds of N. Am., Inc., Philadelphia, PA.

Altman, Bob. 1999. Conservation Strategy for landbirds in coniferous forests of western Oregon and Washington. Version 1.0. Unpublished report prepared for Oregon-Washington Partners in Flight. 83 pp.

Anderson Paul D., David Larson, and Samuel Chan. 2006. Riparian buffer and upslope density management influences on microclimate of young headwater forests of western Oregon. Presented at the 2005 Science and Management of Headwater Streams in the Pacific Northwest, Corvallis, OR. Abstracts from presentations posted: <http://www.headwatersresearch.org/> Page 10.

Andrén, H. 1994. Effects of Habitat Fragmentation on Birds and Mammals in Landscapes with Different Proportions of Suitable Habitat: A Review. *Oikos* 71:355-366.

Andrus, C. 2001a. Inventory of Large Wood Abundance and Transport in Steep Channels of the Elliott State Forest. June 22, 2001. Prepared for the Oregon Department of Forestry.

Andrus, C. 2001b. Summary of Streamside Stand Measurements on the Elliott State Forest. December 14, 2001. Prepared for the Oregon Department of Forestry.

Andrus, C.W. and H.A. Froehlich. 1988. Riparian forest development after logging or fire in the Oregon Coast Range: Wildlife habitat and timber value. *In* Streamside Management: Riparian Wildlife and Forestry Interactions. College of Forest Resources, Seattle, WA.

Anthony, R.G. and F.B. Isaacs. 1989. Characteristics of Bald Eagle Nest Sites in Oregon. *Journal of Wildlife Management*, 53:148-159.

- Anthony, R.G., R.L. Knight, G.T. Allen, B.R. McClelland, and J.I. Hodges.** 1982. Habitat Use by Nesting and Roosting Bald Eagles. Transactions of the North American Wildlife and Natural Resources Conference 47:332-342.
- Anthony, R.G., M.G. Garrett, and C.S. Schuler.** 1993. Environmental Contaminants in Bald Eagles in the Columbia River Estuary. Journal of Wildlife Management, 57:10-19.
- Anthony, R.G., R.W. Frenzel, F.B. Isaacs, and M.G. Garrett.** 1994. Probable causes of nesting failures in Oregon's bald eagle populations. Wildlife Society Bulletin 22:576-582.
- Anthony, R.G., R.J. Steidl, and M.C. Hansen.** 1996. Demographic characteristics of northern spotted owls on lands managed by Oregon Department of Forestry in the northern Coast Ranges and the Elliott State Forest, Oregon. FY 1995 Annual Research Report. Oregon Cooperative Fish and Wildlife Unit, Department of Fisheries and Wildlife, Oregon State University, Corvallis, OR 28 pp.
- Anthony, R.G., M. Hansen, K. Swindle, and A. Ellingson.** 2000a. Demographic Characteristics of Northern Spotted Owls on Lands Managed by Oregon Department of Forestry. Final Research Report. 40 pages. Oregon Department of Forestry, Salem, OR. Revised 3/23/2000.
- Anthony, R.G., B.Glenn, K. Swindle, and M.C. Hansen.** 2000b. Home Range and Habitat Use of Northern Spotted Owls on State Forest Lands in the Oregon Coast Range. Final Research Report. 114 pages. Oregon Department of Forestry, Salem, OR. Revised 3/23/2000.
- Applegarth, J.S.** 1994. Special Status Amphibians and Reptiles of the Eugene District: A Guide for their Conservation. USDI Bureau of Land Management, Eugene, OR. 51 pp.
- Arnett.** 2000. Personal communication to Marnie Allbritten Oregon Department of Fish and Wildlife, Southwest Region Office, 4192 N. Umpqua Highway, Roseburg, OR 97470.
- Arney, J.D.** 1973. Tables for Quantifying the Competitive Stress on Individual Trees. Canadian Forestry Service. Pacific Forest Research Center, Victoria, B.C. Information Report BC-X-78. 45 pp.
- Arney, J.D.** 1999. Personal communication to Jeff Boechler, Oregon Department of Fish and Wildlife, Corvallis, OR.
- Arnqvist, G. and D. Wooster.** 1995. Meta-analysis: Synthesizing Research Findings in Ecology and Evolution. Trends in Ecology and Evolution 10: 236-240.
- Aubry, K.B. and D.B. Houston.** 1992. Distribution and status of the fisher (*Martes pennanti*) in Washington. Northwestern Naturalist, 73:69-79.
- Aubry, K.B. and C.M. Raley.** 2002. Ecological characteristics of fishers (*Martes pennanti*) in the Southwestern Oregon. Progress Report: 1 January 2001 – 1 May 2002. USDA Forest Service, Pacific Northwest Research Station, Olympia, WA. 14 pp.

Aubry, K.B., L.L.C. Jones, and P.A. Hall. 1988. Use of Woody Debris by Plethodontid Salamanders in Douglas-Fir Forests in Washington. *In* Management of Amphibians, Reptiles, and Small Mammals in North America — Proceedings of the Symposium. USDA Forest Service General Technical Report RM-166, Fort Collins, CO.

Aubry, K.B., and C. M. Raley. .2006. Ecological characteristics of fishers (*Martes pennanti*) in the southern Oregon Cascade Range. Update: July 2006 Research Report. USDA Forest Service. Pacific Northwest Research Station, Olympia, WA. 30 pp.

Baichich, Paul J. and Colin J.O. Harrison. 1997. A Guide to the Nests, Eggs, and Nestlings of North American Birds, 2nd ed. San Diego: Academic Press.

Barnes, M.D. 2001. ODF Riparian Vegetation Survey 2001, Elliott State Forest. October 12, 2001. Prepared by Integrated Resource Management, Inc. for the Oregon Department of Forestry.

Bate, L.J. 1995. Monitoring woodpecker abundance and habitat in the central Oregon Cascades. M.S. thesis. University of Idaho. 116 pp.

Beamish, R.J., 1980. Adult biology of the river lamprey (*Lampetra ayresii*) and the Pacific lamprey (*Lampetra tridentate*) from the Pacific coast of Canada. *Can. J. Fish and Aquatic Sci.* 37:1906-1923.

Beamish R.J. and J.H. Youson. 1987. Life history and abundance of young adult *Lampetra ayresii* in the Fraser River and their possible impact on salmon and herring stocks in the Strait of Georgia. *Can. J. Fish and Aquatic Sci.* 44:525-537.

Beckham, S.D. 2001. Oregon History. *In* Oregon Blue Book 2001-2001, Office of the Secretary of State, Salem, OR. pp.338-382.

Beechie T.G. Pess, P. Kennard, R. Bilby, and S. Bolton. 2000. Modeling recovery rates and pathways for woody debris recruitment in Northwestern Washington streams. *North American Journal of Fisheries Management* 20:436-452.

Belden, L.K. and A.R. Blaustein. 2002. Exposure of red-legged frog embryos to ambient UV-B radiation in the field negatively affects larval growth and development. *Oecologia* 130:551-554.

Belt, G.H., J. O’Laughlin, and T. Merrill. 1992. Design of Forest Riparian Buffer Strips for the Protection of Water Quality: Analysis of Scientific Literature. Report No. 8. University of Idaho; Idaho Forest, Wildlife and Range Policy Group; Moscow ID.

Benda L.E. 1990. The Influence of Debris Flows on Channels and Valley Floors in the Oregon Coast Range, USA. *Earth Surface Processes and Landforms* 15: 457-466, Chichester, UK.

- Benda, L.E. and T.W. Cundy.** 1990. Predicting Deposition of Debris Flows in Mountain Channels. *Canadian Geotechnical Journal*, Vol.27, No. 4, pp. 409-417. p. 5-66.
- Benda L. and T. Dunne.** 1997. Stochastic forcing of sediment supply to channel networks from landsliding and debris flow. *Water Resources Research*. Vol 33 No 12. PP 2849-2863
- Benda, L.E. and J.C. Sias.** 1998. Landscape Controls on Wood Abundance in Streams. Earth Systems Institute, Seattle, Washington. (www.Earthsi@aol.com.)
- Benda L., C. Veldhuisen, and J. Black.** 2003. Debris flows as agents of morphological heterogeneity at low-order confluences, Olympic Mountains, Washington. *Geological Society of America Bulletin*. V. 115; no. 9; p. 1110 – 11121.
- Bent, A.C.** 1937. Life histories of North American birds of prey, Part 1. U.S. Natl. Mus. Bull. 167 (Reprinted in 1961 by Dover Publ., New York).
- Bent, Arthur.** 1961. Life Histories of North American Birds of Prey. Dover, New York. 409 pp.).
- Beschta, R.L.** 1978. Long-term patterns of sediment production following road construction and logging in the Oregon Coast Range. *Water Resources Research* 14(6): 1011-1016
- Beschta, R.L., R.E. Bilby, G.W. Brown, L.B. Holtby, and T.D. Hofstra.** 1987. Stream Temperature and Aquatic Habitat: Fisheries and Forestry Interactions. *In* Salo and Cundy, editors; *Streamside Management: Forestry and Fishery Interactions*. University of Washington, Institute of Forest Resources Contribution 57, Seattle, WA.
- Beschta, R.L., M.R. Pyles, A.E. Skaugset, and C.G. Surfleet.** 2000. Peakflow responses to forest practices in the western cascades of Oregon, USA. *Journal of Hydrology*. 233. 102-120.
- Bilby R.E.** 1985a. Contributions of road surface sediment to a Western Washington stream. *Forest Science*. Vol.31. No. 4. pp. 827-838.
- Bilby, R.E.** 1985b. Influence of Stream Size on The Function and Characteristics of Large Organic Debris. *In* Proceedings of the West Coast Meeting of The National Council and Paper Industry for Air and Stream Improvement, Portland, Oregon.
- Bilby, R.E. and P.A. Bisson.** 1998. Function and Distribution of Large Woody Debris. *In* River Ecology and Management: Lessons from The Pacific Coastal Ecosystem. Springer. pp. 324-346.
- Bilby, R.E., K. Sullivan, and S.H. Duncan.** 1989. The Generation and Fate of Road-Surface Sediment in Forested Watersheds in Southwestern Washington. *Forest Science*. Vol. 35, No. 2: 453-468.

Biosystems, Waterwork Consulting, Coos Watershed Association, Alsea Geospatial, Inc. and Karen Bahus. 2003. Elliott State Forest Watershed Analysis. Prepared for the Oregon Department of Forestry. October 2003. 14pp. Website: http://egov.oregon.gov/ODF/STATE_FORESTS/watershed.shtml#Elliott_State_Forest_Analysis

Bisson P.A. and P.E.Reimers. 1977. Geographic variation among Pacific Northwest populations of longnose dace, *Rhinichthys cataractae*. *Copeia*. 1977: 518-522.

Bisson, P.A., R.E. Bilby, M.D. Bryant, C.A. Dolloff, G.B. Grette, R.A. House, M.L. Murphy, K.V. Koski, and J. R. Sedell. 1987. Large Woody Debris in Forested Streams in the Pacific Northwest: Past Present and Future. *In* Salo and Cundy, editors; Streamside Management: Forestry and Fishery Interactions. University of Washington, Institute of Forest Resources Contribution 57, Seattle, WA.

Bisson, P.A., T.P. Quinn, G.H. Reeves, and S.V. Gregory. 1992. Best Management Practices, Cumulative Effects, and Long-Term Trends in Fish Abundance in Pacific Northwest River Systems. *In* R.J. Naiman, editor; Watershed Management: Balancing Sustainability and Environmental Change. Springer-Verlag. 542 pp.

Bjornn, T.C. and D.W. Reiser. 1991. Habitat Requirements of Salmonids in Streams. *In* W.R. Meehan, editor; Influences of Forest and Rangeland Management on Salmonid Fishes and Their Habitats. Special Publication 19. American Fisheries Society, Bethesda, MD.

Blakesley, J.A. 2004. Chapter 5: Habitat Associations. *In* S.P. Courtney, J.A. Blakesley, R.E. Bigley, M. L. Cody, J.P. Dumbacher, R.C. Fleischer, A.B. Franklin, J.F. Franklin, R.J. Gutierrez, J.M. Marzluff, L. Sztukowski. Final Report. Scientific Evaluation of the Status of the Northern Spotted Owl. Sustainable Ecosystems Institute, Portland, Oregon. Pp. 5-1 – 5-26.

Blakesley, J.A., A.B. Franklin, and R.J. Gutierrez. 1992. Spotted Owl Roost and Nest Site Selection in Northwestern California. *Journal of Wildlife Management* 56(2):388-392.

Blaustein, A.R., J.J. Beatty, D. Olson and R.M. Storm. 1995. The Biology of Amphibians and Reptiles in Old-Growth Forests in the Pacific Northwest. U.S. Forest Service General Technical Report PNW-GTR-337.

Boal, C.W. and R.W. Mannan. 1994. Northern Goshawk diets in ponderosa pine forests on the Kiabab Plateau. *In* W.M. Block, M.L. Morrison, and M.H. Resier, eds. The northern goshawk: Ecology of management. *Studies in Avian Biology* 16:97-102.

Botkin, D.B., K. Cummins, T. Dunne, H. Regier, M. Sobel, and L. Talbot. 1995. Status and Future of Salmon of Western Oregon and Northern California: Findings and Options. The Center for the Study of the Environment, P.O. Box 6945, Santa Barbara, CA 93160, research report 951002.

Bourhill, R. 1994. History of Oregon's Timber Harvests and/or Lumber Production: State Data – 1849 to 1992, County Data 1925 to 1992. Oregon Department of Forestry, 2600 State Street, Salem, OR 97310.

Brazier, J.R. and G.W. Brown. 1973. Buffer Strips for Stream Temperature Control. Research Paper 15. Oregon State University, Forest Research Laboratory, Corvallis, OR.

Brown, H.A. 1975. Temperature and Development of the Tailed Frog, *Ascaphus truei*. Comp. Biochem. Physiology 50:397-405..

Brown and Krygier. 1970. Effects of clear-cutting on stream temperature. Water Resource. Res. 6: 1133-1139.

Buchanan, J.B., L.L. Irwin, and E.L. McCutchen. 1993. Characteristics of spotted owl nest trees in the Wenatchee National Forest. Journal of Raptor Research 27:1-7.

Buck, S.G., C. Mullis, A.S. Mossman, I. Show, and C. Coolahan. 1994. Habitat Use by Fishers in Adjoining Heavily and Lightly Harvested Forest. Pp. 368-376. In Buskirk, S.W., A. S. Harestad, M.G. Raphael, and R.A. Powell, editors; Martens, Sables, and Fishers: Biology and Conservation. Cornell University Press, Ithaca, NY. 684 pp.

Bull, E.L. 1987. Ecology of the pileated woodpecker in northeastern Oregon. Journal of Wildlife Management 51(2): 472-481.

Bull, E.L. 2003. Pileated woodpecker. Pp. 372-374. In Birds of Oregon: A General Reference. D.B. Marshall, M.G. Hunter, and A.L. Contreras, Eds. Oregon State University Press, Corvallis, OR.

Bull, E.L. and B.E. Carter. 1996. Tailed Frogs: Distribution, Ecology, and Association with Timber Harvest in Northeastern Oregon. Res. Pap. PNW-RP-497. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 11pp.

Bull, E.L. and J.H. Hohmann. 1994. Breeding biology of Northern Goshawks in northeastern Oregon. Studies in Avian Biology 16:103-105.

Bull, E.L. and R.S. Holthausen. 1993. Habitat use and management of pileated woodpeckers in northeastern Oregon. Journal of Wildlife Management 57(2): 335-345.

Bull, E.L., R.S. Holthausen, and M.G. Henjum. 1992. Roost trees used by pileated woodpeckers in northeastern Oregon. Journal of Wildlife Management 56(4): 786-793.

Burger, Alan E. 1997. Behavior and numbers of marbled murrelets measured with radar. Journal of Field Ornithology 68: 208-223.

Burger, Alan E. 2001. Using radar to estimate populations and assess habitat associations of marbled murrelets. Journal of Wildlife Management 65: 696-715.

Burger, Alan E. 2002. Conservation assessment of marbled murrelets in British Columbia: a review of the biology, populations, habitat associations, and conservation. Technical Report Series Number 387, Canadian Wildlife Service Pacific and Yukon Region, British Columbia.

Bury, B.R. 1983. Differences in Amphibian Populations in Logged and Old Growth Redwood Forest. *Northwest Science* 57:167-178.

Bury, B.R., and P.S. Corn, K.B. Aubry, F.F. Gilbert, L.L.C. Jones. 1991. Aquatic Amphibian Communities in Oregon and Washington. *In* L.F. Ruggiero, and K.B. Aubry, A.B. Carey, M.H. Huff, technical editors; *Wildlife and Vegetation of Unmanaged Douglas-Fir Forests*, pp. 353-362. USDA Forest Service, Pacific Northwest Research Station, Portland, OR. PNW-GTR-285.

Bury, Bruce. 2002. Personal communication to Marnie Allbritten, Oregon Department of Fish and Wildlife, Southwest Region Office, 4192 N. Umpqua Highway, Roseburg, OR 97470, January 28, 2002.

Buskirk, S.W. and R.A. Powell. 1994. Habitat Ecology of Fishers and American Martens. Pp. 283-296. *In* Buskirk, S.W., A.S. Harestad, M.G. Raphael, and R.A. Powell, editors; *Martens, Sables, and Fishers: Biology and Conservation*. Cornell University Press, Ithaca, NY. 684 pp.

Buskirk, S.W., A.S. Harestad, M.G. Raphael, and R.A. Powell (Eds.). 1994. *Martens, Sables, and Fishers: Biology and Conservation*. Cornell University Press, Ithaca, NY. 684 pp.

Carey, A.B., S.P. Horton, and B.L. Biswell. 1992. Northern Spotted Owls: Influence of Prey Base and Landscape Character. *Ecological Monographs* 62(2):223-250.

Carlson, J.Y., C.W. Andrus, and H.A. Froehlich. 1990. Woody debris, channel features, and macroinvertebrates of streams with logged and undisturbed riparian timber in northeastern Oregon, USA. *Canadian Journal of Aquatic Sciences* 47:1103-1111.

Carroll, Carlos, William J. Zielinski, and Reed F. Noss. 1999. Using presence-absence data to build and test spatial habitat models for the fisher in the Klamath Region, U.S.A. *Conservation Biology* 13(6): 1344-1359.

Chamberlin, T.W., R.D. Harr, and F.H. Everest. 1991. Timber Harvesting, Silviculture, and Watershed Processes. *In* W.R. Meehan, editor; *Influences of Forest and Rangeland Management on Salmonid Fishes and Their Habitats*. Special Publication 19. American Fisheries Society, Bethesda, MD.

Chan S.S., P.D. Anderson, R.J. Danehy, and M.L. Reiter. 2006. Riparian microclimate, overstory canopy and understory vegetation and soil moisture relationships to a range of timber harvest prescriptions in headwater forests of the central Oregon coast range. 2006. Presented at the 2005 Science and Management of Headwater Streams in the Pacific Northwest, Corvallis, OR. Abstracts from presentations posted: <http://www.headwatersresearch.org/> Page 18

Coats R.L., L. Collins, J. Forsheim, and D. Kaufman. 1985. Channel changes, sediment transport and fish habitat in a coastal stream. *Environmental Management* 9(1): 35–48.

Compton J.E., M. Robbins Church, S.T. Larned, and W.E. Hogsett. 2003. Nitrogen export from forested watersheds in the Oregon Coast Range: The role of N₂-fixing red alder. *Ecosystems*. 6:8 pp 773-785.

Conroy, Catherine J., Volker Bahn, Michael S. Rodway, Laurie Ainsworth, and Deanna Newson. 2002. Estimating nest densities for marbled murrelets in three habitat suitability categories in the Ursus Valley, Clayoquot Sound. Pages 121-137 *in* Multi-scale studies of populations, distribution and habitat associations of marbled murrelets in Clayoquot Sound, British Columbia. A.E. Burger and T.A. Chatwin (editors). Ministry of Water, Land and Air Protection, Victoria, British Columbia.

Cook, D. and M.R. Jennings. 2001. *Rana aurora draytonii*. Predation. *Herpetological Review* 32:182-183.

Cooper, Brian A. and Kimberly H. Augenfeld. 2001. Radar surveys for marbled murrelets in the Elliott State Forest, Oregon 2001. Unpublished report submitted to Oregon Department of Forestry.

Corkran, C.C. and C. Thoms. 1996. Amphibians of Oregon, Washington and British Columbia. Lone Pine Publishing, Edmonton, Alberta. 175 pp.

Corn, P.S. and R.B. Bury. 1989. Logging in western Oregon: Responses of Headwater Habitats and Stream Amphibians. *Forest Ecology and Management* 29:39-57.

Corn, P.S. and R.B. Bury. 1991. Terrestrial Amphibian Communities in the Oregon Coast Range. *In* L.F. Ruggiero, K.B. Aubry, A.B. Carey, and M.H. Huff, technical coordinators, Wildlife and Vegetation of Unmanaged Douglas-Fir Forests. USDA Forest Service General Technical Report PNW-GTR-285, Portland, OR.

Crocker-Bedford. 1990. Goshawk reproduction and forest management. *Wildlife Society Bulletin* 18:262-269.

Crookham, C.S. 1992. 46 Op. Atty. Gen. 468 (1992), Opinion No. 8223, July 24, 1992. Opinion of Oregon Attorney General Charles S. Crookham on legal mandates for managing Common School Forest Lands.

Csuti, B., A.J. Kimerling, T.A. O’Neil, M.M. Shaughnessy, E.P. Gaines, and M.M.P. Huso. 1997. Atlas of Oregon Wildlife: Distribution, Habitat, and Natural History. Corvallis: Oregon State University Press. 492 pp.

Davidson, C., H.B. Shaffer, and M.R. Jennings. 2001. Declines of the California red-legged frog: climate, UV-B, habitat, and pesticides hypotheses. *Ecological Applications* 11:464-479.

Davidson, C., H.B. Shaffer, and M.R. Jennings. 2002. Spatial tests of the pesticide drift, habitat destruction, UV-B, and climate-change hypotheses for California amphibian declines. *Conservation Biology* 16:1588-1601.

Daw, Sonya K. 1996. Northern Goshawk nest site selection and habitat associations at the post-fledging family area scale in Oregon. M.S. thesis, *Oreg. State Univ., Corvallis.*

DellaSala, D.A., C.L. Thomas, and R.G. Anthony. 1987. Wintering Ecology of Bald Eagles (*Haliaeetus leucocephalus*) in the Willamette Valley and Coburg Hills of Western Oregon. Unpublished report, Bureau of Land Management, Eugene, OR. 22 pp.

deMaynadier, P.G. and M.L. Hunter, Jr. 1995. The Relationship between Forest Management and Amphibian Ecology: A Review of the North American Literature. *Environmental Reviews* 3:230-261.

Dent L. 2001. Harvest effects on riparian function and structure under current forest practice rules. ODF Technical Report #12. Oregon Department of Forestry, 2600 State Street. Salem, Oregon. 97310.

Dent, E.F., K.A. Mills, and J. Robben. 2003. Turbidity Off of Forest Roads in Oregon. In *Conf. Proc.: Total Maximum Daily Load (TMDL) Environmental Regulations—II.* American Society of Agricultural and Biological Engineers. Albuquerque, New Mexico. pp. 191-197.

Desimone, S.M. 1997. Occupancy rates and habitat relationships of Northern Goshawks in historic nesting areas in Oregon. M.S. thesis, *Oreg. State Univ., Corvallis.*

DeStefano, S. and J. McCloskey. 1997. Does Vegetation Structure Limit the Distribution of Northern Goshawks in the Oregon Coast Ranges. *Journal of Raptor Research* 31:34-39.

DeStefano, S., S.K. Daw, S.M. Desimone, and E.C. Meslow. 1994. Density and Productivity of Northern Goshawks: Implications for Monitoring and Management. *Studies in Avian Biology* 16:88-91.

Dietrich and Dunne. 1978. Sediment budget for a small catchment in mountainous terrain. *Z. Geomorph. N.F. Suppl. Bd. 29:* 191-206.

Diller, Lowell V. and Richard L. Wallace. 1996. Distribution and habitat of *Rhyacotriton variegatus* on managed, young growth forests in north coastal California. *Journal of Herpetology* 30(2):184-191.

Duncan, Nancy. 2004. Bureau of Land Management – Roseburg District. Personal communication to Marnie Allbritten, Oregon Department of Fish and Wildlife, Assistant District Wildlife Biologist – Southwest Region Office, 4192 N. Umpqua Hwy, Roseburg, OR 97470..

Duncan, S.H. and E.C. Steinbrenner. 1973. Soil Survey of the Elliott State Forest by S.H. Duncan and E.C. Steinbrenner. 1973. Compiled by Weyerhaeuser Company for the Oregon Department of Forestry.

Duncan, S.H. and J.W. Ward. 1985. The influence of watershed geology and forest roads on the composition of salmon spawning gravel. *Northwest Science*. 59:204-212. Everest et al. 1987.

Dunster, Julian and Katherine Dunster. 1996. Dictionary of Natural Resource Management. UBC Press, University of British Columbia, Vancouver, Canada.

Dupuis, L. and D. Steventon. 1999. Riparian Management and the Tailed Frog in Northern Coastal Forests. *Forest Ecology and Management* 124:35-43.

Eltzroth, E.K. 2003. Western Bluebird. Pp. 474-477. *In* Birds of Oregon: A General Reference. D.B. Marshall, M.G. Hunter, and A.L. Contreras, Eds. Oregon State University Press, Corvallis, OR.

Emmett, R.L., S.L. Stone, S.A. Hinton, and M.E. Monaco. 1991. Distribution and abundance of fishes and invertebrates in West coast estuaries, Volume 2: Species life histories summaries. ELMR Rep. No. 8. NOS/NOAA Strategic Environmental Assessment Division. Rockville, MD.

Erdman, T.C., D.F. Brinker, J.P. Jacobs, J. Wilde, and T.O. Meyer. 1998. Productivity, population trend, and status of northern goshawks, *Accipiter gentiles atricapillus*, in northern Wisconsin. *Canadian Field-Naturalist*.

Evans, D.M., W.P. Ritchie, S.K. Nelson, E. Kuo-Harrison, Peter Harrison, and T.E. Hamer (Eds.). 2000. Methods for surveying marbled murrelets in forests: an update to the protocol for land management and research, by the Pacific Seabird Group Marbled Murrelet Technical Committee. 64 pp.

Everest, F.H., R.L. Beschta, J.C. Scrivener, K.V. Koski, J.R. Sedell and C.J. Cederholm. 1987. Fine Sediment and Salmonid Production: A Paradox. *In* Salo, E.O., editor, Streamside Management, Forestry and Fishery Interactions. College of Forest Resources, Univ. of Washington. Pp 98-142.

Farris, K.L., E.O. Garton, P.J. Heglunch, S. Zack, and P.J. Shea. 2002. Woodpecker foraging and the successional decay of ponderosa pine. Pages 237-246. *In* Laudenslayer Jr., William F., Patrick J. Shea, Bradley E. Valentine, C. Phillip Weatherspoon, and Thomas E. Lisle, technical coordinators. Proceedings of the symposium on the ecology and management of dead wood in western forests. 1999 November 2-4; Reno, NV. Gen Tech. Rep. PSW-GTR-181. Albany, CA; Pacific Southwest Research Station, Forest Service, U.S. Department of Agriculture.

Fick, L. and G. Martin. 1992. The Tillamook Burn: Rehabilitation and Reforestation. Oregon Department of Forestry, Forest Grove District, Forest Grove, OR.

- Fields, R.** 2005. Project Coordinator, Oregon Department of Forestry. Personal communication with Alan Ritchey, Fisheries Biologist, Oregon Department of Fish and Wildlife.
- Folz, R.B.** 1996. Traffic and No-traffic on an Aggregate Surfaced Road: Sediment Production Differences. Paper presented at FAO seminar on Environmentally Sound Forest Road and Wood Transport, Sinaia, Romania.
- Forman, R.T., T.D. Sperling, J.A. Bissonette, A.P. Clevenger, C.D. Cutshall, V.H. Dale, L. Fahrig, R. France, C.R. Goldman, K. Heanue, J.A. Jones, F.J. Swanson, T. Turrentien, and T. Winter.** 2003. Road Ecology. Island Press., Washington, D.C.
- Forsman, E.D., E.C. Meslow, and H.M. Wight.** 1984. Distribution and Biology of the Spotted Owl in Oregon. Wildlife Monographs 87:1-64.
- Franklin, J.F. and C.T. Dyrness.** 1988. Natural Vegetation of Oregon and Washington. Oregon State University Press, Corvallis, OR.
- Frazier, Matthew W.** 1997. Roost site characteristics of the long legged myotis (*Myotis volans*) in the Teanaway River Valley of Washington. M.S. Thesis. University of Washington. 58 pp.
- Frenzel, R.W.** 1984. Environmental Contaminants and Ecology of Bald Eagles in Southcentral Oregon. Unpublished doctoral dissertation, Oregon State University, Corvallis, OR. 143 pp.
- Furness, M.J., T.D. Roelofs, and C.S. Yee.** 1991. Road Construction and Maintenance. *In* W. R. Meehan, editor; Influences of Forest and Rangeland Management on Salmonid Fishes and Their Habitats. Special Publication 19. American Fisheries Society, Bethesda, MD.
- Gabrielson, I.N. and S.G. Jewett.** 1940. Birds of Oregon. *Oreg. State Coll.*, Corvallis. (Reprinted in 1970 as Birds of the Pacific Northwest by Dover Publ., New York.)
- Gallo, Kirsten; Lanigan, Steven H.; Eldred, Peter; Gordon, Sean N.; Moyer, Chris.** 2005. Northwest Forest Plan—The First 10 years (1994–2003): Preliminary Assessment of the Condition of Watersheds. Gen. Tech. Rep. PNW-GTR-647. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 133 p.
- Gardener, R.H., B.T. Milne, M.G. Turner, and R. V. O'Neill.** 1987. Neutral Models for the Analysis of Broad-Scale Landscape Pattern. *Landscape Ecology* 1: 19-28.
- Garrett, M.G., J.W. Watson, and R.G. Anthony.** 1993. Bald Eagle Home Range and Habitat Use in the Columbia River Estuary. *Journal of Wildlife Management*, 57:19-27.
- Gibilisco, C.J.** 1994. Distributional Dynamics of Modern *Martes* in North America. Pp. 59-71. *In* Buskirk, S.W., A.S. Harestad, M.G. Raphael, and R.A. Powell, editors; Martens, Sables, and Fishers: Biology and Conservation. Cornell University Press, Ithaca, NY. 684 pp.

Good, D.A. and D.B. Wake. 1992. Geographic Variation and Speciation in the Torrent Salamanders of Genus *Rhyacotriton* (Caudata: Rhyacotritonidae). University of California Publications, Zool. 126:1-91.

Graham, R.T., R.L. Rodriquez, K.M. Paulin, R.L. Player, A.P. Heap, and R. Williams. 1999. The Northern Goshawk in Utah: habitat assessment and management recommendations. USDA Forest Service, Rocky Mountain Research Station, General Technical Report RMRS-GTR-22. 48 pp.

Grant, G. E. 1987. Assessing effects of peak flow increases on stream channels: a rational approach. *In* Proceedings of the California watershed management conference. 1986 November 18-20. R.Z. Callaham and J.J. DeVries, technical coordinators. Wildland Resources Center Report Number 11. University of California Berkeley, California, USA.

Gregory, S.V., G.A. Lamberti, D.C. Erman, K.V. Koski, M.L. Murphy, and J. R. Sedell. 1987. Influence of Forest Practices on Aquatic Production. *In* Salo and Cundy, editors; Streamside Management: Forestry and Fishery Interactions. University of Washington, Institute of Forest Resources Contribution 57, Seattle, WA.

Groot, C. and L. Margolis. 1991. Pacific Salmon: Life Histories. University of British Columbia Press, Vancouver, British Columbia, Canada.

Gutierrez, R., M. Coyd, S. Clourtney, and D. Kennedy. 2004. Chapter 7: Barred Owls. *In* S.P. Courtney, J.A. Blakesley, R.E. Bigley, M.L. Cody, J.P. Dumbacher, R.C. Fleischer, A.B. Franklin, J.F. Franklin, R.J. Gutierrez, J.M. Marzluff, L. Sztukowski. Final Report. Scientific Evaluation of the Status of the Northern Spotted Owl. Sustainable Ecosystems Institute, Portland, Oregon. Pp. 7-1 – 7-51.

Haines, K.F. 1995. Northern goshawk breeding habitat in conifer stands with natural tree mortality in eastern Oregon. Boise State University. M.S. thesis. 55 pp.

Hamer and Meekins. 1996. A predictive model of habitat suitability for the marbled murrelet and habitat rating strategy for the Elliott State Forest. Report by Hamer Environmental to the Oregon Department of Forestry. 45 pp.

Hamer, T.E. and S.K. Nelson. 1995. Nesting Chronology of the Marbled Murrelet. *In* Ralph, C.J., G.I. Hunt, Jr., M.G. Raphael, and J.F. Piatt, editors; Ecology and Conservation of the Marbled Murrelet. Pacific Southwest Research Station, USDA Forest Service, Albany, California. General Technical Report PSW-GTR-152. Pp. 49-56.

Hamer Environmental. 2002. Marbled murrelet habitat measurements for the Elliott State Forest. Report to the Oregon Department of Forestry. 13 pages.

Hansen, A.J., W.C. McComb, R. Vega, M.G. Raphael, and M. Hunter. 1995. Bird habitat relationships in natural and managed forest in the western Cascades of Oregon. *Ecological Applications* 5:555-569.

- Harestad, A.S. and D.G. Keisker.** 1989. Nest tree use by primary cavity-nesting birds in south central British Columbia. *Can. J. Zool.* 67: 1067-1073.
- Harr, R.D.** 1976. Forest practices and stream flow in western Oregon. USDA Forest Service General Technical Report PNW-49. Pacific Northwest Forest and Range Station, Portland, Oregon, USA.
- Harris, D.D.** 1977. Hydrologic changes after logging in two small Oregon coastal watersheds. United States Geologic Survey Water Supply Paper. 2037. Washington DC, USA.
- Hartwig, C.L., D.S. Eastman, and A.S. Harestad.** 2004. Characteristics of pileated woodpecker (*Dryocopus pileatus*) cavity trees and their patches on southeastern Vancouver Island, British Columbia, Canada. *Forest Ecology and Management* 187: 225-234.
- Hayes, J.P.** 1996. *Arborimus longicaudus*. *Mammalian Species* 532:1-5.
- Hayes, M.P.** 1999. Species Account for *Rhyacotriton kezeri*. Oregon Department of Fish and Wildlife, Salem, OR. 19 pp.
- Hayes J.P., M.D. Adam, D. Bateman, E. Dent, W.H. Emmingham, K.G. Maas, and A.E. Skaugset.** 1996. Integrating research and forest management in riparian areas of the Oregon Coast Range. *West. J. Appl. For.* 11: 85-89.
- Heimann, D.C.** 1988. Recruitment trends and physical characteristics of coarse woody debris in Oregon Coast Range streams. M.S. thesis. Oregon State University.
- Hibbs David E. and Peter A. Giordano.** 1996. Vegetation Characteristics of Alder-Dominated Riparian Buffer Strips in the Oregon Coast Range. *Northwest Science*, Vol. 70, No. 3. pp 213 – 222.
- Hicks, B.J., J.D. Hall, P.A. Bisson, and J.R. Sedell.** 1991. Responses of Salmonids to Habitat Changes. *In* W.R. Meehan, editor; *Influences of Forest and Rangeland Management on Salmonid Fishes and Their Habitats*. Special Publication 19. American Fisheries Society, Bethesda, MD.
- Holling, C.S. (Ed.).** 1978. Adaptive Environmental Assessment and Management. International Series on Applied Systems Analysis 3. John Wiley and Sons, London. 377 pp.
- Hull, Cindy L., Gary W. Kaiser, Cecilia Lougheed, Lynn Lougheed, Sean Boyd, and Fred Cooke.** 2001. Intraspecific variation in commuting distance of marbled murrelets (*Brachyramphus marmoratus*): ecological and energetic consequences of nesting further inland. *Auk* 118: 1036-1046.
- Hunnicut, M.** 1989. Roosting and Staging Activities of Bald Eagles in the Coburg Hills of Western Oregon. Unpublished report, Bureau of Land Management, Eugene, OR. 12 pp.

Hunter, M.G. 1998. Watershed-level Patterns Among Stream Amphibians in the Blue River Watershed, West-Central Cascades of Oregon. M.S. thesis. Oregon State University. 97 pp.

Hunter, M.G. 2003. A review and synthesis of post-1990 literature on the Northern Spotted Owl (*Strix occidentalis caurina*). Unpublished report to Oregon Department of Forestry, Salem, OR. 79 pp.

Independent Multidisciplinary Science Team. 1999. Recovery of Wild Salmonids in Western Oregon Forests: Oregon Forest Practices Act Rules and the Measures in the Oregon Plan for Salmon and Watersheds, Governor's Natural Resources Office, Salem, Oregon.

Ingram, R. 1973. Wolverine, Fisher and Marten in Central Oregon. Central Region Adm., Report 73-2, Oregon State Game Commission, Bend, OR. 41 pp.

Isaacs, F.B. and R.G. Anthony. 1998. Results of the 1998 bald eagle nest survey. Memo to cooperators from Oregon Cooperative Fish and Wildlife Research Unit. October 1, 1998.

Isaacs, F.B. and R.G. Anthony. 2002. Results of the 2002 bald eagle nest survey. Memo to cooperators from Oregon Cooperative Fish and Wildlife Research Unit. October 1, 2002.

Isaacs, F.B. and R.G. Anthony. 2003a. Results of the 2003 bald eagle nest survey. Memo to cooperators from Oregon Cooperative Fish and Wildlife Research Unit. October 1, 2003.

Isaacs, F.B. and R.G. Anthony. 2003b. Bald Eagle. Pp. 140-144 *In* Birds of Oregon: A General Reference. D.B. Marshall, M.G. Hunter, and A.L. Contreras, Eds. Oregon State University Press, Corvallis, OR.

Isaacs, F.B., R.G. Anthony, and R.J. Anderson. 1983. Distribution and Productivity of Nesting Bald Eagles in Oregon, 1978-1982.

Isaacs, F.B., R.G. Anthony, M. Vander Heyden, C.D. Miller, and W. Weatherford. 1996. Habits of Bald Eagles Wintering along the Upper John Day River, Oregon. *Northw. Sci.* 70(1):1-9.

Jackson C.R., D. P. Batzer, S.S. Cross, S. M. Haggerty, and C.A. Sturm. 2006. Abiotic and Biotic Responses of Headwater Streams to Adjacent Timber Harvest: Results of a Four-Year Manipulative Study. Presented at the 2005 Science and Management of Headwater Streams in the Pacific Northwest, Corvallis, OR. Abstracts from presentations posted: <http://www.headwatersresearch.org/> Page 26

Jacobs S.E. and C.X. Cooney. 1997. Oregon coastal salmon spawning surveys, 1994-95. Oregon Department of Fish and Wildlife, Ocean Salmon Management Information Report, Portland, OR.

Jacobs, S., J. Firman, G. Susac, E. Brown, B. Riggers, and K. Tempel. 2000. Status of Oregon coastal stocks of anadromous salmonids. Monitoring Program Report Number OPSW-ODFW-2000-3, Oregon Department of Fish and Wildlife, Portland, Oregon.

- Jenkins, Jeff, Robert Gill, Todd Reinwald, and Dave Vesely.** 2005. Miami Watershed Analysis. Prepared for the Oregon Department of Forestry. August 2005. 27 pp.
- Jobanek, G.** 1993. The European Starling in Oregon. *Oreg. Birds* 19:93-96.
- Jodice, Patrick G.R. and Michael W. Collopy.** 2000. Activity patterns of marbled murrelets in douglas-fir old-growth forests of the Oregon coast range. *Condor* 102: 275-285.
- Johnson, S.** 2004. Factors influencing stream temperature in small streams: substrate and a shading experiment. *Can. J. Fish. Aquat. Sci.* 61:913-923.
- Johnson, D.H. and T.A. O'Neil.** 2001. (Manag. Dirs.) Wildlife – Habitat Relationships in Oregon and Washington. Oregon State University Press, Corvallis, Oregon.
- Jones, J.A. and G.E. Grant.** 1996. Peak flow responses to clear-cutting and roads in small and large basins, western Cascades, Oregon. *Water Resources Research* Vol. 32 No. 4. Pp. 959-974.
- Jones, J.L. and E.O. Garton.** 1994. Selection of Successional Stages by Fishers in North-Central Idaho. Pp. 377-387. *In* Buskirk, S.W., A.S. Harestad, M.G. Raphael, and R.A. Powell, editors; *Martens, Sables, and Fishers: Biology and Conservation*. Cornell University Press, Ithaca, NY. 684 pp.
- Kahler, T.H and T.P. Quinn.** 1998. Juvenile and Resident Salmonid Movement and Passage through Culverts. Fisheries Research Institute. School of Fisheries. 357980. University of Washington. Seattle, Washington. 98195-7980. Prepared for Washington State Transportation Commission.
- Karraker, N.E. and G.S. Beyersdorf.** 1997. A Tailed Frog (*Ascaphus truei*) Nest Site in Northwestern California. *Northwestern Naturalist* 78:110-111.
- Kaufmann P.** 1987. Channel morphology and hydraulic characteristics of torrent-impacted forest streams in the Oregon Coast Range, USA. PhD. Dissertation, Oregon State University, Corvallis, Oregon. 235 pp.
- Kavanagh P.S., K.K. Jones, C.H. Stein, and P.S. Jacobsen.** 2005. Fish Habitat Assessment in the Oregon Department of Forestry Elliott study area. Conservation and Recovery Program, Oregon Department of Fish and Wildlife, 28655 Highway 34, Corvallis OR 97333.
- Keinath D.A.** 2004. Fringed Myotis (*Myotis thysanodes*): A Technical Conservation Assessment [online] www.fs.fed.us/r2/projects/scp/assessments/fringedmyotis.pdf.
- Keister Jr., G.P.** 1981. Characteristics of winter roosts and populations of bald eagles in the Klamath Basin. M.S. Thesis, Oregon State University, Corvallis. 82 pp.
- Keister Jr., G.P. and R.G. Anthony.** 1983. Characteristics of Bald Eagle Communal Roosts in the Klamath Basin, Oregon and California. *Journal of Wildlife Management*, 47:1072-1079.

- Keister, G.P., R.G. Anthony, and E.J. O'Neill.** 1987. Use of Communal Roosts and Foraging Areas by Bald Eagles Wintering in the Klamath Basin. *Journal of Wildlife Management*, 51:415-420.
- Keller, E.A. and F.J. Swanson.** 1979. Effects of Large Organic Material on Channel Form and Fluvial Process. *Earth Surface Processes* 4:361-380.
- Ketcheson G. and H.A. Froehlich.** 1978. Hydrologic factors and environmental impacts of mass soil movements in the Oregon Coast Range. *Waters Resources Research Institute Bulletin* 56, Oregon State University, Corvallis Oregon.
- Kiesecker, J.M. and A.R. Blaustein.** 1998. Effects of introduced bullfrogs and smallmouth bass on microhabitat use, growth, and survival of native red-legged frogs (*RANA AURORA*). *Conservation Biology* 12:776-787.
- King, James E.** 1966. Site index curves for Douglas-fir in the Pacific Northwest. *Weyerhaeuser Forestry Paper No. 8*. Centralia, WA. 49 p.
- King, J.G. and L.C. Tennyson.** 1984. Alteration of streamflow characteristics following road construction in north central Idaho. *Water Resources Research* 20(8):1159-1163.
- Kingfisher Ecological.** 2003. Final Report: Northern spotted owl surveys Elliott State Forest 2003. Unpublished report to Oregon Department of Forestry, Salem, OR; 35 pp.
- Kitzhaber M.D., John (Governor of Oregon).** 1999. Executive Order Number EO 99-01, "The Oregon Plan for Salmon and Watersheds." Office of the Governor, Salem, OR. 18 pp.
- Knutson, K.L. and V.L. Naef.** 1997. Management Recommendations for Washington's Priority Habitats: Riparian. Washington Department of Fish and Wildlife, Olympia, WA.
- Kostow, K.** 2002. Oregon lampreys: Natural history status and analysis of management issues. Oregon Department of Fish and Wildlife, Salem, OR.
- Lamberti G.A., S.V. Gregory, L.R. Ashkenas, R.C. Wildman, and K.M.S. Moore.** 1991. Stream ecosystem recovery following a catastrophic debris flow. *Can. J. Fish. Aq. Sci.* 48 pp 196-208.
- Lee D.S., C.R. Gilbert, C.H. Hocutt, R.E. Jenkins, D.E. McAllister, and J.R. Stauffer.** 1980. Atlas of North American Freshwater Fishes. Publication #1980-12, Northern Carolina Biological Survey. Pp24-34.
- Leonard, W.P., H.A. Brown, L.L.C. Jones, K.R. McAllister, and R.M. Storm.** 1993. Amphibians of Washington and Oregon. Seattle Audubon Society, Seattle, WA. 168 pp.
- Lettman, G.J.** 1995. Timber Management Practices and Land Use Trends on Private Forest Land in Oregon: A Final Report to the Sixty-Eighth Oregon Legislative Assembly. Oregon Department of Forestry, Forest Resources Planning Program, Salem, OR. May 1995.

- Levno, A. and J. Rothacher.** 1967. Increases in Maximum Stream Temperatures After Logging in Old-Growth Douglas-Fir Watersheds. USDA Forest Service Research Note PNW-65 PNW Forest and Range Experiment Station, Portland, Oregon. 12 p.
- Lewis, J.C. and D.W. Stinson.** 1998. Washington State Status Report for the Fisher. Washington Department of Fish and Wildlife, Olympia, WA. 64 pp.
- Lewis, J.C. and G. E. Hayes.** 2004. Feasibility Assessment of Reintroducing Fisher to Washington. Washington Department of Fish and Wildlife, Olympia, WA. 70 pp. Table 4, p.17.
- Lindenmayer, D.B., and J.F. Franklin.** 2002. Conserving Forest Biodiversity: a Comprehensive Multiscaled Approach, Washington, D.C. Island Press. 351 p.
- Long, Colin J., Cathy Whitlock, Patrick J. Bartlein, and Sarah H. Millspaugh.** 1998. A 9000-year fire history from the Oregon Coast Range, based on a high-resolution charcoal study. *Can. J. For. Res.* 28(5): 774-787 (1998).
- Long, Linda L. and C.J. Ralph.** 1998. Regulation and observations of human disturbance near nesting marbled murrelets. Unpublished Report. USDA Forest Service, Pacific Southwest Research Station, Redwood Sciences Laboratory, Arcata, California.
- Loomis, D.R. Anglin and D. Liscia.** 1992. Fish management review Southwest region. Oregon Department of Fish and Wildlife, Roseburg, OR.
- Luce, C., and T. Black.** 1999. Sediment Production from Forest Roads in Western Oregon. *Water Resources Research* 35(8). pp. 2561-2570.
- Lyons J.K. and R.L. Beschta.** 1983. Land use, floods, and channel changes: Upper Middle Fork Willamette River, Oregon (1936 – 1980). *Water Resources Research* 19(2). pp. 463-471.
- Manley, Irene A. and S. Kim Nelson.** 1999. Habitat characteristics associated with nest success and predation at marbled murrelet tree nests. *Pacific Seabirds* 26:40.
- Manley, Irene F., Louise Waterhouse, and Alton S. Harestad.** 1999. Nesting habitat of marbled murrelets on the Sunshine Coast. Forest Research Extension Note EN-002, Vancouver Forest Region, British Columbia Ministry of Forestry, Nanaimo, BC.
- Markle, D., T. Pearsons, and D. Bills.** 1991. Natural History of *Oregonichthys* (Pices: Cyprinidae), with a Description of a New Species from the Umpqua River of Oregon. *Copeia* 1991 (2): 277-293.
- Marr, N.V., W.D. Edge, and R.G. Anthony.** 1995. Sheep Carcass Availability and Use by Bald Eagles. *Wilson Bulletin*, 107:251-257.
- Marshall, D.B.** 1989. The Marbled Murrelet. Audubon Wildlife Report. Pp. 435-455.

Marshall, D.B. 1992. Status of the Northern Goshawk in Oregon and Washington. Audubon Society of Portland, OR.

Marshall, D.B., M.W. Chilcote, and H. Weeks. 1996. Species at Risk: Sensitive, Threatened and Endangered Vertebrates of Oregon. 2nd edition. Oregon Department of Fish and Wildlife, Portland, OR.

Marshall, D.B., M.G. Hunter, and A.L. Contreras, Eds. 2003. Birds of Oregon. :A General Reference. Oregon State University Press, Corvallis, OR.

Maser, C., B.R. Mate, J.F. Franklin, and C.T. Dyrness. 1981. Natural History of Oregon Coast Mammals. USDA For. Serv. Gen. Tech. Rep. PNW-133, 496 pp. Pac. Northwest For. and Range Exp. Stn., Portland, OR.

May C.L. and R.E. Greswell. 2003. Large wood recruitment and redistribution in headwater streams in the southern Oregon Coast Range, U.S.A. Canadian Journal of Forest Research. 33:1352-1362.

McArdle, R.E., W.H. Meyer, and D. Bruce. 1949. The yield of Douglas fir in the Pacific Northwest. U.S. Dept. Agric. Tech. Bull. No. 201 Rev.

McClelland, B.R. and P.T. McClelland. 1999. Pileated woodpecker nest and roost trees in Montana: links with old-growth and forest "health". Wildlife Society Bulletin 27(3): 846-857.

McDade, M.H., F.J. Swanson, W.A. McKee, J.F. Franklin, and J. VanSickle. 1990. Source Distances for Coarse Woody Debris Entering Small Streams in Western Oregon and Washington. Canadian Journal of Forest Resources 20:326-330.

McGarigal, K.R., R.G. Anthony, and F.B. Isaacs. 1991. Interactions of Bald Eagles and Humans in the Columbia River Estuary. Wildlife Monograph No. 115. 47 pp.

McGarry, E.V. 1994. A Quantitative Analysis and Description of the Delivery and Distribution of Large Woody Debris in Cummins Creek, Oregon. Unpublished master's thesis, Oregon State University, Corvallis, OR.

McGrath, M.T. 1997. Northern goshawk habitat analysis in managed forest landscapes. M.S. thesis, Oregon State University, Corvallis, OR.

Meehan, W.R. (Ed.). 1991. Influences of Forest and Rangeland Management on Salmonid Fishes and Their Habitats. Special Publication 19. American Fisheries Society, Bethesda, MD.

Megahan, W.F. and G.L. Ketcheson. 1996. Predicting Downslope Travel of Granitic Sediments from Forest Roads in Idaho. Water Resources Bulletin. Vol. 32, No. 2. pp. 371-381.

Megahan, W.F., W.S. Platts, and B. Kulesza. 1980. Riverbed Improves Over Time: South Fork Salmon. *In* Symposium on Watershed Management, New York, NY; American Society of Civil Engineers; 1980: 380-395.

Mellen, T.K., E.C. Meslow, and R.W. Mannan. 1992. Summertime home range and habitat use of pileated woodpeckers in western Oregon. *Journal of Wildlife Management* 56(1): 96-103.

MFG, Inc. 2005. Effects Analysis For the Draft Western Oregon State Forests Habitat Conservation Plan For Aquatic Species. December 2005. Prepared for: Oregon Department of Forestry, 2600 State Street Salem, OR 97310. Prepared by: MFG, Inc., 875 Crescent Way Arcata, CA 95521

Miller, G.S. 1989. Dispersal of Juvenile Northern Spotted Owls in Western Oregon. Unpublished master's thesis, Oregon State University, Corvallis, OR.

Miller G.S., J. Blakesley, R. Bown, M. Brown, E.D. Forsman, R. Horn, J. Lint, E.C. Meslow, B. Noon, J. Reid, J.A. Thraikill, F.F. Wagner, and C. Zabel. 1990. Standards and Guidelines for Establishment and Implementations of Demographic and Density Study Areas for Spotted Owls. *Oreg. Coop. Wildl. Res. Unit. Unpubl. Rep.* 8pp.

Mills, K., L. Dent, and J. Robben. 2003. Wet Season Road Use Monitoring Project: Final Report. Oregon Department of Forestry.

Minore Don and Howard G. Weatherly. 1994. Riparian Trees, Shrubs and Forest Regeneration in the Coastal Mountains of Oregon. *New Forests* 8: 249-263

Monkkonen, Miko and Pasi Reunanen. 1999. On Critical Thresholds in Landscape Connectivity: A Management Perspective. *Oikos* 84: 302-305.

Murphy, M.L. 1995. Forestry Impacts on Freshwater Habitat of Anadromous Salmonids in the Pacific Northwest and Alaska — Requirements for Protection and Restoration. NMFS Coastal Ocean Program Decision Analysis Series No. 7. NMFS Coastal Ocean Office, Silver Spring, MD.

Murphy, M.L. and K.V. Koski. 1989. Input and Depletion of Woody Debris in Alaska Streams and Implications for Streamside Management. *North American Journal of Fisheries Management* 9:427-436.

Murphy, M.L. and W.R. Meehan. 1991. Stream Ecosystems. *In* W. R. Meehan, editor; Influences of Forest and Rangeland Management on Salmonid Fishes and Their Habitats. Special Publication 19. American Fisheries Society, Bethesda, MD.

Naiman, R.J. (Ed.). 1992. Watershed Management: Balancing Sustainability and Environmental Change. Springer-Verlag, New York, NY.

NatureServe. 2005. NatureServe Explorer: an Online Encyclopedia of Life. Version 4.4. Arlington, Virginia. Accessed May 20, 2005. www.natureserve.org/explorer

Nelson, S. Kim. 1997. Marbled murrelet (*Brachyramphus marmoratus*). *The Birds of North America* No. 276: 1-32.

Nelson, S.K. and T.E. Hamer. 1995. Nest Success and the Effects of Predation on Marbled Murrelets. *In* Ralph, C.J., G.I. Hunt, Jr., M.G. Raphael, and J.F. Piatt, editors; *Ecology and*

Conservation of the Marbled Murrelet. Pacific Southwest Research Station, USDA Forest Service, Albany, CA. General Technical Report PSW-GTR-152. Pp. 89-98.

Nelson, S. Kim and Amanda K. Wilson. 2002. Marbled murrelet habitat characteristics on state lands in western Oregon. Unpublished Report, submitted to Oregon Department of Forestry, Oregon Department of Fish and Wildlife, US Fish and Wildlife Service, and National Council for Air and Stream Improvement.

Nickelson, T.E. 2001. Population assessment: Oregon coast coho salmon ESU. Oregon Department of Fish and Wildlife, Portland, OR.

Nickelson, T.E., J.D. Rodgers, S.L. Johnson, and M.F. Solazzi. 1992a. Seasonal Changes in Habitat Use by Juvenile Coho Salmon (*Oncorhynchus kisutch*) in Oregon Coastal Streams. Canadian Journal of Fisheries and Aquatic Sciences 49:783-789.

Nickelson, T.E., M.F. Solazzi, S.L. Johnson, and J.D. Rodgers. 1992b. Effectiveness of Selected Stream Improvement Techniques to Create Suitable Summer and Winter Rearing Habitat for Juvenile Coho Salmon (*Oncorhynchus kisutch*) in Oregon Coastal Streams. Canadian Journal of Fisheries and Aquatic Sciences 49:790-794.

Nielson, M., K. Lohman, and J. Sullivan. 2001. Phylogeography of the tailed frog (*Ascaphus truei*): implications for the biogeography of the Pacific Northwest. Evolution 55:147-160.

Nierenberg Tara R. and David E. Hibbs. 2000. A Characterization of Unmanaged Riparian Areas in the Central Coast Range of Western Oregon. Forest Ecology and Management. 129. 195-206

Norris, L.A., H.W. Lorz, and S.V. Gregory. 1991. Forest Chemicals. In W. R. Meehan, editor; Influences of Forest and Rangeland Management on Salmonid Fishes and Their Habitats. Special Publication 19. American Fisheries Society, Bethesda, MD.

Nowacki, Gregory., Markc. Kramer. 1998. The effects of wind disturbance on temperate rain forest structure and dynamics in southeast Alsaska. Gen Tech Rep. Pacific Northwest Research Station. 25p.

Nussbaum, R.A. and C.K. Tait. 1977. Aspects of the Life History and Ecology of the Olympic Salamander, *Rhyacotriton olympicus* (Gaiage). Am. Midl. Nat. 98:176-199.

Nussbaum, R.A., E.D. Brodie, Jr., and R.M. Storm. 1983. Amphibians and Reptiles of the Pacific Northwest. University of Idaho Press, Moscow, ID. 332 pp.

O'Donnell, Brian P., Nancy L. Naslund, and C. John Ralph. 1995. Patterns of seasonal variation in activity of marbled murrelets in forested stands. Pages 117-128. In Ecology and conservation of the marbled murrelet. C.J. Ralph, M.G. Raphael, and J.F. Piatt (editors). USDA Forest Service, Pacific Southwest Research Station, General Technical Report PSW-GTR-152.

Olson D.H., C. Rugger, and G. Weaver. 2006. Effects of Headwater Riparian Reserves of Four Widths with Upslope Thinning on Instream and Bank Invertebrates. Presented at the 2005 Science and Management of Headwater Streams in the Pacific Northwest, Corvallis, OR. Abstracts from presentations posted: <http://www.headwatersresearch.org/> Page 39

Olterman, J.H. and B.J. Verts. 1972. Endangered Plants and Animals of Oregon. IV: Mammals. Special Report 364. Agricultural Experiment Station, Oregon State University, Corvallis, OR. 47 pp.

Oregon Climate Service. 2003. Strand Hall, Room 326, Oregon State University, Corvallis, OR 97331. <http://oregonstate.edu/~readw/>.

Oregon Department of Fish and Wildlife. 1992. Protocol for Surveying Proposed Management Activities That May Impact Northern Spotted Owls. Portland, OR. June 29, 1992.

Oregon Department of Fish and Wildlife. 1995. Biennial Report on the Status of Wild Fish in Oregon. December 1995. Portland, OR.

Oregon Department of Fish and Wildlife. 1997. Oregon Department of Fish and Wildlife Sensitive Species. Portland, OR.

Oregon Department of Fish and Wildlife. 1998. Oregon Department of Fish and Wildlife Sensitive Species. Portland, OR.

Oregon Department of Forestry. 1994. Forest Practice Technical Note FP1 — Water Classification. April 1994. Salem, OR. pp. 5-34.

Oregon Department of Forestry. 1999a. Forest Practice Note FP no. 4 – Road Maintenance. July 1999. Salem, OR. pp. 5 .

Oregon Department of Forestry. 1999b. Forest Practice Note FP no. 5 – Ground Based Harvesting. July 1999. Salem, OR. pp. 5.

Oregon Department of Forestry. 1995. Elliott State Forest Habitat Conservation Plan. Oregon Department of Forestry, Salem, OR.

Oregon Department of Forestry. 1996. Marbled Murrelet Management Plan for State Forest Lands. Salem, OR. Unpublished report. 8 pp.

Oregon Department of Forestry. 2000. Forest Roads Manual. Salem, OR. J. Cornell and K. Mills, editors. http://oregon.gov/ODF/STATE_FORESTS/Roads_Manual.shtml

Oregon Department of Forestry. 2001a. Northwest Oregon State Forests Management Plan. Salem, OR.

Oregon Department of Forestry. 2001b. Southwest Oregon State Forest Management Plan. Salem, OR.

Oregon Department of Forestry. 2002a. State Forests Monitoring Program Strategic Plan for the Northwest and Southwest Oregon State Forests Management Plans. Oregon Department of Forestry, Salem, OR. 48pp.

Oregon Department of Forestry. 2002b. Forest Practice Technical Note FP no. 4 – Fish Passage Guidelines for New and Replacement Stream Crossing Structures. Salem, OR. pp. 14.

Oregon Department of Forestry. 2002c. Forest Practice Technical Note FP no. 5 – Determining the 50 – year Peak Flow and Stream Crossing Structure Size for New and Replacement Crossings. Salem, OR. pp. 12.

Oregon Department of Forestry. 2002d. Forest Practice Monitoring Technical Report no. 14 Compliance With Fish Passage and Peak Flow Requirements at Stream Crossings Final Study Results. Salem, OR. pp. 59

Oregon Department of Forestry. 2003a. High Landslide Hazard Locations, Shallow, Rapidly Moving Landslides and Public Safety: Screening and Practices. Forest Practices Technical Note No. 2. Oregon Department of Forestry. 2600 State Street, Salem Oregon 97310. 10p. web site: www.odf.state.or.us/pcf/Pub/fp/LandslideHazNote2.pdf

Oregon Department of Forestry. 2003b. Determination of Rapidly Moving Landslide Impact Rating. Forest Practices Technical Note No. 6. Oregon Department of Forestry. 2600 State Street, Salem Oregon 97310. 12p. web site: www.odf.state.or.us/pcf/Pub/fp/LandslideTechNote6.pdf.

Oregon Department of Forestry. 2004. Marbled Murrelet Operational Policy for State Forest Lands. Oregon Department of Forestry. 2600 State Street, Salem, Oregon 97310.

Oregon Department of Forestry. 2006a. Coos District Implementation Plan. Oregon Department of Forestry. 2600 State Street, Salem, OR 97310.

Oregon Department of Forestry. 2006b. Reforestation Organization Operations Tracking System Manual. Oregon Department of Forestry. 2600 State Street, Salem, OR 97310.

Oregon Department of Forestry and Department of Environmental Quality 2002. Sufficiency Analysis: A statewide evaluation of forest practice act effectiveness in protecting water quality DEQ and ODF. Oregon Forests Practices Act water protection rules: Policy and scientific considerations. Oregon Department of Forestry State Forests Program, 2600 State Street, Salem OR. 97301. 82 pp.

Oregon Department of State Lands. 2006. DSL Asset Management Plan. Oregon Department of State Lands. 775 Summer St. NE Salem, OR 97301-1279

Oregon Fish and Wildlife Commission. 1998. At their December 18, 1998 meeting, they addressed a 5 year review of the State threatened and endangered species list, and recommended removal of the Bald Eagle from the list. Oregon Department of Fish and Wildlife. 3406 Cherry Ave. NE, Salem, OR 97303-4924, phone: 503-947-6000.

Oregon Watershed Enhancement Board. 1999. Oregon Road/Stream Crossing Restoration Guide: June 8, 1999. Oregon Plan for Salmon and Watersheds – Governor’s Watershed Enhancement Board. 255 Capitol St. N.E., Salem, OR. 97310-0203. 28pp.
<http://www.oregon-plan.org/>

Ormsbee, Patricia. 1996. Selection of day roosts by female long-legged myotis (*Myotis volans*) in forests of the central Oregon Cascades. MS. Thesis. Oregon State University. 63 pp.

Pacific Northwest Research Station (PNW). May 2004. Ecological payoffs from red alder in southeast Alaska. *In* Science Findings. Contributing researchers Robert Deal and Mike Wipfli. www.fs.fed.us/pnw USDA Forest Service P.O. Box 3890, Portland, Or. 97208. 6 pp

Pabst, Robert J. and Thomas A. Spies. 1999. Structure and composition of unmanaged riparian forests in the coastal mountains of Oregon, USA. *Canadian Journal of Forest Resources*. Vol. 29. Pp. 1557 -1573

Paragi, T.G. and G.M. Wholecheese. 1994. Marten, *Martes Americana*, predation on a northern goshawk, *Accipiter gentilis*. *Canadian Field-Naturalist* 108:81-82

Pletcher, F.T. 1963. The life history and distribution of lampreys in the Salmon and certain other rivers in British Columbia, Canada. Master of Science Thesis. University of British Columbia, Vancouver B.C.

Powell, R.A. and W.J. Zielinski. 1994. Fisher. Pp. 38-73. *In* Ruggiero, L.F., K.B. Aubry, S.W. Buskirk, L.J. Lyon, and W.J. Zielinski; *The Scientific Basis for Conserving Forest Carnivores: American Marten, Fisher, Lynx, and Wolverine in the Western United States*. USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO. General Technical Report RM-245. 184 pp.

Prescott, H.W. 1980. Causes of decline of the Western Bluebird in Oregon’s Willamette Valley. *Sialia* 2:131-135.

R2 Resource Consultants, Inc. in association with Lee Benda and Associates, Inc. 2005. Upper Nehalem Watershed Analysis. Prepared for the Oregon Department of Forestry. December 2005. 65pp.

Ralph, C.J., S.K. Nelson, M.M. Shaughnessy, S.L. Miller, and T.E. Hamer. 1994. Methods for surveying for marbled murrelets in forests: a protocol for land management and research. Technical Paper 1. Pacific Seabird Group, Marbled Murrelet Technical Committee, Arcata, CA.

Ralph, C.J., G.I. Hunt, Jr., M.G. Raphael, and J.F. Piatt (Technical Eds.). 1995. Ecology and Conservation of the Marbled Murrelet. Pacific Southwest Research Station, USDA Forest Service, Albany, CA. General Technical Report PSW-GTR-152.

Raphael, Martin G., Diane Evans Mack, John M. Marzluff, and John L. Leginbuhl. 2002. Effects of forest fragmentation on populations of the marbled murrelet. *Studies in Avian Biology* No 25: 221-235.

- Rasheed, S.A., P.F.J. Garcia and S.L. Holroyd.** 1995. Status of the Fringed myotis in British Columbia. Ministry of Environment, Lands, and Parks. Wildlife Working Report No. WR-73. 17 pp.
- Reeves, G.H., F.H. Everest, and J.R. Sedell.** 1993. Diversity of Juvenile Anadromous Salmonid Assemblages in Coastal Oregon Basins with Different Levels of Timber Harvest. Transactions of the American Fisheries Society, 122:309-317.
- Reeves G., L. Benda, K. Burnett, P. Bisson, and J. Sedell.** 1995. A Disturbance-Based Ecosystem Approach to Maintaining and Restoring Freshwater Habitats of Evolutionary Significant Units of Anadromous Salmonids in the Pacific Northwest. Evolution, and the Aquatic System: Defining Unique Units in Population Conservation. AM. Fish.Soc.Symp 17.334-349. pp 4-49.
- Reeves G.H., K.M. Burnett, and E.V. McGarry.** 2003. Sources of large wood in the main stem of a fourth-order watershed in coastal Oregon. Canadian Journal of Forest Research 33:1363-1370.
- Reid, L.M. and T. Dunne.** 1984. Sediment Production from Forest Road Surfaces. Water Resources Research. Vol. 20, No. 11: 1753-1761.
- Reimers P., R. Bender, J. Muck, T. Rumreich, R. Smith, and M. Manion** 1993. Fish management review Southwest region. Oregon Department of Fish and Wildlife, Charleston, OR
- Reiser, D.W. and R.G. White.** 1988. Effects of Two Sediment-Size Classes on Steelhead Trout and Chinook Salmon Egg Incubation and Juvenile Quality. North American Journal of Fisheries Management 8:432-437.
- Reynolds, R.T., E.C. Meslow, and H.M. Wight.** 1982. Nesting habits of coexisting Accipiter in Oregon. Journal of Wildlife Management 46:124-31.
- Richards, William H., David O. Wallin, and Nathan Schumaker.** 2002. An Analysis of Late-Seral forest connectivity in Western Oregon, USA. Conservation Biology 16(5): 1409-1421.
- Richardson, J.S.** 1992. Coarse Particulate Detritus Dynamics in Small, Montane Streams of Southwestern British Columbia. Canadian Journal of Fisheries and Aquatic Sciences 49(2):337-346.
- Rissler, L.J.** 1995. Habitat structure analyses of northern goshawk, *Accipiter gentilis atricapillus*, and northern spotted owls, *Strix occidentalis caurina*, nesting stands in the eastern Cascades. Utah State University. M.S. Thesis. 86 pp.
- Robart, G.P.** 1982. Wolverine, fisher, marten sightings in Oregon 1973-1982. Suppl. Rep. Central Reg. Admin. Rep. No. 82-4. Oregon Dept. Fish and Wildlife, Bend. 5 pp.

Robben J. and L. Dent. 2002. Oregon Department of Forestry Best Management Practices Compliance Monitoring Project Final Report. Technical Report #15. Oregon Department of Forestry, Salem, Or. 97310.

http://egov.oregon.gov/ODF/PRIVATE_FORESTS/docs/fp/BMPfinalTR15.pdf

Robison, E.G. and R.L. Beschta. 1990. Identifying Trees in Riparian Areas that Can Provide Coarse Woody Debris to Streams. *Forest Science* 36:790-801.

Robison E.G., K. Mills, J. Paul, L. Dent and A. Skaugset. 1999. Storm Impacts and Landslides of 1996 Final Report. Forest Practices Technical Report 4. Oregon Department of Forestry, 2600 State Street, Salem, OR. 97310. 157 p.

www.odf.state.or.us/pcf/Pub/fp/StormImpactsFinal.pdf

Robison E.G. and John Runyon. 2006. Effects of Different Riparian Management Treatments on Small Headwater Streams at the Upstream Extent of Fish Use for Streams in Western Oregon. Presented at the 2005 Science and Management of Headwater Streams in the Pacific Northwest, Corvallis, OR. Abstracts from presentations posted:

<http://www.headwatersresearch.org/> Page 44

Rosenburg, Kenneth and Martin Raphael. 1986. Effect of Forest Fragmentation on Vertebrates in Douglas Fir Forests. In Verner, J., et. al. (eds.) *Wildlife 2000, Modeling Habitat Relationships of Terrestrial Vertebrates*. University of Wisconsin Press.

Ross, Christine. 1999. Personal communication to Charlie Bruce, Oregon Department of Fish and Wildlife biologist, Corvallis, OR.

Rothacher, J. 1973. Does harvest in west slope Douglas-fir increase peak flow in small streams? USDA Forest Service Research paper PNW-163. Portland Oregon. USA

Ruth, R.H. and R.A. Yoder. 1953. Reducing Wind Damage in the Forests of the Oregon Coast Range. Research Paper Number 7. USDA Forest Service, PNW Experimental Station, Portland, OR.

Salo, E.O. and T.W. Cundy (Eds.). 1987. *Streamside Management: Forestry and Fishery Interactions*. University of Washington, Institute of Forest Resources Contribution 57, Seattle, WA.

Schommer, T. and G. Silovsky. 1994. Goshawk monitoring, management, and research in the Pacific Northwest Region. USDA Forest Service. Portland, OR. 72 pp.

Seaman, D.E., and R.A. Powell. 1996. An Evaluation of the Accuracy of Kernel Density Estimators for Home Range Analysis. *Ecology* 77:2075-2085.

Seaman, D.E., B. Griffith, and R.A. Powell. 1998. KernalHR: A Program for Estimating Animal Home Ranges. *Wildlife Society Bulletin* 1998, 26(1):95-100.

- Sedell, J.R., P.A. Bisson, J.A. June, and R.W. Speaker.** 1982. Ecology and Habitat Requirements of Fish Populations in South Fork Hoh River, Olympic National Park. *In* Starkey, E.E., J.F. Franklin, and J.W. Matthews, Tech. Coords., Ecological Research in National Parks of the Pacific Northwest: Proceedings, 2nd Conference November 1979. Oregon State University, Forest Research Laboratory.
- Sessions, J., K.N. Johnson, J. Beuter, B. Greber, G. Lettman.** 1990. Timber for Oregon's Tomorrow: The 1989 Update. Oregon State University, College of Forestry, Forest Research Laboratory, Corvallis, OR.
- Seymour, R., Malcolm Hunter.** 1999. Principles of forest ecology. Pages 22-61. In M. Hunter (ed.) Maintaining biodiversity in forest ecosystems. Cambridge University Press, New York, NY.
- Shreiber, B. and D.S. deCalesta.** 1992. The relationship between cavity-nesting birds and snags on clearcuts in western Oregon. *Forest Ecology and Management* 50:299-316.
- Sidle, R. C., A. J. Pearce, and C. L. O'Loughlin.** 1985. Hillslope Stability and Land Use. American Geophysical Union Water Resources Monograph 11.
- Skaugset A. E.** 2006. Processes that Influence the Downstream Propagation of Stream Temperature. Presented at the 2005 Science and Management of Headwater Streams in the Pacific Northwest, Corvallis, OR. Abstracts from presentations posted: <http://www.headwatersresearch.org/> Page 49.
- Spence, B.C., G.A. Lomnický, R.M. Hughes, and R.P. Novitzki.** 1996. An Ecosystem Approach to Salmonid Conservation. TR-4501-96-6057. ManTech Environmental Research Services Corp., Corvallis, OR (Available from the National Marine Fisheries Service, Portland, OR).
- Spies, T.A. and J.F. Franklin.** 1991. The Structure of Natural Young, Mature and Old-growth Douglas-fir Forests in Oregon and Washington. *In* Ruggiero, L.F., K.B. Aubry, A.B. Carey, and M.H. Huff, technical coordinators. Wildlife and Vegetation of Unmanaged Douglas-fir Forests. USDA Forest Service, Pacific Northwest Forest and Range Experiment Station, Portland, OR. General Technical Report PNW-GTR-285.
- Squires, J.R. and R.T. Reynolds.** 1997. Northern Goshawk (*Accipiter gentilis*). In A. Poole and F. Gill, editors, The Birds of North America, No. 298. Academy of Natural Sciences, Philadelphia, and American Ornithologists' Union, Washington, DC. 32 pp.
- Stalmaster, M.V., R.L. Knight, B.L. Holder, and R.J. Anderson.** 1985. Bald Eagles. Pages 269-290. *In* E.R. Brown, editor, Management of Wildlife and Fish Habitats in Forests of Western Oregon and Washington, pages 269-290. USDA Forest Service, Publication No. R6-F&WL-192-1985, Portland, OR. 332 pp.
- Stebbins, R.C.** 2003. A field guide to western reptiles and amphibians. Third edition. Houghton Mifflin Company, Boston.
- Steelquist, R.** 1992. Field Guide to the Pacific Salmon. Sasquatch Books, Seattle, WA.

Sullivan K. 1985. Long-term patterns of water quality in managed watershed in Oregon: 1. Suspended sediment. *Water Resources Bulletin*. Volume 21. No. 6. pp. 977-987.

Sullivan, K., T.E. Lisle, C.A. Dolloff, G.E. Grant, and L.M. Reid. 1987. Stream Channels: The Link between Forests and Fishes. *In* Salo and Cundy, editors; *Streamside Management: Forestry and Fishery Interactions*. University of Washington, Institute of Forest Resources Contribution 57, Seattle, WA.

Sullivan, K.D., J. Tooley, K. Doughty, J.E. Caldwell, and P.Knudsen. 1990. Evaluation of prediction models and characterization of stream temperature regimes in Washington. Washington Department of Natural Resources Timber/Fish/Wildlife Report TFW-WQ3-90-006.

Swanson F.J., M.M. Swanson, and C. Woods. 1977. Inventory of mass erosion in the Mapleton Ranger District, Siuslaw National Forest.

Swanson, F.J., L.E. Benda, S.H. Duncan, G.E. Grant, W.F. Megahan, L.M. Reid, and R.R. Ziemer. 1987. Mass Failures and Other Processes of Sediment Production in Pacific Northwest Forest Landscapes. pp. 9-38. *In* Salo, E.O. and T.W. Cundy, editors; *Proceedings of Streamside Management Forestry and Fishery Interactions*. University of Washington, Institute of Forest Resources Contribution No. 57. 471 pp.

Swanston, D.N. 1991. Natural Processes. *In* W.R. Meehan, editor; *Influences of Forest and Rangeland Management on Salmonid Fishes and their Habitats*. Special Publication 19. American Fisheries Society, Bethesda, MD.

Tappeiner, J., Lewis, R., Tesch, S., Thornburgh, D., Weatherspoon, P., Birch, K., Knapp, W., McComb, B., Saunders, B., Spies, T., Zasada, J., Bickford, M., Gara, B., Harrington, C., Hicks, L., Hoyer, J., Leitner, G., Martin, S., Oliver, C., Raettig, T. 1992. Managing Stands for Northern Spotted Owl Habitat. *In* Recovery Plan for the Northern Spotted Owl - DRAFT, Appendix G. Pp. 481 - 525. *In* USDI 1992a.

Tappeiner, J., T. Nierenberg, J. Bailey, and N. Poage. 2000. Characterizing northern spotted owl habitat on state forest lands in the Oregon Coast Range. Report to Oregon Department of Forestry. 114 pp.

Taylor, A.L. and E. D. Forsman. 1976. Recent Range Extensions of the Barred Owl in Western North America, Including the First Records for Oregon. *Condor* 78:560-561.

Taylor, J.A. 1999. Habitat Selection of the Long-Legged Myotis (*Myotis volans*) in a Managed Landscape on the East-Slopes of the Cascade Range. M.Sc. Thesis. Michigan State University, East Lansing, MI.

Thom, B.A., K.K. Jones, and R.L. Flitcroft. 1999. Stream habitat conditions in western Oregon. Monitoring program report 1999-1 to the Oregon Plan for Salmon and Watersheds, Governor's Natural Resource Office, Salem, OR.

Thom, B.A., P.S. Kavanaugh, and K.K. Jones. 2001. Reference Site Selection and Survey Results, 2000. Monitoring Program Report Number OPSW-ODFW-2001-6, Oregon Department of Fish and Wildlife, Portland, Oregon.

Thomas, D.W. 1988. The distribution of bats in different ages of Douglas-fir forests. *The Journal of Wildlife Management*, 52:619-626.

Thomas, J.W., E.D. Forsman, J.B. Lint, E.C. Meslow, B.R. Noon, and J. Verner. 1990. A Conservation Strategy for the Northern Spotted Owl: A Report of the Interagency Scientific Committee to Address the Conservation of the Northern Spotted Owl. USDA Forest Service; USDI Bureau of Land Management, Fish and Wildlife Service, National Park Service, Portland, Oregon.

Thomas, R.B. and W.F. Megahan. 1998. Peak Flow Responses to Clear-cutting and Roads in Small and Large Basins, Western Cascades, Oregon: A Second Opinion. *Water Resources Research*, V.34, No. 12, 3393-3403. December.

Thraillkill, J.A. and L.S. Andrews. 1996. Presence of Breeding Northern Goshawks in the Coast Range of Oregon. *Journal of Raptor Research*, 30:248-249.

Trotter, P.C. 1997. Sea-Run Cutthroat Trout: Life History Profile. Pages 7-15 in J.D. Hall, P.A. Bisson, and R.E. Gresswell, editors. *Sea-run cutthroat trout: biology, management, and future conservation*. Oregon Chapter, American Fisheries Society, Corvallis.

Turnstone Environmental Consultants. 2002. Final Report: Marbled Murrelet Surveys for the Elliott State Forest Habitat conservation Plan. Unpublished report submitted to the Oregon Department of Forestry. 52 pages plus maps.

Ursitti V.L 1990. Riparian vegetation and abundance of woody debris in streams of southwestern Oregon. M.S. thesis. Oregon State University.

USDA Forest Service. 1993. Forest Ecosystem Management: An Ecological, Economic, and Social Assessment. Report of the Forest Ecosystem Management Assessment Team. July 1993. Also known as the FEMAT Report. USDA Forest Service, Pacific Northwest Region, Portland, OR. U.S. Government Printing Office 1993-793-071.

USDA Forest Service, USDI Bureau of Land Management, U.S. Fish and Wildlife Service, National Park Service, US Department of Commerce National Oceanic and Atmospheric Administration National Marine Fisheries Service, and US Environmental Protection Agency. 1994a. Final Supplemental Environmental Impact Statement on Management of Habitat for Late-Successional and Old-Growth Forest Related Species Within the Range of the Northern Spotted Owl. Also known as the Northwest Forest Plan or the Final SEIS. USDA Forest Service, Pacific Northwest Region, Portland, OR. February 1994. libweb.uoregon.edu/govdocs/forestpl.html.

USDA Forest Service, USDI Bureau of Land Management, U.S. Fish and Wildlife Service, National Park Service, US Department of Commerce National Oceanic and Atmospheric Administration National Marine Fisheries Service, and US Environmental Protection Agency. 1994b. Record of Decision for Amendments to Forest Service and Bureau of Land Management Planning Documents Within the Range of the Northern Spotted Owl. USDA Forest Service, Pacific Northwest Region, Portland, OR. April 1994. *[This is the 2nd document of the NW Forest Plan; the 1st is "FEMAT," above. Note – the Standards and Guidelines are Attachment A of this ROD]*

USDA Forest Service, and USDI Bureau of Land Management. 2001. Record of Decision and Standards and Guidelines for Amendments to the Survey and Manage, Protection Buffer, and other Mitigation Measures Standards and Guidelines. USDA Forest Service, Pacific Northwest Region, Portland, OR. January 2001.

USDA Forest Service. 2006. The PNW-FIA Integrated Database, Version 2.0. Developed and compiled by Karen Waddell and Bruce Hiserote, based on data from USDA Forest Service and USDI Bureau of Land Management inventories. USDA Forest Service Pacific Northwest Research Station, Portland Oregon.

U.S. Fish and Wildlife Service. 1986. Pacific Bald Eagle Recovery Plan. U.S. Fish and Wildlife Service, Portland, OR. 160 pp.

U.S. Fish and Wildlife Service. 1990. Procedures Leading to Endangered Species Act Compliance for the Northern Spotted Owl. U.S. Fish and Wildlife Service, Region 1. July 1990.

U.S. Fish and Wildlife Service. 1992a. Final Draft Recovery Plan for the Northern Spotted Owl. USDI, Washington, D.C. USDA Forest Service, Pacific Northwest Region, Portland, OR. 3 documents: Vol. I, 322 pp.; Vol. II, 490 pp., summary booklet, 28 pp.

U.S. Fish and Wildlife Service. 1992b. Protocol for Surveying Proposed Management Activities That May Impact Northern Spotted Owls. U.S. Fish and Wildlife Service, Portland, OR. March 17, 1992.

U.S. Fish and Wildlife Service. 1992c. Endangered and Threatened Wildlife and Plants; Determination of Threatened Status for the Washington, Oregon, and California Population of the Marbled Murrelet. U.S. Federal Register 57:45328-45337.

U.S. Fish and Wildlife Service. 1995. Draft Recovery Plan for the Marbled Murrelet in Washington, Oregon, and California. Region 1, Portland, Oregon.

U.S. Fish and Wildlife Service. 1996. 90-day Finding for a petition to list the fisher in the western United States as Threatened. Federal Register 61(42):8016-8018

U.S. Fish and Wildlife Service and National Marine Fisheries Service. 1996. Handbook for Habitat Conservation Planning and Incidental Take Permit Processing.

U.S. Fish and Wildlife Service. 1997. Recovery Plan for Threatened Marbled Murrelet (*Brachyramphus marmoratus*) in Washington, Oregon, and California. Region 1, Portland, OR.

U.S. Fish and Wildlife Service. 1998a. Announcement for regional workshops to address delisting. Memo, July 24, 1998, Portland, OR.

U.S. Fish and Wildlife Service. 1998b. Federal Register: June 29, 1998, 63(165): 35183-35184).

U.S. Fish and Wildlife Service. 1998c. Northern Goshawk Status Review. Office of Technical Support, Portland, OR.

U.S. Fish and Wildlife Service. 2004a. Northern Spotted Owl, Five-Year Review: Summary and Evaluation. U.S. Fish and Wildlife Service. Portland, OR. November 2004.

U.S. Fish and Wildlife Service. 2004b. Biological Opinion and Letter of Concurrence for Effects to Bald Eagles, Northern Spotted Owls and Marbled Murrelets from the U.S. Department of the Interior, Bureau of Land Management, Eugene District and Salem District, and U.S. Department of Agriculture, Siuslaw National Forest fiscal year 2004/2005 disturbance activities within the North Coast Province (FWS Reference Number 1-7-04-F-1113). U.S. Department of the Interior, U.S. Fish and Wildlife Service, Oregon Fish and Wildlife Office, Portland, Oregon, January 2004. 85 pp.

U.S. Fish and Wildlife Service. 2004c. Federal Register 50 CFR Part 17. Endangered and threatened wildlife and plants: 12 month finding for a petition to list the West Coast Distinct population segment of the fisher (*Martes pennanti*) 18770-18792.
http://frwebgate.access.gpo.gov/cgi-bin/getdoc.cgi?dbname=2004_register&docid=fr08ap04-20.pdf

U.S. Fish and Wildlife Service. 2007. Draft Post-delisting monitoring plan posted:
<http://www.fws.gov/migratorybirds/baldeagle.htm>.

U.S. Fish and Wildlife Service. 2008. Final Recovery Plan for the Northern Spotted Owl, *Strix occidentalis caurina*. U.S. Fish and Wildlife Service, Portland, Oregon. xii + 142 pp.

U.S. Fish and Wildlife Service, Environmental Protection Agency, Department of the Army, and Soil Conservation Service. 1989. Federal Manual for Identifying and Delineating Jurisdictional Wetlands. pp. 5-59.

Van Sickle, J. and S.V. Gregory. 1990. Modeling Inputs of Large Woody Debris to Streams from Falling Trees. Canadian Journal of Forest Research 20:1593-1601.

Verts, B.J., and L.N. Carraway. 1998. Land Mammals of Oregon. University of California Press, Berkeley, CA.

Vesely, D.G. and J. Stamp. 2001. Report on Surveys for Stream-Dwelling Amphibians Conducted on the Elliott State Forest, June–September. Unpublished report to Oregon Department of Forestry. 35 pp.

- Walters, C.J.** 1986. Adaptive Management of Renewable Resources. McGraw-Hill, New York, NY. 374 pp.
- Walters, C.J. and C.S. Holling.** 1990. Large-Scale Management Experiments and Learning by Doing. *Ecology* 71(6): 2060-2068.
- Washington Department of Wildlife.** 1990. Working Implementation Plan for Bald Eagle Recovery in Oregon and Washington. Washington Department of Wildlife, Olympia, WA.
- Washington Forest Practices Board (WFPB).** 1997. Standard methodology for conducting watershed analysis. Version 4.0. Washington Department of Natural Resources, Olympia. November 1997.
- Watson, J.W. and R.G. Anthony.** 1986. Ecology of Bald Eagles in the Tongue Point Area, Lower Columbia River. Final report to the Army Corps of Engineers. No. DACW57-83-C-0100.
- Watson, J.W., M.G. Garrett, and R.A. Anthony.** 1991. Foraging Ecology of Bald Eagles in the Columbia River Estuary. *Journal of Wildlife Management* 55:492-499.
- Weaver, W. E., and D. K. Hagens.** 1994. Handbook for Forest and Ranch Roads. Mendocino County Resource Conservation District in Cooperation with the California Department of Forestry and the U.S. NRCS. 1994.
- Weikel, J.** 2003. A Literature Review on Ecology and Effects of Forest Management on the Marbled Murrelet. Final report submitted to Oregon Department of Forestry. 29 pp.
- Weikel, J. and D. Vesely.** 2001. A report on avian surveys conducted on the Elliott State Forest, May-July, 2001. Final report to the Oregon Department of Forestry. 25 pp.
- Welsh Jr., H.H.** 1990. Relictual Amphibians and Old-Growth Forests. *Conservation Biology* 4:309-319.
- Welsh, C.J.E. and D.E. Capen.** 1992. Availability of nesting sites as a limit to woodpecker populations. *Forest Ecology and Management* 48: 31-41.
- Welsh Jr., H.H. and A.J. Lind.** 1992. Population Ecology of Two Relictual Salamanders from the Klamath Mountains of Northwestern California. Pp. 419-437. *In* McCullough, D.R. and R.H. Barrett (Eds.), *Wildlife 2001: Populations*. Elsevier Applied Science, London.
- Welsh Jr., H.H. and A.J. Lind.** 1996. Habitat Correlates of the Southern Torrent Salamander, *Rhyacotriton variegatus* (Caudata: Rhyacotritonidae), in Northwestern California. *Journal of Herpetology* 30:385-398.
- Welsh Jr., H.,H. and L.M. Ollivier.** 1998. Stream Amphibians as Indicators of Ecosystem Stress: A Case Study from California's Redwoods. *Ecological Applications* 8:1118-1132.
- Wemple, B.C., J.A. Jones, and G.E. Grant.** 1997. Channel network extension by logging roads in two basins, Western Cascades, Oregon. *Water Resources Bulletin* 32(6):1195-1207.

Wilson, J. 1998. Wind Stability in Douglas-fir Plantations. Ph. D. Thesis, University of Washington, Seattle, WA.

Wimberly, M.C., 2000. From Presentation. Assessing Present-Day Forest Landscapes in the Context of Historical Disturbance Regimes. Website:
www.fsl.orst.edu/clams/download/presentations/j02s_wimberly_histvar.pdf

Wimberly, M.C. and T.A. Spies. 2001. Influences of Environment and Disturbance on Forest Patterns in Coastal Oregon Watersheds. *Ecology* 82 (5):1443-1459.

Wimberly, M.C., T.A. Spies, C.J. Long, and C. Whitlock. 2000. Simulating Historical Variability in the Amount of Old Forest in the Oregon Coast Range. *Conservation Biology* 14(1) (February 2000): 167-180.

Wipfli M.S. and D.P. Gregovich. 2002. Export of invertebrates and detritus from fishless headwater streams in southeastern Alaska: implications for downstream salmonid production

With, Kimberly A. and Thomas O. Crist. 1995. Critical Thresholds in Species' Responses to Landscape Structure. *Ecology* 76(8):2446-2459.

With, Kimberly A., Robert H. Gardener and Monica G. Turner. 1997. Landscape Connectivity and Population Distributions in Heterogeneous Environments. *Oikos* 78:151-169.

Worton, B.J. 1995. Using Monte Carlo Simulation to Evaluate Kernel-Based Home Range Estimators. *Journal of Wildlife Management.* 59:794-800.

Wydoski, R.S., and R.R. Whitney. 2003. Inland Fishes of Washington. 2003. American Fisheries Society, Bethesda, Maryland.

Yocum, C.F. and M.T. McCollum. 1973. Status of the Fisher in Northern California, Oregon and Washington. *California Fish and Game* 59(4):305-309.

Zeiner, D.C., W.F. Laudenslayer, Jr., K.E. Mayer, and M. White (Eds.). 1990. California's Wildlife. Volume 2. Birds. State of California, Department of Fish and Game. Sacramento, California. 731 pp.

Zielinski, W.J., T.E. Kucera, and R.H. Barrett. 1995. Current Distribution of the Fisher, *Martes pennanti*, in California. *California Fish and Game* 81(3):104-112.

Ziemer. 1981. Storm flow response to road building and partial cutting in small streams of northern California. *Water Resources Research.* 17:907-917.

Ziemer, R.R. 1998. Proceedings of the conference on coastal watersheds: The Casper Creek Story: 6 May; Ukiah, Ca. Gen. Tech. Rep. PSW-GTR-168. Albany, CA: Pacific Southwest Research Station, Forest Service, U.S. Department of Agriculture. 149 pp.

Ziemer, R.R. and T.E. Lisle. 1998. Chapter 3: Hydrology. *In* R. J. Naiman and R. E. Bilby, editors; River Ecology and Management, Lessons from the Pacific Coastal Ecoregion. Springer-Verlag New York, Inc., 175 Fifth Avenue, New York, NY 10010, USA. ISBN 0-387-98323-6 Springer-Verlag pp 43-69.