



Photo credit: M. Yoklavich, NMFS.

KEY INFORMATION

Areas of Concern

California waters (possibly west coast of North America).

Year Identified as “Species of Concern”

1999

Factors for Decline

- Overfishing
- Poor environmental conditions

Conservation Designations

IUCN: Critically Endangered

Current Status:

Demographic and Genetic Diversity Concerns:

The main reason for listing is the magnitude of decline in abundance. As of the 2005 status assessment, bocaccio were estimated to be between 8.2 and 12.2% of their historical or “unfished” abundance levels (MacCall 2005). Like most long-lived west coast rockfishes (*Sebastes* spp.), productivity is low and the stock is not capable of sustaining “conventional” levels of fishing pressure.

Existing Protections and Conservation Actions:

Although the southern population of bocaccio has substantially declined, NMFS has taken measures to ensure that it will not become endangered within the foreseeable future. The Pacific Fisheries Management Council (Council) has recommended that NMFS eliminate all directed fishing for bocaccio in 2003. The only allowable catch would be bocaccio taken as **bycatch** in other fisheries. In order to ensure that catch levels are not exceeded, the Council recommended that NMFS implement several management measures in 2003, including new depth-based management measures to prohibit bottom trawls, limit entry of fixed gear, and limit open access fishing in the times and areas where bocaccio are expected to occur. Bocaccio has an 80 percent chance of no further declines in 100 years, and the species is expected to rebuild in approximately 170 years with NMFS' implementation of the Council's proposed measures, in addition to measures being implemented by California. In 2004 a formal

Brief Species Description:

Bocaccio is a large (up to 3 feet or 90cm length) piscivorous (fish-eating) rockfish ranging from northern Baja California to Alaska. It is thought to consist of two partially isolated populations: a southern population off California, and a northern population off Washington and British Columbia. They have a distinctively long jaw extending at least to the eye socket. Their back ranges in color from olive- to burnt-orange or brown as adults; its stomach is pink and red. Young bocaccio are light bronze in color and have small brown spots on their sides. Coloring darkens and the spots disappear as they mature. They prefer rocky habitats from 130 to 980 feet (40 to 300 m) deep, but may occur in nearly all habitats. Oil platforms have become somewhat important artificial habitats for this species (Love and York 2006). Young (1-3 yr) bocaccio are relatively **pelagic**, and become more demersal (bottom oriented) with age (maximum age 45 to 50 years). They mature at 4 to 5 years of age. They



Species of Concern

NOAA National Marine Fisheries Service

prefer to eat other rockfishes, but will also eat sablefish, anchovies, lantern fish and squid. They are a component of catches by nearly every fishing gear, and are difficult to avoid. Like other species of their genus, cowcod are internal fertilizers. Mating occurs in the fall (MacCall 2002).

Status Reviews/Research Underway:

On January 31, 2001, NMFS received a petition list the southern population of bocaccio as a threatened species under the Endangered Species Act (ESA). On June 14, 2001, NMFS published its 90-day finding that the petition may be warranted, and announced the initiation of a formal status review as required by the ESA. NMFS' Southwest Fisheries Science Center prepared a comprehensive status review that recognized a northern “**distinct population segment**” (DPS) and a southern DPS for bocaccio, and this is consistent with the current NMFS and Council management of bocaccio. Based on the review, NMFS concluded that listing is not warranted (67 FR 69704, November 19, 2002), but that progress of the Pacific Fishery Management Council’s bocaccio rebuilding program should be monitored.




Figure 1. History of exploitation rates (catch/total biomass) of bocaccio (left), and history of estimated year class strengths (at age 1) of bocaccio (right). NMFS.

Contact Information

For bocaccio, contact:

Dr. Scott Rumsey
NOAA Fisheries, Northwest Region
Protected Resources Division
1201 NE Lloyd, Blvd, #1100
Portland, OR 97232
(503) 872-2791

Scott.Rumsey@noaa.gov

<http://www.nmfs.noaa.gov/pr/species/concern>

For Species of Concern, contact

NOAA Fisheries
Office of Protected Resources
1315 East West Highway
Silver Spring, MD 20910
(301) 713-1401

soc.list@noaa.gov

References:

- Love, M.S. and A. York. 2006. Fishery Bulletin (Seattle) 104:542-549.
- MacCall, A.D. 2002. Status of bocaccio off California in 2002. NMFS, Santa Cruz, CA.
- MacCall, A.D. 2005. Status of bocaccio off California in 2005. NMFS, Santa Cruz, CA.