

TABLE 8

NUMBER OF REPORTED SHIGELLA ISOLATES BY REGION  
FOR 1983-1997

REGION	YEAR																TOTAL
	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997		
NEW ENGLAND	543	354	861	729	645	651	546	646	1757	555	520	414	586	373	486	9666	
MID-ATLANTIC	2050	1094	1607	1263	2331	1179	1310	1595	1263	1071	1215	1562	1758	2240	1874	23412	
EAST NORTH CENTRAL	1881	1863	1494	1550	3816	3597	2673	2096	1643	2154	2816	1970	2105	1190	1457	32305	
WEST NORTH CENTRAL	530	495	387	330	671	1151	1359	630	403	589	828	1368	1391	699	494	11325	
SOUTH ATLANTIC	990	997	683	686	1425	2562	2568	1894	1565	1727	2860	4790	2181	1912	1136	27976	
EAST SOUTH CENTRAL	155	114	188	60	219	683	482	799	915	305	768	1023	802	334	233	7080	
WEST SOUTH CENTRAL	1619	1263	1218	1611	1671	2535	1408	1832	1409	1453	2943	1771	1944	1225	1731	25633	
MOUNTAIN	1525	1150	1223	1137	1174	1692	1161	1655	2248	1174	1348	1436	2713	1665	1238	22539	
PACIFIC	5664	5385	4215	4906	4812	8286	4465	6159	5302	5895	6213	4449	5850	4433	3665	79699	
TOTAL	14957	12715	11876	12272	16764	22336	15972	17306	16505	14923	19511	18783	19330	14071	12314	239635	

TABLE 8A

NUMBER OF REPORTED *S. SONNEI* ISOLATES BY REGION  
FOR 1983-1997

REGION	YEAR																	TOTAL
	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997			
NEW ENGLAND	383	214	713	596	491	481	358	448	1573	435	428	302	456	264	341	7483		
MID-ATLANTIC	1629	823	1128	772	2030	883	1015	1321	1020	880	985	1325	1527	2048	1664	19050		
EAST NORTH CENTRAL	1470	1458	1110	1123	3279	2685	2061	1443	1176	1671	2440	1707	1773	896	1242	25534		
WEST NORTH CENTRAL	414	354	246	147	503	870	1084	446	255	478	698	1166	1281	575	405	8922		
SOUTH ATLANTIC	705	704	450	462	1099	2155	2268	1535	1210	1451	2280	2695	1966	1772	999	21751		
EAST SOUTH CENTRAL	124	103	179	52	136	523	319	513	587	152	341	649	511	232	211	4632		
WEST SOUTH CENTRAL	1071	767	817	1161	1293	2103	1022	1415	1072	1268	2765	1595	1721	1085	1514	20669		
MOUNTAIN	654	575	554	571	470	921	522	1124	1601	722	824	920	2180	1115	695	13448		
PACIFIC	2825	2792	1878	2421	1981	4407	1593	2871	2240	3049	3578	2087	3396	2275	1736	39129		
TOTAL	9275	7790	7075	7305	11282	15028	10242	11116	10734	10106	14339	12446	14811	10262	8807	160618		

TABLE 8B  
 NUMBER OF REPORTED S. FLEXNERI ISOLATES BY REGION  
 FOR 1983-1997

REGION	YEAR																	TOTAL
	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997			
NEW ENGLAND	142	114	128	93	109	127	150	122	94	107	92	106	115	94	123	1716		
MID-ATLANTIC	335	256	377	341	278	257	269	254	227	177	211	213	206	179	188	3768		
EAST NORTH CENTRAL	345	344	326	314	361	422	439	465	339	330	287	238	289	267	185	4951		
WEST NORTH CENTRAL	90	103	108	168	134	156	168	125	115	77	67	121	71	105	82	1690		
SOUTH ATLANTIC	93	102	90	116	154	143	101	156	124	137	173	343	196	122	120	2170		
EAST SOUTH CENTRAL	30	10	8	6	9	8	9	10	8	2	4	18	7	7	20	156		
WEST SOUTH CENTRAL	491	435	346	380	328	376	333	367	280	170	136	141	188	113	184	4268		
MOUNTAIN	774	509	548	496	556	503	472	381	491	371	382	353	464	441	484	7225		
PACIFIC	1924	1879	1602	1653	1831	2346	1741	2151	2034	1879	1709	1568	1483	1376	1187	26363		
TOTAL	4224	3752	3533	3567	3760	4338	3682	4031	3712	3250	3061	3101	3019	2704	2573	52307		

TABLE 8C

NUMBER OF REPORTED S. BOYDII ISOLATES BY REGION  
FOR 1983-1997

REGION	YEAR																	TOTAL
	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997			
NEW ENGLAND	7	10	8	6	14	7	16	17	12	4	.	5	11	6	14	137		
MID-ATLANTIC	8	8	16	16	16	21	17	10	10	6	11	15	12	5	13	184		
EAST NORTH CENTRAL	42	38	35	37	28	64	42	29	20	16	30	19	31	19	23	473		
WEST NORTH CENTRAL	6	4	1	1	2	6	5	5	7	3	.	7	6	3	4	60		
SOUTH ATLANTIC	11	2	3	6	7	5	10	8	8	5	10	27	7	10	7	126		
EAST SOUTH CENTRAL	1	1	.	1	2	2	1	1	.	1	.	.	.	2	.	12		
WEST SOUTH CENTRAL	47	35	33	54	36	39	37	35	44	11	33	24	27	19	27	501		
MOUNTAIN	48	34	66	43	59	52	48	47	30	14	14	18	32	91	41	637		
PACIFIC	246	254	196	209	192	276	290	246	181	164	123	92	103	120	124	2816		
TOTAL	416	386	358	373	356	472	466	398	312	224	221	207	229	275	253	4946		

TABLE 8D

NUMBER OF REPORTED *S. DYSENTERIAE* ISOLATES BY REGION  
FOR 1983-1997

REGION	YEAR																	TOTAL
	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997			
NEW ENGLAND	2	6	5	4	4	12	9	8	6	6	.	.	3	6	6	77		
MID-ATLANTIC	7	7	16	11	7	18	9	9	6	8	2	6	6	6	8	126		
EAST NORTH CENTRAL	22	20	20	7	27	21	23	15	11	21	4	6	11	8	2	218		
WEST NORTH CENTRAL	6	2	3	3	8	8	7	3	1	6	1	2	1	2	2	55		
SOUTH ATLANTIC	15	4	1	2	4	4	2	5	6	4	4	5	2	8	5	71		
EAST SOUTH CENTRAL	.	.	1	1	1	2	.	2	.	1	1	1	.	.	2	12		
WEST SOUTH CENTRAL	8	26	20	13	12	17	15	14	10	2	9	10	5	7	6	174		
MOUNTAIN	20	19	14	9	10	8	25	7	12	5	12	9	6	18	12	186		
PACIFIC	105	95	129	68	159	121	126	118	93	73	72	55	56	48	36	1354		
TOTAL	185	179	209	118	232	211	216	181	145	126	105	94	90	103	79	2273		

TABLE 8E

NUMBER OF REPORTED SHIGELLA ISOLATES WITH UNKNOWN SEROGROUP  
BY REGION FOR 1983-1997

REGION	YEAR																	TOTAL
	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997			
NEW ENGLAND	9	10	7	30	27	24	13	51	72	3	.	1	1	3	2	253		
MID-ATLANTIC	71	.	70	123	.	.	.	1	.	.	6	3	7	2	1	284		
EAST NORTH CENTRAL	2	3	3	69	121	405	108	144	97	116	55	.	1	.	5	1129		
WEST NORTH CENTRAL	14	32	29	11	24	111	95	51	25	25	62	72	32	14	1	598		
SOUTH ATLANTIC	166	185	139	100	161	255	187	190	217	130	393	1720	10	.	5	3858		
EAST SOUTH CENTRAL	.	.	.	.	71	148	153	273	320	149	422	355	284	93	.	2268		
WEST SOUTH CENTRAL	2	.	2	3	2	.	1	1	3	2	.	1	3	1	.	21		
MOUNTAIN	29	13	41	18	79	208	94	96	114	62	116	136	31	.	6	1043		
PACIFIC	564	365	410	555	649	1136	715	773	754	730	731	647	812	614	582	10037		
TOTAL	857	608	701	909	1134	2287	1366	1580	1602	1217	1785	2935	1181	727	602	19491		