

**TABLE 2****Laboratory confirmed *Shigella* isolates reported to the CDC by Subgroup and Serotype in 2002**

<b>Rank</b>	<b>Serotype</b>	<b>Reported</b>	<b>Percent</b>
1	<i>S. sonnei</i>	10851	83.5
2	<i>S. flexneri</i> unspecified	734	5.7
3	<i>S. flexneri</i> 2 unspecified	183	1.4
4	<i>S. flexneri</i> 1 unspecified	110	0.9
5	<i>S. flexneri</i> 2a	101	0.8
6	<i>S. flexneri</i> 4 unspecified	74	0.6
7	<i>S. flexneri</i> 3 unspecified	70	0.5
8	<i>S. flexneri</i> 6	59	0.5
9	<i>S. boydii</i> unspecified	53	0.4
10	<i>S. flexneri</i> 4a	53	0.4
11	<i>S. flexneri</i> 3a	51	0.4
12	<i>S. dysenteriae</i> unspecified	23	0.2
13	<i>S. flexneri</i> 1b	23	0.2
14	<i>S. boydii</i> 2	22	0.2
15	<i>S. flexneri</i> 2b	14	0.1
16	<i>S. flexneri</i> 3b	13	0.1
17	<i>S. boydii</i> 4	10	0.1
18	<i>S. flexneri</i> 1a	9	0.1
19	<i>S. flexneri</i> 5 unspecified	9	0.1
20	<i>S. flexneri</i> variant y	8	0.1
21	<i>S. boydii</i> 1	7	0.1
22	<i>S. dysenteriae</i> 2	5	0.0
23	<i>S. dysenteriae</i> 4	5	0.0
24	<i>S. flexneri</i> 4b	5	0.0
25	<i>S. flexneri</i> variant x	4	0.0
26	<i>S. dysenteriae</i> 9	3	0.0
27	<i>S. flexneri</i> 3c	3	0.0
28	<i>S. boydii</i> 10	2	0.0
29	<i>S. boydii</i> 12	2	0.0
30	<i>S. boydii</i> 6	2	0.0
31	<i>S. dysenteriae</i> 10	2	0.0
32	<i>S. dysenteriae</i> 8	2	0.0
33	<i>S. boydii</i> 14	1	0.0
34	<i>S. boydii</i> 18	1	0.0
35	<i>S. dysenteriae</i> 1	1	0.0
36	<i>S. dysenteriae</i> 3	1	0.0
37	<i>S. dysenteriae</i> 5	1	0.0
	<b>Sub Total</b>	<b>12517</b>	<b>96.4</b>
	Unknown	475	3.7
	<b>Sub Total</b>	<b>475</b>	<b>3.7</b>
	<b>Total</b>	<b>12992</b>	<b>100.1</b>