

Rapid Assessment Reference Condition Model

The Rapid Assessment is a component of the LANDFIRE project. Reference condition models for the Rapid Assessment were created through a series of expert workshops and a peer-review process in 2004 and 2005. For more information, please visit www.landfire.gov. Please direct questions to helpdesk@landfire.gov.

Potential Natural Vegetation Group (PNVG)

R0GFLP Grand Fir/Lodgepole/Larch/Douglas-Fir Mix

General Information

Contributors (additional contributors may be listed under "Model Evolution and Comments")

Modelers

Pat Green pgreen@fs.fed.us
Jason Cole jcole@fs.fed.us
Sue Hagle shagle@fs.fed.us

Reviewers

Cathy Stewart cstewart@fs.fed.us
Steve Barrett sbarrett@mtdig.ne

Vegetation Type

Forested

General Model Sources

- Literature
 Local Data
 Expert Estimate

Rapid Assessment Model Zones

- | | |
|---|--|
| <input type="checkbox"/> California | <input type="checkbox"/> Pacific Northwest |
| <input type="checkbox"/> Great Basin | <input type="checkbox"/> South Central |
| <input type="checkbox"/> Great Lakes | <input type="checkbox"/> Southeast |
| <input type="checkbox"/> Northeast | <input type="checkbox"/> S. Appalachians |
| <input type="checkbox"/> Northern Plains | <input type="checkbox"/> Southwest |
| <input checked="" type="checkbox"/> N-Cent. Rockies | |

Dominant Species*

PICO
PSME
LAOC
ABGR

LANDFIRE Mapping Zones

10	21
19	22
20	29

Geographic Range

This PNVG occurs mostly in Idaho, eastern Washington, eastern Oregon, and western Montana. It is very important in Bailey's section M332.

Biophysical Site Description

Occurs above 4500 feet elevation, just above the grand fir with Douglas fir and larch zone (R0GFDF) and below the spruce-fir zone. Soils are underlain by granitics, metamorphics, and minor volcanic rocks. Most have a volcanic ash influenced loess surface layer.

Vegetation Description

Stands range from relatively open to densely stocked, and are usually dominated by a mix of early to mid seral species, including lodgepole pine, western larch, with lesser amounts of grand fir, Englemann spruce, and ponderosa pine. Grand fir increases markedly during mid to late successional stage, in the absence of fire and in response to pathogens that affect other species, like bark beetles. Stand understories range from moderately open to dense and include beargrass, mountain huckleberry, grouse whortleberry, serviceberry, and snowberry.

Disturbance Description

Fire regime group III, with stand replacing fires sometimes punctuated by mixed severity fires. Root disease and mountain pine beetle are very active in this PNVG.

Adjacency or Identification Concerns

This PNVG represents the warm/moderately moist grand fir habitat types (Pfister et al. 1977). This PNVG grades into grand fir with larch at lower elevations (R0GFDF) and western spruce-fir forest at higher elevations. It may be difficult to differentiate this PNVG from R0GFDF and R0WLLPDF, as the three types commonly overlap. This PNVG typically supports more lodgepole pine than the adjacent (lower elevation) grand fir type. The mosaic of open versus closed canopy is similar between this type and R0GFDF, but the

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

fire return interval is longer.

This PNVG may be similar to the PNVG R#MCONms from the Pacific Northwest model zone.

Scale Description

Sources of Scale Data Literature Local Data Expert Estimate

Terrain is usually rolling hills, convex ridges and mountain slopes with little dissection, so fires spread easily. Large infrequent fires result in large patch sizes, of 100's to 1000's of acres, and some occurrence of 10,000's of acres.

Issues/Problems

Proportion of seral structural stages may fluctuate widely over time because large stand replacing fires can affect 100,000 acres at a time.

Model Evolution and Comments

Workshop code was GFDF2.

Sources on historic composition are derived from Losensky (1993) and Sub-basin Assessments from the 1930s (US Department of Agriculture 1997-2003).

Review comments incorporated on 3/16/2005. As a result of the peer-review process, the mean fire return interval was increased to approximately 70 years (from 55 years) and the proportion of mixed to replacement fire was increased from 55:45 to approximately 70:30.

Succession Classes**

Succession classes are the equivalent of "Vegetation Fuel Classes" as defined in the Interagency FRCC Guidebook (www.frcc.gov).

Class A 15 %

Early1 PostRep

Description

Post stand-replacing fire, lasting about 30 years. This class is initially dominated by resprouting forbs and shrubs, and transitions to seedling and sapling-dominated. Lodgepole pine is a frequent early seral dominant. Douglas fir and larch are common, while ponderosa pine and grand fir are less common. Residual, large western larch often survive all but the most severe fire to serve as seed sources.

Dominant Species* and Canopy Position

XETE
VAGL
PICO
PSME

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	Min	Max
Cover	0 %	100 %
Height	no data	no data
Tree Size Class	no data	

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Class B 15%

Mid1 Closed

Description

Pole and immature forest (or mature lodgepole) of 30 to 100 years. Tree canopy cover of 40 percent or more. Lodgepole pine is the most common dominant. Douglas-fir and western larch are secondary dominants. Larch may be reduced by grand fir competition, in the absence of fire.

Dominant Species* and Canopy Position

PICO
PSME
LAOC
ABGR

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	Min	Max
Cover	40 %	100 %
Height	no data	no data
Tree Size Class	no data	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class C 25%

Mid1 Open

Description

Pole and immature forest (or mature lodgepole) of 30 to 100 years. Tree canopy less than 40 percent. These are usually created by mixed fire, root disease activity, or mountain pine beetle activity in mixed conifer stands.

Dominant Species* and Canopy Position

PICO
ABGR
PSME
LAOC

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	Min	Max
Cover	0 %	40 %
Height	no data	no data
Tree Size Class	no data	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class D 20%

Late1 Open

Description

Mature forest of 100 years or more. Tree canopy less than 40 percent. These are usually the result of mixed severity fire, leaving an overstory of larch, Douglas fir, with some residual grand fir or ponderosa pine and lodgepole. They may also occur as a result of insect or pathogen activity removing a Douglas fir, lodgepole or grand fir understory.

Dominant Species* and Canopy Position

LAOC
PSME
PIPO
PICO

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	Min	Max
Cover	0 %	40 %
Height	no data	no data
Tree Size Class	no data	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Class E 25%

Late I Closed

Description

Mature forest of 100 years or more. Tree canopy cover greater than 40 percent. These are usually the result of uninterrupted succession in areas of low root disease occurrence or in areas of larch dominance.

Dominant Species* and Canopy Position

LAOC
ABGR
PSME
PICO

Structure Data (for upper layer lifeform)

	Min	Max
Cover	40 %	100 %
Height	no data	no data
Tree Size Class	no data	

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Fuel Model no data

Disturbances

Disturbances Modeled

- Fire
- Insects/Disease
- Wind/Weather/Stress
- Native Grazing
- Competition
- Other:
- Other

Fire Regime Group: 3

- I: 0-35 year frequency, low and mixed severity
- II: 0-35 year frequency, replacement severity
- III: 35-200 year frequency, low and mixed severity
- IV: 35-200 year frequency, replacement severity
- V: 200+ year frequency, replacement severity

Fire Intervals (FI)

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class. All values are estimates and not precise.

Historical Fire Size (acres)

Avg: no data
Min: no data
Max: no data

Sources of Fire Regime Data

- Literature
- Local Data
- Expert Estimate

	Avg FI	Min FI	Max FI	Probability	Percent of All Fires
Replacement	220	50	250	0.00455	31
Mixed	100	35	150	0.01	69
Surface					
All Fires	69			0.01456	

References

Ager, A., D. Scott; C. Schmitt. 1995. UPEST: Insect and disease risk calculator for the forests of the Blue Mountains. File document. Pendleton, OR: USDA Forest Service, Pacific Northwest Region, Umatilla and Wallowa-Whimian National Forests. 25 p.

Amman, G.D. 1977. The role of mountain pine beetle in lodgepole pine ecosystems: impact on succession. In: W.J. Mattson, ed. The role of arthropods in forest ecosystems. Springer-Verlag, New York, New York, USA.

Barrett, S. W. 2004. Altered fire intervals and fire cycles in the Northern Rockies. Fire Management Today 64(3): 25-29.

Barrett, S. W. 2004. Fire Regimes in the Northern Rockies. Fire Management Today 64(2): 32-38.

Byler, J. W. ; Marsden, M. A.; Hagle, S.K. 1992. The probability of root disease on the Lolo national Forest, Montana. Can. J. For. Res. 20:987-994.

Byler, J. W. and S. K. Hagle. 2000. Succession Functions of Pathogens and Insects. Ecoregion sections M332a and M333d in northern Idaho and western Montana. Summary. USDA Forest Service, State and Private Forestry. R1-FHP 00-09. 37 pp.

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Filip, G.M.; Goheen, D.J. 1984. Root diseases cause severe mortality in white and grand fir stands of the Pacific Northwest. *Forest Science* 30: 138-142.

Furniss, M.M.; Livingston, R.L.; McGregor, M.D. 1981. Development of a stand susceptibility classification for Douglas-fir beetle (*Dendroctonus pseudotsugae*). In: Hedden, R.L. ; Barres, S.J.; Coster, J.E., tech. coords. Hazard rating systems in forest insect pest management. Symposium proceedings; 1980 July 31-August 1; Athens, Georgia. Gen. Tech. Rep. WO-27. Washington, D.C.: U.S.D.A.- Forest Service: 115-128.

Goheen, D.J.; E.M. Hansen. 1993. Effects of pathogens and bark beetles on forests. In: Beetle- pathogen interactions in conifer forests. Academic Press Ltd. PP. 176-196.

Hagle, S.; J. Schwandt; T. Johnson; S. Kegley; C. Bell Randall; J. Taylor; I. B. Lockman; N. Sturdevant; M. Marsden. 2000. Successional functions of pathogens and insects; Ecoregion sections M332a and M333d in northern Idaho and western Montana. Volume 2: Results. USDA Forest Service, State and Private Forestry, Northern Region. R1-FHP 00-11. 262 pp. Appendices.

Hagle, S.; T. Johnson; M. Marsden; L. Lewis; L. Stipe; J. Schwandt; J. Byler; S. Kegley; C. Bell Randall; J. Taylor; I. B. Lockman; N. Sturdevant; S. Williams. 2000. Successional functions of pathogens and insects; Ecoregion sections M332a and M333d in northern Idaho and western Montana. Volume 1: Methods. USDA Forest Service, State and Private Forestry, Northern Region. R1-FHP 00-10. 97 pp.

Hagle, S.K.; Byler, J.W. 1993. Root diseases and natural disease regimes in a forest of western U.S.A. In: Johansson, M.; Stenlid, J., eds., Proceedings of the Eighth International Conference on Root and Butt Rots, Wik, Sweden and Haikko, Finland, August 9-16, 1993. pp. 606-617.

Hagle, S.K.; Byler, J.W.; Jeheber-Matthews, S.; Barth, R.; Stock, J.; Hansen, B.; Hubbard, C. 1994. Root disease in the Coeur d'Alene river basin: An assessment. In: Interior Cedar-Hemlock-White pine forests: Ecology and Management, 1993, March 2-4; Spokane, WA: Washington State University, Pullman, WA. Pp. 335-344.

Haig, I. T.; K. P. Davis; R. H. Weidman. 1941. Natural regeneration in the western white pine type. USDA Tech. Bull. 767. Washington, DC: 99 p.

Holah, J. C.; M.V. Wilson; E.M. Hansen. Impacts of a native root-rotting pathogen on successional development of old-growth Douglas-fir forests. *Oecologia* (1977) 111: 429-433.

Kurz, W.A.; Beukema, S.J.; Robinson, D.C.E. 1994. Assessment of the role of insect and pathogen disturbance in the Columbia River Basin: a working document. Prepared by ESSA Technologies, Ltd., Vancouver, B.C. USDA Forest Service, Coeur d'Alene, ID, 56 pp.

Leiberg, J.B. 1900. The Bitterroot Forest Preserve. Dept of Interior, US Geological survey. 19th Annual Report, Part V. Forest Reserves. Washington, D.C. Pp 217-252.

Losensky, B.J. 1993. Historical vegetation in Region One by climatic section. USDA forest Service, Northern Region. Draft report on file at Nez Perce National forest.

Morgan, P. and Parsons, R. 2001. Historical range of variability of forests of the Idaho southern batholith Ecosystem. Final report to Boise Cascade Corporation, Boise, ID. On file at Nez Perce Forest Headquarters. 34 pp.

Morgan, P., SC bunting, AE Black, T Merrill, and S. Barrett. 1996. Fire regimes in the interior Columbia river Basin: past and Present. Final Report submitted to the Intermountain fire Sciences laboratory, Intermountain Research Station, Missoula, MT. 35 pp.

Pfister, R. D., B. L. Kovalchik, S. F. Arno, and R. C. Presby. 1977. Forest habitat types of Montana. USDA Forest Service, Intermountain Forest and Range Experiment Station, General Technical Report, INT-34.

Smith, Jane Kapler; Fischer, William C. 1997. Fire ecology of the forest habitat types of northern Idaho. Gen. Tech. Rep. INT-GTR-363. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station. 142 p.

Swetnam, T.W.; Lynch, A. 1989. A tree-ring reconstruction of western spruce budworm history in the southern Rocky Mountains. *For. Sci.* 35:962-986.

US Department of Agriculture. 1911. Extensive land survey. Nez Perce National Forest. On file at Forest headquarters.

US Department of Agriculture. 1914. Extensive land survey. Selway National Forest. On file at Forest headquarters.

US Department of Agriculture. 1938. Forest Statistics: Idaho County, Idaho. Forest Survey Release Number 15. A September progress Report. USDA Forest Service. Northern Rocky Mountain Forest and Range Experiment Station, Missoula MT. 31 pp.

US Department of Agriculture. 1954. Timber type map for the Nez Perce National Forest. On file at Forest headquarters.

US Department of Agriculture. 1997-2003. Subbasin Assessments: 1930s vegetation mapping

Veblen, T.T.; Hadley, K. S.; Reid, M. S.; Rebertus, A. J. 1991. The response of subalpine forests to spruce beetle outbreak in Colorado. *Ecology* 72(1): 213-231.

Williams, C.B.; Azuma, D.L.; Ferrell, G.T. 1992. Incidence and effects of endemic populations of forest pests in young mixed-conifer forest of the Sierra Nevada. USDA- Forest Service, Research Paper PSW-RP-212, Pacific Southwest Research Station, 8 p.