

Rapid Assessment Reference Condition Model

The Rapid Assessment is a component of the LANDFIRE project. Reference condition models for the Rapid Assessment were created through a series of expert workshops and a peer-review process in 2004 and 2005. For more information, please visit www.landfire.gov. Please direct questions to helpdesk@landfire.gov.

Potential Natural Vegetation Group (PNVG)

R0PIPObl Ponderosa Pine-Black Hills-Low Elevation

General Information

Contributors (additional contributors may be listed under "Model Evolution and Comments")

Modelers

Kelly Pohl kpohl@tnc.org
Cody Wienk cody_wienk@nps.gov
Carolyn Sieg csieg@fs.fed.us

Reviewers

Vegetation Type

Forested

General Model Sources

- Literature
 Local Data
 Expert Estimate

Rapid Assessment Model Zones

- California Pacific Northwest
 Great Basin South Central
 Great Lakes Southeast
 Northeast S. Appalachians
 Northern Plains Southwest
 N-Cent. Rockies

Dominant Species*

PIPO
PRVI
QUMA
ORAS

LANDFIRE Mapping Zones

10	21
19	22
20	29

Geographic Range

Black Hills region of Wyoming and South Dakota.

Biophysical Site Description

This PNVG is found below Ponderosa Pine Black Hills high elevation and above Ponderosa Pine Northern Plains (generally 4000-6000 ft), predominately on the lower limestone plateau and material weathered from metamorphic rocks. This type is generally on sites with sandy loam to clayey loam soils.

Vegetation Description

Ponderosa pine, bur oak (in northern Hills and Bear Lodge Mts.), chokecherry, Saskatoon serviceberry, aspen, Ribes species, rose species, ironwood, hawthorn, Oregon grape, raspberry, roughleaf ricegrass, Canada wildrye, needlegrasses, sideoats gramma, sedges.

Disturbance Description

Generally frequent fire return interval with surface fire. Mixed severity fire occurs if fire return intervals are missed, and stand replacement fire is very infrequent (300+ years). Precipitation is concentrated in April through June, but occurs throughout the growing season, resulting in good pine regeneration and dense patches of saplings. Elk, and to a lesser extent, bison, were important ungulates. Windthrow, storm damage, and mountain pine beetles were important disturbances in this type, especially when stands reached high densities.

Adjacency or Identification Concerns

This type occurs at elevations above Ponderosa Pine Northern Plains and at elevations below Ponderosa Pine Black Hills High Elevation. This type differs from Ponderosa Pine Black Hills High Elevation because it has more frequent surface fires, less frequent replacement fires, and less closed canopy forest.

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Scale Description

Sources of Scale Data Literature Local Data Expert Estimate

Patch size probably ranged from 10's to 1000's of acres. Most disturbances would have been relatively small and patchy in nature.

Issues/Problems

There is considerable debate over the role of mixed severity and surface fires in the historical range of variability in this and other ponderosa pine forests in the northern and central Rockies (Baker and Ehle 2001, 2003; Barrett 2004; Veblen et al. 2000).

Model Evolution and Comments

Quantitative model was developed post-workshop by Kelly Pohl with input from Cody Wienk and Carolyn Sieg. Additional input was provided during the workshop by Deanna Reyher, Blaine Cook, Bill Baker and factored into the model development. Because of the model's late development it received no peer review.

Succession Classes**

Succession classes are the equivalent of "Vegetation Fuel Classes" as defined in the Interagency FRCC Guidebook (www.frcc.gov).

Class A 5%

Early1 PostRep

Description

Herbaceous/shrubby post-replacement class, persists 0-15 years. In Bear Lodge this stage dominated by bur oak. In the Black Hills proper, lower limestone, it is dominated by grass/forb with chokecherry, serviceberry, leadplant, raspberry, rose, and current present.

Dominant Species* and Canopy Position

QUMA

PRVI

AMAL

PIPO

Upper Layer Lifeform

- Herbaceous
 Shrub
 Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	Min	Max
Cover	0 %	100 %
Height	no data	no data
Tree Size Class	no data	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class B 15%

Mid1 Closed

Description

Pole (dog hair), persists 15-50 years. Very few understory species present due to canopy closure.

Dominant Species* and Canopy Position

PIPO

Upper Layer Lifeform

- Herbaceous
 Shrub
 Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	Min	Max
Cover	50 %	100 %
Height	no data	no data
Tree Size Class	no data	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Class C 20%

Mid1 Open
Description

Open pole with patches of 100+ year old trees, persists 15-50 years. In Bear Lodge Mountains, bur oak persists, particularly in open canopy stands.

Dominant Species* and Canopy Position

PIPO
AUMA
ORAS
PRVI

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	Min	Max
Cover	0 %	50 %
Height	no data	no data
Tree Size Class	no data	

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class D 50%

Late1 Open
Description

Open canopy stand; persists 50+ years. Patches of dense doghair and 200+ year old trees persist. Bur oak mostly restricted to northern Black Hills and Bear Lodge. Common juniper and rough leaf ricegrass common.

Dominant Species* and Canopy Position

PIPO
ORAS
JUCO
QUMA

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	Min	Max
Cover	0 %	60 %
Height	no data	no data
Tree Size Class	no data	

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class E 10%

Late1 Closed
Description

Closed canopy, multi-layer stand, persists 50+ years. At >70% canopy closure, mountain pine beetle outbreaks occur, opening up the canopy. Ironwood and bur oak in northern Black Hills and Bear Lodge Mountains.

Dominant Species* and Canopy Position

PIPO
JUCO
OSVI
QUMA

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	Min	Max
Cover	60 %	100 %
Height	no data	no data
Tree Size Class	no data	

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Disturbances

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Disturbances Modeled

- Fire
- Insects/Disease
- Wind/Weather/Stress
- Native Grazing
- Competition
- Other:
- Other

Historical Fire Size (acres)

Avg: no data
 Min: no data
 Max: no data

Sources of Fire Regime Data

- Literature
- Local Data
- Expert Estimate

Fire Regime Group: 1

- I: 0-35 year frequency, low and mixed severity
- II: 0-35 year frequency, replacement severity
- III: 35-200 year frequency, low and mixed severity
- IV: 35-200 year frequency, replacement severity
- V: 200+ year frequency, replacement severity

Fire Intervals (FI)

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class. All values are estimates and not precise.

	<i>Avg FI</i>	<i>Min FI</i>	<i>Max FI</i>	<i>Probability</i>	<i>Percent of All Fires</i>
<i>Replacement</i>	300	200	400	0.00333	7
<i>Mixed</i>	100	50	400	0.01	21
<i>Surface</i>	30	5	50	0.03333	71
<i>All Fires</i>	21			0.04667	

References

Baker, William L., Ehle, Donna S. 2001. Uncertainty in surface-fire history: The case of ponderosa pine forests in the western United States. *Canadian Journal of Forest Research* 31: 1205-1226.

Baker, William L.; Ehle, Donna S. 2003. Uncertainty in fire history and restoration of ponderosa pine forests in the western United States. In: Omi, Philip N.; Joyce, Linda A., tech. eds. *Fire, fuel treatments, and ecological restoration: conference proceedings; 2002 April 16-18; Fort Collins, CO. Proceedings RMRS-P-29. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station: 319-333.*

Barrett, S. W. 2004. Altered fire intervals and fire cycles in the Northern Rockies. *Fire Management Today* 64(3): 25-29.

Barrett, S. W. 2004. Fire Regimes in the Northern Rockies. *Fire Management Today* 64(2): 32-38.

Brown and Sieg. 1996. Fire history in interior ponderosa pine communities of the Black Hills, South Dakota, USA. *International Journal of Wildland Fire*. 6:97-105.

Brown, P.M., M.G. Ryan, and T.G. Andrews. 2000. Historical surface fire frequency in ponderosa pine stands in Research Natural Areas, Central Rocky Mountains and Black Hills, USA. *Natural Areas Journal* 20:133-139.

Graves, H.S. 1899. The Black Hills Forest Reserve. In: the 19th Annual Report of the Survey, 1897-1898. Part V. Forest Reserves. Washington, D.C. U.S. G.S. 67-164.

Parrish, J.B., D.J. Herman, D.J. Reyher. 1996. A century of change in Black Hills forest and riparian ecosystems. U.S. Forest Service and South Dakota Agricultural Experiment Station B722, South Dakota State University, Brookings, SD.

Progulske, D.R. 1974. Yellow ore, yellow hair, yellow pine: A photographic study of a century of forest ecology. *Agricultural Experiment Station Bulletin* 616, South Dakota State University, Brookings, SD.

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Sheppard, W.D., and M.A. Bettaglia. 2002. Ecology, silviculture and management of Black Hills Ponderosa Pine. RMRS GTR-97, Rocky Mtn. Research Station, Fort Collins, CO.

Shinneman, D. J., and W. L. Baker. 1997. Nonequilibrium dynamics between catastrophic disturbances and old-growth forests in ponderosa pine landscapes of the Black Hills. *Conservation Biology* 11:1276-1288.

Uresk, D.W., and K.E. Severson. 1989. Understory-overstory relationships in ponderosa pine forests, Black Hills, SD. *Journal of Range Management* 42: 203-208.

Veblen, Thomas T.; Kitzberger, Thomas; Donnegan, Joseph. 2000. Climatic and human influences on fire regimes in ponderosa pine forests in the Colorado Front Range. *Ecological Applications*. 10(4): 1178-1195.

Wienk, C. L., C. H. Sieg and G. R. McPherson. 2004. Evaluating the role of cutting treatments, fire and soil seed banks in an experimental framework in ponderosa pine forest of the Black Hills, South Dakota. *Forest Ecology and Management* 192: 375-393.