

Rapid Tornadogenesis in Nelson County, ND - 18 July 2001

David J. Kellenbenz
National Weather Service - Eastern North Dakota

1. INTRODUCTION

A supercell thunderstorm developed rapidly in southwest Nelson County, ND during the afternoon of 18 July 2001, and produced a long lived weak F3 tornado. The tornado touchdown occurred in an area with numerous intersecting surface boundaries. The tornadic supercell's mid level rotation persisted for only 10-15 minutes before producing the tornado. Research done by Burgess et. al (1993) on supercell thunderstorms indicate mesocyclones were present on average at least 36 minutes prior to tornadogenesis. The Nelson county tornado also occurred very close to three distinct boundaries. Research by Atkins et al. (1999) has shown that mesocyclonic low level circulations were "stronger and more persistent" in the vicinity of boundaries, than when boundaries were not present. During Project VORTEX, 70% of F2 or greater tornadoes occurred near low level boundaries (Markowski et al. 1998a). In addition, the tornado occurred on the cool side of a boundary, which could have lead to lower lifted condensation levels (LCLs) in the local tornadogenesis region. Recent research has indicated that lower LCLs may aid in tornadogenesis, which occurs on the cool side of boundaries (Markowski 2003).

This paper will examine the rapid tornadogenesis as seen from the WSR-88D and mesoanalysis point of view. In addition, hypotheses are presented as to why tornadogenesis occurred so quickly, and if this tornado could have been anticipated sooner with the environmental conditions that were present.

2. SYNOPTIC OVERVIEW

A broad west to southwest flow was present in the upper troposphere present, as a 35 m/s speed max was approaching the Northern Plains at 300 hPa, along with divergence at 300 hPa (Fig. 1). Dry air was being advected into the region in the mid levels, as 700mb dewpoint depressions rose around 2 degrees C from 1200 UTC to 0000 UTC near Bismarck, ND (KBIS). A very unstable low-level environment was present with convective available potential energy (CAPE) around 6,000 J/Kg were present in northeastern ND during the afternoon hours. Storm relative helicity (SRH) and shear values were in the moderate category with 0-3km SRH values around $200 \text{ m}^2/\text{s}^2$, and around 35kts of shear in the 0-6 km layer. Abundant moisture was available as well, with surface dewpoints in the low and middle 70s and precipitable water amounts of 1.80 inches.




Figure 1. 300 hPa analysis for 00 UTC 19 July 2001.

3. MESOSCALE ANALYSIS

Numerous surface boundaries and a meso-low were present on the surface analysis near the eventual tornadogenesis region around tornadogenesis time at 2300 UTC (Fig. 2). These discontinuity lines were also seen on the KVMX WSR 88-D base reflectivity imagery, as they were oriented from south-southwest to north-northeast, from south-southeast to north-northwest, in addition to horizontal convective rolls oriented southeast to northwest at 2238 UTC in base reflectivity imagery (Fig. 3).




Figure 2. Surface analysis for 2300 UTC 18 July 2001.




Figure 3. KMXV WSR-88D 0.5 degree base reflectivity image at 2238 UTC on 18 July 2001.

The movement of the surface boundaries throughout the afternoon were seen in the surface analyses, as well as in the radar reflectivity imagery. A meso-low circulation developed around 2200 UTC in northeastern North Dakota, shortly before tornadogenesis commenced around 2300 UTC. This focused boundary layer convergence near the intersection of the mesoscale

boundaries. 0-3km SRH near Grand Forks at 2100 UTC on 18 July 2001 from the 12Z ETA model was predicted to be around $200 \text{ m}^2/\text{s}^2$. It is believed that the SRH was significantly enhanced on the supercellular scale, as stated by Rasmussen and Markowski (2000). The hodograph from the Dimmitt, TX tornado of 2 June 1995 revealed a SRH of 1100 J/kg . It may be common for an increase of SRH locally near a supercell updraft due to boundary interaction. It is hypothesized that augmentation of the local SRH occurred in the vicinity of the tornado, which was near numerous surface boundary intersections. In addition, the rapid stretching of horizontal vorticity into vertical vorticity from the powerful updraft also lead to rapid tornado formation once the supercell crossed onto the “cold” side of the boundary (Markowski et al. 1998a).

4. RADAR ANALYSIS OF THE TORNADIC SUPERCELL

Horizontal convective rolls, N-S and SW-NE oriented boundaries were present in southwest Nelson County before convective initiation occurred around 2213 UTC . A cross section of reflectivity imagery (Figs. 4 and 5) shows the rapid growth of the supercell from 2246 UTC until 2300 UTC , and the attendant strong updraft with a pronounced Bounded Weak Echo Region (BWER).




Figure 4. KMXV WSR-88D reflectivity cross section at 2246 UTC 18 July 2001.




Figure 5. KMXV reflectivity cross section at 2323 UTC on 18 July 2001 with pronounced bounded weak echo region.

Mid level rotation in the KMXV WSR-88D storm relative motion (SRM) data increased from a very weak mesocyclone circulation at 2240 UTC (Fig. 6), to a moderate mesocyclonic circulation 15 minutes later at 2255 UTC (Fig. 7). The circulation aloft rapidly translated to the surface, with the first tornado reported around 2255 UTC. Therefore, the mesocyclone persisted for about 10-15 minutes before producing a tornado. Research done by Burgess et al. (1993) indicate mesocyclones were present, on average at least 36 minutes before tornadogenesis occurred. This tornado developed shortly after a mesocyclonic circulation in the mid-levels was first detected.




Figure 6. KMOVX 4.3 degree Storm Relative Motion image at 2240 UTC 18 July 2001.




Figure 7. KMVX 5.2 degree Storm Relative Motion image at 2255 UTC 18 July 2001.

Once the three surface boundaries intersected (evident with the base reflectivity imagery), a tornado was produced. At 2253 UTC (Fig. 8), surface boundaries were distinctly separated, but once the supercell crossed the boundary intersection, tornadogenesis occurred around 2258 UTC (Fig. 9). The tornado touched down on the cool side of the boundary intersection, which has been linked to persistent and stronger tornadoes than those forming along or on the warm side of boundaries (Houston and Wilhelsom 1998). This may have also lead to lower LCLs in a potentially warm-type rear flank downdraft (RFD), which can then be easily ingested into the attendant strong updraft. Visual clues of a notable clear slot with the potential warm-type RFD

were also noted by storm chasers following this supercell (Markowski 2003).




Figure 8. KMVX 0.5 degree base reflectivity image just before tornadogenesis at 2253 UTC 18 July 2001.




Figure 9. KMVX 0.5 degree base reflectivity image around tornadogenesis time at 2258 UTC 18 July 2001 as boundaries intersect in extreme southwest Nelson County.

5. SUMMARY AND DISCUSSION

Utilization of surface and radar data played a key role in identifying the many mesoscale boundaries present near the tornadogenesis region. Once the mesocyclone formed, the tornado formed rapidly in 10-15 minutes. Video of the storm from a College of Dupage (IL) chase team showed a classic, well defined tornado, wall cloud and RFD wrapping around the parent mesocyclone, evident in the clear slot behind the parent updraft. This is uncommon as most mesocyclones persist for at least 20 to 30 minutes before producing the classic, non-landspout tornado.

The strong updraft ingested enhanced SRH, which can be augmented by an order of magnitude on the tornado scale. The strong updraft was able to rapidly convert horizontal into vertical vorticity as it crossed the boundary intersection. Therefore, mesoscale boundaries appear to play a key role in increasing SRH on the mesoscale, so close attention must be paid to them in a region of moist, deep convection. The “exceedingly variability” of SRH may explain why some storms in the same mesoscale environment produce tornadoes and some fail. This is a topic of ongoing research and will be better understood once the resolution of mesoscale models improve. (Markowski et al. 1998b). Research is now showing that this process from mesocyclone to tornadic development can take as little as 10 minutes (Rasmussen 2003). This timescale was comparable to the tornado highlighted in this paper.

There was no lead time for this particular tornadogenesis event. A severe thunderstorm warning was in effect for this supercell before reports of the tornado touchdown were received. An upgrade to a tornado warning occurred when the report of the tornado was received. The local storm-scale 0-1km SRH values were likely greatly enhanced near the boundary intersection at tornadogenesis time. This interaction would have caused a local increase in horizontal vorticity, which was then rapidly stretched into the vertical with the intense updraft speed of the supercell. A common misconception for warning operations is the required persistence of a long-lived mesocyclone (more than 20 minutes) prior to tornadogenesis. Although this may be true for the majority of F2 or greater tornadoes, this case demonstrates how a supercell can rapidly become tornadic in about 10 minutes. On-going mesoanalysis of boundary intersection in conjunction with the thermodynamic conditions (extreme instability and low LCL heights) and WSR-88D trends may have led to earlier anticipation of tornado formation.

6. ACKNOWLEDGMENTS

The author would like to thank Bradley Bramer (SOO FGF) for providing his review, comments and valuable suggestions on the manuscript.

7. REFERENCES

Atkins, N. T, M. L. Weisman, and L. J. Wicker, 1999: The influence of pre-existing boundaries on supercell evolution. *Mon. Wea. Rev.*, **127**, 2910-2927.

Burgess, D. W, R. J. Donaldson, and P. R. Desrochers, 1993: Tornado Detection and Warning by Radar. *The Tornado: Its Structure, Dynamics, Prediction, and Hazards, Geophys. Monogr., No. 79, Amer. Geophys. Union*, 203-221.

Houston, L. A, and R. B. Wilhelmson, 1998: Numerically modeled interactions between supercells and boundaries. *19th Conference on Severe Local Storms Minneapolis MN, Amer. Meteor. Soc.*, 332-333.

Markowski, P. M., E. N. Rasmussen, and J. M. Straka, 1998a: The occurrence of tornadoes in supercells interacting with boundaries during VORTEX-95. *Wea. Forecasting*, **13**, 852-859.

Markowski, P. M., J. M. Straka, and E. N. Rasmussen, D. O. Blanchard, 1998b: Variability of Storm-Relative Helicity during VORTEX. *Mon. Wea. Rev.*, **126**, 2959-2971.

Markowski, P. M., 2003: Current Research on RFD/LCL characteristics. URL found at <http://www.meteo.psu.edu/~marko/research.html>.

Rasmussen E. N. and P. M. Markowski, 2000: Tornado Forecasting. URL found at <http://www.nssl.noaa.gov/~erik/www/SSR>.