

Table 93. Contraceptive Use by Women, 15 to 44 Years of Age, Who Have Ever Had Sexual Intercourse: 1982 to 2002

[46,684 represents 46,684,000. Based on the National Survey of Family Growth, see Appendix III]

Contraceptive status and method	1982	1995	2002			
			Total ¹	White only, Non-Hispanic	Black only, Non-Hispanic	Hispanic ²
Women who have had sexual intercourse (1,000)...	46,684	53,800	54,190	34,999	7,403	7,887
PERCENT DISTRIBUTION						
Ever used any method	94.8	98.2	98.2	98.8	97.3	96.5
Female sterilization	22.3	23.4	20.7	19.2	26.9	24.1
Male sterilization	10.1	14.6	13.0	16.7	5.0	4.4
Pill	76.3	82.2	82.3	87.0	79.8	68.5
Norplant implant	(X)	2.1	2.1	1.4	3.3	4.0
1-month injectable (Lunelle TM)	(X)	(X)	0.9	0.5	1.0	3.1
3-month injectable (Depo-Provera TM)	(X)	4.5	16.8	13.6	23.3	24.3
Emergency contraception	(X)	0.8	4.2	4.2	4.1	3.8
Contraceptive patch	(X)	(X)	0.9	0.7	1.4	1.1
Today TM	(X)	12.0	7.3	8.8	6.3	2.7
Intrauterine device	18.4	10.0	5.8	4.7	5.4	10.0
Diaphragm	17.1	15.2	8.5	10.4	6.9	3.4
Condom.	51.8	82.0	89.7	92.2	91.8	78.2
Female condom.	(X)	1.2	1.9	1.2	5.4	1.4
Periodic abstinence—calendar rhythm	17.0	24.3	16.2	16.6	13.7	16.2
Periodic abstinence—natural family planning.	2.3	4.2	3.5	3.6	1.7	4.7
Withdrawal	24.5	40.6	56.1	59.7	50.5	47.2
Foam alone	24.9	18.3	12.1	12.5	15.5	8.2
Jelly/cream alone.	5.8	9.1	7.3	7.7	9.6	3.4
Suppository/insert	9.7	10.6	7.5	7.9	8.8	4.5
Other methods	9.3	0.3	1.0	0.8	1.3	1.3

X Not applicable. ¹ Includes other races, not shown separately. ² Persons of Hispanic origin may be any race.

Source: U.S. National Center for Health Statistics, *Advance Data*, Number 350, "Use of Contraception and Use of Family Planning Services in the United States: 1982–2002," December 10, 2004.