

Alabama, 2005

Primary and secondary syphilis

- CASES: Alabama reported 169 cases in 2005.
- RATES: Alabama ranked 13 among 50 states + DC + 3 territories: 3.7 cases per 100,000 population compared to the U.S. rate of 3.0 cases per 100,000 population.

Male-female rates of primary and secondary syphilis

- In Alabama, the rate among males was 4.6 per 100,000 population compared to the U.S. male rate of 5.1 per 100,000.
- The rate among females was 3.0 per 100,000 compared to the U.S. female rate of 0.9 per 100,000.

Racial disparities of primary and secondary syphilis


In Alabama, the race/ethnicity adjusted rates per 100,000 population were 1.1 among whites, 11.0 among blacks, 0.0 among Hispanics, 2.5 among Asian/Pacific Islanders, and 4.1 among American Indians/Alaska Natives.

- The rate among blacks was 10 times that of whites.
- The rate among Asian/Pacific Islanders was 2.3 times that of whites.
- The rate among American Indians/Alaska Natives was 3.7 times that of whites.


Congenital syphilis

- CASES: Alabama reported 5 cases in 2005.
- RATES: Alabama ranked 14 among 28 states + 2 territories reporting at least one congenital syphilis case: 8.4 cases per 100,000 live births compared to the U.S. rate of 8.0 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Alabama, 1996-2005



Total P&S Syphilis Rates by County, Alabama, 2005



Rate (per 100,000 population)

