

New York, 2005

Primary and secondary syphilis

- **CASES:** New York reported 705 cases in 2005.
- **RATES:** New York ranked 11 among 50 states + DC + 3 territories: 3.7 cases per 100,000 population compared to the U.S. rate of 3.0 cases per 100,000 population.

Male-female rates of primary and secondary syphilis

- In New York, the rate among males was 7.3 per 100,000 population compared to the U.S. male rate of 5.1 per 100,000.
- The rate among females was 0.3 per 100,000 compared to the U.S. female rate of 0.9 per 100,000.

Racial disparities of primary and secondary syphilis


In New York, the race/ethnicity adjusted rates per 100,000 population were 1.9 among whites, 10.9 among blacks, 4.8 among Hispanics, 0.9 among Asian/Pacific Islanders, and 2.8 among American Indians/Alaska Natives.

- The rate among blacks was 5.7 times that of whites.
- The rate among Hispanics was 2.5 times that of whites.
- The rate among American Indians/Alaska Natives was 1.5 times that of whites.


Congenital syphilis

- **CASES:** New York reported 8 cases in 2005.
- **RATES:** New York ranked 19 among 28 states + 2 territories reporting at least one congenital syphilis case: 3.2 cases per 100,000 live births compared to the U.S. rate of 8.0 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, New York, 1996-2005



Total P&S Syphilis Rates by County, New York, 2005



Rate (per 100,000 population)

