

Mutations in Retroviral Genes Associated with Drug Resistance

Urvi Parikh,¹ Charles Calef,² Brendan Larder,³ Raymond Schinazi,⁴
and John W. Mellors¹

¹ *University of Pittsburgh, Scarfe Hall, Suite 818, 3550 Terrace St., Pittsburgh, PA 15261.*

² *T10, MS K710, Los Alamos National Laboratory, Los Alamos, NM 87545.*

³ *Virco, U.K., 162A Cambridge Science Park, Milton Road, Cambridge CB4 4GH, U.K.*

⁴ *Emory University/VAMC, 1670 Clairmont Rd., Decatur, GA 30033.*

Introduction

Drug resistance is the inevitable consequence of incomplete suppression of HIV replication. The rapid replication rate of HIV and its inherent genetic variation have led to the identification of many HIV variants that exhibit altered drug susceptibility. The growing number of drug resistance mutations listed in this revised table stands as a testimony to the genetic flexibility of HIV. This table, updated in March 2003, lists 647 HIV-1 mutation/drug combinations, of which 179 occur in protease, 1 in integrase, 305 in RT, and 139 in Env. Although the tables are quite comprehensive, the reader should be reminded that the mutations described are predominantly found in clade B virus and not in other HIV genotypes. The revised table also includes drug resistance mutations that have been identified for SIV and FIV.

In the table the phrase “Enzyme resist” refers to inhibition assays done just with a mutated enzyme. Instead of introducing the mutations into a virus and testing the susceptibility of the mutant virus to a drug, researchers introduce the mutation(s) into the enzyme and determine their effect by running enzyme activity assays. This type of susceptibility testing does not take into account changes in other viral proteins (like Gag) that would also help confer resistance, which is the reason for distinguishing enzyme resistance from whole virus resistance. In the “Amino Acid Change” column a + means amino acids have been inserted into the sequence, while a Δ indicates a deletion. In the “Class of Drug” column the abbreviation MN stands for “Multiple Nucleoside” and refers to resistance to combinations of nucleoside RTIs. Other abbreviations used in the table are listed in a separate Abbreviations Table on page 158.

All of the information contained in these printed tables and other useful tools are available at our Web site: http://resdb.lanl.gov/Resist_DB.

Acknowledgments

The authors would like to gratefully acknowledge their colleagues for assistance in assembling this table. This work was supported in part by the National Institutes of Health and the Department of Veterans Affairs.

Contents

HIV1 Gag	95
HIV1 RT	96
HIV1 Protease	118
HIV1 Integrase	138
HIV1 Envelope	139
SIV RT	163
FIV RT	164
Table of Abbreviations	165
Table of Compounds	165
References	170

Mutations in HIV Gag that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
E 12 K		PI	VX-478 (amprenavir)	Y				in vitro selection in MT-2 cells, passage 31	Gatanaga02
V 35 I		PI	VX-478 (amprenavir)	Y				in vitro selection in MT-2 cells, passage 10	Gatanaga02
L 75 R		PI	VX-478 (amprenavir)	Y				in vitro selection in MT-2 cells, passage 10	Gatanaga02
H 219 Q		PI	JE-2147	Y				in vitro selection in MT-2 cells, passage 6	Gatanaga02
H 219 Q		PI	KNI-272	Y				in vitro selection in MT-2 cells	Gatanaga02
H 219 Q		PI	UIC-94003	Y				in vitro selection in MT-2 cells	Gatanaga02
H 219 Q		PI	VX-478 (amprenavir)	Y				in vitro selection in MT-2 cells, passage 10	Gatanaga02
V 390 A		PI	JE-2147	Y				in vitro selection in MT-2 cells, passage 6	Gatanaga02
V 390 D		PI	VX-478 (amprenavir)	Y				in vitro selection in MT-2 cells, passage 31	Gatanaga02
R 409 K		PI	JE-2147	Y				in vitro selection in MT-2 cells, passage 33	Gatanaga02
R 409 K		PI	KNI-272	Y				in vitro selection in MT-2 cells	Gatanaga02
R 409 K		PI	UIC-94003	Y				in vitro selection in MT-2 cells	Gatanaga02
R 409 K		PI	VX-478 (amprenavir)	Y				in vitro selection in MT-2 cells, passage 10	Gatanaga02
L 449 F		PI	JE-2147	Y				in vitro selection in MT-2 cells, passage 33	Gatanaga02
L 449 F		PI	VX-478 (amprenavir)	Y				in vitro selection in MT-2 cells, passage 20	Gatanaga02
E 468 K		PI	VX-478 (amprenavir)	Y				in vitro selection in MT-2 cells, passage 20	Gatanaga02

Mutations in HIV RT that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
M 41 L	ATG → TTG/CTG	NRTI	AZT (zidovudine)	?	Y	4		M41L/T215Y: 60–70-fold; M41L/D67N/K70R/T215Y: 180-fold.	Larder89, Larder91, Kellam92
E 44 A	GAA → GCA	NRTI	3TC (lamivudine)	N	Y			Seen in NRTI-experienced patients but not in antiretroviral naïve patients.	Montes02
E 44 D	GAA → GAC	MN	AZT (zidovudine) + 3TC (lamivudine)	N	Y	1.0		Confers moderate levels of resistance to 3TC (4 to 50-fold) when present in an AZT-resistant genetic background without diminishing AZT resistance.	Hertogs00
A 62 V	GCC → GTC	MN		N	Y	Nil		A62V alone has no effect, but in combination with mutations at 75, 77, 116, 151 causes multi NRTI resistance.	Iversen96, Shirasaka95
I 63 M	ATA → ATG	NNRTI	BHAP U-90152 (delavirdine)	Y	N			Compensatory mutation restoring replication competence to W229Y mutant. I63M/V189I/W229Y/E396G mutant 2.5-fold resistant to delavirdine with respect to WT	Pelemans01
I 63 M	ATA → ATG	NNRTI	BL-RG-587 (nevirapine)	Y	N			Compensatory mutation restoring replication competence to W229Y mutant. I63M/V189I/W229Y/E396G mutant 23-fold resistant to nevirapine with respect to WT	Pelemans01
I 63 M	ATA → ATG	NNRTI	GW 420867X	Y	N			Compensatory mutation restoring replication competence to W229Y mutant. I63M/V189I/W229Y/E396G mutant 8-fold resistant to GW 420867X with respect to WT	Pelemans01
I 63 M	ATA → ATG	NNRTI	HBV 097	Y	N			Compensatory mutation restoring replication competence to W229Y mutant. I63M/V189I/W229Y/E396G mutant 5-fold resistant to HBV 097 with respect to WT	Pelemans01

Mutations in HIV RT that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
I 63 M	ATA → ATG	NNRTI	I-EBU (emivirine)	Y	N			Compensatory mutation restoring replication competence to W229Y mutant. I63M/V189I/W229Y/E396G mutant 7-fold resistant to emivirine with respect to WT	Pelemans01
I 63 M	ATA → ATG	NNRTI	UC-781	Y	N			Compensatory mutation restoring replication competence to W229Y mutant. I63M/V189I/W229Y/E396G mutant 2.5-fold resistant to UC-781 with respect to WT	Pelemans01
K 65 R	AAA → AGA	NRTI	1592U89 (abacavir)	Y	N	3		K65R/L74Y: 3.6-fold; K65R/M184V: 7-fold; K65R/L74V/M184V: 10.2-fold	Tisdale97
K 65 R	AAA → AGA	NRTI	ddC (zalcitabine)	Y	Y	4-10			Zhang94, Gu94
K 65 R	AAA → AGA	NRTI	ddI (didanosine)	Y	Y	4-10	ddC; PMEA; 3TC5	Infrequently observed in patients receiving ddI or ddC	Zhang94
K 65 R	AAA → AGA	NRTI	dOTC	Y	?			K65R/M184V: 4.2-fold.	Rando99
K 65 R	AAA → AGA	NRTI	DXG	Y	?	8	other dioxolane derivatives	Reverses AZT resistance in D67N/K70R/T215Y/K219Q background	Mellors96
K 65 R	AAA → AGA	NRTI	PMEA (adefovir)	Y	N	10-25			Gu95, Foli96
K 65 R	AAA → AGA	NRTI	PMPA (tenofovir)	Y	?	3.5			Cherrington97
67	deletion	MN	AZT (zidovudine) + ddI (didanosine)	N	Y	AZT: 1.2, ddI: 3.8, ddC:18.0		Δ67/T69G/AZT resistance mutations: 445-fold AZT resistance. When NNRTI resistance mutations are also added, AZT resistance increases to 1,813-fold.	Imamichi00a, Imamichi00b, Imamichi01
D 67 E	GAC → GAG	MN		N	Y				Larder99
D 67 G	GAC → GGC	MN		N	Y				Larder99
D 67 G	GAC → GAG	NNRTI	(-)dOTFC	Y	?	4.5	(-)dOTFC: 5.2-fold		Richard00
D 67 N	GAC → AAC	NRTI	AZT (zidovudine)	Y	Y			D67N/K70R/T215Y/K219Q: 120-fold; M41L/D67N/K70R/T215Y: 180-fold.	Larder89, Larder91, Kellam92
D 67 S	GAC → AAC	MN		N	Y				Larder99

Mutations in HIV RT that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
S 68 G	AGT→GGT	MN		?	Y			Frequently associated with other multi-ddN resistance mutations V75I, F77L, F116Y and Q151M.	Schmitz98
S 68 N	AGT→AAT	MN		N	Y				Larder99
S 68 N	AGT→AAT	MN		N	Y				Larder99
S 68 S + GGG	AGT→AAT	MN		N	Y				Larder99
S 68 S + SS	AGT→AAT	MN		N	Y				Larder99
S 68 S + SSG	AGT→AAT	MN		N	Y				Larder99
S 68 S + ST	AGT→AAT	MN		N	Y				Larder99
S 68 S + SV	AGT→AAT	MN		N	Y				Larder99
S 68 Y	AGT→TAT	MN		N	Y				Larder99
ins 69 TRVMG	ACT+→ACG AGA GTG ATG GGG	NRTI	1592U89 (abacavir)	Y	Y	32			Lobato02
ins 69 TRVMG	ACT+→ACG AGA GTG ATG GGG	NRTI	3TC (lamivudine)	Y	Y	84			Lobato02
ins 69 TRVMG	ACT+→ACG AGA GTG ATG GGG	NRTI	AZT (zidovudine)	Y	Y	371			Lobato02
ins 69 TRVMG	ACT+→ACG AGA GTG ATG GGG	NRTI	d4T (stavudine)	Y	Y	15			Lobato02
ins 69 TRVMG	ACT+→ACG AGA GTG ATG GGG	NRTI	ddC (zalcitabine)	Y	Y	4			Lobato02
ins 69 TRVMG	ACT+→ACG AGA GTG ATG GGG	NRTI	ddI (didanosine)	Y	Y	12			Lobato02
T 69 A	ACT→GCT	MN	3TC (lamivudine) + d4T (stavudine)	?	Y			Seen in one patient on 3TC + d4T combination therapy.	Lawrence99
T 69 A + SG	ACT→GCT + AGT GGT	MN		?	Y	Confers >4-fold resistance to: AZT, ddI, ddC, 3TC and PMEA.		Seen in heavily treated patients.	Winters98
T 69 D	ACT→GAT	MN	AZT (zidovudine) + 3TC (lamivudine)	?	Y			Seen in one patient on AZT + 3TC combination therapy.	Lawrence99
T 69 D	ACT→GAT	NRTI	ddC (zalcitabine)	N	Y	5			Fitzgibbon92
T 69 G	ACT→GGT	MN	AZT (zidovudine) + ddI (didanosine)	N	Y	AZT: 1.5, ddC: 11.0, ddI: 10.0		Δ67/T69G/AZT resistance mutations: 445-fold AZT resistance. When NNRTI resistance mutations are also added, AZT resistance increases to 1,813-fold.	Imamichi00a, Imamichi00b, Imamichi01

Mutations in HIV RT that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
T 69 N	ACT→AAT	MN	3TC (lamivudine) + ddI (stavudine)	?	Y			Seen in two patients on 3TC + ddI combination therapy.	Lawrence99
T 69 S + AG	ACT→AAT	MN		N	Y				Larder99
T 69 S + EA	ACT→AGT + AGA GCA	MN		?	Y	Confers >4-fold resistance to: AZT, ddI, ddC, 3TC and PMEA.		Seen in heavily treated patients.	Winters98
T 69 S + EE	ACT→AGT + AGA GCA	MN		N	Y				Larder99
T 69 S + RA	ACT→AGT + AGA GCA	MN		?	Y	Confers >4-fold resistance to: AZT, ddI, ddC, 3TC and PMEA.		Seen in heavily treated patients.	Winters98
T 69 S + SA	ACT→AGC + AGC GCT	MN		?	Y	Confers >4-fold resistance to: AZT, ddI, ddC and PMEA.		Seen in heavily treated patients.	Winters98
T 69 S + SA	ACT→TCT + AGT GCT	MN		?	Y	Confers >4-fold resistance to: AZT, ddI, ddC and PMEA.		Seen in heavily treated patients.	Winters98
T 69 S + SA	ACT→AGT + AGC GCT	MN		?	Y	Confers >4-fold resistance to: AZT, ddI, ddC and PMEA.		Seen in heavily treated patients.	Winters98
T 69 S + SG	ACT→AGT + AGT GGT	MN		?	Y	Confers >4-fold resistance to: AZT, ddI, ddC and PMEA.		Seen in heavily treated patients.	Winters98
T 69 S + SG	ACT→AGT + AGT GGT	NNRTI	ddI (didanosine) + hydroxyurea	?	Y			Seen in one patient.	DeAntoni97

Mutations in HIV RT that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
T 69 S + SS	ACT→TCT + AGC TCT	MN		?	Y	Confers >4-fold resistance to: AZT, ddI, ddC, 3TC and PMEA.		Seen in heavily treated patients.	Winters98
T 69 S + SS	ACT→TCT + AGT TCT	MN		?	Y	Confers >4-fold resistance to: AZT, ddI, ddC, 3TC and PMEA.		Seen in heavily treated patients.	Winters98
T 69 S + SS	ACT→AGT + AGT AGT	NNRTI	ddI (didanosine) + hydroxyurea	?	Y			Seen in one patient.	DeAntoni97
T 69 S + TS	ACT→TCT + ACC TCT	MN		?	Y	Confers >4-fold resistance to: AZT, ddI, ddC, 3TC and PMEA.		Seen in heavily treated patients.	Winters98
T 69 S + VG	ACT→TCT + ACC TCT	MN		N	Y				Larder99
K 70 E	AAA→GAA	NRTI	PMEA (adefovir)	Y	Y	9	3TC (7); PFA: 2-fold hypersusceptibility		Cherrington96, Mulato97
K 70 R	AAA→AGA	NRTI	AZT (zidovudine)	Y	Y			D67N/K70R/T215Y/K219Q: 120-fold	Larder89, Larder91, Kellam92
K 70 S	AAA→AGA	MN	ddI (didanosine) + d4T (stavudine)	?	Y			Seen in one patient on ddC + d4T combination therapy.	
L 74 I	TTA→ATA	NNRTI	HBV 097	Y	?				Kleim96
L 74 V	TTA→GTA	NRTI	1592U89 (abacavir)	Y	N	4		K65R/L74V: 3.6-fold; K65R/L74V/M184V: 10.2-fold	Tisdale97
L 74 V	TTA→GTA	NRTI	ddI (didanosine)	N	Y	5-10	ddC (4)	Can reverse effect of T215Y AZT resistance mutation	StClair91
L 74 V	TTA→GTA	NRTI	DXG	Y	?	4			Mellors96
L 74 V	TTA→GTA	NNRTI	HBV 097	Y	?				Kleim96
V 75 I	GTA→ATA	MN		N	Y	Nil		V75I alone has no effect, but in combination with mutations at 62, 77, 116, 151 causes multi NRTI resistance.	Iversen96, Shirasaka95
V 75 I	GTA→TTA	NNRTI	HBV 097	Y	?			Compensates for negative effect of G190E mutation on RT activity	Kleim96

Mutations in HIV RT that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
V 75 L	GTA → TTA	NNRTI	HBV 097	Y	?	?		Seen in one patient on ddC + d4T combination therapy.	Kleim96
V 75 M	GTA → ATG	MN	ddC (zalcitabine) + d4T (stavudine)	?	Y	Y		Observed with d4T selection in vitro, rarely in patients receiving d4T	Lawrence99
V 75 T	GTA → ACA	NRTI	d4T (stavudine)	Y	Y	7	ddI; ddC; d4C; (-)-FTC	Observed after selection with AZT and PFA; suppresses effects of AZT mutations	Lacey94, Schinazi96
F 77 L	TTC → CTC	MN		N	Y	Nil		F77L alone has no effect, but in combination with mutations at 62, 75, 116, 151 causes multi NRTI resistance.	Iversen96, Shirasaka95
W 88 G	TGG → GGG	PARTI	PFA (foscarnet)	Y	Y	5	Hypersusceptibility to AZT	Observed after selection with AZT and PFA; suppresses effects of AZT mutations	Mellors95, Tachedjian95, Tachedjian96
W 88 S	TGG → TCG	PARTI	PFA (foscarnet)	N	Y	2-4	Wild-type susceptibility to AZT.	Partially suppresses effects of AZT resistance mutations	Mellors95, Tachedjian95, Tachedjian96
E 89 G	GAA → GGA	PARTI	PFA (foscarnet)	Y	N	14		Isolated by screening RT clones for ddGTP resistance	Prasad91
E 89 K	GAA → GGA	PARTI	PFA (foscarnet)	Y	N	> 16		Suppresses effects of AZT resistance mutations	Tachedjian95, Tachedjian96
L 92 I	TTA → ATA	PARTI	PFA (foscarnet)	Y	N	8		Partially suppresses effects of AZT resistance mutations	Tachedjian95, Tachedjian96
A 98 G	GCA → GGA	NNRTI	BI-RG-587 (nevirapine)	N	Y	Y			Richman94
A 98 G	GCA → GGA	NNRTI	L-697,661	N	Y	8			Byrnes93
L 100 I	TTA → ATA	NNRTI	BHAP U-88204E	Y	?	?			Balzarini93d, Vasudevachari92
L 100 I	TTA → ATA	NNRTI	BI-RG-587 (nevirapine)	N	Y	Y			Richman93
L 100 I	TTA → ATA	NNRTI	DMP-266 (efavirenz)	Y	?	8-11		Combinations of mutations needed for high-level resistance; L100I/V108I: 1,000-fold; L100I/V179D/Y181C: 1,000-fold	Young95, Winslow96
L 100 I	TTA → ATA	NNRTI	DMP-266 (efavirenz)	Y	Y	Y			Bachelert00
L 100 I	TTA → ATA	NNRTI	L-697,661	Y	N	2			Byrnes93
L 100 I	TTA → ATA	NNRTI	TIBO R82150	Y	?	> 100		Suppresses effects of AZT resistance mutations	Mellors93, Balzarini93c, Byrnes93a
L 100 I	TTA → ATA	NNRTI	TIBO R82913	Y	?	?		Found in combination with E138K	Larder92

Mutations in HIV RT that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
L 100 I	TTA → ATA	NNRTI	UC-68	Y	?	70			Balzarini95
L 100 I	TTA → ATA	NNRTI	UC-70	Y	?	758			Buckheit95a
L 100 I	TTA → ATA	NNRTI	UC-781	Y	?	20		Activity of UC-781 versus L100I, K103N, V106A, E138K, Y181C and Y188L reduced by 2-, 7-, 1.5-, 1.5-, 5- and 150-fold, respectively, compared to wild type	Balzarini96a, Balzarini96b
L 100 I	TTA → ATA	NNRTI	UC-84	Y	?	> 40, > 33			Buckheit95a, Buckheit95b
K 101 E	AAA → GAA	NNRTI	8-Chloro-TIBO (tivrapiene)	?	Y				Moeremans95
K 101 E	AAA → GAA	NNRTI	ADAMII	Y	?	30		Not selected for in vitro, resistance determined against a panel of mutants. Viruses with the L100I mutation show an enhanced sensitivity to ADAMII.	Cushman98
K 101 E	AAA → GAA	MN	AZT (zidovudine) + BHAP U-8720IE (ateviridine)	?	Y			Seen in one patient on atevirdine + AZT combination therapy. Found in association with K103N.	Demeter98
K 101 E	AAA → GAA	NNRTI	DMP-266 (efavirenz)	Y	?	1,000			Young95
K 101 E	AAA → GAA	NNRTI	DMP-266 (efavirenz)	Y	Y				Bachelier00
K 101 E	AAA → GAA	NNRTI	L-697,661	N	Y	8			Byrnes93
K 101 E	AAA → GAA	NNRTI	UC-10	Y	?	12		K101E/Y181C: 200-fold	Buckheit95a, Buckheit97
K 101 E	AAA → GAA	NNRTI	UC-38	Y	N			K101E/G190E: > 100-fold	Balzarini95a, Balzarini95
K 101 E	AAA → GAA	NNRTI	UC-57	Y	?			K101E/Y181C: 58-fold	Buckheit95a
K 101 E	AAA → GAA	NNRTI	UC-781	Y	?	7	UC040 (18); Nevirapine (15)	V108I/Y181C: 55-fold; K101E/V108I/Y181C: 500-fold.	Buckheit97
K 101 I	AAA → ATA	NNRTI	UC-16	Y	N	10		K101I/G141E: 10-fold	Balzarini95
K 101 Q	AAA → CAA	NNRTI	DMP-266 (efavirenz)	N	Y				Bachelier00
K 101 Q	AAA → CAA	NNRTI	LY-300046 HCl (troviridine)	Y	?			Found in combination with V108I	Zhang95, Vrang93
K 103 E	AAA → GAA	NRTI	BHAP U-8720IE (ateviridine)	?	Y			Found in association with Y181C in one patient on monotherapy. K103E, K103N and Y181C observed with monotherapy.	Demeter98
K 103 N	AAA → AAC	NNRTI	8-Chloro-TIBO (tivrapiene)	?	Y				Moeremans95

Mutations in HIV RT that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
K 103 N	AAA → AAC	NNRTI	ADAMII	Y	?	>28		Not selected for in vitro, resistance determined against a panel of mutants. Viruses with the L100I mutation show an enhanced sensitivity to ADAMII.	Cushman98
K 103 N	AAA → AAC	NNRTI	α -APA (loviride)	Y	Y				Staszewski96a
K 103 N	AAA → AAC	NRTI	BHAP U-87201E (ateviridine)	?	Y			Found in association with Y181C in several patients on monotherapy. Also seen in patients on ATV + AZT combination therapy.	Demeter98
K 103 N	AAA → AAC	NNRTI	BHAP U-90152 (delavirdine)	?	Y			K103N/Y181C seen separately and in combination in patients	Demeter95
K 103 N	AAA → AAC	NNRTI	BI-RG-587 (nevirapine)	N	Y				Richman93
K 103 N	AAA → AAC	NNRTI	DMP-266 (efavirenz)	Y	Y	67		Predominant mutation in vivo	Winslow96
K 103 N	AAA → AAC	NNRTI	DMP-266 (efavirenz)	Y	Y				Bachelert00
K 103 N	AAA → AAC	NRTI	GW 420867X	Y	?				Kleim99
K 103 N	AAA → AAC	NNRTI	I-EBU (emivirine)	Y	?			Predominant mutation in vivo	Seki95
K 103 N	AAA → AAC	NNRTI	L-697,593	Y	?	20		K103N/Y181C: > 1,000-fold	Nunberg91
K 103 N	AAA → AAC	NNRTI	L-697,661	Y	Y	8		K103N and Y181C most common with monotherapy	Byrnes93, Saag93
K 103 N	AAA → AAC	NNRTI	TIBO R82913	Y	?	> 100		K103N/Y181C: > 1,000-fold	Balzarini93d
K 103 N	AAA → AAC	NNRTI	UC-10	Y	N	5			Balzarini95
K 103 N	AAA → AAC	NNRTI	UC-81	Y	?				Balzarini95, Yang97
K 103 Q	AAA → CAA	NNRTI	L-697,661	N	Y	8			Saag93
K 103 R	AAA → AGA	NNRTI	I-EBU (emivirine)	Y	Y				BorritoEsoda97
K 103 R	AAA → AGA	NNRTI	LY-300046 HCl (troviridine)	Y	?			K103R/V179D: 500-fold; Found in combination with V179D or Y181C	Zhang95, Vrang93
K 103 T	AAA → ACA	NNRTI	BHAP U-90152 (delavirdine)	?	Y				Demeter95
K 103 T	AAA → ACA	NRTI	S-1153	Y	?				Fujiwara98
K 103 T	AAA → ACA	NNRTI	UC-42	Y	N	100			Balzarini95

Mutations in HIV RT that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
V 106 A	GTA→GCA	NNRTI	ADAMII	Y	?	7.13		Not selected for in vitro, resistance determined against a panel of mutants. Viruses with the L100I mutation show an enhanced sensitivity to ADAMII.	Cushman98
V 106 A	GTA→GCA	NNRTI	BHAP U-88204E	Y	?				Vasudevachari92
V 106 A	GTA→GCA	NNRTI	BI-RG-587 (nevirapine)	Y	Y	100		No effect on AZT resistance	Richman94, Larder92, Richman93, Balzarini93d
V 106 A	GTA→GCA	NNRTI	E-EBU-dM	Y	?				Balzarini93
V 106 A	GTA→GCA	NRTI	GW 420867X	Y	?			V106A/Y181C: 400-fold resistance	Klein99
V 106 A	GTA→GCA	NRTI	S-1153	Y	?	4.5		V106A + F227L: 387-fold	Fujiwara98
V 106 A	GTA→GCA	NNRTI	S-2720 (Quinoxaline)	Y	?			P225H follows V106A. Also seen with L101I and Y181C. Double and triple mutants highly resistant to other NNRTI's, including MKC442	Pelermans97
V 106 A	GTA→GCA	NNRTI	TIBO R82913	Y	?	100			Larder92
V 106 A	GTA→GCA	NNRTI	UC-69	Y	?			V106A/V181C: 166-fold	Buckheit95a
V 106 A	GTA→GCA	NNRTI	UC-82	Y	?	13		Activity of UC-82 versus L100I, K103N, V106A, E138K, Y181C and Y188L reduced by 2-, 6-, 1.5-, 2-, 4- and 200-fold, respectively, compared to wild type	Balzarini96b, Balzarini96a
V 106 I	GTA→ATA	NNRTI	HBV 097					Appears under lowered drug concentration selection	Klein97
V 106 M	GTG→ATG	NNRTI	BHAP U-90152 (delavirdine)	Y			1000		Brenner02
V 106 M	GTG→ATG	NNRTI	BI-RG-587 (nevirapine)	Y			1000		Brenner02
V 106 M	GTG→ATG	NNRTI	DMP-266 (efavirenze)	Y	Y			selected in vitro under efavirenze pressure in Clade C virus. Also developed in 3/6 efavirenze-treated patients with Clade C infection.	Brenner02
V 108 I	GTA→ATA	NNRTI	ADAMII	Y	?	6.74		Not selected for in vitro, resistance determined against a panel of mutants. Viruses with the L100I mutation show an enhanced sensitivity to ADAMII.	Cushman98

Mutations in HIV RT that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
V 108 I	GTA → ATA	NNRTI	α-APA (loviride)	Y	?				Staszewski96a
V 108 I	GTA → ATA	NNRTI	BI-RG-587 (nevirapine)	N	Y				Richman93
V 108 I	GTA → ATA	NNRTI	DMP-266 (efavirenz)	Y	?			L100I/V108I: 1,000-fold	Winslow96
V 108 I	GTA → ATA	NNRTI	DMP-266 (efavirenz)	Y	Y				Bachelier00
V 108 I	GTA → GCA	NNRTI	I-EBU (emivirine)	Y	?				Seki95
V 108 I	GTA → GCA	NNRTI	L-697.661	Y	Y	4			Byrnes93
V 108 I	GTA → ATA	NNRTI	LY-300046 HCl (troviridine)	Y	?			Found in combination with K101Q	Zhang95
V 108 I	GTT → GAT	NNRTI	TIBO R82913	N	Y	> 100	R82150 (> 100)		Vandamme94a
V 108 I	GTA → ATA	NNRTI	UC-781	Y	?		V108I/Y181C: 55 fold. K101E/V108I/Y181C: 500 fold.		Buckheit97
Y 115 F	TAT → TTT	NRTI	I592U89 (abacavir)	Y	N	2		K65R/L74V and/or Y115F with M184V: 10 fold; L74V/Y115F/M184V: 11-fold	Tisdale97
F 116 Y	TTT → TAT	MN		N	Y	Nil		F116Y alone has no effect, but in combination with mutations at 62, 75, 77, 151 causes multi NRTI resistance.	Iversen96, Shirasaka95
V 118 I	GTT → ATT	MN	AZT (zidovudine) + 3TC (lamivudine)	N	Y	2.0		Confers moderate levels of resistance to 3TC (4 to 50-fold) when present in an AZT-resistant genetic background without diminishing AZT resistance.	Hertogs00
P 119 S	CCC → TCC	NRTI	F-ddA (lodenosine)	Y	?	4.6		Found with V179D and/or L214F, which are possibly compensatory	Tamaka97
I 135 L	ATA → AAAA	NNRTI		N	Y	Nil		Mutation identified by logistic regression analysis. I135L/L283I: 5.0-fold	Brown00
I 135 M	ATA → ATG	NNRTI		N	Y	Nil		Delavirdine resistance, 4.2-fold Nevirapine resistance, 4.1-fold Efavirenze resistance.	
								Mutation identified by logistic regression analysis. I135L/L283I: 4.0-fold	Brown00
								Delavirdine resistance, 4.5-fold Nevirapine resistance, 3.2-fold Efavirenze resistance.	

Mutations in HIV RT that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
I 135 T	ATA→ACA	NNRTI		N	Y	Nil		Mutation identified by logistic regression analysis. I135L/L283I: 3.4-fold Nevirapine resistance.	Brown00
E 138 A	GAG→GCG	NNRTI	TSAO	N	Y			Mutation reducing susceptibility to TSAO in TSAO therapy naive patients.	VanLaethem00
E 138 K	GAG→AAG	NNRTI	I-EBU (emivirine)	Y	N			Obtained in the concomitant presence of low 3TC concentrations	Balzarini96c
E 138 K	GAG→AAG	NNRTI	TIBO R82913	Y	?			Found in combination with L100I	Balzarini93c
E 138 K	GAG→AAG	NNRTI	TSAO	Y	?	> 100		E138A (GAG to GCG) in TSAO-naive patients confers TSAO viral resistance	Balzarini93a, Balzarini93b, Vandamme96
E 138 K	GAG→AAG	NNRTI	UC-82	Y	?	5		Activity of UC-82 versus L100I, K103N, V106A, E138K, Y181C and Y188L reduced by 2-, 6-, 1.5-, 2-, 4- and 200-fold, respectively, compared to wild type	Balzarini96b, Balzarini96a
E 138 K	GAG→AAG	NNRTI	UC-84	Y	?	> 100	TSAOs		Balzarini95, Balzarini95b
T 139 I	ACA→ATA	NNRTI	ADAMII	Y	?	38		Not selected for in vitro, resistance determined against a panel of mutants. Viruses with the L100I mutation show an enhanced sensitivity to ADAMII.	Cushman98
T 139 I	ACA→ATA	NNRTI	Calanolide A	Y	?	> 70			Buckheit95c
G 141 E	GGG→GAG	NNRTI	UC-16	Y	N				Balzarini95
Q 151 M	CAG→ATG	MN		N	Y	AZT: 10; ddI/ddC: 5		K101I/G141E: 10-fold Pivotal multi nucleoside RTI resistance mutation (first to occur), found in association with combinations of four other mutations: A62V/V75I/F77L/F116Y/Q151M: AZT 190-fold; ddI 50-fold; ddC 20-fold; d4T > 10-fold	Iversen96, Shirasaka95, Schmit96
S 156 A	TCA→GCA	PARTI	PFA (foscarnet)	Y	N	4.5			Tachedjian95
P 157 S	CCA→TCA	NRTI	3TC (lamivudine)			5		Mutation increases susceptibility to AZT and PMPA	Smith99

Mutations in HIV RT that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
Q 161 L	CAA → CTA	PARTI	PFA (foscarnet)	Y	Y	5		Q161L/H208Y: 9-fold; Q161L/H208Y suppresses effects of AZT mutations	Mellors95
V 179 D	GTT → GAT	NNRTI	ADAMII	Y	?	28		Not selected for in vitro, resistance determined against a panel of mutants. Viruses with the L100I mutation show an enhanced sensitivity to ADAMII.	Cushman98
V 179 D	GTT → GAT	NNRTI	DMP-266 (efavirenz)	Y	?			L100I/V179D/Y181C: 1,000-fold	Winslow96
V 179 D	GTT → GAT	NNRTI	L-697,661	N	Y	4			Byrnes93
V 179 D	GTT → GAT	NNRTI	LY-300046 HCl (troviridine)	Y	?			Found in combination with K103R or Y181C; V179D/Y181C: > 1,000-fold	Zhang95
V 179 D	GTT → GAT	NNRTI	QM96521	Y	?	10	Other TDD derivative: 15-140-fold; 8-chloro-TIBO: 10-fold		Witvrouw98
V 179 D	GTT → GAT	NNRTI	TIBO R82913	N	Y	20	R82150 (20)		Vandamme94
V 179 D	GTT → GAT	NNRTI	UC-10	Y	?	16			Balzarini95, Balzarini96a
V 179 E	GTT → GAG	NNRTI	L-697,661	N	Y	8			Byrnes93
Y 181 C	TAT → TGT	NNRTI	1737	Y	?	20		Y181C also confers resistance to numerous other tetrahydronaphthalene derivatives.	Hara97
Y 181 C	TAT → TGT	NNRTI	ADAMII	Y	?	>28		Not selected for in vitro, resistance determined against a panel of mutants. Viruses with the L100I mutation show an enhanced sensitivity to ADAMII.	Cushman98
Y 181 C	TAT → TGT	NNRTI	α-APA (loviride)	Y	?				deBethune93
Y 181 C	TAT → TGT	NNRTI	α-APA (loviride)	?	Y				Staszewski96
Y 181 C	TAT → TGT	NNRTI	BHAP U-87201E (ateviridine)	N	Y			K103E, K103N and Y181C observed with monotherapy	Demeter95, Demeter98
Y 181 C	TAT → TGT	NNRTI	BHAP U-88204E	Y	?				Vasudevachari92
Y 181 C	TAT → TGT	NNRTI	BHAP U-90152 (delavirdine)	?	Y			K103N/Y181C seen separately and in combination in vivo	Demeter95
Y 181 C	TAT → TGT	NNRTI	BI-RG-587 (nevirapine)	Y	Y	> 100	Other NNRTIs	Can suppress effects of AZT mutations	Richman94, Richman91, Mellors92

Mutations in HIV RT that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
Y 181 C	TAT→TGT	NNRTI	BM+51,0836	Y	?	?			Maass93
Y 181 C	TAT→TGT	NNRTI	DMP-266 (efavirenz)	Y	?	4		L100I/V179D/Y181C: 1,000-fold; uncommon in vivo	Winslow96, Young95
Y 181 C	TAT→TGT	NNRTI	E-BPTU	Y	?	160	Other NNRTIs		Buckheit95c
Y 181 C	TAT→TGT	NNRTI	E-EBU	Y	?	?			Balzarini93
Y 181 C	TAT→TGT	NNRTI	E-EPSeU	Y	?	> 50		Y188C confers greater resistance than Y181C	Nguyen94
Y 181 C	TAT→TGT	NNRTI	E-EPU	Y	?	> 95		Y188C confers greater resistance than Y181C	Nguyen94
Y 181 C	TAT→TGT	NRTI	GW 420867X	Y	?	?		V106A/Y181C: 400-fold resistance	Kleim99
Y 181 C	TAT→TGT	NNRTI	I-EBU (emivirine)	?	Y	?			BorotoEsoda97
Y 181 C	TAT→TGT	NNRTI	L-697,593	Y	?	> 100		K103N/Y181C: > 1,000-fold	Nunberg91
Y 181 C	TAT→TGT	NNRTI	L-697,661	Y	Y	> 30		K103N and Y181C most common with monotherapy	Byrnes93, Saag93
Y 181 C	TAT→TGT	NNRTI	LY-300046 HCl (trovirdine)	Y	?	?	Nevirapine; 9-chloro-TIBO	V179D/Y181C: > 1,000-fold; Found in combination with K103R or V179D	Zhang95, Vrang93
Y 181 C	TAT→TGT	NNRTI	TIBO R82913	Y	?	> 100		K103N/Y181C: > 1,000-fold	Larder92
Y 181 C	TAT→TGT	NNRTI	UC-10	Y	?	6		K101E/Y181C: 200-fold	Buckheit95a
Y 181 C	TAT→TGT	NNRTI	UC-32	Y	?	38			Buckheit95a
Y 181 C	TAT→TGT	NNRTI	UC-38	Y	?	8-149	Other NNRTIs	K101E/Y181C: 58-fold	Buckheit95a
Y 181 C	TAT→TGT	NNRTI	UC-57	Y	?	?			Buckheit95a
Y 181 C	TAT→TGT	NNRTI	UC-68	Y	?	5			Buckheit95a
Y 181 C	TAT→TGT	NNRTI	UC-69	Y	?	?		V106A/V181C: 166-fold	Buckheit95a
Y 181 C	TAT→TGT	NNRTI	UC-781	Y	?	13		V108/Y181C: 55 fold; K101E/V108I/Y181C: 500 fold. 42	Balzarini98, Buckheit97
Y 181 C	TAT→TGT	NNRTI	UC-80	Y	?	18			Buckheit95a
Y 181 C	TAT→TGT	NNRTI	UC-81	Y	?	53			Balzarini95, Yang97
Y 181 C	TAT→TGT	NNRTI	UC-84	Y	?	> 118			Buckheit95a

Mutations in HIV RT that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
Y 181 I	TGT→ATT	NNRTI	BHAP U-88204E	Y	Y			Appeared after treatment of Y181C-mutated virus with BHAP; high-level resistance to BHAP, nevirapine and TIBO; observed in one nevirapine-treated patient	Balzarini94
Y 181 I	TGT→ATT	NNRTI	BI-RG-587 (nevirapine)	N	Y	High-level		Observed in one patient	Shaw94
Y 181 I	TAT→ATT	NNRTI	I-EBU (emivirine)	Y	N	1,000			Balzarini96c
M 184 I	ATG→ATA	NRTI	3TC (lamivudine)	Y	Y			M184V and M184I can suppress effects of AZT resistance mutations	Schinazi93, Tisdale93, Gao93
M 184 I	ATG→ATA	NNRTI	QYL-685	Y	?	9	QYL-609	Additional passage of virus did not select M184V	Yoshimura99a
M 184 T	ATG→ACG	NRTI	3TC (lamivudine)	Y	?			Reduced replication capacity and RT activity	Larder95, Keulen96
M 184 V	ATG→GTG	NRTI	I592U89 (abacavir)	Y	N	3		K65R/L74V and/or Y115F with M184V: 10-fold; K65R/M184V: 8-fold; L74V/M184V: 9-fold resistance; L74V/Y115F/M184V: 11-fold	Tisdale97
M 184 V	ATG→GTG	NRTI	3TC (lamivudine)	Y	Y	>100	ddI; ddC; (-)-FTC	M184V and M184I can suppress effects of AZT resistance mutations; GTA seen in cell culture	Schinazi93, Tisdale93, Gao93
M 184 V	ATG→GTG	NRTI	ddC (zalcitabine)	Y	Y	2-5		Rarely observed in patients receiving ddI	Gu92
M 184 V	ATG→GTG	NRTI	ddI (didanosine)	Y	Y	2-5			Gu92
M 184 V	ATG→GTG	NRTI	dOTC	Y	?			K65R/M184V: 4.2-fold.	Rando99
M 184 V	ATG→GTG	NRTI	(-)dOTC	Y	?	nil			Rando99
M 184 V	ATG→GTG	NRTI	(+)dOTC	Y	?				Rando99
M 184 V	ATG→GTG	NNRTI	(-)dOTFC	Y	?	13			Richard00
M 184 V	ATG→GTG	NNRTI	(-)dOTFC	Y	?	>15.0			Richard00
M 184 V	ATG→GTG	NRTI	(-)FTC (emtricitabine)	Y	?	>100		M184V can suppress effects of AZT mutations	Schinazi93, Tisdale93
M 184 V	ATG→GTG	NRTI	L-FddC	Y	?	>100			Schinazi95

Mutations in HIV RT that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
Y 188 C	TAT→TGT	NNRTI	ADAMIII	Y	?	6.07		Not selected for in vitro, resistance determined against a panel of mutants. Viruses with the L100I mutation show an enhanced sensitivity to ADAMIII.	Cushman98
Y 188 C	TAT→TGT	NNRTI	BI-RG-587 (nevirapine)	N	Y				Richman93
Y 188 C	TAT→TGT	NNRTI	E-EPSeU	Y	?	> 250		Y188C is the predominant mutation for E-EPSeU; Y188C confers greater resistance than Y181C	Nguyen94
Y 188 C	TAT→TGT	NNRTI	E-EPU	Y	?	> 250		Y188C confers greater resistance than Y181C	Nguyen94
Y 188 C	TAT→TGT	NNRTI	HEPT	Y	?				Balzarini93
Y 188 H	TAT→CAT	NNRTI	ADAMIII	Y	?	>128		Not selected for in vitro, resistance determined against a panel of mutants. Viruses with the L100I mutation show an enhanced sensitivity to ADAMIII.	Cushman98
Y 188 H	TAT→CAT	MN	AZT (zidovudine) + BHAP U-87201E (atevirdine)	?	Y			Found in two patients on atevirdine + AZT combination therapy.	Demeter98
Y 188 H	TAT→CAT	NNRTI	DMP-266 (efavirenz)	N	Y				Bachelier00
Y 188 H	TAT→CAT	NNRTI	TIBO R82913	Y	?				Balzarini93c
Y 188 H/L	TAT→CAT/CTT	NNRTI	α-APA (loviride)	?	Y				Staszewski96
Y 188 L	TAT→TTA	NNRTI	DMP-266 (efavirenz)	Y	?	1,000			Winslow96
Y 188 L	TAT	NNRTI	DMP-266 (efavirenz)		Y				Bachelier00
Y 188 L	TAT→TTA	NNRTI	TIBO R82913	N	Y				Vandamme94
V 189 I	GTA→ATA	NNRTI	BHAP U-90152 (delavirdine)	Y	N			Compensatory mutation restoring replication competence to W229Y mutant. I63M/V189I/W229Y/E396G mutant 2.5-fold resistant to delavirdine with respect to WT	Pelemans01

Mutations in HIV RT that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
V 189 I	GTA → ATA	NNRTI	BI-RG-587 (nevirapine)	Y	N			Compensatory mutation restoring replication competence to W229Y mutant. I63M/V189I/W229Y/E396G mutant 2.3-fold resistant to nevirapine with respect to WT	Pelemans01
V 189 I	GTA → ATA	NNRTI	GW 420867X	Y	N			Compensatory mutation restoring replication competence to W229Y mutant. I63M/V189I/W229Y/E396G mutant 8-fold resistant to GW 420867X with respect to WT	Pelemans01
V 189 I	GTA → ATA	NNRTI	HBV 097	Y	?	2	Other NNRTIs (2-6)		Kleim96
V 189 I	GTA → ATA	NNRTI	HBV 097	Y	N			Compensatory mutation restoring replication competence to W229Y mutant. I63M/V189I/W229Y/E396G mutant 5-fold resistant to HBV 097 with respect to WT	Pelemans01
V 189 I	GTA → ATA	NNRTI	I-EBU (emivirine)	Y	N			Compensatory mutation restoring replication competence to W229Y mutant. I63M/V189I/W229Y/E396G mutant 7-fold resistant to emivirine with respect to WT	Pelemans01
V 189 I	GTA → ATA	NNRTI	UC-781	Y	N			Compensatory mutation restoring replication competence to W229Y mutant. I63M/V189I/W229Y/E396G mutant 2.5-fold resistant to UC-781 with respect to WT	Pelemans01
G 190 A	GGA → GCA	NNRTI	α-APA (loviride)	?	Y				Moeremans95
G 190 A	GGA → GCA	NNRTI	BI-RG-587 (nevirapine)	N	Y				Richman94
G 190 A	GGA → GCA	NNRTI	DMP-266 (efavirenz)	N	Y				Bachelor00

Mutations in HIV RT that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
G 190 E	GGA → GAA	NNRTI	AAP-BHAP (U-104489)	Y	?	>100		TI39I/G190E/T200A/L214F: >100. Additional mutations possibly restore the replication capacity of the G190E mutant	Olmsted96
G 190 E	GGA → GAA	NNRTI	DMP-266 (efavirenz)	N	Y				Bachelert00
G 190 E	GGA → GAA	NRTI	GW 420867X	Y	?				Kleim99
G 190 E	GGA → GAA	NNRTI	HBV 097	Y	?		Other NNRTIs	Reduces enzymatic activity of RT and viral replication competency	Kleim95
G 190 E	GGA → GAA	NNRTI	S-2720 (Quinoxaline)	Y	?				Kleim93
G 190 E	GGA → GAA	NNRTI	UC-38	Y	N			K101E/G190E: > 100-fold; cross resistance to: TSAO-m3T, Nev, TIBO R82913, BHAP U88204; susceptible to L697,661	Balzarni95a
G 190 Q	GGA → CAA	NNRTI	HBV 097	Y	?		Other NNRTIs	Appears exclusively in connection with V179D	Kleim96
G 190 S	GGA → TCA	NNRTI	DMP-266 (efavirenz)	N	Y				Bachelert00
G 190 T	GGA → ACA	NNRTI	HBV 097	Y	?			Appears during selection with low drug concentrations.	Kleim97
H 208 Y	CAT → TAT	MN	AZT (zidovudine) + 3TC (lamivudine)	?	Y			Polymorphism facilitating AZT+3TC dual resistance	Kemp98
H 208 Y	CAT → TAT	PARTI	PFA (foscarnet)	Y	Y	2		Q161L/H208Y: 9-fold; increased susceptibility to AZT 100-fold, nevirapine (20-fold) and TIBO R82150 (30-fold); Q161L/H208Y suppresses effects of AZT mutations	Mellors95
L 210 W	TTG → TGG	NRTI	AZT (zidovudine)	Y	Y			210W/215Y: 42-fold 41L/210W/215Y; 49-fold 41L/67N/70R/210W/215Y: 366-fold Mutation arises after prolonged AZT therapy.	Gurusinghe95, Harrigan96, Hooker96
R 211 K	AGG → AAG	MN	AZT (zidovudine) + 3TC (lamivudine)	?	Y			Polymorphism facilitating AZT+3TC dual resistance in association with M184V and other AZT resistance mutations.	Kemp98

Mutations in HIV RT that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
F 214 L	TTT→CTT	MN	AZT (zidovudine) + 3TC (lamivudine)	?	Y			Polymorphism facilitating AZT+3TC dual resistance in association with M184V and other AZT resistance mutations.	Stuyver97
T 215 F	ACC→TTC	NRTI	AZT (zidovudine)	?	Y			K67N/K70R/T215Y/K219Q: 120-fold	Larder89, Larder91, Kellam92
T 215 Y	ACC→TAC	NRTI	AZT (zidovudine)	Y	Y			M41L/T215Y: 60–70-fold; K67N/K70R/T215Y/K219Q: 120-fold. Effect of T215Y is reversed by a ddl mutation (L74V), NNRTI mutations (L100I;Y181C) or (-)-FTC/3TC mutations (M184I/V)	Larder89, Larder91, Kellam92
Y 215 C	TTC→TGC	NRTI	ddC (zalcitabine)	N	Y	4		Arises on background of T215Y AZT resistance	Slade93
K 219 E	AAA→GAA	NRTI	AZT (zidovudine)	Y	N				Larder89, Larder91, Kellam92
K 219 Q	AAA→CAA	NRTI	AZT (zidovudine)	?	Y			K67N/K70R/T215Y/K219Q: 120-fold	Larder89, Larder91, Kellam92
K 219 R	AAA→AGA	MN	3TC (lamivudine) + d4T (stavudine)	?	Y			Seen in two patient on 3TC + d4T combination therapy.	Lawrence99
K 219 R	AAA→AGA	MN	AZT (zidovudine) + 3TC (lamivudine)	?	Y			Seen in two patient on AZT + 3TC combination therapy.	Lawrence99
K 219 W	AAA→TGG	MN	ddC (zalcitabine) + d4T (stavudine)	?	Y			Seen in one patient on ddC + d4T combination therapy.	Lawrence99
P 225 H	CCT→CAT	NNRTI	DMP-266 (efavirenz)	N	Y				Bachelier00
P 225 H	CCT→CAT	NNRTI	S-2720 (Quinoxaline)	Y	?	4.0	MKC-442 (5.7); HBY-097 (4.0); UC-781 (3.7)	P225H follows V106A. Also seen with L101I and Y181C. Double and triple mutants highly resistant to other NNRTI's, including MKC442. The presence of P225H in a V106A background restores sensitivity to BHAP U-90152.	Pelemans97, Pelemans98
F 227 L	TTA→CTC	NRTI	S-1153	Y	?	nil		V106A + F227L: 387-fold. This mutation confers hypersensitivity to delavirdine.	Fujitwara98

Mutations in HIV RT that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
F 227 L	TTA → CTC	NNRTI	UC-781	Y	?			V106A/F227L: 10-fold. Found with V106A, K101I, Y181C and L100I. Appears in a V106A background following dose-escalating UC-781 treatment.	Balzarini98
W 229 Y	TGG → TAC	NNRTI	BHAP U-90152 (delavirdine)	Y	N			Compensatory mutation restoring replication competence to W229Y mutant. I63M/V189I/W229Y/E396G mutant 2.5-fold resistant to delavirdine with respect to WT	Pelemans01
W 229 Y	TGG → TAC	NNRTI	BI-RG-587 (nevirapine)	Y	N			Compensatory mutation restoring replication competence to W229Y mutant. I63M/V189I/W229Y/E396G mutant 23-fold resistant to nevirapine with respect to WT	Pelemans01
W 229 Y	TGG → TAC	NNRTI	GW 420867X	Y	N			Compensatory mutation restoring replication competence to W229Y mutant. I63M/V189I/W229Y/E396G mutant 8-fold resistant to GW 420867X with respect to WT	Pelemans01
W 229 Y	TGG → TAC	NNRTI	HBV 097	Y	N			Compensatory mutation restoring replication competence to W229Y mutant. I63M/V189I/W229Y/E396G mutant 5-fold resistant to HBV 097 with respect to WT	Pelemans01
W 229 Y	TGG → TAC	NNRTI	I-EBU (emivirine)	Y	N			Compensatory mutation restoring replication competence to W229Y mutant. I63M/V189I/W229Y/E396G mutant 7-fold resistant to emivirine with respect to WT	Pelemans01

Mutations in HIV RT that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
W 229 Y	TGG→TAC	NNRTI	UC-781	Y	N			Compensatory mutation restoring replication competence to W229Y mutant. I63M/V189I/W229Y/E396G mutant 2.5-fold resistant to UC-781 with respect to WT	Pelemans01
V 233 E	GAA→GTA	NNRTI	AZT (zidovudine) + BHAP U-87201E (atevirdine)	N	Y			Seen in 1 patient. K101E, Y188H and K238T also seen in patients on ATV/AZT combination therapy.	Demeter98
L 234 I	CTC→ATC	NRTI	S-1153	Y	?	22		This mutation confers hypersensitivity to Loviride.	Fujiwara98
P 236 L	CCT→CTT	NNRTI	BHAP U-87201E (atevirdine)	Y	N				Dueweke93
P 236 L	CCT→CTT	NNRTI	BHAP U-90152 (delavirdine)	Y	Y			Sensitizes RT 10-fold to nevirapine, TIBO R829I3 and L-697,661	Dueweke93
P 236 L	CCT→CTT	NNRTI	HEPT	Y	?				Buckheit95c
K 238 T	AAA→ACA	MN	AZT (zidovudine) + BHAP U-87201E (atevirdine)	N	Y			Seen in 1 patient. K101E, K103N, Y188H, and V233E also observed with ATV/AZT combination therapy.	Demeter98
K 238 T	AAA→ACA	MN	AZT (zidovudine) + BHAP U-87201E (atevirdine)	N	Y			Seen in 1 patient. K101E, K103N, Y188H and E233V also seen in patients on ATV/AZT combination therapy.	Demeter98
L 283 I	CTT→ACT	NNRTI		N	Y	Nil		Mutation identified by logistic regression analysis. Confers resistance to Delavirdine, Nevirapine, and Efavirenze in conjunction with mutations at codon 135.	Brown00
G 333 D	GGC→GAC	MN	AZT (zidovudine) + 3TC (lamivudine)	Y	Y			Facilitates dual resistance to AZT+3TC in association with M184V and standard AZT resistance mutations.	Kemp98
G 333 D	GGC→GAC	MN	AZT (zidovudine) + 3TC (lamivudine) + 1592U89 (abacavir)	?	Y				Caride00
G 333 E	GGC→GAG	MN	AZT (zidovudine) + 3TC (lamivudine)	Y	Y			Facilitates dual resistance to AZT+3TC in association with M184V and standard AZT resistance mutations.	Kemp98

Mutations in HIV RT that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
G 333 E	GGC→GAG	MN	AZT (zidovudine) + 3TC (lamivudine) + 1592U89 (abacavir)	?	Y				Caride00
T 386 I	ACT→ATT	MN	AZT (zidovudine) + 3TC (lamivudine) + 1592U89 (abacavir)	?	Y			Abrogates M184V suppression of L210W and L210W/G333D/E	Caride00
E 396 G	GAA→GGA	NNRTI	BHAP U-90152 (delavirdine)	Y	N			Compensatory mutation restoring replication competence to W229Y mutant. I63M/V189I/W229Y/E396G mutant 2.5-fold resistant to delavirdine with respect to WT	Pelemans01
E 396 G	GAA→GGA	NNRTI	BI-RG-587 (nevirapine)	Y	N			Compensatory mutation restoring replication competence to W229Y mutant. I63M/V189I/W229Y/E396G mutant 23-fold resistant to nevirapine with respect to WT	Pelemans01
E 396 G	GAA→GGA	NNRTI	GW 420867X	Y	N			Compensatory mutation restoring replication competence to W229Y mutant. I63M/V189I/W229Y/E396G mutant 8-fold resistant to GW 420867X with respect to WT	Pelemans01
E 396 G	GAA→GGA	NNRTI	HBV 097	Y	N			Compensatory mutation restoring replication competence to W229Y mutant. I63M/V189I/W229Y/E396G mutant 5-fold resistant to HBV 097 with respect to WT	Pelemans01
E 396 G	GAA→GGA	NNRTI	I-EBU (emivirine)	Y	N			Compensatory mutation restoring replication competence to W229Y mutant. I63M/V189I/W229Y/E396G mutant 7-fold resistant to emivirine with respect to WT	Pelemans01

Mutations in HIV RT that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
E 396 G	GAA → GGA	NNRTI	UC-781	Y	N			Compensatory mutation restoring replication competence to W229Y mutant. I63M/V189I/W229Y/E396G mutant 2.5-fold resistant to UC-781 with respect to WT	Pelemans01

Mutations in HIV PROTEASE that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
R 8 K	CGA → AAA	PI	A-77003	Y	?	10		R8K/M46I/G48V: 20-fold	Ho94, Tisdale94
R 8 Q	CGA → CAA	PI	A-77003	Y	?	10		M46I improves replication competency of R8Q mutant	Ho94, Kaplan94
L 10 F	CTC → TTC	PI	ABT-378 (lopinavir)	Y	?		Passage 17 virus: ritonavir, 21-fold; saquinavir, 4-fold	I84V/L10F/M46I: 4 fold, I84V/L10F/M46I/T91S: 12 fold, I84V/L10F/M46I/T91S/V32I/I47V: 46 fold, Passage 17 virus: I84V/L10F/M46I/T91S/V32I/I47V/V47A/G16E/H69Y: 338 fold (in presence of p7/p1 (AN/F to VN/F) cleavage-site mutation and p1/p6 (F/L to F/F) cleavage-site mutation).	Carrillo98
L 10 F	CTC → TTC	PI	ABT-378 (lopinavir)	N	Y			In conjunction with other mutations, susceptibility to lopinavir was reduced by mutations at positions 82,54,10,63,71 and 84 by 4–10-fold, K20M/R by >20-fold and F53L by >40-fold. IC50 of lopinavir against isolates with 0–3 mutations was 0.8-fold higher	Kempf01
L 10 F	CTC → TTC	PI	BILA 2185 BS	Y	?		BILA 1906 BS (360)	L10F/L23I/V32I/M46I/I47V/I54M/A71V/I84V: 1,500-fold. Associated Gag mutations: p1/p6 cleavage site (L to F (CTT to TTT at P1'); p7/p1 cleavage site (Q to R (CAG to CGG) at P3', A to V (GCT to CTT) at P2').	Croteau97
L 10 F	CTC → TTC	PI	BMS 232632	Y	?			V32L/L33F/M46I/A71V/I84V/N88S: 183-fold., L10Y,F/I50L/LG3P/A71V/N88S: 93-fold., V32I/M46I/I84V/L89M: 96-fold.	Gong00
L 10 F	CTC → TTC	PI	DMP 450	Y	?			Probably compensatory	Otto95, Winslow95
L 10 F	CTC → TTC	PI	DMP-323	Y	?			L10F/V82A: 2-fold; L10F/K45I/I84V: 50-fold	King95

Mutations in HIV PROTEASE that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
L 10 F	CTC→TTC	PI	JE-2147	Y	?		KNI-272: 7-fold; Ritonavir: 9-fold	L10F/I47V/I84V: 19-fold. L10F/M46I/I47V/I84V: 28-fold. >50 passages required for isolation of resistant virus.	Yoshimura99
L 10 F	CTC→TTC	PI	SC-55389A	Y	?	2.8	Not SC-52151	N88S/L10F: 25-fold	Potts94, Pillay96, Smidt97
L 10 F	CTC→TTC	PI	VB 11,328	Y	?			L10F/I84V: 8-fold	Partaledis95
L 10 F	CTC→TTC	PI	VX-478 (amprenavir)	Y	?				Tisdale96
L 10 I	CTC→ATC	PI	ABT-378 (lopinavir)	N	Y			In conjunction with other mutations, susceptibility to lopinavir was reduced by mutations at positions 82,54,10,63,71 and 84 by 4–10-fold, K20M/R by >20-fold and F53L by >40-fold. IC50 of lopinavir against isolates with 0–3 mutations was 0.8-fold higher	Kempf01
L 10 I	CTC→ATC	PI	MK-639 (indinavir)	?	Y			Found in combination with G48V in vivo.	Condra96
L 10 I	CTC→ATC	PI	Ro 31–8959 (saquinavir)		Y				Schapiro96
L 10 R	CTC→CGC	PI	ABT-378 (lopinavir)	N	Y			In conjunction with other mutations, susceptibility to lopinavir was reduced by mutations at positions 82,54,10,63,71 and 84 by 4–10-fold, K20M/R by >20-fold and F53L by >40-fold. IC50 of lopinavir against isolates with 0–3 mutations was 0.8-fold higher	Kempf01
L 10 R	CTC→CGC	PI	MK-639 (indinavir)	N	Y		XM-323 (15)	L10R/M46I/L63P/V82T: 4-fold; L10R/M46I/L63P/V82T/I84V: 8-fold	Condra96, Condra95
L 10 V	CTC→GTC	PI	ABT-378 (lopinavir)	N	Y			In conjunction with other mutations, susceptibility to lopinavir was reduced by mutations at positions 82,54,10,63,71 and 84 by 4–10-fold, K20M/R by >20-fold and F53L by >40-fold. IC50 of lopinavir against isolates with 0–3 mutations was 0.8-fold higher	Kempf01

Mutations in HIV PROTEASE that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
L 10 V	CTC → GTC	PI	MK-639 (indinavir)	?	Y		A-80987 (4)	V32L/L33F/M46I/A71V/I84V/N88S: 183-fold., L10Y,F/I50L/LG3P/A71V/N88S: 93-fold., V32I/M46I/I84V/L89M: 96-fold.	Condra96, Condra95 Gong00
L 10 Y	CTC → TAC	PI	BMS 232632	Y	?				
I 15 V	CTC → TAC	PI	PNU-140690 (tipranavir)	?	Y			Passage 17 virus: I84V/L10F/M46I/T91S/V32I/I47V/V47A/G16E/H69Y: 338 fold (in presence of p7/p1 (AN/F to VN/F) cleavage-site mutation and p1/p6 (F/L to F/F) cleavage-site mutation).	Rusconi00
G 16 E	GGG → GAG	PI	ABT-378 (lopinavir)	Y	?		Passage 17 virus: ritonavir, 21-fold; saquinavir, 4-fold	In conjunction with other mutations, susceptibility to lopinavir was reduced by mutations at positions 82,54,10,63,71 and 84 by 4–10-fold, K20M/R by >20-fold and F53L by >40-fold. IC50 of lopinavir against isolates with 0–3 mutations was 0.8-fold higher	Carrillo98
K 20 M	AAG → ATG	PI	ABT-378 (lopinavir)	N	Y			In conjunction with other mutations, susceptibility to lopinavir was reduced by mutations at positions 82,54,10,63,71 and 84 by 4–10-fold, K20M/R by >20-fold and F53L by >40-fold. IC50 of lopinavir against isolates with 0–3 mutations was 0.8-fold higher	Kempf01
K 20 M	AAG → ATG	PI	AG1343 (nelfinavir)	?	Y			Seen in two patients following a switch from saquinavir. Associated with reduced susceptibility to both saquinavir and nelfinavir.	Lawrence99
K 20 M	AAG → ATG	PI	MK-639 (indinavir)	?	Y		VX-478 (8)	In conjunction with other mutations, susceptibility to lopinavir was reduced by mutations at positions 82,54,10,63,71 and 84 by 4–10-fold, K20M/R by >20-fold and F53L by >40-fold. IC50 of lopinavir against isolates with 0–3 mutations was 0.8-fold higher	Condra96
K 20 R	AAG → AGG	PI	ABT-378 (lopinavir)	N	Y				Kempf01
K 20 R	AAG → AGG	PI	ABT-538 (ritonavir)	N	Y				Molla96
K 20 R	AAG → AGG	PI	MK-639 (indinavir)	?	Y		Ro-31-8959 (8);		Condra96

Mutations in HIV PROTEASE that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
L 23 I	CTA → ATA	PI	BILA 2185 BS	Y	?		Ro-31-8959 (50); L-735,524 (80); BILA 1906 BS (360)	L10F/L23I/V32I/M46I/I47V/I54M/A71V/I84V: 1,500-fold. Associated Gag mutations: p1/p6 cleavage site (L to F (CTT to TTT at P1'); p7/p1 cleavage site (Q to R (CAG to CGG) at P3', A to V (GCT to CTT) at P2').	Croteau97, Doyon96
L 24 I	TTA → ATA	PI	ABT-378 (lopinavir)	N	Y			In conjunction with other mutations, susceptibility to lopinavir was reduced by mutations at positions 82,54,10,63,71 and 84 by 4–10-fold, K20M/R by >20-fold and F53L by >40-fold. IC50 of lopinavir against isolates with 0–3 mutations was 0.8-fold higher	Kempf01
L 24 I	TTA → ATA	PI	MK-639 (indinavir)	?	Y		SC-52151 (8)		Condra96, Condra95
L 24 V	TTA → GTA	PI	SC-52151 (telinavir)	Y	?	10–20	SC55389A	L24V/G48V/A71V/V75I/P81T: 1000-fold	Potts94, Pillay96
D 30 N	GAT → AAT	PI	AG1343 (nelfinavir)	Y	Y			D30N/A71V: 7-fold; D30N and N88D are most common in vivo after 24 weeks of therapy; they do not cause cross-resistance to other protease inhibitors	Patrick96, Patrick97
V 32 I	GTA → ATA	PI	A-77003	Y	?	7 (enzyme resist.)		V32I appears first; progression to V32I/M46V and V32I/M46V/A71V/V82A occurs even in the absence of drug	Kaplan94
V 32 I	GTA → ATA	PI	ABT-378 (lopinavir)	Y	?		Passage 17 virus: ritonavir, 21-fold; saquinavir, 4-fold	I84V/L10F/M46I/T91S/V32I/I47V: 46 fold, Passage 17 virus: I84V/L10F/M46I/T91S/V32I/I47V/V47A/G16E/H69Y: 338 fold (in presence of p7/p1 (AN/F to VN/F) cleavage-site mutation and p1/p6 (F/L to F/F) cleavage-site mutation).	Carrillo98
V 32 I	GTA → ATA	PI	ABT-538 (ritonavir)	Y	?	40		V32I and V82I are synergistic mutations yielding 20-fold enzyme resistance	Molla96

Mutations in HIV PROTEASE that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
V 32 I	GTA →ATA	PI	BILA 1906 BS	Y	?	?			V32I/A71V: 3-fold; V32I/M46I/L/A71V/I84V: 5-fold; V32I/M46I/L/A71V/I84A: 520-fold. 32I/46L/71V/84A are functionally impaired. Associated Gag mutations: p1/p6 cleavage site (L to F (CTT to TTT at P1'))	Lamarre94, Croteau97
V 32 I	GTA →ATA	PI	BILA 2011 (palinavir)	Y	?	?	1200	BILA 1906 (1400)	Other mutations found in p1/p6 cleavage site	Lamarre95
V 32 I	GTA →ATA	PI	BILA 2185 BS	Y	?	?		BILA 1906 (360)	L10F/L23I/V32I/M46I/I47V/I54M/A71V/I84V: 1,500-fold. Associated Gag mutations: p1/p6 cleavage site (L to F (CTT to TTT at P1')); p7/p1 cleavage site (Q to R (CAG to CGG) at P3', A to V (GCT to CTT) at P2')	Croteau97
V 32 I	GTA →ATA	PI	BMS 232632	Y	?	?			V32L/L33F/M46I/A71V/I84V/N88S: 183-fold., L10YF/I50L/L63P/A71V/N88S: 93-fold., V32I/M46I/I84V/L89M: 96-fold.	Gong00
V 32 I	GTA →ATA	PI	JE-2147	Y					in vitro selection in MT-2 cells, passage 33	Gatanaga02
V 32 I	GTA →ATA	PI	KNI-272	Y	?	?	2		V32I/M46I/I84V: 37-fold; V32I/L33F/K45I/F53L/A71V/I84V/L89M: 130-fold	Gulnik95
V 32 I	GTA →ATA	PI	MK-639 (indinavir)	Y	Y	Y			V32I/M46I/V82A: 3-fold; V32I/M46I/A71V/V82A: 14-fold	Condra96, Condra95
V 32 I	GTA →ATA	PI	VX-478 (amprenavir)	Y	Y	Y			in vitro selection in MT-2 cells, passage 10	Gatanaga02
L 33 F	TTA →TTC	PI	ABT-538 (ritonavir)	N	Y	Y			M36I/I54V/A71V/V82I: 8-fold; K20R/M36I/I54V/V82A: 41-fold. In vivo, V82A/F/T/S occurs first, often followed by changes at 54,71 and 36	Molla96

Mutations in HIV PROTEASE that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
L 33 F	TTA → TTC	PI	BMS 232632	Y	?			V32L/L33F/M46I/A71V/I84V/N88S: 183-fold., L10Y/F150L/L63P/A71V/N88S: 93-fold., V32I/M46I/I84V/L89M: 96-fold.	Gong00
E 35 D	TTA → TTC	PI	PNU-140690 (tipranavir)	?	Y			Seen in 60% of patients receiving Tipranavir therapy.	Rusconi00
M 36 I	ATG → ATA	PI	ABT-538 (ritonavir)	N	Y			In vivo, V82 occurs first, often followed by changes at 54, 71 and 36	Molla96
M 36 I	ATG → ATA	PI	AG1343 (nelmivir)		Y			Seen in 30% of patients receiving Tipranavir therapy.	Patck96
N 37 D	ATG → ATA	PI	PNU-140690 (tipranavir)	?	Y			Seen in 20% of patients receiving Tipranavir therapy.	Rusconi00
R 41 K	ATG → ATA	PI	PNU-140690 (tipranavir)	?	Y			Seen in 20% of patients receiving Tipranavir therapy.	Rusconi00
K 45 I	AAA → ATA	PI	DMP-323					L10F/K45I/I84V: 50-fold	Tisdale94
M 46 F	ATG → TTC	PI	A-77003	Y	?	4 (enzyme resist.)		Seen with V82A	Kaplan94
M 46 I	ATG → ATA	PI	A-77003	Y	?			No effect on susceptibility but improves replication competency of R8Q mutant; R8K/M46I/G48Y: 20-fold	Ho94, Kaplan94
M 46 I	ATG → ATA	PI	ABT-378 (lopinavir)	Y			Passage 17 virus: ritonavir, 21-fold; saquinavir, 4-fold	I84V/L10F/M46I: 4 fold, I84V/L10F/M46I/T91S: 12 fold, I84V/L10F/M46I/T91S/V32I/I47V: 46 fold, Passage 17 virus: I84V/L10F/M46I/T91S/V32I/I47V/A/G16E/H69Y: 338 fold (in presence of p7/p1 (AN/F to VN/F) cleavage-site mutation and p1/p6 (F/L to F/F) cleavage-site mutation).	Carrillo98
M 46 I	ATG → ATA	PI	ABT-538 (ritonavir)	Y	Y			M46I/L63P/A71V/V82F/I84V: 27-fold	Molla96
M 46 I	ATG → ATA	PI	AG1343 (nelmivir)	Y	Y				Patck96

Mutations in HIV PROTEASE that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
M 46 I	ATG→ATA	PI	BILA 1906 BS	Y	?		L 735,524 (60)	V32I/A71V: 3-fold; V32I/M46I/L/A71V/I84V: 5-fold; V32I/M46I/L/A71V/I84A: 520-fold. V32I/M46I/A71V/I84A is functionally impaired. Associated Gag mutations: p1/p6 cleavage site (L to F (CTT to TTT at P1')	Croteau97, Doyon96, Lamarre94, Lamarre95
M 46 I	ATG→ATA	PI	BILA 2185 BS	Y	?		BILA 1906 (360)	L10F/L23I/V32I/M46I/I47V/I54M/A71V/I84V: 1,500-fold. Associated Gag mutations: p1/p6 cleavage site (L to F (CTT to TTT at P1'); p7/p1 cleavage site (Q to R (CAG to CCG) at P3', A to V (GCT to CTT) at P2').	Croteau97
M 46 I	ATG→ATA	PI	BMS 232632	Y	?			V32L/L33F/M46I/A71V/I84V/N88S: 183-fold., L10YF/I50L/L63P/A71V/N88S: 93-fold., V32I/M46I/I84V/L89M: 96-fold.	Gong00
M 46 I	ATG→ATA	PI	DMP 450	Y	?			Probably compensatory	Otto95, Winslow95
M 46 I	ATG→ATA	PI	JE-2147	Y	?		KN1-272: 7-fold; Ritonavir: 9-fold	L10F/M46I/I47V/I84V: 28-fold. >50 passages required for isolation of resistant virus.	Yoshimura99
M 46 I	ATG→?	PI	JE-2147	Y				in vitro selection in MT-2 cells, passage 6	Gatanaga02
M 46 I	ATG→ATA	PI	MK-639 (indinavir)	N	Y			M46I/L63P/V82T: 4-fold; L10R/M46I/L63P/V82T: 4-fold; L10R/M46I/L63P/V82T/I84V: 8-fold	Condra96, Condra95
M 46 I	ATG→ATA	PI	VB 11,328	Y	?			I50V/M46I/I47V: 20-fold	Tisdale94, Partaledis95
M 46 I	ATG→ATA	PI	VX-478 (amprenavir)	Y	?	Nil			Partaledis95
M 46 L	ATG→TTC	PI	A-77003	Y	?	2-3 (enzyme resist.)			Kaplan94

Mutations in HIV PROTEASE that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
M 46 L	ATG→TTG	PI	ABT-378 (lopinavir)	N	Y			In conjunction with other mutations, susceptibility to lopinavir was reduced by mutations at positions 82,54,10,63,71 and 84 by 4–10-fold, K20M/R by >20-fold and F53L by >40-fold. IC50 of lopinavir against isolates with 0–3 mutations was 0.8-fold higher	Kempf01
M 46 L	ATG→TTG	PI	BILA 1906 BS	Y	?			Associated p1/p6 cleavage site mutation (L to F (CTT to TTT) at P1')	Croteau97, Doyon96, Lamarre94, Lamarre95
M 46 L	ATG→CTG	PI	DMP-323	Y	?			V82A/M46L: 7-fold; V82A/M46L/L97V: 11-fold	King95
M 46 L	ATG→TTG	PI	MK-639 (indinavir)	Y	Y			V32I/M46L/A71V/V82A: 14-fold; V32I/M46L/V82A: 3-fold	Tisdale94
M 46 V	ATG→GTG	PI	A-77003	Y	?			V32I appears first; progression to V32I/M46V and V32I/M46V/A71V/V82A occurs even in the absence of drug.	Tisdale94
I 47 V	ATA→GTA	PI	ABT-378 (lopinavir)	Y	?		Passage 17 virus: ritonavir, 21-fold; saquinavir, 4-fold	I84V/L10F/M46I/T91S/V32I/I47V: 46 fold, Passage 17 virus: I84V/L10F/M46I/T91S/V32I/I47V/V47A/G16E/H69Y: 338 fold (in presence of p7/p1 (AN/F to VN/F) cleavage-site mutation and p1/p6 (F/L to F/F) cleavage-site mutation).	Carrillo98
I 47 V	ATA→CTA	PI	BILA 2185 BS	Y	?		BILA 1906 (360)	L10F/L23I/V32I/M46I/I47V/I54M/A71V/I84V: 1,500-fold. Associated Gag mutations: p1/p6 cleavage site (L to F (CTT to TTT at P1'); p7/p1 cleavage site (Q to R (CAG to CGG) at P3', A to V (GCT to CTT) at P2').	Croteau97
I 47 V	ATA→CTA	PI	JE-2147	Y	?		KNI-272: 7-fold; Ritonavir: 9-fold	L10F/I47V/I84V: 19-fold. L10F/M46I/I47V/I84V: 28-fold. >50 passages required for isolation of resistant virus.	Yoshimura99

Mutations in HIV PROTEASE that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
I 47 V	ATA → GTA	PI	JE-2147	Y				in vitro selection in MT-2 cells, passage 33	Gatanaga02
I 47 V	ATA → CTA	PI	VB 11,328	Y	?			I50V/M46I/I47V: 20-fold	Partaledis95
I 47 V	ATA → CTA	PI	VX-478 (amprenavir)	Y	?	Nil			Partaledis95
V 47 A	GTA → TAT	PI	ABT-378 (lopinavir)	Y	?		Passage 17 virus: ritonavir, 21-fold; saquinavir, 4-fold	I84V/L10F/M46I/T91S/V32I/I47V: 46 fold, Passage 17 virus: I84V/L10F/M46I/T91S/V32I/I47V/V47A/G16E/H69Y: 338 fold (in presence of p7/p1 (AN/F to VN/F) cleavage-site mutation and p1/p6 (F/L to F/F) cleavage-site mutation).	Carrillo98
G 48 V	GGG → GTG	PI	A-77003	Y	?			R8K/M46I/G48V: 20-fold; G48V/I82T: 100-fold	Borman95
G 48 V	GGG → GTG	PI	MK-639 (indinavir)	?	Y				Vasudevachari96
G 48 V	GGG → GTG	PI	MP-167	Y	?	20	MP-134(5) SC-52151(16) Ro31-8959(5) (Fold increase in IC90s).	L10F/G48V: 20-fold	Mo96
G 48 V	GGG → GTG	PI	Ro 31-8959 (saquinavir)	Y	Y			Found in comb. with L10I in vivo; G48V/I84V/L90M: 30-fold; G48V/L90M: >100-fold enzyme resistance; G48V/L90M/I54V: > 50-fold (subtype B or O)	Jacobsen94, Eberle95
G 48 V	GGG → GTG	PI	SC-52151 (telinavir)	Y	?		Ro 31-8959	G48V/V82A, G48V/L63P/V82A or I54T: 10- to 20-fold; L24V/G48V/A71V/V75I/P81T: 1000-fold	Potts94, Pillay96
I 50 L	ATT → CTT	PI	BMS 232632	Y	?			V32L/L33F/M46I/A71V/I84V/N88S: 183-fold, L10Y/F/I50L/L63P/A71V/N88S: 93-fold., V32I/M46I/I84V/L89M: 96-fold.	Gong00
I 50 V	ATT → GTT	PI	VB 11,328	Y	?	3		I50V/M46I/I47V: 20-fold	Tisdale94, Partaledis95
I 50 V	ATT → GTT	PI	VX-478 (amprenavir)	Y	?	3		D30N/G52S: 93-fold	Partaledis95, Rao96
G 52 S	GGT → AGT	PI	AG1343 (nelfinavir)	?	Y				Patrick98

Mutations in HIV PROTEASE that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
F 53 L	TTT→?	PI	ABT-378 (lopinavir)	N	Y			In conjunction with other mutations, susceptibility to lopinavir was reduced by mutations at positions 82,54,10,63,71 and 84 by 4–10-fold, K20M/R by >20-fold and F53L by >40-fold. IC50 of lopinavir against isolates with 0–3 mutations was 0.8-fold higher	Kempf01
I 54 L	ATC→CTC	PI	ABT-378 (lopinavir)	N	Y			In conjunction with other mutations, susceptibility to lopinavir was reduced by mutations at positions 82,54,10,63,71 and 84 by 4–10-fold, K20M/R by >20-fold and F53L by >40-fold. IC50 of lopinavir against isolates with 0–3 mutations was 0.8-fold higher	Kempf01
I 54 M	ATT→ATG	PI	BILA 2185 BS	Y	?		BILA 1906 (360)	L10F/L23I/V32I/M46I/I47V/I54M/A71V/I84V: 1,500-fold. Associated Gag mutations: p1/p6 cleavage site (L to F (CTT to TTT at P1'); p7/p1 cleavage site (Q to R (CAG to CGG) at P3'; A to V (GCT to CTT) at P2').	Croteau97
I 54 M	ATC→ATG	PI	VX-478 (amprenavir)	Y				in vitro selection in MT-2 cells, passage 31	Gatanaga02
I 54 T	ATC→ACC	PI	ABT-378 (lopinavir)	N	Y			In conjunction with other mutations, susceptibility to lopinavir was reduced by mutations at positions 82,54,10,63,71 and 84 by 4–10-fold, K20M/R by >20-fold and F53L by >40-fold. IC50 of lopinavir against isolates with 0–3 mutations was 0.8-fold higher	Kempf01

Mutations in HIV PROTEASE that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
I 54 V	ATC → GTC	PI	ABT-378 (lopinavir)	N	Y			In conjunction with other mutations, susceptibility to lopinavir was reduced by mutations at positions 82,54,10,63,71 and 84 by 4–10-fold, K20M/R by >20-fold and F53L by >40-fold. IC50 of lopinavir against isolates with 0–3 mutations was 0.8-fold higher	Kempf01
I 54 V	ATC → GTC	PI	ABT-538 (ritonavir)	N	Y			I54V/V82T: 9-fold; K20R/M36I/I54V/V82A: 41-fold; M36I/I54V/A71V/V82T: 8-fold; I54V/A71V/V82A/L90N: 7-fold; In vivo, V82A/F/T/S occurs first, followed by changes at 54, 71 and 36	Molla96
I 54 V	ATC → GTC	PI	MIK-639 (indinavir)	?	Y			In subtype O and B	Lamarre94
I 54 V	ATA → GTA	PI	Ro 31-8959 (saquinavir)	Y				Seen in one patient following a switch from saquinavir. Associated with reduced susceptibility to both saquinavir and nelfinavir.	Jacobsen94, Eberle95
K 55 R	AAA → AGA	PI	AG1343 (nelfinavir)	?	Y				Lawrence99
R 57 K	AGA → AAA	PI	AG1343 (nelfinavir)	?	Y			Seen in one patient following a switch from saquinavir. Associated with reduced susceptibility to both saquinavir and nelfinavir.	Lawrence99
D 60 E	GAT → GAA	PI	DMP 450	Y	?			Probably compensatory	Otto95, Winslow95
D 60 E	GAT → GAA	PI	PNU-140690 (tipranavir)	?	Y			Seen in 30% of patients receiving Tipranavir therapy.	Rusconi00
L 63 P	CTC → CCC	PI	ABT-378 (lopinavir)	N	Y			In conjunction with other mutations, susceptibility to lopinavir was reduced by mutations at positions 82,54,10,63,71 and 84 by 4–10-fold, K20M/R by >20-fold and F53L by >40-fold. IC50 of lopinavir against isolates with 0–3 mutations was 0.8-fold higher	Kempf01

Mutations in HIV PROTEASE that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
L 63 P	CTC→CCC	PI	AG1343 (nelfinavir)	?	Y			D30N/M36I/L63P: 60-fold	Patick98
L 63 P	CTC→CCC	PI	BMS 232632	Y	?			V32L/L33F/M46I/A71V/I84V/N88S: 183-fold., L10Y/F150L/L63P/A71V/N88S: 93-fold., V32I/M46I/I84V/L89M: 96-fold.	Gong00
L 63 P	CTC→CCC	PI	MK-639 (indinavir)	N	Y			M46I/L63P/V82T: 4-fold; L10R/M46I/L63P/V82T/I84V: 8-fold; L10R/M46I/L63P/V82T: 4-fold	Condra96, Condra95
H 69 Y	CAT→TAT	PI	ABT-378 (lopinavir)	Y	?		Passage 17 virus: ritonavir, 21-fold; saquinavir, 4-fold	Passage 17 virus: I84V/L10F/M46I/T91S/V32I/I47V/V47A/G16E/H69Y: 338 fold (in presence of p7/p1 (AN/ F to V/N/F) cleavage-site mutation and p1/p6 (F/L to F/F) cleavage-site mutation).	Carrillo98
A 71 I	GCT→ATT	PI	ABT-378 (lopinavir)	N	Y			In conjunction with other mutations, susceptibility to lopinavir was reduced by mutations at positions 82,54,10,63,71 and 84 by 4–10-fold, K20M/R by >20-fold and F53L by >40-fold. IC50 of lopinavir against isolates with 0–3 mutations was 0.8-fold higher	Kempf01
A 71 L	GCT→CTC	PI	ABT-378 (lopinavir)	N	Y			In conjunction with other mutations, susceptibility to lopinavir was reduced by mutations at positions 82,54,10,63,71 and 84 by 4–10-fold, K20M/R by >20-fold and F53L by >40-fold. IC50 of lopinavir against isolates with 0–3 mutations was 0.8-fold higher	Kempf01

Mutations in HIV PROTEASE that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
A 71 T	GCT→ACT	PI	ABT-378 (lopinavir)	N	Y			In conjunction with other mutations, susceptibility to lopinavir was reduced by mutations at positions 82,54,10,63,71 and 84 by 4–10-fold, K20M/R by >20-fold and F53L by >40-fold. IC50 of lopinavir against isolates with 0–3 mutations was 0.8-fold higher	Kempf01
A 71 T	GCT→ACT	PI	BMS 186318	Y	?			A71T/V82A: 15-fold	Patrick95, Rose94
A 71 T	GCT→ACT	PI	MK-639 (indinavir)	?	Y				Condra96, Condra95
A 71 T	GCT→ACT	PI	PNU-140690 (tipranavir)	?	Y				Ruscioni00
A 71 V	GCT→GTT	PI	A-77003	Y	?			V32I appears first; progression to V32I/M46V and V32I/M46V/A71V/V82A occurs even in the absence of drug;	Tisdale94, King95
A 71 V	GCT→GTC	PI	ABT-378 (lopinavir)	N	Y			M46I/L63P/A71V/V82F/I84V: 27-fold	Kempf01
A 71 V	GCT→GTT	PI	ABT-538 (ritonavir)	Y	Y			In conjunction with other mutations, susceptibility to lopinavir was reduced by mutations at positions 82,54,10,63,71 and 84 by 4–10-fold, K20M/R by >20-fold and F53L by >40-fold. IC50 of lopinavir against isolates with 0–3 mutations was 0.8-fold higher	Molla96
A 71 V	GCT→GTT	PI	AG1343 (nelfinavir)	Y	?	5		D30N/A71V: 7-fold; M46I/L63P/A71V/I84V: 30-fold	Patrick98
A 71 V	GCT→GTT	PI	BILA 1906 BS	Y	?			V32I/A71V: 3-fold; V32I/M46I,L/A71V/I84V: 5-fold; V32I/M46I,L/A71V/I84A: 520-fold. 32I/46L/71V/84A are functionally impaired. Associated Gag mutations: p1/p6 cleavage site (L to F (CTT to TTT at P1')	Croteau97, Doyon96, Lamarre94, Lamarre95
A 71 V	GCT→GTT	PI	BILA 2011 (palinavir)	Y	?		BILA 2185: 30-fold		Lamarre94

Mutations in HIV PROTEASE that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
A 71 V	GCT→GTT	PI	BILA 2185 BS	Y	?		BILA 1906 (360)	L10F/L23I/V32I/M46I/I47V/I54M/A71V/I84V: 1,500-fold. Associated Gag mutations: p1/p6 cleavage site (L to F (CTT to TTT at P1'); p7/p1 cleavage site (Q to R (CAG to CGG) at P3'; A to V (GCT to CTT) at P2').	Croteau97
A 71 V	GCT→GTT	PI	BMS 232632	Y	?			V32L/L33F/M46I/A71V/I84V/N88S: 183-fold., L10YF/I50L/L63P/A71V/N88S: 93-fold., V32I/M46I/I84V/L89M: 96-fold.	Gong00
A 71 V	GCT→GTT	PI	MK-639 (indinavir)	Y	Y			V32I/M46L/A71V/V82A: 14-fold	Tisdale94
A 71 V	GCT→GTT	PI	SC-52151 (telinavir)	Y	?		Not L-735,524	A71V/V75I/P81T: 20- to 30-fold; L24V/G48V/A71V/V75I/P81T: 1000-fold; N88D or I11V/M46I/F53L/A71V/N88D: 10- to 20-fold	Potts94, Pillay96
A 71 V	GCT→GTT	PI	VX-478 (amprenavir)	Y	?			in vitro selection in MT-2 cells, passage 31	Gatanaga02
G 73 S	GGT→AGT	PI	AG1343 (nelfinavir)	?	Y			Seen in two patients following a switch from saquinavir. Associated with reduced susceptibility to both saquinavir and nelfinavir.	Lawrence99
G 73 S	GGT→GCT	PI	MK-639 (indinavir)	?	Y			Emerges following a switch from saquinavir to indinavir.	Dultoust97
V 75 I	GTA→ATA	PI	SC-52151 (telinavir)	Y	?			L24V/G48V/A71V/V75I/P81T: 1000-fold; A71V/V75I/P81T: 20- to 30-fold; L24V/G48V/A71V/V75I/P81T: 1000-fold	Potts94, Pillay96
V 77 I	GTA→ATA	PI	AG1343 (nelfinavir)	Y	Y				Patrick98
P 81 T	CCT→ACT	PI	SC-52151 (telinavir)	Y	?			A71V/V75I/P81T: 20- to 30-fold; L24V/G48V/A71V/V75I/P81T: 1000-fold	Potts94, Pillay96
I 82 T	ATC→ACC	PI	A-77003	Y	?			G48V/I82T: 100-fold 82T was derived from in vitro passage of 82I)	Swanstrom94

Mutations in HIV PROTEASE that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
V 82 A	GTC→GCC	PI	A-77003	Y	?			Rare; seen with M46F; V32I appears first; progression to V32I/M46V and V32I/M46V/A71V/V82A occurs even in the absence of drug	Tisdale94, Borman95, Swanstrom94
V 82 A	GTC→GCC	PI	ABT-378 (lopinavir)	N	Y			In conjunction with other mutations, susceptibility to lopinavir was reduced by mutations at positions 82,54,10,63,71 and 84 by 4–10-fold, K20M/R by >20-fold and F53L by >40-fold. IC50 of lopinavir against isolates with 0–3 mutations was 0.8-fold higher	Kempf01
V 82 A	GTC→GCC	PI	ABT-538 (ritonavir)	N	Y	2		In vivo, V82 occurs first, often followed by changes at I54, A71 and M36	Molla96
V 82 A	GTC→GCC	PI	AG1343 (nelfinavir)	?	Y				Lawrence99
V 82 A	GTC→GCC	PI	BMS 186318	Y	?		A-77003 (4)	A71T/V82A: 15-fold	Patrick95, Rose94
V 82 A	GTC→GCC	PI	DMP-323	Y	?			V82A/M46L: 7-fold; V82A/M46L/L97V: 11-fold; L10F/V82A: 2-fold; ; V82A/L97V: 3-fold	King95
V 82 A	GTC→GCC	PI	MK-639 (indinavir)	Y	Y			V32I/M46L/V82A: 3-fold; V32I/M46L/A71V/V82A: 14-fold	Condra96, Condra95
V 82 A	GTC→GCC	PI	P9941	Y	?	6–8			Otto93
V 82 A	GTC→GCC	PI	Ro 31-8959 (saquinavir)	?	Y			Follows G48V during saquinavir therapy or after a switch to nelfinavir or indinavir.	Winters97, Eastman97, Schapiro97
V 82 A	GTC→GCC	PI	SC-52151 (telinavir)	Y	?			G48V/V82A, G48V/L63P/V82A or I54T: 10- to 20-fold	Potts94, Pillay96
V 82 A	GTC→GCC	PI	SKF108922	Y	?				Shao95

Mutations in HIV PROTEASE that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
V 82 F	GTC→TTC	PI	ABT-378 (lopinavir)	N	Y			In conjunction with other mutations, susceptibility to lopinavir was reduced by mutations at positions 82,54,10,63,71 and 84 by 4–10-fold, K20M/R by >20-fold and F53L by >40-fold. IC50 of lopinavir against isolates with 0–3 mutations was 0.8-fold higher	Kempf01
V 82 F	GTC→TTC	PI	ABT-538 (ritonavir)	Y	Y			V82F/I84V: 8- to 10-fold; M46I/L63P/A71V/V82F/I84V: 27-fold	Molla96
V 82 F	GTC→TTC	PI	DMP-323	Y	?			V82F/I84V: 92-fold	King95
V 82 F	GTC→TTC	PI	MK-639 (indinavir)	?	Y			No resistance alone but V32I and V82I are synergistic mutations yielding 20-fold enzyme resistance 82T was derived from in vitro passage of 82I	Partaledis94 Kaplan94
V 82 I	GTC→ATC	PI	A-77003	Y	?				King95
V 82 I	GTC→ATC	PI	DMP-323	Y	?	< 2			King95
V 82 I	GTC→ATC	PI	JE-2147	Y				in vitro selection in MT-2 cells, passage 33	Gatanaga02
V 82 S	GTC→TCC	PI	ABT-538 (ritonavir)	N	Y	6		In vivo, V82 occurs first, often followed by changes at I54, A71 and M36	Molla96
V 82 T	GTC→ACC	PI	ABT-378 (lopinavir)	N	Y			In conjunction with other mutations, susceptibility to lopinavir was reduced by mutations at positions 82,54,10,63,71 and 84 by 4–10-fold, K20M/R by >20-fold and F53L by >40-fold. IC50 of lopinavir against isolates with 0–3 mutations was 0.8-fold higher	Kempf01
V 82 T	GTC→ACC	PI	ABT-538 (ritonavir)	N	Y	3		In vivo, V82 occurs first, often followed by changes at I54, A71 and M36; V82T has reduced replication efficacy in natural background	Molla96

Mutations in HIV PROTEASE that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
V 82 T	GTC → ACC	PI	MK-639 (indinavir)	N	Y			M46I/L63P/V82T: 4-fold; L10R/M46I/L63P/V82T: 4-fold; L10R/M46I/L63P/V82T/V84V: 8-fold	Condra96, Condra95
V 82 T	GTC → ACC	PI	SKF108842	Y	?				Shao95
V 82 T	GTC → ACC	PI	SKF108922	Y	?				Shao95
I 84 A	ATA → GCA	PI	BILA 1906 BS	Y	?		BILA 2185 BS (200)	V32I/A71V: 3-fold; V32I/M46I,L/A71V/184V: 5-fold; V32I/M46I,L/A71V/184A: 520-fold. 32I/46L/71V/84A are functionally impaired. Associated Gag mutations: p1/p6 cleavage site (L to F (CTT to TTT at P1'))	Croteau97, Doyon96, Lamarre94, Lamarre95
I 84 A	ATG → ATA	PI	BILA 2011 (palmavir)	Y	?		Ro, 31-8959 (400);	184A is the most common mutation	Lamarre94
I 84 V	ATA → GTA	PI	ABT-378 (lopinavir)	Y	?		Passage 17 virus: ritonavir, 21-fold; saquinavir, 4-fold	184V/L10F/M46I: 4 fold, 184V/L10F/M46I/T91S: 12 fold, 184V/L10F/M46I/T91S/V32I/47V: 46 fold, Passage 17 virus: 184V/L10F/M46I/T91S/V32I/147V/V47A/G16E/H69Y: 338 fold (in presence of p7/p1 (AN/F to VN/F) cleavage-site mutation and p1/p6 (F/L to F/F) cleavage-site mutation).	Carrillo98
I 84 V	ATA → GTA	PI	ABT-538 (ritonavir)	Y	Y			M46I/L63P/A71V/V82F/184V: 27-fold; V82F/184V: 8- to 10-fold; M46I/L63P/A71V/V82F/184V: 27-fold	Molla96
I 84 V	ATA → GTA	PI	AG1343 (nelfinavir)		?			M46I/L63P/A71V/184V: 30-fold	Patrick96
I 84 V	ATA → GTA	PI	BILA 1906 BS	Y	?		BILA 2185 BS(200)	V32I/A71V: 3-fold; V32I/M46I,L/A71V/184V: 5-fold; V32I/M46I,L/A71V/184A: 520-fold. 32I/46L/71V/84A are functionally impaired. Associated Gag mutations: p1/p6 cleavage site (L to F (CTT to TTT at P1'))	Croteau97, Doyon96, Lamarre94, Lamarre95

Mutations in HIV PROTEASE that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
I 84 V	ATA→GTA	PI	BILA 2185 BS	Y	?		BILA 1906 BS(360)	L10F/L23I/V32I/M46I/I47V/I54M/A71V/I84V: 1,500-fold. Associated Gag mutations: p1/p6 cleavage site (L to F (CTT to TTT at P1'); p7/p1 cleavage site (Q to R (CAG to CGG) at P3', A to V (GCT to CTT) at P2').	Croteau97
I 84 V	ATA→GTA	PI	BMS 232632	Y	?			Minor resistance mutation for BMS-232632.	Gong99
I 84 V	ATA→GTA	PI	BMS 232632	Y	?			V32L/L33F/M46I/A71V/I84V/N88S: 183-fold., L10Y,F/I50L/L63P/A71V/N88S: 93-fold., V32I/M46I/I84V/L89M: 96-fold.	Gong00
I 84 V	ATA→GTA	PI	DMP 450	Y	?			V82F/I84V: 92-fold; L10F/K45I/I84V: 50-fold	Otto95, Winslow95
I 84 V	ATA→GTA	PI	DMP-323	Y	?	12	P9941; not A-77003 or Ro 31-8959		Tisdale94, King95
I 84 V	ATA→GTA	PI	JE-2147	Y	?			L10F/I47V/I84V: 19-fold. L10F/M46I/I47V/I84V: 28-fold. >50 passages required for isolation of resistant virus.	Yoshimura99
I 84 V	ATA→GTA	PI	MK-639 (indinavir)	N	Y			G48V/I84V/L90M: 30-fold; L10R/M46I/L63P/V82T/I84V: 8-fold	Condra96, Condra95
I 84 V	ATA→GTA	PI	MP-134	Y	?	10	MP-167(5) ABT-538(10) MK-639(8) SC-52151(8) Ro31-895(2) (IC90 data)		Mo96
I 84 V	ATA→GTA	PI	Ro 31-8959 (saquinavir)	Y	?				Tisdale94
I 84 V	ATA→GTA	PI	RPL-312	Y	?	5			el-Farrash94
I 84 V	ATA→GTA	PI	SKF108842	Y	?				Shao95
I 84 V	ATA→GTA	PI	VB 11,328	Y	?			L10F/I84V: 8-fold	Partaledis95
I 84 V	ATA→GTA	PI	VX-478 (amprenavir)	Y	?				Partaledis95
N 88 D	AAT→GAT	PI	AG1343 (nelfinavir)	Y	Y			D30N and N88D are most common in vivo after 24 weeks of therapy; they do not cause cross-resistance to other protease inhibitors.	Patrick96

Mutations in HIV PROTEASE that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
N 88 D	AAT→GAT	PI	SC-52151 (telinavir)	Y	?			N88D compensatory, no resistance alone	Potts94, Pillay96
N 88 S	AAT→AGT	PI	BMS 232632	Y	?			Major resistance mutation for BMS-232632.	Gong99
N 88 S	AAT→AGT	PI	BMS 232632	Y	?			V32L/L33F/M46I/A71V/I84V/N88S: 183-fold., L10Y/F150L/L63P/A71V/N88S: 93-fold., V32I/M46I/I84V/L89M: 96-fold.	Gong00
N 88 S	AAT→AGT	PI	SC-55389A	Y	?	20	L735,524 (3); not SC-52151	N88S/L10F: 25	Smidt97
N 88 S	AAT→AGT	PI	VX-478 (amprenavir)	Y	Y			Confers >2.5-fold hypersusceptibility to Amprenavir.	Ziermann00
L 89 M	AAT→AGT	PI	BMS 232632	Y	?			V32L/L33F/M46I/A71V/I84V/N88S: 183-fold., L10Y/F150L/L63P/A71V/N88S: 93-fold., V32I/M46I/I84V/L89M: 96-fold.	Gong00
L 90 M	TTG→ATG	PI	ABT-378 (lopinavir)	N	Y			In conjunction with other mutations, susceptibility to lopinavir was reduced by mutations at positions 82,54,10,63,71 and 84 by 4–10-fold, K20M/R by >20-fold and F53L by >40-fold. IC50 of lopinavir against isolates with 0–3 mutations was 0.8-fold higher	Kempf01
L 90 M	TTG→ATG	PI	ABT-538 (ritonavir)	N	Y			82A/54V/I71V/90L/M: 7-fold	Molla96
L 90 M	TTG→ATG	PI	AG1343 (nelfinavir)	N	Y			Rare in patients	Patrick96
L 90 M	TTG→ATG	PI	MK-639 (indinavir)	?	Y				Condra96
L 90 M	TTG→ATG	PI	Ro 31–8959 (saquinavir)	Y	Y			G48V/L90M: >100-fold enzyme resistance; double mutant rare in vivo; L90M most common in vivo; G48V/I84V/L90M: 30-fold	Jacobsen94

Mutations in HIV PROTEASE that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
T 91 S	ACT→TCT	PI	ABT-378 (lopinavir)	Y	?		Passage 17 virus: ritonavir, 21-fold; saquinavir, 4-fold	I84V/L10F/M46I/T91S: 12 fold, I84V/L10F/M46I/T91S/V32I/I47V: 46 fold, Passage 17 virus: I84V/L10F/M46I/T91S/V32I/I47V/V47A/G16E/H69Y: 338 fold (in presence of p7/p1 (AN/F to VN/F) cleavage-site mutation and p1/p6 (F/L to F/F) cleavage-site mutation).	Carrillo98
L 97 V	TTA→GTA	PI	DMP-323	Y	?			No resistance alone; V82A/L97V: 3-fold; V82A/M46L/L97V: 11-fold	King95

Mutations in HIV INTEGRASE that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
G 140 S	GGC→AGC	II	L-Chicoric Acid	Y	?	156-fold		Mutation located in the catalytic core of integrase. Mildly attenuates virus growth.	King98

Mutations in HIV ENV that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
R 22 A	AGG→AGA	F/BI	RPR103611	Y	?			found in combination with G36D in patients	Labrosse97
Q 32 H	AGG→AGA	F/BI	T20 (pentafuside)		Y			found in combination with G36D in patients	Wei02
Q 32 R	AGG→AGA	F/BI	T20 (pentafuside)		Y			Resistance lost when R122G substitution is present in HR2 domain	Wei02
G 36 D	GGT→GAT	F/BI	T20 (pentafuside)	Y	Y	13.6		Both G36S and V38M mutations must be present to confer resistance.	Rimsky98
G 36 S	GGT→AGT	F/BI	T20 (pentafuside)	Y	?				Wei02
I 37 V	GGT→AGT	F/BI	T20 (pentafuside)		Y				Wei02
V 38 A	GTG→GCG	F/BI	T20 (pentafuside)	Y	Y	45			Rimsky98
V 38 M	GTG→ATG	F/BI	T20 (pentafuside)	Y	?				Wei02
Q 39 R	GTC→ATG	F/BI	T20 (pentafuside)		Y			represents a conservative change that is present in the HIV-1 LAI consensus sequence	Wei02
R 46 M	AGG→ATG	F/BI	T20 (pentafuside)		Y				Wei02
V 69 I	GTC→ATC	F/BI	T20 (pentafuside)		Y				Wei02
I 84 S	ATC→AGC	F/BI	RPR103611	Y	?				Labrosse97
L 91 H	ATC→AGC	F/BI	RPR103611	Y	?				Labrosse00
N 106 K	AAT→AAG	F/BI	SDF-1 α	Y	?		SDF-1 β : 15-fold; AMB2763: 3-fold.		Schols98
S 113 N	AGT→AAT	F/BI	DS (Dextran sulphate)	Y	?				Este96a, Este97
S 134 N	AGC→AAC	F/BI	DS (Dextran sulphate)	Y	?				Este97, Este96a
S 134 N	AGC→AAC	F/BI	SDF-1 α	Y	?		SDF-1 β : 15-fold; AMB2763: 3-fold.		Schols98

Mutations in HIV ENV that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
F 145 L	TTC→TTA	F/BI	JM-3100	Y	?			Combination of mutations: 2- to 100-fold	DeVreese96, DeVreese96a
F 145 L	TTC→TTA	F/BI	SDF-1 α	Y	?		SDF-1 β : 15-fold; AMB2763: 3-fold.	106K/134N/145L/245I/269E/278H/288V/293D/364-367Deletion/387T: 15-fold.	Schols98
N 188 K	AAT→AAA	F/BI	Siamycin I	Y	?			NI88K/G332E/N351D/A550T/N633D/L762S: 9-fold	Lin96
I 228 V	ATA→GTA	F/BI	JM-2763	Y	?			Combination of mutations	DeVreese96a
G 237 R	ATA→GTA	F/BI	IC9564 (emvirine)	Y	?				Holz-Smith01
F 245 I	TTC→ATC	F/BI	SDF-1 α	Y	?		SDF-1 β : 15-fold; AMB2763: 3-fold.	106K/134N/145L/245I/269E/278H/288V/293D/364-367Deletion/387T: 15-fold.	Schols98
R 252 K	TTC→ATC	F/BI	IC9564 (emvirine)	Y	?				Holz-Smith01
K 269 E	AAA→GAA	F/BI	DS (Dextran sulphate)	Y	?			V3 loop region; S113N/S134N/K269E/Q278E/N293D/N323S/R387I: 250-fold	Este97, Este96a
N 269 E	AAC→GAA	F/BI	SDF-1 α	Y	?		SDF-1 β : 15-fold; AMB2763: 3-fold.	106K/134N/145L/245I/269E/278H/288V/293D/364-367Deletion/387T: 15-fold.	Schols98
N 269 K	AAC→?	F/BI	ALX40-4C	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 145-fold cross-resistant to ALX40-4C. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
N 269 K	AAC→?	F/BI	AMD3100	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 15-fold cross-resistant to AMD3100. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01

Mutations in HIV ENV that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
N 269 K	AAC→?	F/BI	SDF-1	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 26-fold cross-resistant to SDF-1. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
N 269 K	AAC→?	F/BI	T134	Y	N			In vitro selected virus (p145 of HIV-INL4-3, contains mutations N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275 in V3 loop of gp120) 15-fold resistant to T134. Role of each mutation not confirmed by site	Kanbara01
N 269 K	AAC→?	F/BI	T140	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 21-fold cross-resistant to T140. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
N 269 K	AAC→?	F/BI	vMIP-II	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 42-fold cross-resistant to vMIP II. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
N 270 S	AAT→AGT	F/BI	JM-3100	Y	?				DeVreese96, DeVreese96a
R 272 T	AGA→ACA	F/BI	JM-3100	Y	?				DeVreese96, DeVreese96a

Mutations in HIV ENV that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
S 274 del	AGA → ACA	F/BI	ALX40-4C	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 145-fold cross-resistant to ALX40-4C. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
S 274 del	AGA → ACA	F/BI	AMD3100	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 15-fold cross-resistant to AMD3100. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
S 274 del	AGA → ACA	F/BI	SDF-1	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 26-fold cross-resistant to SDF-1. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
S 274 del	AGA → ACA	F/BI	T134	Y	N			In vitro selected virus (p145 of HIV-INL4-3, contains mutations N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275 in V3 loop of gp120) 15-fold resistant to T134. Role of each mutation not confirmed by site	Kanbara01

Mutations in HIV ENV that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
S 274 del	AGA → ACA	F/BI	T140	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 21-fold cross-resistant to T140. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
S 274 del	AGA → ACA	F/BI	vMIP-II	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 42-fold cross-resistant to vMIP II. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
S 274 R	AGT → AGA	F/BI	JM-2763	Y	?			Combination of mutations: 95- to 792-fold	DeVreese96, DeVreese96a
S 274 R	AGT → AGA	F/BI	JM-3100	Y	?	DS (> 7 to 6.667)			DeVreese96, DeVreese96a
I 275 del	AGT → AGA	F/BI	ALX40–4C	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 145-fold cross-resistant to ALX40–4C. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
I 275 del	AGT → AGA	F/BI	AMD3100	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 15-fold cross-resistant to AMD3100. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01

Mutations in HIV ENV that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
I 275 del	AGT→AGA	F/BI	SDF-1	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 26-fold cross-resistant to SDF-1. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
I 275 del	AGT→AGA	F/BI	T134	Y	N			In vitro selected virus (p145 of HIV-INL4-3, contains mutations N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275 in V3 loop of gp120) 15-fold resistant to T134. Role of each mutation not confirmed by site	Kanbara01
I 275 del	AGT→AGA	F/BI	T140	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 21-fold cross-resistant to T140. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
I 275 del	AGT→AGA	F/BI	vMIP-II	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 42-fold cross-resistant to vMIP II. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
Q 278 H	CAG→CAT	F/BI	DS (Dextran sulphate)	Y	?			V3 loop region; S113N/S134N/K269E/Q278E/N293D/N323S/R387I: 250-fold	Este97, Este96a
Q 278 H	CAG→CAT	F/BI	JM-2763	Y	?				DeVreese96, DeVreese96a
Q 278 H	CAG→CAC	F/BI	JM-3100	Y	?				DeVreese96, DeVreese96a

Mutations in HIV ENV that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
Q 278 H	CAG→CAT	F/BI	SDF-1 α	Y	?		SDF-1 β : 15-fold; AMB2763: 3-fold.	106K/134N/145L/245I/269E/278H/288V/293D/364-367Deletion/387T: 15-fold.	Schols98
Q 278 T	CAG→ACG	F/BI	ALX40-4C	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 145-fold cross-resistant to ALX40-4C. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
Q 278 T	CAG→ACG	F/BI	AMD3100	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 15-fold cross-resistant to AMD3100. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
Q 278 T	CAG→ACG	F/BI	SDF-1	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 26-fold cross-resistant to SDF-1. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
Q 278 T	CAG→ACG	F/BI	T134	Y	N			In vitro selected virus (p145 of HIV-INL4-3, contains mutations N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275 in V3 loop of gp120) 15-fold resistant to T134. Role of each mutation not confirmed by site	Kanbara01

Mutations in HIV ENV that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
Q 278 T	CAG→ACG	F/BI	T140	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 21-fold cross-resistant to T140. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
Q 278 T	CAG→ACG	F/BI	vMIP-II	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 42-fold cross-resistant to vMIP II. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
R 279 K	CAG→ACG	F/BI	ALX40–4C	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 145-fold cross-resistant to ALX40–4C. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
R 279 K	CAG→ACG	F/BI	AMD3100	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 15-fold cross-resistant to AMD3100. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01

Mutations in HIV ENV that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
R 279 K	CAG→ACG	F/BI	SDF-1	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 26-fold cross-resistant to SDF-1. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
R 279 K	CAG→ACG	F/BI	T134	Y	N			In vitro selected virus (p145 of HIV-INL4-3, contains mutations N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275 in V3 loop of gp120) 15-fold resistant to T134. Role of each mutation not confirmed by site	Kanbara01
R 279 K	CAG→ACG	F/BI	T140	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 21-fold cross-resistant to T140. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
R 279 K	CAG→ACG	F/BI	vMIP-II	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 42-fold cross-resistant to vMIP II. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01

Mutations in HIV ENV that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
A 284 V	CAG→ACG	F/BI	ALX40-4C	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 145-fold cross-resistant to ALX40-4C. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
A 284 V	CAG→ACG	F/BI	AMD3100	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 15-fold cross-resistant to AMD3100. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
A 284 V	CAG→ACG	F/BI	SDF-1	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 26-fold cross-resistant to SDF-1. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
A 284 V	CAG→ACG	F/BI	T134	Y	N			In vitro selected virus (p145 of HIV-INL4-3, contains mutations N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275 in V3 loop of gp120) 15-fold resistant to T134. Role of each mutation not confirmed by site	Kanbara01

Mutations in HIV ENV that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
A 284 V	CAG→ACG	F/BI	T140	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 21-fold cross-resistant to T140. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
A 284 V	CAG→ACG	F/BI	vMIP-II	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 42-fold cross-resistant to vMIP II. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
F 285 L	CAG→ACG	F/BI	ALX40–4C	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 145-fold cross-resistant to ALX40–4C. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
F 285 L	CAG→ACG	F/BI	AMD3100	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 15-fold cross-resistant to AMD3100. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01

Mutations in HIV ENV that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
F 285 L	CAG→ACG	F/BI	SDF-1	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 26-fold cross-resistant to SDF-1. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
F 285 L	CAG→ACG	F/BI	T134	Y	N			In vitro selected virus (p145 of HIV-INL4-3, contains mutations N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275 in V3 loop of gp120) 15-fold resistant to T134. Role of each mutation not confirmed by site	Kanbara01
F 285 L	CAG→ACG	F/BI	T140	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 21-fold cross-resistant to T140. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
F 285 L	CAG→ACG	F/BI	vMIP-II	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 42-fold cross-resistant to vMIP II. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01

Mutations in HIV ENV that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
V 286 Y	CAG→ACG	F/BI	ALX40-4C	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 145-fold cross-resistant to ALX40-4C. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
V 286 Y	CAG→ACG	F/BI	AMD3100	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 15-fold cross-resistant to AMD3100. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
V 286 Y	CAG→ACG	F/BI	SDF-1	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 26-fold cross-resistant to SDF-1. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
V 286 Y	CAG→ACG	F/BI	T134	Y	N			In vitro selected virus (p145 of HIV-INL4-3, contains mutations N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275 in V3 loop of gp120) 15-fold resistant to T134. Role of each mutation not confirmed by site	Kanbara01

Mutations in HIV ENV that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
V 286 Y	CAG→ACG	F/BI	T140	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 21-fold cross-resistant to T140. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
V 286 Y	CAG→ACG	F/BI	vMIP-II	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 42-fold cross-resistant to vMIP II. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
I 288 T	ATA→ACA	F/BI	ALX40–4C	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 145-fold cross-resistant to ALX40–4C. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
I 288 T	ATA→ACA	F/BI	AMD3100	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 15-fold cross-resistant to AMD3100. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01

Mutations in HIV ENV that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
I 288 T	ATA→ACA	F/BI	SDF-I	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 26-fold cross-resistant to SDF-1. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
I 288 T	ATA→ACA	F/BI	T134	Y	N			In vitro selected virus (p145 of HIV-INL4-3, contains mutations N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275 in V3 loop of gp120) 15-fold resistant to T134. Role of each mutation not confirmed by site	Kanbara01
I 288 T	ATA→ACA	F/BI	T140	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 21-fold cross-resistant to T140. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
I 288 T	ATA→ACA	F/BI	vMIP-II	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 42-fold cross-resistant to vMIP II. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
I 288 V	ATA→GTA	F/BI	JM-3100	Y	?				DeVreese96, DeVreese96a
I 288 V	ATA→GTC	F/BI	SDF-I α	Y	?		SDF-1 β : 1.5-fold; AMB2763: 3- fold.	106K/134N/145L/245I/269E/278H/ 288V/293D/364–367Deletion/387T: 15-fold.	Schols98

Mutations in HIV ENV that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
ins 290 T	ATA→GTC	F/BI	ALX40-4C	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 145-fold cross-resistant to ALX40-4C. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
ins 290 T	ATA→GTC	F/BI	AMD3100	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 15-fold cross-resistant to AMD3100. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
ins 290 T	ATA→GTC	F/BI	SDF-1	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 26-fold cross-resistant to SDF-1. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
ins 290 T	ATA→GTC	F/BI	T134	Y	N			In vitro selected virus (p145 of HIV-1NL4-3, contains mutations N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275 in V3 loop of gp120) 15-fold resistant to T134. Role of each mutation not confirmed by site	Kanbara01

Mutations in HIV ENV that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
ins 290 T	ATA→GTC	F/BI	T140	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 21-fold cross-resistant to T140. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
ins 290 T	ATA→GTC	F/BI	vMIP-II	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 42-fold cross-resistant to vMIP II. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
K 290 E	ATA→GTC	F/BI	ALX40-4C	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 145-fold cross-resistant to ALX40-4C. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
K 290 E	ATA→GTC	F/BI	AMD3100	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 15-fold cross-resistant to AMD3100. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01

Mutations in HIV ENV that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
K 290 E	ATA→GTC	F/BI	SDF-I	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 26-fold cross-resistant to SDF-1. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
K 290 E	ATA→GTC	F/BI	T134	Y	N			In vitro selected virus (p145 of HIV-1NL4-3, contains mutations N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275 in V3 loop of gp120) 15-fold resistant to T134. Role of each mutation not confirmed by site	Kanbara01
K 290 E	ATA→GTC	F/BI	T140	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 21-fold cross-resistant to T140. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
K 290 E	ATA→GTC	F/BI	vMIP-II	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 42-fold cross-resistant to vMIP II. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01

Mutations in HIV ENV that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
N 293 D	AAT→GAT	F/BI	ALX40-4C	Y	N			TI134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 145-fold cross-resistant to ALX40-4C. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
N 293 D	AAT→GAT	F/BI	AMD3100	Y	N			TI134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 15-fold cross-resistant to AMD3100. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
N 293 D	AAT→GAT	F/BI	DS (Dextran sulphate)	Y	?			V3 loop region; S113N/S134N/K269E/Q278E/N293D/N323S/R387I: 250-fold	Este97, Este96a
N 293 D	AAT→GAT	F/BI	SDF-I	Y	N			TI134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 26-fold cross-resistant to SDF-I. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
N 293 D	AAT→GAT	F/BI	SDF-1 α	Y	?		SDF-1 β : 15-fold; AMB2763: 3-fold.	106K/134N/145L/245I/269E/278H/288V/293D/364-367Deletion/387T: 15-fold.	Schols98

Mutations in HIV ENV that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
N 293 D	AAT→GAT	F/BI	T134	Y	N			In vitro selected virus (p145 of HIV-1NL4-3, contains mutations N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275 in V3 loop of gp120) 15-fold resistant to T134. Role of each mutation not confirmed by site	Kanbara01
N 293 D	AAT→GAT	F/BI	T140	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 21-fold cross-resistant to T140. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
N 293 D	AAT→GAT	F/BI	vMIP-II	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 42-fold cross-resistant to vMIP II. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
N 293 H	AAT→CAT	F/BI	JM-3100	Y	?				DeVreese96, DeVreese96a
M 294 I	AAT→CAT	F/BI	ALX40-4C	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 145-fold cross-resistant to ALX40-4C. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01

Mutations in HIV ENV that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
M 294 I	AAT→CAT	F/BI	AMD3100	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 15-fold cross-resistant to AMD3100. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
M 294 I	AAT→CAT	F/BI	SDF-I	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 26-fold cross-resistant to SDF-1. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
M 294 I	AAT→CAT	F/BI	T134	Y	N			In vitro selected virus (p145 of HIV-1NL4–3, contains mutations N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275 in V3 loop of gp120) 15-fold resistant to T134. Role of each mutation not confirmed by site	Kanbara01
M 294 I	AAT→CAT	F/BI	T140	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 21-fold cross-resistant to T140. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01

Mutations in HIV ENV that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
M 294 I	AAT→CAT	F/BI	vMIP-II	Y	N			TI134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 42-fold cross-resistant to vMIP II. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
Q 296 K	AAT→CAT	F/BI	ALX40-4C	Y	N			TI134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 145-fold cross-resistant to ALX40-4C. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
Q 296 K	AAT→CAT	F/BI	AMD3100	Y	N			TI134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 15-fold cross-resistant to AMD3100. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
Q 296 K	AAT→CAT	F/BI	SDF-1	Y	N			TI134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274–275) 26-fold cross-resistant to SDF-1. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01

Mutations in HIV ENV that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
Q 296 K	AAT→CAT	F/BI	T134	Y	N			In vitro selected virus (p145 of HIV-INL4-3, contains mutations N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275 in V3 loop of gp120) 15-fold resistant to T134. Role of each mutation not confirmed by site	Kanbara01
Q 296 K	AAT→CAT	F/BI	T140	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 21-fold cross-resistant to T140. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
Q 296 K	AAT→CAT	F/BI	vMIP-II	Y	N			T134-resistant virus (selected in vitro, contains N269K, Q278T, R279K, A284V, F285L, V286Y, I288T, K290E, N293D, M294I, and Q296K; 290Tins, D274-275) 42-fold cross-resistant to vMIP II. Role of each mutation not confirmed by site-directed mutagenesis.	Kanbara01
A 297 T	GCA→ACA	F/BI	JM-2763	Y	?				DeVreese96, DeVreese96a
A 297 T	GCA→ACA	F/BI	JM-3100	Y	?				Este97, Este96a
N 323 S	AAT→AGT	F/BI	DS (Dextran sulphate)	Y	?			C3 region; S113N/S134N/K269E/Q278E/N293D/N323S/R387I: 250-fold	Lin96
G 332 E	GGA→GAA	F/BI	Siamycin I	Y	?			N188K/G332E/N351D/A550T/N633D/L762S: 9-fold	Lin96
N 351 D	AAT→GAT	F/BI	Siamycin I	Y	?			N188K/G332E/N351D/A550T/N633D/L762S: 9-fold	Lin96
P 385 L	CCA→CTA	F/BI	JM-2763	Y	?				DeVreese96, DeVreese96a
P 385 L	CCA→CTA	F/BI	JM-3100	Y	?				DeVreese96, DeVreese96a

Mutations in HIV ENV that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
R 387 I	AGA→ACA	F/BI	DS (Dextran sulphate)	Y	?	?		CD4 binding region; S113N/S134N/K269E/Q278E/N293D/N323S/R387I: 250-fold	Este97, Este96a
R 387 T	AGA→ACA	F/BI	SDF-1 α	Y	?	?	SDF-1 β : 15-fold; 106K/134N/145L/245I/269E/278H/AMB2763: 3-288V/293D/364-367Deletion/387T: 15-fold.		Schols98
Q 410 E	CAA→GAA	F/BI	JM-3100	Y	?	?			DeVreese96, DeVreese96a
S 433 P	TCC→CCC	F/BI	JM-3100	Y	?	?			DeVreese96, DeVreese96a
V 457 I	GTA→ATA	F/BI	JM-3100	Y	?	?			DeVreese96, DeVreese96a
A 550 T	GCC→ACC	F/BI	Siamycin I	Y	?	?		N188K/G332E/N351D/A550T/N633D/L762S: 9-fold	Lin96
N 633 D	AAT→GAT	F/BI	Siamycin I	Y	?	?		N188K/G332E/N351D/A550T/N633D/L762S: 9-fold	Lin96
L 762 S	TTG→TCG	F/BI	Siamycin I	Y	?	?		N188K/G332E/N351D/A550T/N633D/L762S: 9-fold	Lin96
FNSTW 364-368	Deletion	F/BI	SDF-1 α	Y	?	?	SDF-1 β : 15-fold; 106K/134N/145L/245I/269E/278H/AMB2763: 3-288V/293D/364-367Deletion/387T: 15-fold.		Schols98

Mutations in HIV SIVRT that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	-Fold -resist	Cross-resist (-fold)	Comments	Refs
K 65 R	AAA→AGA	F/BI	PMPA (tenofovir)	?	Y	5	3TC (80); ddi; ddC; d4T; PMEA	K65R appears first, followed by N69S and I118V. Observed changes at N69S and I118V do not result in increased resistance.	VanRompay96, Cherrington96a, VanRompay97a
Q 151 M	CAG→ATG	F/BI	AZT (zidovudine)	?	Y	>100	ddi; ddC; d4T; 3TC		VanRompay97
M 184 V	ATG→GTG	F/BI	(-)-FTC (emtricitabine)	Y	?				Schinazi95

Mutations in HIV FIVRT that confer drug resistance, ordered by position.

Amino Acid Change	Codon Change	Drug Class	Compound	In vitro	In vivo	In vivo -resist	-Fold	Cross-resist (-fold)	Comments	Refs
D 3 H	GAT→CAT	FIV NRTI	ddC (zalcitabine)	Y	?	4		ddI; PFA		Medlin96, Zhu96
V 47 I	GTA→ATA	FIV NRTI	d4T (stavudine)	Y	?	4-6		PFA (>50); AZT; ddI; PMEA		Smith96
P 156 S	CCA→TCA	FIV NRTI	3TC (lamivudine)	Y	?	7		AZT (4), AZT + 3TC (6)		Smith98
M 183 T	ATG→ACG	FIV NRTI	(-)-FTC (emtricitabine)	Y	?	10		ddC	Corresponds to 184 in HIV; M183V recombinant displays 10-fold resistance to 3TC or (-)-FTC.	Smith97

Abbreviations used in tables

Amino acids

A	alanine
C	cysteine
D	aspartate
E	glutamate
F	phenylalanine
G	glycine
H	histidine
I	isoleucine
K	lysine
L	leucine
M	methionine
N	asparagine
P	proline
Q	glutamine
R	arginine
S	serine
T	threonine
V	valine
W	tryptophan
Y	tyrosine

Drug class

F/BI	Fusion/Binding Inhibitor
II	Integrase Inhibitor
MN	Multiple Nucleoside
NRTI	Nucleoside Reverse Transcriptase Inhibitor
NNRTI	HIV-1 Specific Nonnucleoside RT Inhibitor
PI	Protease Inhibitor
PARTI	Pyrophosphate Analogue RTI
SIV RTI	SIV Nucleoside RTI

Compounds

Compound	Other Names (Company)	Chemical Name or Description
1737		Tetrahydronaphthalene lignan derivative
(-)-dOTC	BCH-10652	(-)-2'-deoxy-3'-oxa-4'-thiocytidine
(-)-dOTFC		(-)-2'-deoxy-3'-oxa-4'-thio-5-fluorocytidine
(-)-FTC	Emtricitabine, Coviracil (Triangle Pharmaceuticals)	(-)-(2 <i>R</i> ,5 <i>S</i>)-5-fluoro-1-[2-(hydroxymethyl)-1,3-oxathiolan-5-yl]cytosine
(+)-dOTC		(+)-2'-deoxy-3'-oxa-4'-thiocytidine
(+)-dOTFC		(+)-2'-deoxy-3'-oxa-4'-thio-5-fluorocytidine
1592U89	Abacavir, Ziagen (Glaxo Wellcome)	(1 <i>S</i> ,4 <i>R</i>)-4-[2-amino-6-cyclopropyl-amino]-9 <i>H</i> -purin-9-yl]-2-cyclopentene-1-methanol succinate
3TC	(-)-BCH-189, Lamivudine, Efavir (Glaxo Wellcome)	(-)- β -L-2',3'-dideoxy-3'-thiacytidine
8-chloro-TIBO	RO91767, R86183, tivrapipe	(+)-(S)-4,5,6,7-Tetrahydro-8-chloro-5-methyl-6-(3-methyl-2-butenyl)imidazol[4,5,1-jk][1,4]benzodiazepine
A-77003	C2 symmetry-based protease inhibitor (Abbott)	2PyridCH ₂ NCH ₃ CO-Val-NHCH(Bz)]CHOHCHOH
AAP-BHAP	U-104489 (Pharmacia & Upjohn)	1-[(5-Methanesulfonamidoindol-2-yl)carbonyl]-4-[N-ethyl-N-[3-(1,1-dimethylamino)-2-pyridinyl]amino]piperidine
ABT-378	Aluviran, Lopinavir (Abbott)	N-[(1 <i>S</i> ,3 <i>S</i> ,4 <i>S</i>)-4-[[2,6-dimethylphenoxy]acetyl]amino]-3-hydroxy-5-phenyl-1-(phenylmethyl)pentyl]tetrahydro- α -(1-methylethyl)-2-oxo-1(2 <i>H</i>)-pyrimidineacetamide
ABT-538	Ritonovir, Norvir (Abbott)	10-Hydroxy-2-methyl-5-(1-methylethyl)-1-[2-(1-methylethyl)-4-thiazolyl]-3,6-dioxo-8,11-bis(phenylmethyl)-2,4,7,12-tetraazatridecan-13-oic acid, 5-thiazolylmethyl ester

Abbreviations (cont)

Compounds (cont)

ADAMII		Methyl 3',3''-dichloro-4',4''-dimethoxy-5',5''-
AG1343	Nelfinavir, Viracept (Agouron)	bis(methoxycarbonyl)-6,6-diphenyl-5-hexenoate (3 <i>S</i> ,4 <i>aS</i> ,8 <i>aS</i>)- <i>N</i> -tert-Butyl-2-[(2 <i>R</i> ,3 <i>R</i>)-3-(3,2-cresotamido)- 2-hydroxy-4-(phenylthio)butyl]decahydro-3-isoquinoline- carboxamide monomethanesulfonate
ALX40-4C		a polypeptide of nine d-Arg residues
AMD3100		octahydrochloride dihydrate of 1,19-[1,4-phenylene-bis- (methylene)]-bis-1,4,8,11-tetra-azacyclotetradecane
AZT	zidovudine (Glaxo Wellcome)	3'-azido-3'-deoxythymidine
BHAP U-87201E	Atevirdine (Pharmacia Upjohn)	1-[(5-Methoxyindol-2-yl)carbonyl]-4-[3-(ethylamino)-2- pyridyl]piperazine
BHAP U-88204E		1-(Indolyl-2-carbonyl)-4-[3-[(1-methylethyl)amino]pyridyl- piperazine
BHAP U-90152	Delavirdine, Rescriptor (Pharmacia Upjohn)	1-(5-Methanesulphonamido)-1 <i>H</i> -indol-2-yl-carbonyl)-4-[3- (isopropylamino)-2-pyridinyl]piperazine
BHAP U-90153		bisheteroaryl piperidinyl derivative
BHAP U-90154		bisheteroaryl piperidinyl derivative
BHAP U-90155		bisheteroaryl piperidinyl derivative
BILA 1906 BS	(Bio-Mega/Boehringer Ingelheim)	<i>N</i> -{1 <i>S</i> -[[[3-[2 <i>S</i> -(1,1-dimethylethyl)amino]carbonyl-4 <i>R</i> - 3-pyridinylmethyl)thio]-1-piperidinyl]-2 <i>R</i> -hydroxy-1 <i>S</i> - (phenylmethyl)propyl]amino]carbonyl}-2-methylpropyl}-2- quinolinecarboxamide
BILA 2011	Palinavir (Bio- Mega/Boehringer Ingelheim)	<i>N</i> -{1 <i>S</i> -[[[3-[2 <i>S</i> -(1,1-dimethylethyl)amino]carbonyl]-4 <i>R</i> - [4-pyridinylmethyl)oxy]-1-piperidinyl]-2 <i>R</i> -hydroxy-1 <i>S</i> - (phenylmethyl)propyl]amino]carbonyl}-2-methylpropyl}-2- quinolinecarboxamide
BILA 2185 BS	(Bio-Mega/Boehringer Ingelheim)	<i>N</i> -(1,1-dimethylethyl)-1-[2 <i>S</i> -[[2-2,6-dimethylphenoxy)-1- oxoethyl]amino]-2 <i>R</i> -hydroxy-4-phenylbutyl]4 <i>R</i> -pyridinylthio)- 2-piperidine-carboxamide
BI-RG-587	Nevaripine, Viramune (Boehringer Ingelheim)	11-Cyclopropyl-4-methyl-5,11-dihydro-6 <i>H</i> -dipyrido[3,2- <i>b</i> :2',3'- e]-[1,4]diazepin-6-one
BM+51.0836		thiazolo-isoindolinone derivative
BMS 186318	(Bristol-Myers Squibb)	[1 <i>S</i> -[1 <i>R</i> *,2 <i>S</i> *(2 <i>S</i> *,3 <i>R</i> *)]]-[3-[[3-[[(1,1-Dimethylethoxy)- carbonyl]amino]-2-hydroxy-4-[4-[2-(4-morpholinyl)- 2-oxoethoxy]phenyl]butyl]amino]-2-hydroxy-1- (phenylmethyl)propyl]carbamic Acid, 1,1-dimethylethyl-ester
BMS 232632		azapeptide protease inhibitor
Calanolide A	NSC675451	a dipyrano-coumarin
d4T	Stavudine, Zerit (Bristol- Myers Squibb)	2',3'-didehydro-3'-deoxythymidine
ddC	Zalcitabine, Hivid (Roche)	2',3'-dideoxycytidine
ddI	Didanosine, Videx (Bristol-Myers Squibb)	2',3'-dideoxyinosine
DMP-266	Efavirenze, Sustiva (Dupont Merck)	(-)-6-Chloro-4-cyclopropylethynyl-4-trifluoromethyl-1,4,- dihydro-2 <i>H</i> -3,1-benzoxazin-one
DMP-323	XM-323 (Dupont Merck)	[4 <i>R</i> -(4- α ,5- α ,6- β ,7- β)]-hexahydro-5,6-dihydroxy-1,3-bis[(4- hydroxymethyl)phenyl]methyl]-4,7-bis(phenylmethyl)-2 <i>H</i> -1,3- diazepin-2-one

Abbreviations (cont)

Compounds (cont)

DMP-450	(Avid Therapeutics)	[4 <i>R</i> -(4- α ,5- α ,6- β ,7- β)]-hexahydro-5,6-bis(hydroxy)-1,3-bis(3-amino)phenyl]methyl)-4,7-bis(phenylmethyl)-2 <i>H</i> -1,3-diazepin-2-onebismesylate
DS		dextran sulfate
DXG	(-)- β -dioxolane-G	(-)-(2 <i>R</i> ,4 <i>R</i>)-9-[2-(Hydroxymethyl)-1,3-dioxolan-4-yl]guanine
E-BPTU	NSC 648400	1-benzyloxymethyl-5-ethyl-6-(2-pyridylthio)uracil
EBU-dM		5-ethyl-1-ethoxymethyl-6-(3,5-dimethylbenzyl)uracil
E-EBU		5-ethyl-1-ethoxymethyl-6-benzyluracil
E-EPSeU		1-(ethoxymethyl)-(6-phenylselenyl)-5-ethyluracil
E-EPU		1-(ethoxymethyl)-(6-phenyl-thio)-5-ethyluracil
F-ddA	Lodenosine	2'-fluoro-2',3'-dideoxyadenosine
GW420867X		S-3-ethyl-6-fluoro-4-isopropoxycarbonyl-3,4-dihydro-quinoxalin-2(1 <i>H</i>)-one
HBY 097		(<i>S</i>)-4-isopropoxycarbonyl-6-methoxy-3-(methylthio-methyl)-3,4-dihydroquinoxalin-2(1 <i>H</i>)-thione
HEPT		1-[(2-hydroxyethoxy)methyl]6-(phenylthio)thymine
IC9564	Betulinic acid derivative	4 <i>S</i> -[8-(28 betuliniyl) amino-octanoylamino]-3 <i>R</i> -hydroxy-6-methylheptanoic acid
I-EBU	MKC-442, emivirine, coactinon (Triangle Pharmaceuticals)	6-benzyl-1-ethoxymethyl-5-isopropyluracil (I-EBU, Triangle Pharmaceuticals/
JE-2147		an allophenylnorstatine-containing dipeptide protease inhibitor
JM-2763	(Johnson Matthey)	1,10-(1,3-propanediyl)-bis-1,4,8,11-tetraazacyclo-tetradecane
JM-3100	SID791 (Johnson Matthey)	1,10-[1,4-phenylenebis-(methylene)]bis-(1,4,8,11-tetraazacyclotetradecane)octahydrochloride dihydrate
KNI-272	Kynostatin 272	(2 <i>S</i> ,3 <i>S</i>)-3-amino-2-hydroxy-4-phenylbutyric acid-containing tripeptide
L-697,593		5-ethyl-6-methyl-3-(2-phthalimido-ethyl)pyridin-2(1 <i>H</i>)-one
L-697,661		3-[-(4,7-dichloro-1,3-benzoxazol-2-yl)methyl]amino-5-ethyl-6-methylpyridin-2(1 <i>H</i>)-one
L-Chicoric acid		[<i>S</i> -(<i>R</i> *, <i>R</i> *)]-2,3-Bis[[3-(3,4-dihydroxyphenyl)-1-oxo-2-propenyl]oxy]butanedioic acid
L-FddC		(-)- β -L-5-fluoro-2',3'-dideoxy-cytidine
LY-300046 HCl	Trovirdine (Lilly/Medivir/Abbott)	N-[2-(2-pyridylethyl)- <i>N</i> '-[2-(5-bromopyridyl)thiourea,hydrochloride
MK-639	Indinavir, Crixivan, L-735,524 (Merck)	[1(1 <i>S</i> ,2 <i>R</i>),5(<i>S</i>)]-2,3,5-Trideoxy-N-(2,3-dihydro-2-hydroxy-1 <i>H</i> -inden-1-yl)-5-[2-[[1,1-dimethylethyl]amino]carbonyl]-4-(3-pyridinylmethyl)-1-piperazinyl]-2-(phenylmethyl)-D-erythro-pentonamide sulfate
MP-134		C2 symmetry-based protease inhibitor
P9941	(Dupont Merck)	[2-pyridylacetyl-IlePheAla-y(CHOH)] ₂
PFA	Foscarnat (Astra)	phosphonofornate
PMEA	(Gilead Sciences)	9-(2-phosphonylmethoxyethyl)adenine
PMPA	(Gilead Sciences)	(<i>R</i>)-9-(2-phosphonyl-methoxypropyl)adenine
PNU-140690	Tipranavir, U-140690 (Pharmacia & Upjohn)	(6 <i>R</i>)-3-(1 <i>R</i>)-1-[3-([Trifluoromethyl](2-pyridyl)]sulfonylamino-phenyl]propyl-4-hydroxy-6-(2-phenylethyl)-6-propyl-5,6-dihydro-2 <i>H</i> -pyran-2-one

Abbreviations (cont)

Compounds (cont)

QM96521		1,1,3-trioxo-2H,4H-thieno[2,4-3][1,2,4]thiadiazine derivative (TTD)
QYL-685		methylenecyclopropane nucleoside analog with a phenylphosphoralaninate moiety
Ro 31-8959	Saquinavir, Invirase, Fortovase (Roche)	N(1)-[3-[3-[[[(1,1-dimethylethyl)amino]carbonyl]octahydro-2(1H)-isoquinolinyl]-2-hydroxy-1-(phenylmethyl)propyl]-2-[(2-quinolinylcarbonyl)amino]-, [3S-[2[1R*(R*),2S*],3 α ,4 α , β ,8 α , β]]-, monomethanesulfonate
RPI-312		1-[(3S)-3-(n-alpha-benzyloxycarbonyl)-1-asparginyl]-amino-2-hydroxy-4-phenyl-butyl]-n-tert-butyl-L-proline amide (peptidyl protease inhibitor)
RPR103611		a triterpene betulinic acid derivative
S-1153		5-(3,5-dichlorophenyl)thio-4-isopropyl-1-(4-pyridyl)methyl-1Himidazol-2-yl methyl carbamate
S-2720		6-chloro-3,3-dimethyl-4-(isopropenyl-oxycarbonyl)-3,4-dihydroquinoxalin-2(1H)thione
SC-52151	Telnavir	N-tert-butyl-N'-isobutyl-N'-[2(R)-hydroxy-4-phenyl-3(S)-[4-amino-1,4-dioxo-2(S)-(2-quinolinylcarboxamido)butyl-amino]butyl]urea
SC-55389A	(Searle)	hydroxyethyl-urea isostere protease inhibitor
SDF-1		Stromal derived factor 1
SDF-1 α		Stromal cell-derived factor 1 α
Siamycin I		21-residue tricyclic peptide
SKF108842		protease inhibitor
T134		[Tyr5,12, Lys7]-polyphemusin II-derivative with amino acid sequence R-R-W-C-Y-R-K-DK-P-Y-R-Ci-C-R-COOH
T140		[Tyr5,12, Lys7]-polyphemusin II-derivative
T20	DP-178, Pentafuside (Trimeris)	Ac-YTSLIHSLEESQNQQEKNEQELLELDKWASLWNWF-NH ₂
TIBO R82150	(Janssen)	(+)-(5S)-4,5,6,7-tetrahydro-5-methyl-6-(3-methyl-2-butenyl)-imidazo[4,5,1-jk][1,4]-benzodiazepin-2(1H)-thione
TIBO R82913	(Janssen)	(+)-(5S)-4,5,6,7-tetrahydro-9-chloro-5-methyl-6-(3-methyl-2-butenyl)-imidazo-[4,5,1-jk]-[1,4]benzo-diazepin-2(1H)-thione
TSAO		[2',5'-bis-O-(tert-butyl dimethylsilyl)-3'-spiro-5''-(4''-amino-1'',2''-oxathiole-2'',2''-dioxide)]- β -D-pentofuranosyl derivative
UC-10	NSC 645129 (Uniroyal Chemical Co)	thiocarboxanilide derivative
UC-16	(Uniroyal Chemical Co)	thiocarboxanilide derivative
UC-32	NSC 645542 (Uniroyal Chemical Co)	thiocarboxanilide derivative
UC-38	NSC 629243 (Uniroyal Chemical Co)	4-chloro-3-(isopropoxycarbonyl)phenylcarbamothioic acid, O-isopropyl ester
UC-42	(Uniroyal Chemical Co)	thiocarboxanilide derivative
UC-57	NSC 647014 (Uniroyal Chemical Co)	thiocarboxanilide derivative

Abbreviations (cont)**Compounds (cont)**

UC-68	NSC 638532 (Uniroyal Chemical Co)	thiocarboxanilide derivative
UC-69	NSC 646989 (Uniroyal Chemical Co)	thiocarboxanilide derivative
UC-70	NSC 638534 (Uniroyal Chemical Co)	thiocarboxanilide derivative
UC-781	(Uniroyal Chemical Co)	N-[4-chloro-3-(3-methyl-2-butenyloxy)phenyl]-2-methyl-3-furancarbothioamide
UC-80	NSC 639475 (Uniroyal Chemical Co)	thiocarboxanilide derivative
UC-81	NSC 615727 (Uniroyal Chemical Co)	thiocarboxanilide derivative
UC-82	(Uniroyal Chemical Co)	N-[4-chloro-3-(3-methyl-2-butenyloxy)phenyl]-2-methyl-3-thiophenecarbothioamide
UC-84	NSC 615985 (Uniroyal Chemical Co)	thiocarboxanilide derivative
VB 11,328	(Vertex)	Carbamic acid, [3-[[4-methoxyphenyl)sulfonyl](cyclopentylmethyl)amino]-2-hydroxy-1-(phenylmethyl)propyl]-,tetrahydro-3-furanyl ester
vMIP-II		viral macrophage inflammatory protein II
VX-478	141W94, Amprenavir, Agenerase	Carbamic acid, ((1 <i>S</i> ,2 <i>R</i>)-3-(((4-aminophenyl)sulfonyl)(2-methylpropyl)amino)-2-hydroxy-1-(phenylmethyl)propyl)-, (3 <i>S</i>)-tetrahydro-3-furanyl ester
α -APA	R18893, loviride analog	(+)-2,6-Dichloro- α -[(2-acetyl-5-methylphenyl)amino]benzamide

- Bachelor00 Bachelor LT, Anton ED, Kudish P, Baker D, Bunville J, Krakowski K, Bolling L, Aujay M, Wang XV, Ellis D, Becker MF, Lasut AL, George HJ, Spalding DR, Hollis G, Abremski K., Human immunodeficiency virus type 1 mutations selected in patients failing efavirenz combination therapy., *Antimicrobial Agents and Chemotherapy*, **44**, 2475–2484, 2000, Medline: 10952598
- Balzarini93 J. Balzarini, A. Karlsson, E. De Clercq, Human immunodeficiency virus type 1 drug-resistance patterns with different 1-[(2-hydroxyethoxy)methyl]-6-(phenylthio)thymine derivatives., *Mol Pharmacol*, **44**, 694–701, 1993, Medline: 94049697
- Balzarini93a J. Balzarini, S. Velazquez, A. San-Felix, A. Karlsson, M. J. Perez-Perez, M. J. Camarasa, E. De Clercq, Human immunodeficiency virus type 1-specific [2',5'-bis-O-(tert-butylidimethylsilyl)-beta-D-ribofuranosyl]-3'-spiro-5-(4-amino-1,2-oxathiole-2,2-dioxide)-purine analogues show a resistance spectrum that is different from that of the human immunodeficiency virus type 1- specific non-nucleoside analogues., *Mol Pharmacol*, **43**, 109–14, 1993, Medline: 93140699
- Balzarini93b J. Balzarini, A. Karlsson, E. De Clercq, J. Balzarini, A. Karlsson, A. M. Vandamme, M. J. Perez-Perez, H. Zhang, L. Vrang, B. Oberg, K. Backbro, T. Unge, A. San-Felix, et al, Human immunodeficiency virus type 1 (HIV-1) strains selected for resistance against the HIV-1-specific [2',5'-bis-O-(tert-butylidimethylsilyl)-3'-spiro-5''-(4''-amino-1'',2''-oxathiole-2'',2''-dioxide)]-beta-D-pentofurano syl (TSAO) nucleoside analogues retain sensitivity to HIV-1-specific nonnucleoside inhibitors., *Proc Natl Acad Sci U S A*, **90**, 6952–6, 1993, Medline: 93348190
- Balzarini93c J. Balzarini, A. Karlsson, M. J. Perez-Perez, L. Vrang, J. Walbers, H. Zhang, B. Oberg, A. M. Vandamme, M. J. Camarasa, E. De Clercq, HIV-1-specific reverse transcriptase inhibitors show differential activity against HIV-1 mutant strains containing different amino acid substitutions in the reverse transcriptase., *Virology*, **192**, 246–53, 1993, Medline: 93297111
- Balzarini93d J. Balzarini, A. Karlsson, M. J. Perez-Perez, M. J. Camarasa, W. G. Tarpley, E. De Clercq, Treatment of human immunodeficiency virus type 1 (HIV-1)-infected cells with combinations of HIV-1-specific inhibitors results in a different resistance pattern than does treatment with single-drug therapy., *J Virol*, **67**, 5353–9, 1993, Medline: 93353611
- Balzarini94 J. Balzarini, A. Karlsson, V. V. Sardana, E. A. Emini, M. J. Camarasa, E. De Clercq, Human immunodeficiency virus 1 (HIV-1)-specific reverse transcriptase (RT) inhibitors may suppress the replication of specific drug-resistant (E138K)RT HIV-1 mutants or select for highly resistant (Y181C- >C181I)RT HIV-1 mutants., *Proc Natl Acad Sci U S A*, **91**, 6599–603, 1994, Medline: 94294426
- Balzarini95 J. Balzarini, M. J. Perez-Perez, S. Velazquez, A. San-Felix, M. J. Camarasa, E. De Clercq, A. Karlsson, Suppression of the breakthrough of human immunodeficiency virus type 1 (HIV-1) in cell culture by thiocarboxanilide derivatives when used individually or in combination with other HIV-1-specific inhibitors (i.e., TSAO derivatives), *Proc Natl Acad Sci U S A*, **92**, 5470–4, 1995, Medline: 95296332
- Balzarini95a J. Balzarini, W. G. Brouwer, E. E. Felauer, E. De Clercq, A. Karlsson, Activity of various thiocarboxanilide derivatives against wild-type and several mutant human immunodeficiency virus type 1 strains., *Antiviral Res*, **27**, 219–36, 1995, Medline: 96145332
- Balzarini95b J. Balzarini, H. Jonckheere, W.A. Harrison, D.C. Dao, J. Anne, E. De Clercq, A. Karlsson, Oxathiin carboxanilide derivatives: a class of non-nucleoside HIV-1-specific reverse transcriptase inhibitors (NNRTIs) that are active against mutant HIV-1 strains resistant to other NNRTIs, *Antiviral Chemistry and Chemotherapy*, **6**, 169–78
- Balzarini96a J. Balzarini, H. Pelemans, S. Aquaro, C. F. Perno, M. Witvrouw, D. Schols, E. De Clercq, A. Karlsson, Highly favorable antiviral activity and resistance profile of the novel thiocarboxanilide pentenyloxy ether derivatives UC-781 and UC-82 as inhibitors of human immunodeficiency virus type 1 replication., *Mol Pharmacol*, **50**, 394–401, 1996, Medline: 96319790
- Balzarini96b J. Balzarini, W. G. Brouwer, D. C. Dao, E. M. Osika, E. De Clercq, Identification of novel thiocarboxanilide derivatives that suppress a variety of drug-resistant mutant human immunodeficiency virus type 1 strains at a potency similar to that for wild-type virus., *Antimicrob Agents Chemother*, **40**, 1454–66, 1996, Medline: 96338367
- Balzarini96c Balzarini J, Pelemans H, Perez-Perez MJ, San-Felix A, Camarasa MJ, De Clercq E, Karlsson A, Marked inhibitory activity of non-nucleoside reverse transcriptase inhibitors against

- human immunodeficiency virus type 1 when combined with (-)2',3'-dideoxy-3'-thiacytidine. , *Mol Pharmacol* , **49**(5), 882–90, 1996, Medline: 96212950
- Balzarini98 Balzarini J, Pelemans H, Esnouf R, De Clercq E., A novel mutation (F227L) arises in the reverse transcriptase of human immunodeficiency virus type 1 on dose-escalating treatment of HIV type 1-infected cell cultures with the nonnucleoside reverse transcriptase inhibitor thiocarbozanilide UC-781., *AIDS Res Hum Retroviruses*, **14**(3), 255–260, 1998, Medline: 98150880
- Borman95 A.M. Borman, S. Paulous, Clavel F, Continued accumulation of protease inhibitor resistance mutations in culture in the absence of the drug, *Fourth International Workshop on HIV Drug Resistance Sardinia, Italy*
- BorrotoEsoda97 Borroto-K. Esoda, D.S. Noel, C.P. Moxham, Furman P.A., Preliminary genotypic analysis of HIV-1 in plasma from volunteers receiving repeated multiple doses of MKC-442, *Sixth International Workshop on HIV Drug Resistance St. Petersburg, FL, USA*
- Brenner02 Brenner B, Turner D, Oliveira M, Moisi D, Detorio M, Carobene M, Marlink RG, Schapiro J, Roger M, Wainberg MA, A V106M mutation in HIV-1 clade C viruses exposed to efavirenz confers cross-resistance to non-nucleoside reverse transcriptase inhibitors, *AIDS*, **17**(1), F1–5, 2002, Medline: 12478089
- Brown00 Brown AJ, Precious HM, Whitcomb JM, Wong JK, Quigg M, Huang W, Daar ES, D'Aquila RT, Keiser PH, Connick E, Hellmann NS, Petropoulos CJ, Richman DD, Little SJ., Reduced susceptibility of human immunodeficiency virus type 1 (HIV-1) from patients with primary HIV infection to nonnucleoside reverse transcriptase inhibitors is associated with variation at novel amino acid sites., *Journal of Virology*, **74**, 10269–10273, 2000, Medline: 11044070
- Buckheit95a R. W. Buckheit, T. L. Kinjerski, V. Fliakas-Boltz, J. D. Russell, T. L. Stup, L. A. Pallansch, W. G. Brouwer, D. C. Dao, W. A. Harrison, R. J. Schultz, et al, Structure-activity and cross-resistance evaluations of a series of human immunodeficiency virus type-1-specific compounds related to oxathiin carboxanilide., *Antimicrob Agents Chemother*, **39**, 2718–27, 1995, Medline: 96161287
- Buckheit95b R. W. Buckheit, V. Fliakas-Boltz, W. D. Decker, J. L. Roberson, T. L. Stup, C. A. Pyle, E. L. White, J. B. McMahon, M. J. Currens, M. R. Boyd, et al, Comparative anti-HIV evaluation of diverse HIV-1-specific reverse transcriptase inhibitor-resistant virus isolates demonstrates the existence of distinct phenotypic subgroups., *Antiviral Res*, **26**, 117–32, 1995, Medline: 95328856
- Buckheit95c Buckheit RW Jr, Fliakas-Boltz V, Yeagy-Bargo S, Weislow O, Mayers DL, Boyer PL, Hughes SH, Pan BC, Chu SH, Bader JP, Resistance to 1-[(2-hydroxyethoxy)methyl]-6-(phenylthio)thymine derivatives is generated by mutations at multiple sites in the HIV-1 reverse transcriptase, *Virology*, **210**, 186–193, 1995, Medline: 95313352
- Buckheit97 Buckheit RW Jr, Snow MJ, Fliakas-Boltz V, Kinjerski TL, Russell JD, Pallansch LA, Brouwer WG, Yang SS, Highly potent oxathiin carboxanilide derivatives with efficacy against nonnucleoside reverse transcriptase inhibitor-resistant human immunodeficiency virus isolates. , *Antimicrob Agents Chemother*, **41**, 831–837, 1997, Medline: 97242500
- Byrnes93 V. W. Byrnes, V. V. Sardana, W. A. Schleif, J. H. Condra, J. A. Waterbury, J. A. Wolfgang, W. J. Long, C. L. Schneider, A. J. Schlabach, B. S. Wolanskii, Comprehensive mutant enzyme and viral variant assessment of human immunodeficiency virus type 1 reverse transcriptase resistance to nonnucleoside inhibitors., *Antimicrob Agents Chemother*, **37**, 1576–9, 1993, Medline: 94028780
- Byrnes93a V. Byrnes, O. Blahy, J. Condra, L. Gotlib, D. Graham, W. Long, J. Quintero, A. Rhodes, E. Roth, V. Sardana, A. Schlabach, W. Schleif, C. Schneider, D. Titus, B. Wolanski, J. Wolfgang, E. Emini, Phenotypic susceptibility of human immunodeficiency virus type 1 RT containing substitutions which engender resistance to nucleoside and non-nucleoside inhibitors, *Third Workshop on Viral Resistance Gaithersburg, MD, USA*
- Caride00 Caride E, Brindeiro R, Hertogs K, Larder B, Dehertogh P, Machado E, de Sa CA, Eyer-Silva WA, Sion FS, Passioni LF, Menezes JA, Calazans AR, Tanuri A., Drug-resistant reverse transcriptase genotyping and phenotyping of B and non-B subtypes (F and A) of human immunodeficiency virus type I found in Brazilian patients failing HAART., *Virology*, **275**, 107–115, 2000, Medline: 11017792
- Carrillo98 Carrillo A, Stewart KD, Norbeck DW, Kohlbrenner WE, Leonard JM, Kempf DJ, Molla A., In vitro selection and characterization of human immunodeficiency virus type 1 variants with

- increased resistance to ABT-378, a novel protease inhibitor, *J Virol*, **72**, 7532–7541, 1998, Medline: 98362159
- Cherrington96 J. M. Cherrington, A. S. Mulato, M. D. Fuller, M. S. Chen, Novel mutation (K70E) in human immunodeficiency virus type 1 reverse transcriptase confers decreased susceptibility to 9-[2-(phosphonomethoxy)ethyl]adenine in vitro., *Antimicrob Agents Chemother*, **40**, 2212–6, 1996, Medline: 97032863
- Cherrington96a J.M. Cherrington, K.K.A. Van Rompay, A.S. Mulato, M.L. Marthas, C.J. Berardi, S. Telm, N. Bischofberger, N.C. Pedersen, Phenotypic and genotypic characterization of simian immunodeficiency viruses (SIV) with reduced susceptibility to PMPA isolated after PMPA therapy, *Fifth International Workshop on HIV Drug Resistance Whistler, BC, Canada*
- Cherrington97 J.M. Cherrington, R. Chandok, A.S. Mulato, P.D. Lamy, H. Mitsuya, M. Wainberg, In vitro selection and characterization of HIV-1 variants with reduced susceptibility to PMPA, *Sixth International Workshop on HIV Drug Resistance, St. Petersburg, FL, USA*
- Condra95 J. H. Condra, W. A. Schleif, O. M. Blahy, L. J. Gabryelski, D. J. Graham, J. C. Quintero, A. Rhodes, H. L. Robbins, E. Roth, M. Shivaprakash, et al, In vivo emergence of HIV-1 variants resistant to multiple protease inhibitors, *Nature*, **374**, 569–71, 1995, Medline: 95214785
- Condra96 Condra JH, Holder DJ, Schleif WA, Blahy OM, Danovich RM, Gabryelski LJ, Graham DJ, Laird D, Quintero JC, Rhodes A, Robbins HL, Roth E, Shivaprakash M, Yang T, Chodakewitz JA, Deutsch PJ, Leavitt RY, Massari FE, Mellors JW, Squires KE, Steigbigel RT, Teppler H, Emini EA, Genetic correlates of in vivo viral resistance to indinavir, a human immunodeficiency virus type 1 protease inhibitor, *J Virol*, **70**(12), 8270–6, 1996, Medline: 97126022
- Croteau97 G. Croteau, L. Doyon, D. Thibeault, G. McKercher, L. Pilote, D. Lamarre, Impaired fitness of human immunodeficiency virus type 1 variants with high-level resistance to protease inhibitors., *J Virol*, **71**, 1089–96, 1997, Medline: 97151093
- Cushman98 Cushman M, Casimiro-Barcia A, Hejchman E, Ruell JA, Huang M, Schaeffer CA, Williamson K, Rice WG, Buckheit Jr. RW., New alkenyldiarylmethanes with enhanced potencies as anti-HIV agents which act as non-nucleoside reverse transcriptase inhibitors., *J Med Chem*, **41**, 2076–2089, 1998, Medline: 98285673
- DeAntoni97 A. De Antoni, A. Foli, J. Lisziewicz, F. Lori, Mutations in the pol gene of human immunodeficiency virus type 1 in infected patients receiving didanosine and hydroxyurea combination therapy, *J Infect Dis*, **176**, 899–903, 97, Medline: 97472322
- deBethune93 M-P. de Bethune, R. Pauwels, K. Andries, A.M. Vandamme, M. Peeters, R. Colebunders, P. Stoffels, E. De Clercq, J. Desmyter, AZT resistance reversal by the non-nucleoside reverse transcriptase inhibitor α -APA R18893 in a symptomatic HIV-infected individual, *Second HIV Drug Resistance Workshop, Noordwijk, The Netherlands*
- Demeter95 L.M. Demeter, R.W. Shafer, M. Para, G. Morse, W. Freimuth, T.C. Merigan, R.C. Reichman, Delavirdine (DLV) susceptibility of HIV-1 isolates obtained from patients receiving DLV monotherapy (ACTG 260), *J Acquir Immune Defic Syndrom Hum Retrovir*, **10**(S3), 23
- Demeter98 L. M. Demeter, P. M. Meehan, G. Morse, M. A. Fischl, M. Para, W. Powderly, J. Leedom, J. Holden-Wiltse, C. Greisberger, K. Wood, J. Timpone, L. K. Wathen, T. Nevin, L. Resnick, D. H. Batts, R. C. Reichman, Phase I study of atevirdine mesylate (U-87201E) monotherapy in HIV-1-infected patients, *J Acquir Immune Defic Syndr Hum Retrovir*, **19**, 135–44, 98, Medline: 98439558
- DeVreese96 K. De Vreese, D. Reymen, P. Griffin, A. Steinkasserer, G. Werner, G. J. Bridger, J. Este, W. James, G. W. Henson, J. Desmyter, J. Anne, I. De Clercq, The bicyclams, a new class of potent human immunodeficiency virus inhibitors, block viral entry after binding., *Antiviral Res*, **29**, 209–19, 1996, Medline: 96315998
- DeVreese96a de Vreese K, Kofler-Mongold V, Leutgeb C, Weber V, Vermeire K, Schacht S, Anne J, de Clercq E, Datema R, Werner G, The molecular target of bicyclams, potent inhibitors of human immunodeficiency virus replication, *J Virol*, **70**(2), 689–96, 1996, Medline: 96135175
- Doyon96 L. Doyon, G. Croteau, D. Thibeault, F. Poulin, L. Pilote, D. Lamarre, Second locus involved in human immunodeficiency virus type 1 resistance to protease inhibitors., *J Virol*, **70**, 3763–9, 1996, Medline: 96211509
- Dueweke93 T. J. Dueweke, T. Pushkarskaya, S. M. Poppe, S. M. Swaney, J. Q. Zhao, I. S. Chen, M. Stevenson, W. G. Tarpley, A mutation in reverse transcriptase of bis(heteroaryl)piperazine- resistant

- human immunodeficiency virus type 1 that confers increased sensitivity to other nonnucleoside inhibitors., *Proc Natl Acad Sci U S A*, **90**, 4713–7, 1993, Medline: 93281649
- Dulioust97 A. Dulioust, S. Paulous, L. Guillemot, F. Boue, P. Galanaud, Clavel F, Selection of saquinavir-resistant mutants by indinavir following a switch from saquinavir, *Sixth International Workshop on HIV Drug Resistance St. Petersburg, FL, USA*
- Eastman97 P.S. Eastman, I.B. Duncan, C. Gee, Race E, Acquisition of genotypic mutations associated with reduced susceptibility to protease inhibitors during saquinavir monotherapy, *Sixth International Workshop on HIV Drug Resistance St. Petersburg, FL, USA*
- Eberle95 J. Eberle, B. Bechowsky, D. Rose, U. Hauser, K. von der Helm, L. Gurtler, H. Nitschko, Resistance of HIV type 1 to proteinase inhibitor Ro 31–8959., *AIDS Res Hum Retroviruses*, **11**, 671–6, 1995, Medline: 96078227
- el-Farrash94 M. A. el-Farrash, M. J. Kuroda, T. Kitazaki, T. Masuda, K. Kato, M. Hatanaka, S. Harada, Generation and characterization of a human immunodeficiency virus type 1 (HIV-1) mutant resistant to an HIV-1 protease inhibitor., *J Virol*, **68**, 233–9, 1994, Medline: 94076412
- Este96 J. A. Este, K. De Vreese, M. Witvrouw, J. C. Schmit, A. M. Vandamme, J. Anne, J. Desmyter, G. W. Henson, G. Bridger, E. De Clercq, Antiviral activity of the bicyclam derivative JM3100 against drug-resistant strains of human immunodeficiency virus type 1., *Antiviral Res*, **29**, 297–307, 1996, Medline: 96316006
- Este96a J.A. Este, K. Van Laethem, A.M. Vandamme, J. Desmyter, E. De Clercq, Resistant phenotype of human immunodeficiency virus type 1 to dextran sulfate is conferred by specific amino acid substitutions in the gp120 molecule, *Fifth International Workshop on HIV Drug Resistance Whistler, BC, Canada*
- Este97 J.A. Este, D. Schols, K. De Vreese, D. Van Laethem, A.M. Vandamme, J. Desmyter, E. De Clercq, Development of resistance of human immunodeficiency virus type 1 to dextran sulfate associated with the emergence of specific mutations in the envelope gp120 glycoprotein, *Molecular Pharmacology*, **52**, 98–104, 1997, Medline: 97368051
- Fitzgibbon01 Fitzgibbon JE, Gaur S, Walsman SM, Janahi M, Whitley-Williams P, John JF Jr., Emergence of drug resistance mutations in a group of HIV-infected children taking nelfinavir-containing regimens., *AIDS Res Hum Retroviruses*, **17**(14), 1321–8, 2001, Medline: 11602042
- Fitzgibbon92 J. E. Fitzgibbon, R. M. Howell, C. A. Haberkzettel, S. J. Sperber, D. J. Gocke, D. T. Dubin, Human immunodeficiency virus type 1 pol gene mutations which cause decreased susceptibility to 2',3'-dideoxycytidine., *Antimicrob Agents Chemother*, **36**, 153–7, 1992, Medline: 92272541
- Foli96 A. Foli, K. M. Sogocio, B. Anderson, M. Kavlick, M. W. Saville, M. A. Wainberg, Z. Gu, J. M. Cherrington, H. Mitsuya, R. Yarchoan, In vitro selection and molecular characterization of human immunodeficiency virus type 1 with reduced sensitivity to 9-[2-(phosphonomethoxy)ethyl]adenine (PMEA)., *Antiviral Res*, **32**, 91–8, 1996, Medline: 97046247
- Fujiwara98 T. Fujiwara, A. Sato, M. el-Farrash, S. Miki, K. Abe, Y. Isaka, M. Kodama, Y. Wu, L. B. Chen, H. Harada, H. Sugimoto, M. Hatanaka, Y. Hinuma, S-1153 inhibits replication of known drug-resistant strains of human immunodeficiency virus type 1, *Antimicrob Agents Chemother*, **42**, 1340–5, 98, Medline: 98287568
- Gao92 Q. Gao, Z. X. Gu, M. A. Parniak, X. G. Li, M. A. Wainberg, In vitro selection of variants of human immunodeficiency virus type 1 resistant to 3'-azido-3'-deoxythymidine and 2',3'-dideoxyinosine., *J Virol*, **66**, 12–9, 1992, Medline: 92085373
- Gao93 Q. Gao, Z. Gu, M. A. Parniak, J. Cameron, N. Cammack, C. Boucher, M. A. Wainberg, The same mutation that encodes low-level human immunodeficiency virus type 1 resistance to 2',3'-dideoxyinosine and 2',3'-dideoxycytidine confers high-level resistance to the (-) enantiomer of 2',3'-dideoxy-3'-thiacytidine., *Antimicrob Agents Chemother*, **37**, 1390–2, 1993, Medline: 93319281
- Gatanaga02 Gatanaga H, Suzuki Y, Tsang H, Yoshimura K, Kavlick MF, Nagashima K, Gorelick RJ, Mardy S, Tang C, Summers MF, Mitsuya H, Amino acid substitutions in Gag protein at non-cleavage sites are indispensable for the development of a high multitude of HIV-1 resistance against protease inhibitors, *J Biol Chem*, **277**(8), 5952–61, 2002, Medline: 11741936

- Gong00 Gong YF, Robinson BS, Rose RE, Deminie C, Spicer TP, Stock D, Colonno RJ, Lin PF. In vitro resistance profile of the human immunodeficiency virus type 1 protease inhibitor BMS-232632. *Antimicrobial Agents and Chemotherapy*, **44**, 2319–2326, 2000, Medline: 10952574
- Gong99 Y. Gong, B. Robinson, R. Rose, K. Riccardi, C. Deminie, D. Stock, T. Spicer, F. Djang, J. Cross, R. Colonno, P-F. Lin. , Resistance profile and drug combination studies of an HIV-1 protease inhibitor BMS-232632. *6th Conference on Retroviruses and Opportunistic Infections Chicago, IL, USA*, Abstract 603
- Gu92 Z. Gu, Q. Gao, X. Li, M. A. Parniak, M. A. Wainberg, Novel mutation in the human immunodeficiency virus type 1 reverse transcriptase gene that encodes cross-resistance to 2',3'-dideoxyinosine and 2',3'-dideoxycytidine. *J Virol*, **66**, 7128–35, 1992, Medline: 93059660
- Gu94 Z. Gu, Q. Gao, H. Fang, H. Salomon, M. A. Parniak, E. Goldberg, J. Cameron, M. A. Wainberg, Identification of a mutation at codon 65 in the IKKK motif of reverse transcriptase that encodes human immunodeficiency virus resistance to 2',3'-dideoxycytidine and 2',3'-dideoxy-3'-thiacytidine. *Antimicrob Agents Chemother*, **38**, 275–81, 1994, Medline: 94250000
- Gu95 Z. Gu, H. Salomon, J. M. Cherrington, A. S. Mulato, M. S. Chen, R. Yarchoan, A. Folli, K. M. Sogocio, M. A. Wainberg, K65R mutation of human immunodeficiency virus type 1 reverse transcriptase encodes cross-resistance to 9-(2-phosphonylmethoxyethyl)adenine. *Antimicrob Agents Chemother*, **39**, 1888–91, 1995, Medline: 96100773
- Gulnik95 S. V. Gulnik, L. I. Suvorov, B. Liu, B. Yu, B. Anderson, H. Mitsuya, J. W. Erickson, Kinetic characterization and cross-resistance patterns of HIV-1 protease mutants selected under drug pressure. *Biochemistry*, **34**, 9282–7, 1995, Medline: 95352609
- Gurusinghe95 A. D. Gurusinghe, S. A. Land, C. Birch, C. McGavin, D. J. Hooker, G. Tachedjian, R. Doherty, N. J. Deacon, Reverse transcriptase mutations in sequential HIV-1 isolates in a patient with AIDS. *J Med Virol*, **46**, 238–43, 1995, Medline: 96028714
- Hara97 Hara H, Fujihashi T, Sakata T, Kaji A, Kaji H, Tetrahydronaphthalene lignan compounds as potent anti-HIV type 1 agents. *AIDS Res Hum Retroviruses*, **13**, 695–705, 1997, Medline: 97311521
- Harrigan96 P. R. Harrigan, I. Kinghorn, S. Bloor, S. D. Kemp, I. Najera, A. Kohli, B. A. Larder, Significance of amino acid variation at human immunodeficiency virus type 1 reverse transcriptase residue 210 for zidovudine susceptibility. *J Virol*, **70**, 5930–4, 1996, Medline: 96323108
- Hertogs00 Hertogs K, Bloor S, De Vroey V, van Den Eynde C, Dehertogh P, van Cauwenberge A, Sturmer M, Alcorn T, Wegner S, van Houtte M, Miller V, Larder BA. A novel human immunodeficiency virus type 1 reverse transcriptase mutational pattern confers phenotypic lamivudine resistance in the absence of mutation 184V. *Antimicrobial Agents and Chemotherapy*, **44**, 568–573, 2000, Medline: 10681319
- Ho94 D. D. Ho, T. Toyoshima, H. Mo, D. J. Kempf, D. Norbeck, C. M. Chen, N. E. Wideburg, S. K. Burt, J. W. Erickson, M. K. Singh, Characterization of human immunodeficiency virus type 1 variants with increased resistance to a C2-symmetric protease inhibitor. *J Virol*, **68**, 2016–20, 1994, Medline: 94149902
- Holz-Smith01 Holz-Smith SL, Sun IC, Jin L, Matthews TJ, Lee KH, Chen CH. Role of Human Immunodeficiency Virus (HIV) Type 1 Envelope in the Anti-HIV Activity of the Betulinic Acid Derivative IC9564. *Antimicrobial Agents and Chemotherapy*, **45**, 60–66, 2001, Medline: 11120945
- Hooker96 D. J. Hooker, G. Tachedjian, A. E. Solomon, A. D. Gurusinghe, S. Land, C. Birch, J. L. Anderson, B. M. Roy, E. Arnold, N. J. Deacon, An in vivo mutation from leucine to tryptophan at position 210 in human immunodeficiency virus type 1 reverse transcriptase contributes to high-level resistance to 3'-azido-3'-deoxythymidine. *J Virol*, **70**, 8010–8, 1996, Medline: 97048084
- Imamichi00a Imamichi T, Sinha T, Imamichi H, Zhang YM, Metcalf JA, Falloon J, Lane HC. High-level resistance to 3'-azido-3'-deoxythymidine due to a deletion in the reverse transcriptase gene of human immunodeficiency virus type 1. *Journal of Virology*, **74**, 1023–1028, 2000, Medline: 10623768
- Imamichi00b Imamichi T, Berg SC, Imamichi H, Lopez JC, Metcalf JA, Falloon J, Lane HC. Relative replication fitness of a high-level 3'-azido-3'-deoxythymidine-resistant variant of human immunodeficiency virus type 1 possessing an amino acid deletion at codon 67 and a novel substitution (Thr→Gly) at codon 69. *Journal of Virology*, **74**, 10958–10964, 2000, Medline: 11069990

- Imamichi01 Imamichi, T., Murphy, M.A., Imamichi, H., Lane, H.C., Amino Acid Deletion at Codon 67 and Thr-to-Gly Change at Codon 69 of Human Immunodeficiency Virus Type 1 Reverse Transcriptase Confer Novel Drug Resistance Profiles, *Journal of Virology*, **75**, 3988–3992
- Ingate95 S. Ingate, M. J. Perez-Perez, E. De Clercq, J. Balzarini, M. J. Camarasa, Synthesis and anti-HIV-1 activity of novel TSAO-T derivatives modified at the 2'- and 5'-positions of the sugar moiety., *Antiviral Res*, **27**, 281–99, 1995, Medline: 96145337
- Iversen96 A. K. Iversen, R. W. Shafer, K. Wehrly, M. A. Winters, J. I. Mullins, B. Chesebro, T. C. Merigan, Multidrug-resistant human immunodeficiency virus type 1 strains resulting from combination antiretroviral therapy., *J Virol*, **70**, 1086–90, 1996, Medline: 96135222
- Jacobsen94 H. Jacobsen, Brun-F. Vezinet, I. Duncan, M. Hanggi, M. Ott, S. Vella, J. Weber, Mous J, Genotypic characterization of HIV-1 from patients after prolonged treatment with proteinase inhibitor saquinavir, *Third International Workshop on HIV Drug Resistance Kauai, HI, USA*
- Kanbara01 Kanbara K, Sato S, Tanuma J, Tamamura H, Gotoh K, Yoshimori M, Kanamoto T, Kitano M, Fujii N, Nakashima H., Biological and genetic characterization of a human immunodeficiency virus strain resistant to CXCR4 antagonist T134., *AIDS Res Hum Retroviruses*, **17**(7), 615–22, 2001, Medline: 11375057
- Kaplan94 A. H. Kaplan, S. F. Michael, R. S. Wehbie, M. F. Knigge, D. A. Paul, L. Everitt, D. J. Kempf, D. W. Norbeck, J. W. Erickson, R. Swanstrom, Selection of multiple human immunodeficiency virus type 1 variants that encode viral proteases with decreased sensitivity to an inhibitor of the viral protease., *Proc Natl Acad Sci U S A*, **91**, 5597–601, 1994, Medline: 94261633
- Kellam92 P. Kellam, C. A. Boucher, B. A. Larder, Fifth mutation in human immunodeficiency virus type 1 reverse transcriptase contributes to the development of high-level resistance to zidovudine., *Proc Natl Acad Sci U S A*, **89**, 1934–8, 1992, Medline: 92179296
- Kemp98 S. D. Kemp, C. Shi, S. Bloor, P. R. Harrigan, J. W. Mellors, B. A. Larder, A novel polymorphism at codon 333 of human immunodeficiency virus type 1 reverse transcriptase can facilitate dual resistance to zidovudine and L-2',3'-dideoxy-3'-thiacytidine., *J Virol*, **72**, 5093–8, 1998, Medline: 98241751
- Kempf01 Kempf DJ, Isaacson JD, King MS, Brun SC, Xu Y, Real K, Bernstein BM, Japour AJ, Sun E, Rode RA., Identification of genotypic changes in human immunodeficiency virus protease that correlate with reduced susceptibility to the protease inhibitor lopinavir among viral isolates from protease inhibitor-experienced patients., *J Virol.*, **75**(16), 7462–9, 2001, Medline: 11462018
- Keulen96 W. Keulen, A. van Wijk, C. Boucher, B. Berkhout, Initial appearance of 184Ile variant in 3TC-treated patients can be explained by the mutation bias of the HIV-1 RT enzyme, *Fifth International Workshop on HIV Drug Resistance Whistler, BC, Canada*
- King95 R.W. King, S. Garber, D.L. Winslow, C. Reid, L.T. Bachelier, E. Anton, M.J. Otto, Multiple mutations in the human immunodeficiency virus protease gene are responsible for decreased susceptibility to protease inhibitors., *Antiviral Chemistry and Chemotherapy*, **669**(9), 80–88
- King98 P.J. King, W. E. Robinson Jr. , Resistance to the anti-human immunodeficiency virus type 1 compound L-chicoric acid results from a single mutation at amino acid 140 of integrase., *Journal of Virology*, **72**, 8420–8424
- Kleim93 J. P. Kleim, R. Bender, U. M. Billhardt, C. Meichsner, G. Riess, M. Rosner, I. Winkler, A. Paessens, Activity of a novel quinoxaline derivative against human immunodeficiency virus type 1 reverse transcriptase and viral replication., *Antimicrob Agents Chemother*, **37**, 1659–64, 1993, Medline: 94028795
- Kleim95 J. P. Kleim, R. Bender, R. Kirsch, C. Meichsner, A. Paessens, M. Rosner, H. Rubsamens-Waigmann, R. Kaiser, M. Wichers, K. E. Schneeweis, et al, Preclinical evaluation of HBY 097, a new nonnucleoside reverse transcriptase inhibitor of human immunodeficiency virus type 1 replication., *Antimicrob Agents Chemother*, **39**, 2253–7, 1995, Medline: 96109422
- Kleim96 J. P. Kleim, M. Rosner, I. Winkler, A. Paessens, R. Kirsch, Y. Hsiou, E. Arnold, G. Riess, Selective pressure of a quinoxaline nonnucleoside inhibitor of human immunodeficiency virus type 1 (HIV-1) reverse transcriptase (RT) on HIV-1 replication results in the emergence of nucleoside RT-inhibitor-specific (RT Leu-74→Val or Ile and Val-75→Leu or Ile) HIV-1 mutants., *Proc Natl Acad Sci U S A*, **93**, 34–8, 1996, Medline: 96133872

- Kleim97 J. P. Kleim, I. Winkler, M. Rosner, R. Kirsch, H. Rubsamen-Waigmann, A. Paessens, G. Riess, In vitro selection for different mutational patterns in the HIV-1 reverse transcriptase using high and low selective pressure of the nonnucleoside reverse transcriptase inhibitor HBY 097., *Virology*, **231**, 112–8, 1997, Medline: 97288331
- Kleim99 J-P. Kleim, V. Burt, M. Maguire, R. Ferris, R.J. Hazen, G. Roberts, M. St. Clair. , NNRTI GW420867X: Comparative evaluation of the in vitro resistance profile. , *6th Conference on Retroviruses and Opportunistic Infections, Chicago, IL, USA.* , Abstract 600
- Labrosse00 Labrosse B, Treboute C, Alizon M., Sensitivity to a nonpeptidic compound (RPR103611) blocking human immunodeficiency virus type 1 Env-mediated fusion depends on sequence and accessibility of the gp41 loop region., *Journal of Virology*, **74**(5), 2142–50, 2000, Medline: 10666243
- Labrosse97 B. Labrosse, O. Pleskoff, N. Sol, C. Jones, Y. Henin, M. Alizon, Antiviral and resistance studies of RPR103611, an inhibitor of HIV replication, *Sixth International Workshop on HIV Drug Resistance, St. Petersburg, FL, USA*
- Lacey94 S. F. Lacey, B. A. Larder, Novel mutation (V75T) in human immunodeficiency virus type 1 reverse transcriptase confers resistance to 2',3'-didehydro-2',3'- dideoxythymidine in cell culture., *Antimicrob Agents Chemother*, **38**, 1428–32, 1994, Medline: 94379807
- Lamarre94 D. Lamarre, G. Croteau, L. Pilote, P. Rousseau, Doyon L, Molecular characterization of HIV-1 variants resistant to specific viral protease inhibitors, *Third International Workshop on HIV Drug Resistance Kauai, HI, USA*
- Lamarre95 D. Lamarre, L. Doyon, G. Croteau, L. Pilote, Thibeault D, Molecular basis of HIV-1 resistance to protease inhibitors Structural flexibility of the protease and second-site compensatory mutations in cleavage sites, *Fourth International Workshop on HIV Drug Resistance Sardinia, Italy*
- Larder89 B. A. Larder, S. D. Kemp, Multiple mutations in HIV-1 reverse transcriptase confer high-level resistance to zidovudine (AZT)., *Science* , **246**, 1155–8, 1989, Medline: 90069587
- Larder91 B. A. Larder, K. E. Coates, S. D. Kemp, Zidovudine-resistant human immunodeficiency virus selected by passage in cell culture., *J Virol*, **65**, 5232–6, 1991, Medline: 91374572
- Larder92 B. A. Larder, 3'-Azido-3'-deoxythymidine resistance suppressed by a mutation conferring human immunodeficiency virus type 1 resistance to nonnucleoside reverse transcriptase inhibitors., *Antimicrob Agents Chemother*, **36**, 2664–9, 1992, Medline: 93128874
- Larder95 B. A. Larder, S. D. Kemp, P. R. Harrigan, Potential mechanism for sustained antiretroviral efficacy of AZT-3TC combination therapy., *Science*, **269**, 696–9, 1995, Medline: 95350663
- Larder99 Larder BA, Bloor S, Kemp SD, Hertogs K, Desmet RL, Miller V, Sturmer M, Staszewski S, Ren J, Stammers DK, Stuart DI, Pauwels R., A family of insertion mutations between codons 67 and 70 of human immunodeficiency virus type 1 reverse transcriptase confer multinucleoside analog resistance., *Antimicrobial Agents and Chemotherapy*, **43**, 1961–1967, 1999, Medline: 10428920
- Lawrence99 J. Lawrence, J. Schapiro, M. Winters, J. Montoya, A. Zolopa, R. Pesano, B. Efron, D. Winslow, T. C. Merigan, Clinical resistance patterns and responses to two sequential protease inhibitor regimens in saquinavir and reverse transcriptase inhibitor- experienced persons, *J Infect Dis*, **179**, 1356–64, 99, Medline: 99246335
- Lin96 P. F. Lin, H. Samanta, C. M. Bechtold, C. A. Deminie, A. K. Patick, M. Alam, K. Riccardi, R. E. Rose, R. J. White, R. J. Colonno, Characterization of siamycin I, a human immunodeficiency virus fusion inhibitor., *Antimicrob Agents Chemother*, **40**, 133–8, 1996, Medline: 96379881
- Lobato02 Lobato RL, Kim EY, Kagan RM, Merigan TC, Genotypic and phenotypic analysis of a novel 15-base insertion occurring between codons 69 and 70 of HIV type 1 reverse transcriptase, *AIDS Res Hum Retroviruse*, **18**(10), 733–6, 2002, Medline: 12167282
- Maass93 G. Maass, U. Immendoerfer, B. Koenig, U. Leser, B. Mueller, R. Goody, E. Pfaff, Viral resistance to the thiazolo-iso-indolinones, a new class of nonnucleoside inhibitors of human immunodeficiency virus type 1 reverse transcriptase., *Antimicrob Agents Chemother*, **37**, 2612–7, 1993, Medline: 94153035
- Medlin96 H. K. Medlin, Y. Q. Zhu, K. M. Remington, T. R. Phillips, T. W. North , Selection and characterization of a mutant of feline immunodeficiency virus resistant to 2',3'-dideoxycytidine., *Antimicrob Agents Chemotherapy*, **40**, 953–7, 1996, Medline: 96254566
- Mellors92 J. W. Mellors, G. E. Dutschman, G. J. Im, E. Tramontano, S. R. Winkler, Y. C. Cheng, In vitro selection and molecular characterization of human immunodeficiency virus-1 resistant to

- non-nucleoside inhibitors of reverse transcriptase [published erratum appears in *Mol Pharmacol* 1992 Jul;42(1):174], *Mol Pharmacol*, **41**, 446–51, 1992, Medline: 92186808
- Mellors93 J. W. Mellors, G. J. Im, E. Tramontano, S. R. Winkler, D. J. Medina, G. E. Dutschman, H. Z. Bazmi, G. Piras, C. J. Gonzalez, Y. C. Cheng, A single conservative amino acid substitution in the reverse transcriptase of human immunodeficiency virus-1 confers resistance to (+)-(5S)-4,5,6,7-tetrahydro-5-methyl-6-(3-methyl-2-butenyl)imidazo[4,5, 1- jk][1,4]benzodiazepin-2(1H)-thione (TIBO R82150)., *Mol Pharmacol*, **43** **192**, 11–6 246–53, 1993 1993 1993, Medline: 93140700
- Mellors95 J. W. Mellors, H. Z. Bazmi, R. F. Schinazi, B. M. Roy, Y. Hsiou, E. Arnold, J. Weir, D. L. Mayers, Novel mutations in reverse transcriptase of human immunodeficiency virus type 1 reduce susceptibility to foscarnet in laboratory and clinical isolates., *Antimicrob Agents Chemother*, **39**, 1087–92, 1995, Medline: 95351747
- Mellors96 J.W. Mellors, H. Bazmi, C.K. Chu, Schinazi R.F., K65R mutation in HIV-1 reverse transcriptase causes resistance to (-)- β -D-dioxolane-guanine and reverses AZT resistance, *Fifth International Workshop on HIV Drug Resistance Whistler, BC, Canada*
- Mo96 H. Mo, M. Markowitz, P. Majer, S. K. Burt, S. V. Gulnik, L. I. Suvorov, J. W. Erickson, D. D. Ho, Design, synthesis, and resistance patterns of MP-134 and MP-167, two novel inhibitors of HIV type 1 protease., *AIDS Res Hum Retroviruses*, **12**, 55–61, 1996, Medline: 96423018
- Moeremans95 M. Moeremans, M. De Raeymaeker, R. Van den Broeck, P. Stoffels, M. De Brabander, J. De Cree, K. Hertogs, R. Pauwels, S. Staszewski, K. Andries, Virological analysis of HIV-1 isolates in patients treated with the non-nucleoside reverse transcriptase inhibitor RO91767, 8-chloro-TIBO., *Fourth International Workshop on HIV Drug Resistance Sardinia, Italy*
- Moeremans95a M. Moeremans, M. De Raeymaeker, R. Van den Broeck, P. Stoffels, K. Andries, Genotypic analysis of HIV-1 isolates from patients receiving zidovudine alone or in combination with nucleoside reverse transcriptase inhibitor, *Fourth International Workshop on HIV Drug Resistance Sardinia, Italy*
- Molla96 A. Molla, M. Korneyeva, Q. Gao, S. Vasavanonda, P. J. Schipper, H. M. Mo, M. Markowitz, T. Chernyavskiy, P. Niu, N. Lyons, A. Hsu, G. R. Granneman, D. D. Ho, C. A. Boucher, J. M. Leonard, D. W. Norbeck, D. J. Kempf, Ordered accumulation of mutations in HIV protease confers resistance to ritonavir., *Nat Med*, **2**, 760–6, 1996, Medline: 96266327
- Montes02 Montes B, Segondy M., Prevalence of the mutational pattern E44D/A and/or V118I in the reverse transcriptase (RT) gene of HIV-1 in relation to treatment with nucleoside analogue RT inhibitors., *J Med Virol.*, **66**(3), 299–303, 2002, Medline: 11793380
- Mulato97 A.S. Mulato, P.L. Lamy, W. Li, M.D. Miller, J.M. Cherrington, Genotypic characterization of HIV-1 variants isolated from AIDS patients treated with adefovir dipivoxil (bis-POM PMEA)., *Sixth International Workshop on HIV Drug Resistance St. Petersburg, FL, USA*
- Nguyen94 M. H. Nguyen, R. F. Schinazi, C. Shi, N. M. Goudgaon, P. M. McKenna, J. W. Mellors, Resistance of human immunodeficiency virus type 1 to acyclic 6- phenylselenenyl- and 6-phenylthiopyrimidines., *Antimicrob Agents Chemother*, **38**, 2409–14, 1994, Medline: 95142586
- Nunberg91 J. H. Nunberg, W. A. Schleif, E. J. Boots, J. A. O'Brien, J. C. Quintero, J. M. Hoffman, E. A. Emini, M. E. Goldman, Viral resistance to human immunodeficiency virus type 1-specific pyridinone reverse transcriptase inhibitors., *J Virol*, **65**, 4887–92, 1991, Medline: 91333034
- Olmsted96 R. A. Olmsted, D. E. Slade, L. A. Kopta, S. M. Poppe, T. J. Poel, S. W. Newport, K. B. Rank, C. Biles, R. A. Morge, T. J. Dueweke, Y. Yagi, D. L. Romero, R. C. Thomas, S. K. Sharma, W. G. Tarpley, (Alkylamino) piperidine bis(heteroaryl)piperazine analogs are potent, broad-spectrum non-nucleoside reverse transcriptase inhibitors of drug-resistant isolates of human immunodeficiency virus type 1 (HIV-1) and select for drug-resistant variants of HIV-1III B with reduced replication phenotypes., *J Virol*, **70**, 3698–705, 1996, Medline: 96211502
- Otto93 M. J. Otto, S. Garber, D. L. Winslow, C. D. Reid, P. Aldrich, P. K. Jadhav, C. E. Patterson, C. N. Hodge, Y. S. Cheng, In vitro isolation and identification of human immunodeficiency virus (HIV) variants with reduced sensitivity to C-2 symmetrical inhibitors of HIV type 1 protease., *Proc Natl Acad Sci U S A*, **90**, 7543–7, 1993, Medline: 93361483
- Otto95 M.J. Otto, C.D. Reid, R.W. King, S. Garber, D.B. Baker, E. Anton, Winslow D.L., Exposure of chronically infected PBMCs to DMP 450 can completely suppress virus replication or select

- resistant variants depending upon the dose of compound, *Second National Conference on Human Retroviruses and Related Infections Washington, DC, USA*
- Partaledis94 J.A. Partaledis, K. Yamaguchi, Byrn R.A., In vitro selection and characterization of HIV-1 viral isolates with reduced sensitivity to inhibitors of HIV protease, *Third International Workshop on HIV Drug Resistance Kauai, HI, USA*
- Partaledis95 Partaledis JA, Yamaguchi K, Tisdale M, Blair EE, Falcione C, Maschera B, Myers RE, Pazhanisamy S, Futer O, Cullinan AB, Stuver CM, Byrn RA, Livingston DJ., In vitro selection and characterization of human immunodeficiency virus type 1 (HIV-1) isolates with reduced sensitivity to hydroxyethylamino sulfonamide inhibitors of HIV-1 aspartyl protease., *J Virol*, **69**(9), 5228–5235, 1995, Medline: 95363927
- Patick95 A. K. Patick, R. Rose, J. Greytok, C. M. Bechtold, M. A. Hermsmeier, P. T. Chen, J. C. Barrish, R. Zahler, R. J. Colonno, P. F. Lin, Characterization of a human immunodeficiency virus type 1 variant with reduced sensitivity to an aminodiol protease inhibitor., *J Virol*, **69**, 2148–52, 1995, Medline: 95190985
- Patick96 A. K. Patick, H. Mo, M. Markowitz, K. Appelt, B. Wu, L. Musick, V. Kalish, S. Kaldor, S. Reich, D. Ho, S. Webber, Antiviral and resistance studies of AG1343, an orally bioavailable inhibitor of human immunodeficiency virus protease [published erratum appears in *Antimicrob Agents Chemother* 1996 Jun;40(6):1575], *Antimicrob Agents Chemother*, **40**, 292–7, 1996, Medline: 96431786
- Patick97 A.K. Patick, D. Kuritzkes, V.A. Johnson, D. Shugarts, M. Bakhtiari, K.E. Potts, A. Farnsworth, R. Anderson, J.L. Koel, J.D. Hazelwood, C.D. Nail, M. Duran, M. Markowitz, Ho D. Richman D, Genotypic and phenotypic analyses of HIV-1 variants isolated from patients treated with nelfinavir and other HIV-1 protease inhibitors, *Sixth International Workshop on HIV Drug Resistance St. Petersburg, FL, USA*
- Patick98 Patick AK, Duran M, Cao Y, Shugarts D, Keller MR, Mazabel E, Knowles M, Chapman S, Kuritzkes DR, Markowitz M, Genotypic and phenotypic characterization of human immunodeficiency virus type 1 variants isolated from patients treated with the protease inhibitor nelfinavir, *Antimicrob Agents Chemother*, **42**(10), 2637–44, 1998, Medline: 98443459
- Pelemans01 Pelemans H, Esnouf R, Min KL, Parniak M, De Clercq E, Balzarini J., Mutations at amino acid positions 63, 189, and 396 of human immunodeficiency virus type 1 reverse transcriptase (RT) partially restore the DNA polymerase activity of a Trp229Tyr mutant RT., *Virology*, **287**(1), 143–50, 2001, Medline: 11504549
- Pelemans97 H. Pelemans, R. Esnouf, A. Dunkler, M.A. Parniak, A-M. Vandamme, A. Karlsson, E. De Clercq, J-P. Kleim, J. Balzarini. , Characteristics of the Pro225His mutation in human immunodeficiency virus type 1 (HIV-1) reverse transcriptase that appears under selective pressure of dose-escalating quinoxaline treatment of HIV-1., *J Virol*, **71**(11), 8195–8203, 1997, Medline: 98001335
- Pelemans98 Pelemans H, Esnouf RM, Parniak MA, Vandamme AM, De Clercq E, Balzarini J., A proline-to-hisitidine substitution at position 225 of human immunodeficiency virus type 1 (HIV-1) reverse transcriptase (RT) sensitizes HIV-1 RT to BHAP U-90152., *Journal of General Virology*, **79**(6), 1347–52, 1998, Medline: 98295831
- Pillay96 D. Pillay, M.L. Smidt, K.E. Potts, M.L. Bryant, D.D. Richman, In vitro selection of protease inhibitors resistant human immunodeficiency virus type 1 (HIV-1) strains., *Thirty fourth Interscience Conference on Antimicrobial Agents and Chemotherapy, Orlando, FL, USA*
- Potts94 Potts KE, Smidt ML, Stallings WC, Clare M, Pillay D, Richman DD, and Bryant ML, In vitro selection and characterization of human immunodeficiency virus type 1 (HIV-1) variants with decreased sensitivity to hydroxyethylurea isostere containing protease inhibitors, *Third International Workshop on HIV Drug Resistance, Kauai, Hawaii, USA, 2–5 August, 1994. Abstract 4*
- Prasad91 V. R. Prasad, I. Lowy, T. de los Santos, L. Chiang, S. P. Goff, Isolation and characterization of a dideoxyguanosine triphosphate- resistant mutant of human immunodeficiency virus reverse transcriptase., *Proc Natl Acad Sci U S A*, **88**, 11363–7, 1991, Medline: 92107950
- Rando99 R.F. Rando, D.L. Taylor, L.A. Kelly, L.A. Wood, A.S. Tyms, J.M. McCune, C.A. Stoddart, and J. Dedard. , Anti-HIV activity, drug combination, and resistance studies of dOTC (BCH-10652). , *Second International Workshop on Salvage Therapy for HIV Infection, Toronto, Canada*

- Rao96 B.G. Rao, M.D. Dwyer, J.A. Thomson, C.T. Baker, D.D. Deiningner, M.A. Murcko, R.D. Tung, M.A. Navia, Kim E.E., Structural and modelling analysis of the basis of viral resistance to VX-478, *Fifth International Workshop on HIV Drug Resistance, Whistler, BC, Canada*, **69**(9), 5228–5235
- Richard00 Richard N, Salomon H, Rando R, Mansour T, Bowlin TL, Wainberg MA., Selection and characterization of human immunodeficiency virus type 1 variants resistant to the (+) and (-) enantiomers of 2'-deoxy-3'-oxa-4'-thio-5-fluorocytidine., *Antimicrobial Agents and Chemotherapy*, **44**, 1127–1131, 2000, Medline: 10770740
- Richman91 D. Richman, C. K. Shih, I. Lowy, J. Rose, P. Prodanovich, S. Goff, J. Griffin, Human immunodeficiency virus type 1 mutants resistant to nonnucleoside inhibitors of reverse transcriptase arise in tissue culture., *Proc Natl Acad Sci U S A*, **88**, 11241–5, 1991, Medline: 92107925
- Richman93 D. D. Richman, Resistance of clinical isolates of human immunodeficiency virus to antiretroviral agents., *Antimicrob Agents Chemother*, **37**, 1207–13, 1993, Medline: 93319246
- Richman94 D. D. Richman, D. Havlir, J. Corbeil, D. Looney, C. Ignacio, S. A. Spector, J. Sullivan, S. Cheeseman, K. Barringer, D. Pauletti, et al, Nevirapine resistance mutations of human immunodeficiency virus type 1 selected during therapy., *J Virol*, **68**, 1660–6, 1994, Medline: 94149857
- Rimsky98 Rimsky LT, Shugars DC, Matthews TJ., Determinants of human immunodeficiency virus type 1 resistance to gp41-derived inhibitor peptides., *J Virol*, **72**(2), 986–993, 1998, Medline: 98105736
- Rose94 B. Rose, J. Greytok, C. Bechtold, M. Alam, B. Terry, Gong Y.F. DeK. Vore, A. Patrick, R. Colono, Lin P, Combination therapy with two protease inhibitors as an approach to antiviral therapy, *Third International Workshop on HIV Drug Resistance Kauai, HI, USA*
- Rusconi00 Rusconi S, La Seta Catamancio S, Citterio P, Kurtagic S, Violin M, Balotta C, Moroni M, Galli M, d'Arminio-Monforte A., Susceptibility to PNU-140690 (Tipranavir) of human immunodeficiency virus type 1 isolates derived from patients with multidrug resistance to other protease inhibitors., *Antimicrobial Agents and Chemotherapy*, **44**, 1328–1332, 2000, Medline: 10770770
- Saag93 M. S. Saag, E. A. Emini, O. L. Laskin, J. Douglas, W. I. Lapidus, W. A. Schleif, R. J. Whitley, C. Hildebrand, V. W. Byrnes, J. C. Kappes, et al, A short-term clinical evaluation of L-697,661, a non-nucleoside inhibitor of HIV-1 reverse transcriptase. L-697,661 Working Group., *N Engl J Med*, **329**, 1065–72, 1993, Medline: 93382466
- Schapiro96 J.M. Schapiro, Winters M.A. M. Vierra, H. Jacobsen, J. Mous, Merigan T.C., Resistance mutations in patients receiving saquinavir: simultaneous appearance in lymph nodes, peripheral blood mononuclears (PBM) and plasma, *Fifth International Workshop on HIV Drug Resistance Whistler, BC, Canada*
- Schapiro97 J.M. Schapiro, M. Winters, J. Lawrence, J. Norris, T.C. Merigan, Clinical and genotypic cross-resistance between the protease inhibitors saquinavir and indinavir, *Sixth International Workshop on HIV Drug Resistance, St. Petersburg, FL, USA*
- Schinazi93 R. F. Schinazi, R. M. Lloyd, M. H. Nguyen, D. L. Cannon, A. McMillan, N. Ilksoy, C. K. Chu, D. C. Liotta, H. Z. Bazmi, J. W. Mellors, Characterization of human immunodeficiency viruses resistant to oxathiolane-cytosine nucleosides., *Antimicrob Agents Chemother*, **37**, 875–81, 1993, Medline: 93263665
- Schinazi95 R.F. Schinazi, R.M. Lloyd, McA. Millan, G. Gosselin, J.L. Imbach, Sommadossi J-P, Development of HIV-1 and SIV resistant to β -L-2',3'-dideoxycytidine analogues, *Fourth International Workshop on HIV Drug Resistance Sardinia, Italy*
- Schinazi96 R.F. Schinazi, L. Stuyver, A. Wyseur, R.M. Lloyd, L. Hough, A. Rombout, R. Rossau, D. Rimland, Proviral and plasma virus genotyping using a line probe assay in nucleoside treated HIV infected Veterans Affairs patients, *Fifth International Workshop on HIV Drug Resistance, Whistler, BC, Canada*
- Schinazi97 R.F. Schinazi, L. Stuyver, A. Wyseur, R.M. Lloyd, Hough L. A. Rombout, R. Rossau, and Rimland D, Genotypic characterization of HIV-1 variants isolated from AIDS patients treated with adefovir dipivoxil (bis-POM PMEA), *Sixth International Workshop on HIV Drug Resistance St. Petersburg, FL, USA*

- Schmit96 J.C. Schmit, I. Vanderlinden, L. Ruiz, B. Clotet, P. Hermans, S. Sprecher, Arendt V. W. Peetermans, T. Harrer, D. Vaira, J. Desmyter, E. De Clercq, A.M. Vandamme, Prevalence of multi-drug resistance to dideoxynucleoside (ddN) analogues in patients on ddN combination therapy, *Fifth International Workshop on HIV Drug Resistance Whistler, BC, Canada*
- Schmit98 J. C. Schmit, K. Van Laethem, L. Ruiz, P. Hermans, S. Sprecher, A. Sonnerborg, M. Leal, T. Harrer, B. Clotet, V. Arendt, E. Lissen, M. Witvrouw, J. Desmyter, E. De Clercq, A. M. Vandamme, Multiple dideoxynucleoside analogue-resistant (MddNR) HIV-1 strains isolated from patients from different European countries, *AIDS*, **12**, 2007–15, 98, Medline: 99030034
- Schols98 Schols D, Este JA, Cabrera C, Cabrera C, De Clercq E., T-cell-line-tropic human immunodeficiency virus type 1 that is made resistant to stromal cell derived factor 1a contains mutations in envelope gp120 but does not show a switch in coreceptor use., *J Virol*, **72**(5), 4032–4037, 1998, Medline: 98216769
- Seki95 M. Seki, Y. Sadakata, S. Yuasa, M. Baba, Isolation and characterization of human immunodeficiency virus type-1 mutants resistant to the non-nucleoside reverse transcriptase inhibitor MKC-442, *Antiviral Chemistry and Chemotherapy*, **6**, 73–9
- Shao95 Shao W, Smith T, Swanstrom R., Selection and analysis of HIV-1 variants with increased resistance to SKF108842 and SKF108922, two protease inhibitors., *Fourth International Workshop on HIV Drug Resistance, Sardinia, Italy, 6–9 July*
- Shaw94 G. Shaw, X. Wei, Johnson V, M. Taylor, J. Decker, M. Kilby, J. Lifson, B. Hahn, Saag M, Nucleotide sequence analysis of HIV-1 RNA and DNA from plasma and PBMCs of patients treated with ZDV, ddI and nevirapine: rapid turnover and resistance development in vivo, *Third International Workshop on HIV Drug Resistance Kauai, HI, USA*
- Shirasaka95 T. Shirasaka, M. F. Kavlick, T. Ueno, W. Y. Gao, E. Kojima, M. L. Alcaide, S. Chokekijchai, B. M. Roy, E. Arnold, R. Yarchoan, et al, Emergence of human immunodeficiency virus type 1 variants with resistance to multiple dideoxynucleosides in patients receiving therapy with dideoxynucleosides., *Proc Natl Acad Sci U S A*, **92**, 2398–402, 1995, Medline: 95199357
- Slade93 D.E. Slade, C.L. Vavro, J.T. Stapelton, N. Swack, St Clair M.H., A cysteine at codon 215 of HIV RT confers resistance to ddC, *Second HIV Drug Resistance Workshop Noordwijk, The Netherlands*
- Smidt97 M. L. Smidt, K. E. Potts, S. P. Tucker, L. Blystone, T. R. Stiebel, W. C. Stallings, J. J. McDonald, D. Pillay, D. D. Richman, M. L. Bryant, A mutation in human immunodeficiency virus type 1 protease at position 88, located outside the active site, confers resistance to the hydroxyethylurea inhibitor SC-55389A., *Antimicrob Agents Chemother*, **41**, 515–22, 1997, Medline: 97209043
- Smith96 R. Smith, K. Remington, R. Lloyd, R. Schinazi, North T, Mutants of feline immunodeficiency virus resistant to FTC and 3TC, *Third International Feline Retrovirus Research Symposium Ft. Collins, CO, USA*
- Smith97 R. A. Smith, K. M. Remington, R. M. Lloyd, R. F. Schinazi, T. W. North, A novel Met-to-Thr mutation in the YMDD motif of reverse transcriptase from feline immunodeficiency virus confers resistance to oxathiolane nucleosides., *J Virol*, **71**, 2357–62, 1997, Medline: 97184570
- Smith98 R. A. Smith, K. M. Remington, B. D. Preston, R. F. Schinazi, T. W. North, A novel point mutation at position 156 of reverse transcriptase from feline immunodeficiency virus confers resistance to the combination of (-)-beta-2',3'-dideoxy-3'-thiacytidine and 3'-azido-3'-deoxythymidine., *J Virol*, **72**, 2335–40, 1998, Medline: 97184570
- Smith99 Smith RA, Klarmann GJ, Stray KM, von Schwedler UK, Schinazi RF, Preston BD, North TW., A new point mutation (P157S) in the reverse transcriptase of human immunodeficiency virus type 1 confers low-level resistance to (-)-beta-2',3'-dideoxy-3'-thiacytidine., *Antimicrobial Agents and Chemotherapy*, **43**, 2077–2080, 1999, Medline: 10428942
- Staszewski96 S. Staszewski, V. Miller, S. Rehmet, T. Stark, J. De Cree, M. De Brabander, M. Peeters, K. Andries, M. Moeremans, M. De Raeymaeker, G. Pearce, R. Van den Broeck, W. Verbiest, P. Stoffels, Virological and immunological analysis of a triple combination pilot study with zidovudine, lamivudine and zalcitabine in HIV-1-infected patients., *AIDS*, **10**, F1–7, 1996, Medline: 96314090
- Staszewski96a S. Staszewski, V. Miller, A. Kober, R. Colebunders, B. Vandercam, J. Delescluse, N. Clumeck, F. Van Wanzele, M. De Brabander, J De Cree, M. Moeremans, K. Andries, C. Boucher,

- P. Stoffels, P.A.J. Janssen, Evaluation of the efficacy and tolerance of RO18893, RO89439 (loviride) and placebo in asymptomatic HIV-1-infected patients, *Antiviral Therapy*, **1**, 42–50
- StClair91 M. H. St Clair, J. L. Martin, G. Tudor-Williams, M. C. Bach, C. L. Vavro, D. M. King, P. Kellam, S. D. Kemp, B. A. Larder, Resistance to ddI and sensitivity to AZT induced by a mutation in HIV-1 reverse transcriptase., *Science*, **253**, 1557–9, 1991, Medline: 91376665
- Stuyver97 L. Stuyver, A. Wyseur, A. Rombout, J. Louwagie, T. Scarcez, C. Verhofstede, D. Rimland, R. F. Schinazi, R. Rossau, Line probe assay for rapid detection of drug-selected mutations in the human immunodeficiency virus type 1 reverse transcriptase gene., *Antimicrob Agents Chemother*, **41**, 284–91, 1997, Medline: 97173275
- Swanstrom94 R. Swanstrom, T. Smith, S. Petit, D. Irlbeck, W. Shao, Wehbie R. R. Sawhney, L. Everitt, Erickson I, Multiple sequence changes within HIV-1 protease confer reduced sensitivity to a symmetric protease inhibitor, *Third International Workshop on HIV Drug Resistance Kauai, HI, USA*
- Tachedjian95 G. Tachedjian, D. J. Hooker, A. D. Gurusinghe, H. Bazmi, N. J. Deacon, J. Mellors, C. Birch, J. Mills, Characterisation of foscarnet-resistant strains of human immunodeficiency virus type 1., *Virology*, **212**, 58–68, 1995, Medline: 95407116
- Tachedjian96 G. Tachedjian, J. Mellors, H. Bazmi, C. Birch, J. Mills, Zidovudine resistance is suppressed by mutations conferring resistance of human immunodeficiency virus type 1 to foscarnet., *J Virol*, **70**, 7171–81, 1996, Medline: 96386614
- Tachedjian98 G. Tachedjian, M. French, J. Mills, Coresistance to zidovudine and foscarnet is associated with multiple mutations in the human immunodeficiency virus type 1 reverse transcriptase, *Antimicrob Agents Chemother*, **42**, 3038–43, 98, Medline: 99013598
- Tanaka97 M. Tanaka, R. V. Srinivas, T. Ueno, M. F. Kavlick, F. K. Hui, A. Fridland, J. S. Driscoll, H. Mitsuya, In vitro induction of human immunodeficiency virus type 1 variants resistant to 2'-beta-Fluoro-2',3'-dideoxyadenosine., *Antimicrob Agents Chemother*, **41**, 1313–8, 1997, Medline: 97316916
- Tisdale93 M. Tisdale, S. D. Kemp, N. R. Parry, B. A. Larder, Rapid in vitro selection of human immunodeficiency virus type 1 resistant to 3'-thiacytidine inhibitors due to a mutation in the YMDD region of reverse transcriptase., *Proc Natl Acad Sci U S A*, **90**, 5653–6, 1993, Medline: 93296196
- Tisdale94 M. Tisdale, R. Myers, N.R. Parry, N. Oliver, B. Machera, Blair E, Comprehensive analysis of HIV-1 variants individually selected for resistance to six HIV protease inhibitors, *Third International Workshop on HIV Drug Resistance Kauai, HI, USA*
- Tisdale96 M. Tisdale, R. Myers, I. Najera, A. Kohli, Kemp S. and Larder B.A., Analysis of resistance interactions with 141W94 (VX-478) and other HIV-1 protease inhibitors, *Fifth International Workshop on HIV Drug Resistance Whistler, BC, Canada*, **69**(9), 5228–5235
- Tisdale97 M. Tisdale, T. Alnadaf, D. Cousens, Combination of mutations in human immunodeficiency virus type 1 reverse transcriptase required for resistance to the carbocyclic nucleoside 1592U89., *Antimicrob Agents Chemother*, **41**, 1094–8, 1997, Medline: 97291261
- Vandamme94 A. M. Vandamme, Z. Debyser, R. Pauwels, K. De Vreese, P. Goubau, M. Youle, B. Gazzard, P. A. Stoffels, G. F. Cauwenbergh, J. Anne, et al, Characterization of HIV-1 strains isolated from patients treated with TIBO R82913., *AIDS Res Hum Retroviruses*, **10**, 39–46, 1994, Medline: 94235372
- Vandamme94a A.-M. Vandamme, Polymerase chain reaction (PCR) as a diagnostic tool in HIV infection, *Verhandelingen van de Koninklijke Academie voor Geneeskunde van Belgie*, **56**, 231–265
- Vandamme96 A.-M. Vandamme, J.C. Schmit, J. Balzarini, K. Van Laethem, M. Witvrouw, P. Hermans, S. Sprecher, J. Martinez-Picado, B. Clotet, W. Peetermans, J. Desmyter, E. De Clercq, Presence of TSAO-resistant virus strains in non-experienced patients, *Fifth International Workshop on HIV Drug Resistance Whistler, BC, Canada*
- VanLaethem00 Van Laethem K, Schmit JC, Pelemans H, Balzarini J, Witvrouw M, Perez-Perez MJ, Camarasa MJ, Esnouf RM, Aquaro S, Cenci A, Perno CF, Hermans P, Sprecher S, Ruiz L, Clotet B, Van Wijngaerden E, Van Ranst M, Desmyter J, De Clercq E, Vandamme

- AM., Presence of 2',5'-Bis-O-(tert-butyl dimethylsilyl)-3'-spiro-5''-(4''-amino-1'',2''-oxathiole-2'',2''-dioxide) (TSAO)-resistant virus strains in TSAO-inexperienced HIV patients., *AIDS Research and Human Retroviruses*, **16**, 825–833, 2000, Medline: 10875608
- VanRompay96 K. K. Van Rompay, J. M. Cherrington, M. L. Marthas, C. J. Berardi, A. S. Mulato, A. Spinner, R. P. Tarara, D. R. Canfield, S. Telm, N. Bischofberger, N. C. Pedersen, 9-[2-(Phosphonomethoxy)propyl]adenine therapy of established simian immunodeficiency virus infection in infant rhesus macaques., *Antimicrob Agents Chemother*, **40**, 2586–91, 1996, Medline: 97070544
- VanRompay97 Van Rompay KK, Greenier JL, Marthas ML, Otsyula MG, Tarara RP, Miller CJ, Pedersen NC, A zidovudine-resistant simian immunodeficiency virus mutant with a Q151M mutation in reverse transcriptase causes AIDS in newborn macaques, *Antimicrob Agents Chemother*, **41**, 278–83, 1997, Medline: 97173274
- VanRompay97a K. Van Rompay, J. Cherrington, M. Marthas, E. Agatep, Z. Dehqanzada, P. Lamy, C. Berardi, N. Bischofberger, N. Pedersen, Therapeutic efficacy of PMPA treatment for infant macaques infected with PMPA-resistant simian immunodeficiency virus, *Sixth International Workshop on HIV Drug Resistance St. Petersburg, FL, USA*
- Vasudevachari92 Vasudevachari MB, Battista C, Lane HC, Psallidopoulos MC, Zhao B, Cook J, Palmer JR, Romero DL, Tarpley WG, Salzman NP, Prevention of the spread of HIV-1 infection with nonnucleoside reverse transcriptase inhibitors, *Virology*, **190**(1), 269–77, 1992, Medline: 92410603
- Vasudevachari96 M. B. Vasudevachari, Y. M. Zhang, H. Imamichi, T. Imamichi, J. Falloon, N. P. Salzman, Emergence of protease inhibitor resistance mutations in human immunodeficiency virus type 1 isolates from patients and rapid screening procedure for their detection., *Antimicrob Agents Chemother*, **40**, 2535–41, 1996, Medline: 97070533
- Vrang93 L. Vrang, C. Rydergard, C. Ahgren, P. Engelhardt, M. Hogberg, N.G. Johansson, J. Kangasmetsa, P. Lind, R. Noreen, C. Sahlberg, X. X. Zhou, A. Karlsson, C. Lopez, J.M. Morin, R.J. Ternansky, F.W. Bell, C.L. Jordan, M.D. Kinnick, J.A. Palkowitz, C.A. Parrish, P. Pranc, R.T. Vasileff, S.J. West, B.Oberg, Comparative rates of in vitro resistance development of HIV-1 to non-nucleoside analog RT inhibitors, *Antiviral Res*, **20** (S1), 77
- Wei02 Wei X, Decker JM, Liu H, Zhang Z, Arani RB, Kilby JM, Saag MS, Wu X, Shaw GM, Kappes JC, Emergence of resistant human immunodeficiency virus type 1 in patients receiving fusion inhibitor (T-20) monotherapy, *Antimicrob Agents Chemother*, **46**(6), 1896–905, 2002, Medline: 12019106
- Winslow95 D.L. Winslow, S. Garber, C. Reid, E. Anton, Otto M.J., DMP 450, a new cyclic urea inhibitor of HIV protease with potent in vitro antiviral activity, *Eighth International Conference on Antiviral Research Santa Fe, NM, USA*
- Winslow96 D. L. Winslow, S. Garber, C. Reid, H. Scarnati, D. Baker, M. M. Rayner, E. D. Anton, Selection conditions affect the evolution of specific mutations in the reverse transcriptase gene associated with resistance to DMP 266., *AIDS*, **10**, 1205–9, 1996, Medline: 97037953
- Winters97 M.A. Winters, J.M. Schapiro, J. Lawrence, Merigan T.C., Genotypic and phenotypic analysis of the protease gene in HIV-1-infected patients that failed long-term saquinavir therapy and switched to other protease inhibitors, *Sixth International Workshop on HIV Drug Resistance St. Petersburg, FL, USA*
- Winters98 M. A. Winters, K. L. Coolley, Y. A. Girard, D. J. Levee, H. Hamdan, R. W. Shafer, D. A. Katzenstein, T. C. Merigan, A 6-basepair insert in the reverse transcriptase gene of human immunodeficiency virus type 1 confers resistance to multiple nucleoside inhibitors, *J Clin Invest*, **102**, 1769–75, 98, Medline: 99038179
- Witvrouw98 Witvrouw M, Arranz ME, Pannecouque C, Declercq R, Jonckheere H, Schmit J-C, Vandamme A-M, Diaz JA, Ingate ST, Desmyter J, Esnouf R, Van Meervelt L, Vega S, Balzarini J, De Clercq E., 1,1,3-Trioxo-2H,4H-thieno[3,4-e][1,2,4]thiadiazine (TDD) derivatives: a new class of nonnucleoside human immunodeficiency virus type 1 (HIV-1) reverse transcriptase inhibitors with anti-HIV-1 activity., *Antimicrob Agents Chemother*, **42**(3), 618–623, 1998, Medline: 98177121

- Yang97 Yang SS, Pattabiraman N, Gussio R, Pallansch L, Buckheit RW Jr, Bader JP, Cross-resistance analysis and molecular modeling of nonnucleoside reverse transcriptase inhibitors targeting drug-resistance mutations in the reverse transcriptase of human immunodeficiency virus. , *Leukemia*, **11 S3**, 89–92 , 1997, Medline: 97353050
- Yoshimura99 K. Yoshimura, R. Kato, K. Yusa, M. F. Kavlick, V. Maroun, A. Nguyen, T. Mimoto, T. Ueno, M. Shintani, J. Falloon, H. Masur, H. Hayashi, J. Erickson, H. Mitsuya, JE-2147: a dipeptide protease inhibitor (PI) that potently inhibits multi-PI-resistant HIV-1, *Proc Natl Acad Sci U S A*, **96**, 8675–80, 99, Medline: 99342077
- Yoshimura99a Yoshimura K, Feldman R, Kodama E, Kavlick MF, Qiu YL, Zemlicka J, Mitsuya H., In vitro induction of human immunodeficiency virus type 1 variants resistant to phosphoralaninate prodrugs of Z-methylenecyclopropane nucleoside analogues., *Antimicrobial Agents and Chemotherapy*, **43**, 2479–2483, 1999, Medline: 10508028
- Young95 S. D. Young, S. F. Britcher, L. O. Tran, L. S. Payne, W. C. Lumma, T. A. Lyle, J. R. Huff, P. S. Anderson, D. B. Olsen, S. S. Carroll, et al, L-743, 726 (DMP-266): a novel, highly potent nonnucleoside inhibitor of the human immunodeficiency virus type 1 reverse transcriptase., *Antimicrob Agents Chemother*, **39**, 2602–5, 1995, Medline: 96161265
- Zhang94 D. Zhang, A. M. Caliendo, J. J. Eron, K. M. DeVore, J. C. Kaplan, M. S. Hirsch, R. T. D'Aquila, Resistance to 2',3'-dideoxycytidine conferred by a mutation in codon 65 of the human immunodeficiency virus type 1 reverse transcriptase., *Antimicrob Agents Chemother*, **38**, 282–7, 1994, Medline: 94250001
- Zhang95 H. Zhang, L. Vrang, K. Backbro, P. Lind, C. Sahlberg, T. Unge, B. Oberg, Inhibition of human immunodeficiency virus type 1 wild-type and mutant reverse transcriptases by the phenyl ethyl thiazolyl thiourea derivatives trovirdine and MSC-127., *Antiviral Res*, **28**, 331–42, 1995, Medline: 96264013
- Zhu96 Y. Q. Zhu, K. M. Remington, T. W. North, Mutants of feline immunodeficiency virus resistant to 2',3'-dideoxy-2',3'-didehydrothymidine., *Antimicrob Agents Chemother*, 1983-, 1996, Medline: 97032819
- Ziermann00 Ziermann R, Limoli K, Das K, Arnold E, Petropoulos CJ, Parkin NT., A mutation in human immunodeficiency virus type 1 protease, N88S, that causes in vitro hypersensitivity to amprenavir., *Journal of Virology*, **74**, 4414–4419, 2000, Medline: 10756056