

Table 2. Estimated numbers of cases of HIV/AIDS, by race/ethnicity, sex, and age group, 2001–2004—35 areas with confidential name-based HIV infection reporting

Age at diagnosis (yrs)	Year of diagnosis			
	2001	2002	2003	2004
	White, not Hispanic			
Male				
<13	19	12	15	4
13–14	4	3	4	0
15–19	80	80	77	93
20–24	485	517	593	634
25–29	890	917	908	1,029
30–34	1,624	1,632	1,473	1,448
35–39	2,175	2,189	2,050	1,977
40–44	1,636	1,743	1,864	1,970
45–49	1,152	1,145	1,102	1,266
50–54	612	641	673	744
55–59	340	352	329	461
60–64	156	171	161	221
≥65	147	126	112	171
Subtotal	9,321	9,529	9,360	10,018
Female				
<13	26	15	15	5
13–14	3	0	5	3
15–19	78	82	78	68
20–24	188	183	160	178
25–29	248	226	193	212
30–34	344	282	239	249
35–39	360	341	321	289
40–44	281	285	279	287
45–49	190	194	217	222
50–54	103	112	122	130
55–59	53	48	58	78
60–64	27	31	31	31
≥65	19	23	19	35
Subtotal	1,920	1,823	1,737	1,788
Total	11,242	11,352	11,097	11,806

Table 2. Estimated numbers of cases of HIV/AIDS, by race/ethnicity, sex, and age group, 2001–2004—35 areas with confidential name-based HIV infection reporting (cont)

Age at diagnosis (yrs)	Year of diagnosis			
	2001	2002	2003	2004
	Black, not Hispanic			
Male				
<13	118	84	68	63
13–14	9	12	8	10
15–19	314	344	376	463
20–24	1,057	996	1,141	1,406
25–29	1,280	1,277	1,187	1,323
30–34	1,887	1,776	1,578	1,515
35–39	2,460	2,375	2,138	1,912
40–44	2,271	2,065	1,982	1,908
45–49	1,691	1,573	1,577	1,503
50–54	1,030	951	960	987
55–59	530	489	514	523
60–64	259	283	264	268
≥65	265	232	229	246
Subtotal	13,170	12,458	12,023	12,126
Female				
<13	121	117	67	63
13–14	32	28	29	17
15–19	399	329	352	300
20–24	863	820	771	790
25–29	1,070	1,017	948	894
30–34	1,356	1,198	1,084	1,046
35–39	1,541	1,303	1,161	1,061
40–44	1,212	1,155	1,108	1,072
45–49	845	823	792	813
50–54	477	495	494	474
55–59	212	232	234	278
60–64	136	140	116	142
≥65	121	124	131	130
Subtotal	8,386	7,779	7,287	7,080
Total	21,556	20,237	19,310	19,206

Table 2. Estimated numbers of cases of HIV/AIDS, by race/ethnicity, sex, and age group, 2001–2004—35 areas with confidential name-based HIV infection reporting (cont)

Age at diagnosis (yrs)	Year of diagnosis			
	2001	2002	2003	2004
	Hispanic			
Male				
<13	34	24	16	21
13–14	3	1	3	6
15–19	77	110	82	99
20–24	420	418	447	534
25–29	734	757	756	850
30–34	1,076	1,021	958	938
35–39	1,232	1,097	1,050	1,114
40–44	891	767	903	781
45–49	572	514	566	495
50–54	318	283	329	311
55–59	190	144	187	183
60–64	105	89	90	103
≥65	89	88	99	108
Subtotal	5,741	5,314	5,488	5,544
Female				
<13	38	26	16	16
13–14	1	1	2	5
15–19	55	62	48	41
20–24	177	158	155	154
25–29	261	203	204	185
30–34	334	249	231	196
35–39	365	283	268	243
40–44	293	261	246	218
45–49	184	165	179	160
50–54	141	126	120	102
55–59	67	49	62	49
60–64	35	37	34	25
≥65	23	30	25	33
Subtotal	1,973	1,650	1,590	1,425
Total	7,714	6,964	7,078	6,970

Table 2. Estimated numbers of cases of HIV/AIDS, by race/ethnicity, sex, and age group, 2001–2004—35 areas with confidential name-based HIV infection reporting (cont)

Age at diagnosis (yrs)	Year of diagnosis			
	2001	2002	2003	2004
	Asian/Pacific Islander			
Male				
<13	2	1	1	0
13–14	0	1	0	0
15–19	0	1	1	3
20–24	13	13	25	25
25–29	44	36	38	51
30–34	51	59	56	60
35–39	33	52	68	64
40–44	34	39	38	44
45–49	23	25	29	31
50–54	18	12	12	11
55–59	3	1	11	4
60–64	2	5	2	4
≥65	1	1	2	4
Subtotal	223	247	285	300
Female				
<13	0	2	1	0
13–14	0	0	1	0
15–19	2	2	1	2
20–24	6	7	7	7
25–29	9	18	11	16
30–34	10	15	17	26
35–39	4	7	11	11
40–44	8	5	11	14
45–49	4	5	10	4
50–54	6	8	3	6
55–59	0	2	5	5
60–64	1	0	2	3
≥65	4	0	1	3
Subtotal	56	72	83	94
Total	279	319	367	394

Table 2. Estimated numbers of cases of HIV/AIDS, by race/ethnicity, sex, and age group, 2001–2004—35 areas with confidential name-based HIV infection reporting (cont)

Age at diagnosis (yrs)	Year of diagnosis			
	2001	2002	2003	2004
American Indian/Alaska Native				
Male				
<13	0	3	0	1
13–14	0	0	0	0
15–19	2	3	2	3
20–24	9	12	13	12
25–29	25	17	20	24
30–34	26	24	23	18
35–39	30	33	17	37
40–44	15	21	19	28
45–49	11	13	23	18
50–54	5	1	8	4
55–59	2	4	3	4
60–64	1	3	2	1
≥65	1	1	0	0
Subtotal	127	137	130	151
Female				
<13	0	0	0	0
13–14	0	0	0	0
15–19	1	2	4	3
20–24	5	8	10	0
25–29	9	7	6	9
30–34	5	12	3	10
35–39	3	11	9	17
40–44	7	11	12	5
45–49	7	7	9	8
50–54	3	5	1	1
55–59	2	1	1	1
60–64	0	0	1	1
≥65	0	0	1	0
Subtotal	44	66	56	57
Total	171	202	187	208

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts are adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor. The estimates do not include adjustment for incomplete reporting.

Data include persons with a diagnosis of HIV infection, a diagnosis of HIV infection and a later diagnosis of AIDS, or concurrent diagnoses of HIV infection and AIDS.

Since 2000, the following 35 areas have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Alaska, Arizona, Arkansas, Colorado, Florida, Idaho, Indiana, Iowa, Kansas, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, West Virginia, Wisconsin, Wyoming, Guam, and the U.S. Virgin Islands. Since July 1997, Florida has had confidential name-based HIV infection reporting for new diagnoses only.

Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.