

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All causes												
All races 3/, both sexes												
2004.....	816.5	685.2	29.9	16.8	80.1	102.1	193.5	427.0	910.3	2,164.6	5,275.1	13,823.5
2003.....	841.9	700.0	31.5	17.0	81.5	103.6	201.6	433.2	940.9	2,255.0	5,463.1	14,593.3
2002.....	847.3	695.0	31.2	17.4	81.4	103.6	202.9	430.1	952.4	2,314.7	5,556.9	14,828.3
2001.....	848.5	683.4	33.3	17.3	80.7	105.2	203.6	428.9	964.6	2,353.3	5,582.4	15,112.8
2000.....	854.0	736.7	32.4	18.0	79.9	101.4	198.9	425.6	992.2	2,399.1	5,666.5	15,524.4
1999.....	857.0	736.0	34.2	18.6	79.3	102.2	198.0	418.2	1,005.0	2,457.3	5,714.5	15,554.6
All races, male												
2004.....	817.6	753.7	32.4	19.2	114.7	139.5	243.6	543.5	1,128.8	2,644.8	6,394.3	15,031.1
2003.....	840.3	777.4	35.1	19.8	116.5	141.4	255.0	552.2	1,165.5	2,771.7	6,641.8	15,794.0
2002.....	846.6	761.5	35.2	20.0	117.3	142.2	257.5	547.5	1,184.0	2,855.3	6,760.5	16,254.5
2001.....	846.4	749.8	37.0	19.8	117.0	143.7	259.6	545.1	1,192.7	2,911.5	6,833.0	16,744.8
2000.....	853.0	806.5	35.9	20.9	114.9	138.6	255.2	542.8	1,230.7	2,979.6	6,972.6	17,501.4
1999.....	859.2	805.0	37.9	21.5	113.1	139.7	254.9	533.1	1,252.0	3,073.7	7,083.3	17,597.2
All races, female												
2004.....	815.4	613.4	27.3	14.3	43.6	63.5	143.5	314.3	707.4	1,761.4	4,521.8	13,280.3
2003.....	843.4	619.1	27.8	14.0	44.4	64.6	148.5	318.4	732.7	1,823.0	4,675.5	14,062.5
2002.....	848.0	625.3	27.0	14.7	43.7	64.0	148.8	316.9	738.0	1,864.7	4,757.9	14,209.6
2001.....	850.4	613.9	29.5	14.6	42.6	66.0	148.2	316.8	754.0	1,890.8	4,760.5	14,429.9
2000.....	855.0	663.4	28.7	15.0	43.1	63.5	143.2	312.5	772.2	1,921.2	4,814.7	14,719.2
1999.....	854.9	663.6	30.3	15.6	43.7	64.1	141.8	307.6	777.6	1,952.3	4,825.4	14,731.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All causes												
White, both sexes												
2004.....	863.2	574.0	27.0	15.5	76.4	92.9	179.8	394.0	863.5	2,120.4	5,272.8	14,010.1
2003.....	890.1	591.3	28.5	15.8	77.1	94.3	185.7	397.5	892.2	2,207.0	5,457.8	14,792.3
2002.....	895.7	586.7	28.1	16.1	77.1	93.5	186.6	395.3	902.5	2,267.6	5,544.2	15,015.9
2001.....	895.1	570.9	30.7	16.2	75.5	95.3	185.6	393.2	913.5	2,298.9	5,560.9	15,287.5
2000.....	900.2	610.6	29.2	17.0	74.5	90.3	179.9	388.3	940.2	2,341.5	5,637.7	15,707.0
1999.....	901.4	606.1	30.8	17.2	73.2	91.4	177.3	380.4	948.1	2,396.2	5,679.5	15,718.0
White, male												
2004.....	854.2	631.6	29.4	17.9	108.3	127.0	229.1	504.0	1,065.9	2,584.0	6,384.8	15,250.7
2003.....	877.6	658.7	31.5	18.4	108.9	128.8	237.6	509.2	1,102.3	2,707.3	6,620.6	16,037.9
2002.....	884.0	650.9	31.5	18.4	109.7	128.3	239.3	505.4	1,118.6	2,795.4	6,738.8	16,473.2
2001.....	881.9	627.6	34.2	18.4	108.0	130.3	239.7	501.3	1,127.5	2,842.3	6,799.7	16,935.4
2000.....	887.8	667.6	32.6	19.8	105.8	124.1	233.6	496.9	1,163.3	2,905.7	6,933.1	17,716.4
1999.....	892.1	667.0	33.9	19.7	102.8	125.4	230.8	484.6	1,179.7	2,998.7	7,040.1	17,752.9
White, female												
2004.....	871.9	513.6	24.4	13.0	42.4	56.8	129.5	285.4	671.8	1,723.6	4,514.4	13,450.9
2003.....	902.3	520.7	25.5	13.1	43.2	58.0	132.9	287.2	693.7	1,780.6	4,671.6	14,240.6
2002.....	907.0	519.4	24.5	13.7	42.4	56.9	133.2	286.8	698.7	1,819.7	4,742.5	14,382.8
2001.....	907.9	511.5	27.1	13.9	40.8	58.6	131.0	286.8	712.2	1,840.2	4,738.5	14,597.6
2000.....	912.3	550.5	25.5	14.1	41.1	55.1	125.7	281.4	730.9	1,868.3	4,785.3	14,890.7
1999.....	910.4	542.0	27.5	14.6	41.5	56.2	123.2	277.9	731.0	1,893.9	4,787.8	14,900.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All causes												
Black, both sexes												
2004.....	744.3	1,284.7	44.8	23.6	109.4	179.3	322.3	744.8	1,499.6	2,982.6	6,170.4	13,338.5
2003.....	763.6	1,273.8	46.8	22.9	113.1	181.6	343.6	771.4	1,549.9	3,108.4	6,353.9	13,976.7
2002.....	768.4	1,263.6	47.1	24.5	113.7	186.8	348.6	765.8	1,565.2	3,159.3	6,518.9	14,260.8
2001.....	773.5	1,304.5	47.5	23.3	117.7	185.0	356.5	771.9	1,576.8	3,257.6	6,585.6	14,550.8
2000.....	781.1	1,426.1	49.9	24.2	119.4	187.8	356.9	786.9	1,607.8	3,335.6	6,701.3	14,714.3
1999.....	788.1	1,457.3	52.6	26.5	121.2	184.4	368.4	782.4	1,655.6	3,413.6	6,753.3	14,757.9
Black, male												
2004.....	792.6	1,414.2	48.6	26.0	164.3	252.3	397.0	954.9	1,960.8	3,818.3	7,710.3	14,452.5
2003.....	813.7	1,410.0	53.7	26.8	171.3	256.2	426.7	991.4	2,011.2	3,981.3	8,067.0	14,903.4
2002.....	816.7	1,351.5	54.4	28.9	172.6	264.5	434.7	983.0	2,039.2	4,024.5	8,169.6	15,635.5
2001.....	823.9	1,426.4	53.2	27.7	180.7	259.2	445.0	996.1	2,025.3	4,166.6	8,355.9	16,439.9
2000.....	834.1	1,567.6	54.5	28.2	181.4	261.0	453.0	1,017.7	2,080.1	4,253.5	8,486.0	16,791.0
1999.....	847.4	1,592.8	59.0	32.0	184.6	258.6	469.2	1,030.7	2,145.6	4,352.3	8,559.1	17,304.5
Black, female												
2004.....	700.3	1,149.9	40.9	21.1	53.7	112.3	256.0	564.1	1,128.6	2,386.1	5,300.0	12,896.9
2003.....	717.9	1,132.2	39.7	18.9	54.0	113.5	270.0	582.3	1,178.8	2,487.2	5,385.5	13,616.7
2002.....	724.4	1,172.0	39.5	19.9	54.4	116.4	272.3	579.4	1,184.2	2,545.0	5,584.4	13,734.2
2001.....	727.7	1,176.3	41.7	18.7	54.9	117.9	278.0	579.1	1,215.5	2,616.1	5,591.6	13,849.9
2000.....	733.0	1,279.8	45.3	20.0	58.3	121.8	271.9	588.3	1,227.2	2,689.6	5,696.5	13,941.3
1999.....	734.3	1,317.4	46.1	20.9	58.6	117.6	279.4	569.8	1,262.7	2,751.5	5,742.4	13,805.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Certain other intestinal infections (A04,A07-A09)

All races 3/, both sexes

2004.....	1.5	*	*	*	*	*	0.1	0.2	0.9	3.3	12.8	34.3
2003.....	1.1	*	*	*	*	*	*	0.2	0.6	2.4	9.3	24.8
2002.....	0.9	*	*	*	*	*	*	0.1	0.5	2.0	7.0	21.0
2001.....	0.5	*	*	*	*	*	*	0.1	0.3	1.3	4.5	12.8
2000.....	0.5	*	*	*	*	*	*	0.1	0.3	1.2	3.7	11.8
1999.....	0.4	*	*	*	*	*	*	0.1	0.2	1.0	3.2	10.3

All races, male

2004.....	1.1	*	*	*	*	*	*	0.2	0.9	3.1	12.7	32.0
2003.....	0.8	*	*	*	*	*	*	0.2	0.6	2.2	9.3	21.4
2002.....	0.7	*	*	*	*	*	*	0.1	0.6	2.2	7.0	20.4
2001.....	0.4	*	*	*	*	*	*	*	0.3	1.3	4.6	12.1
2000.....	0.4	*	*	*	*	*	*	*	0.3	1.3	4.0	11.4
1999.....	0.3	*	*	*	*	*	*	*	0.2	0.9	3.5	9.5

All races, female

2004.....	1.8	*	*	*	*	*	*	0.2	0.8	3.4	12.9	35.4
2003.....	1.3	*	*	*	*	*	*	0.2	0.6	2.6	9.3	26.3
2002.....	1.0	*	*	*	*	*	*	0.1	0.5	1.8	6.9	21.3
2001.....	0.7	*	*	*	*	*	*	0.1	0.3	1.3	4.5	13.1
2000.....	0.6	*	*	*	*	*	*	*	0.2	1.2	3.6	11.9
1999.....	0.5	*	*	*	*	*	*	*	0.2	1.0	3.0	10.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year	2/	years	years	years	years	years	years	years	years	and over

Certain other intestinal infections (A04,A07-A09)

White, both sexes

2004.....	1.7	*	*	*	*	*	0.1	0.2	0.9	3.5	13.5	36.6
2003.....	1.2	*	*	*	*	*	*	0.1	0.6	2.6	9.7	26.1
2002.....	1.0	*	*	*	*	*	*	0.1	0.5	2.1	7.4	22.0
2001.....	0.6	*	*	*	*	*	*	0.1	0.3	1.3	4.7	13.5
2000.....	0.5	*	*	*	*	*	*	0.1	0.3	1.3	4.0	12.2
1999.....	0.4	*	*	*	*	*	*	*	0.2	1.0	3.3	10.9

White, male

2004.....	1.3	*	*	*	*	*	*	0.2	0.9	3.2	13.4	33.9
2003.....	0.9	*	*	*	*	*	*	0.2	0.6	2.4	9.8	22.7
2002.....	0.8	*	*	*	*	*	*	0.2	0.5	2.3	7.4	21.4
2001.....	0.5	*	*	*	*	*	*	*	0.3	1.3	4.6	12.7
2000.....	0.4	*	*	*	*	*	*	*	0.3	1.3	4.2	11.9
1999.....	0.3	*	*	*	*	*	*	*	0.2	1.0	3.7	9.9

White, female

2004.....	2.1	*	*	*	*	*	*	0.2	0.9	3.7	13.6	37.8
2003.....	1.5	*	*	*	*	*	*	0.1	0.7	2.8	9.7	27.6
2002.....	1.2	*	*	*	*	*	*	0.1	0.5	1.9	7.4	22.3
2001.....	0.8	*	*	*	*	*	*	0.2	0.3	1.3	4.7	13.9
2000.....	0.6	*	*	*	*	*	*	*	0.3	1.2	3.8	12.3
1999.....	0.5	*	*	*	*	*	*	*	*	1.0	3.1	11.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Tuberculosis (A16-A19)

All races 3/, both sexes

2004.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.7	1.3	2.4
2003.....	0.2	*	*	*	*	*	0.1	0.3	0.4	0.6	1.7	2.3
2002.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.9	1.8	2.6
2001.....	0.3	*	*	*	*	0.1	0.1	0.2	0.5	0.8	1.6	2.8
2000.....	0.3	*	*	*	*	0.1	0.1	0.2	0.3	0.8	2.0	2.7
1999.....	0.3	*	*	*	*	*	0.1	0.3	0.5	0.9	2.1	4.2

All races, male

2004.....	0.3	*	*	*	*	*	0.2	0.3	0.4	1.1	1.6	3.8
2003.....	0.3	*	*	*	*	*	0.1	0.4	0.5	0.9	2.3	3.1
2002.....	0.3	*	*	*	*	*	0.2	0.3	0.6	1.2	2.4	4.0
2001.....	0.3	*	*	*	*	*	0.1	0.3	0.7	1.2	2.0	4.4
2000.....	0.3	*	*	*	*	*	0.1	0.4	0.5	1.0	2.8	4.6
1999.....	0.4	*	*	*	*	*	0.2	0.5	0.8	1.4	2.8	7.1

All races, female

2004.....	0.2	*	*	*	*	*	*	0.1	*	0.4	1.0	1.7
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.3	1.9
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.5	2.1
2001.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	2.2
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.7	1.5	2.0
1999.....	0.2	*	*	*	*	*	*	*	0.3	0.6	1.6	3.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Tuberculosis (A16-A19)

White, both sexes

2004.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.0	1.7
2003.....	0.2	*	*	*	*	*	*	0.1	0.3	0.5	1.3	1.8
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.6	1.4	2.1
2001.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.6	1.2	2.2
2000.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	2.2
1999.....	0.2	*	*	*	*	*	0.1	0.2	0.4	0.7	1.6	3.2

White, male

2004.....	0.2	*	*	*	*	*	*	0.2	0.3	0.7	1.2	2.8
2003.....	0.2	*	*	*	*	*	*	0.2	0.4	0.6	1.7	2.4
2002.....	0.2	*	*	*	*	*	*	0.2	0.3	0.9	1.8	3.2
2001.....	0.2	*	*	*	*	*	*	0.2	0.5	0.7	1.5	3.1
2000.....	0.2	*	*	*	*	*	*	0.2	0.2	0.7	1.8	3.8
1999.....	0.3	*	*	*	*	*	*	0.3	0.5	1.0	2.1	5.4

White, female

2004.....	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.3
2003.....	0.2	*	*	*	*	*	*	*	*	0.3	1.1	1.6
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.3	1.2	1.6
2001.....	0.2	*	*	*	*	*	*	*	*	0.4	1.0	1.8
2000.....	0.2	*	*	*	*	*	*	*	*	0.4	1.1	1.6
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.2	2.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All Under 1-4 5-14 15-24 25-34 35-44 45-54 55-64 65-74 75-84 85 years
ages 1/ 1 year 2/ years years years years years years years years years and over

Respiratory tuberculosis (A16)

All races 3/, both sexes

2004.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	0.9	2.2
2003.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.5	1.4	1.8
2002.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.7	1.5	2.3
2001.....	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	1.3	2.4
2000.....	0.2	*	*	*	*	*	0.1	0.2	0.2	0.6	1.7	2.3
1999.....	0.3	*	*	*	*	*	0.1	0.2	0.5	0.8	1.8	3.7

All races, male

2004.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.9	1.2	3.7
2003.....	0.2	*	*	*	*	*	*	0.3	0.4	0.7	1.9	2.5
2002.....	0.3	*	*	*	*	*	0.1	0.2	0.5	1.0	1.9	3.5
2001.....	0.3	*	*	*	*	*	*	0.3	0.6	0.9	1.7	3.8
2000.....	0.3	*	*	*	*	*	0.1	0.3	0.4	0.9	2.4	4.1
1999.....	0.4	*	*	*	*	*	0.1	0.4	0.7	1.2	2.5	6.4

All races, female

2004.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.5
2003.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	1.0	1.5
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.3	1.8
2001.....	0.2	*	*	*	*	*	*	*	*	0.4	1.1	1.8
2000.....	0.2	*	*	*	*	*	*	*	*	0.5	1.2	1.6
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.3	2.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Respiratory tuberculosis (A16)												
White, both sexes												
2004.....	0.1	*	*	*	*	*	0.1	0.1	0.1	0.4	0.7	1.6
2003.....	0.1	*	*	*	*	*	*	0.1	0.2	0.4	1.1	1.4
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.1	1.8
2001.....	0.2	*	*	*	*	*	*	0.1	0.3	0.4	1.0	1.8
2000.....	0.2	*	*	*	*	*	*	0.1	0.1	0.4	1.2	1.9
1999.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	1.3	3.0
White, male												
2004.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.0	2.6
2003.....	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.4	1.8
2002.....	0.2	*	*	*	*	*	*	0.1	0.3	0.7	1.4	2.9
2001.....	0.2	*	*	*	*	*	*	0.1	0.5	0.6	1.3	2.6
2000.....	0.2	*	*	*	*	*	*	0.2	0.2	0.6	1.5	3.4
1999.....	0.3	*	*	*	*	*	*	0.2	0.5	0.8	1.8	5.2
White, female												
2004.....	0.1	*	*	*	*	*	*	*	*	0.2	0.5	1.1
2003.....	0.1	*	*	*	*	*	*	*	*	*	0.9	1.2
2002.....	0.1	*	*	*	*	*	*	*	*	*	1.0	1.3
2001.....	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.4
2000.....	0.1	*	*	*	*	*	*	*	*	0.3	1.0	1.2
1999.....	0.2	*	*	*	*	*	*	*	*	0.3	1.0	2.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Meningococcal infection (A39)

All races 3/, both sexes

2004.....	0.0	*	*	*	0.1	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2002.....	0.1	*	*	0.0	0.1	*	*	*	*	*	*	*
2001.....	0.1	*	*	0.1	0.1	*	*	*	*	*	*	*
2000.....	0.1	0.6	*	*	0.1	*	*	0.1	*	*	*	*
1999.....	0.1	0.8	0.2	0.1	0.1	0.1	0.1	*	*	*	*	*

All races, male

2004.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	0.2	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*	*	*

All races, female

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	0.1	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	0.1	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over
Septicemia (A40-A41)												
All races 3/, both sexes												
2004.....	11.4	6.6	0.5	0.2	0.3	0.8	1.9	5.4	12.9	32.4	81.6	186.7
2003.....	11.7	6.9	0.5	0.2	0.4	0.8	2.1	5.3	13.1	32.6	85.0	202.5
2002.....	11.7	7.3	0.5	0.2	0.3	0.8	1.9	5.2	12.6	34.7	86.5	203.0
2001.....	11.3	7.7	0.7	0.2	0.3	0.7	1.8	5.0	12.3	32.8	82.3	205.9
2000.....	11.1	7.2	0.6	0.2	0.3	0.7	1.9	4.9	11.9	31.0	80.4	215.7
1999.....	11.0	7.5	0.6	0.2	0.3	0.7	1.8	4.6	11.4	31.2	79.4	220.7
All races, male												
2004.....	10.4	8.0	0.5	0.1	0.3	0.8	2.0	5.9	13.8	35.8	93.3	203.9
2003.....	10.5	7.4	0.7	0.2	0.4	0.8	2.1	5.8	14.1	36.2	95.0	213.8
2002.....	10.6	7.7	0.5	0.2	0.3	0.8	2.0	5.7	13.7	38.0	99.0	213.9
2001.....	10.2	8.4	0.8	0.2	0.4	0.7	2.0	5.4	13.2	36.7	95.5	224.3
2000.....	9.8	7.1	0.7	0.2	0.3	0.8	2.0	5.6	13.0	34.5	91.9	223.0
1999.....	9.8	9.0	0.6	0.2	0.3	0.7	2.1	5.1	12.4	35.2	91.9	233.5
All races, female												
2004.....	12.3	5.2	0.6	0.2	0.3	0.8	1.9	4.9	12.0	29.6	73.7	179.0
2003.....	12.9	6.5	0.4	0.2	0.4	0.8	2.0	4.8	12.2	29.5	78.3	197.5
2002.....	12.9	7.0	0.5	0.2	0.3	0.8	1.8	4.7	11.7	31.9	78.2	198.3
2001.....	12.4	7.0	0.6	0.1	0.3	0.7	1.7	4.7	11.4	29.5	73.7	198.3
2000.....	12.3	7.3	0.6	0.1	0.2	0.6	1.9	4.2	11.0	28.1	72.9	212.7
1999.....	12.2	5.8	0.5	0.2	0.3	0.6	1.6	4.1	10.5	28.0	71.3	215.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Septicemia (A40-A41)												
White, both sexes												
2004.....	11.2	5.2	0.5	0.2	0.3	0.7	1.6	4.4	11.0	29.1	76.6	178.1
2003.....	11.5	4.6	0.4	0.2	0.3	0.6	1.7	4.3	11.2	28.9	80.1	191.1
2002.....	11.5	6.1	0.5	0.2	0.3	0.6	1.6	4.2	10.6	31.1	80.7	191.9
2001.....	11.1	5.4	0.7	0.2	0.3	0.5	1.4	4.0	10.3	29.3	77.1	195.3
2000.....	10.8	5.7	0.6	0.2	0.2	0.5	1.6	3.9	9.9	27.4	73.9	203.0
1999.....	10.6	5.5	0.5	0.2	0.3	0.6	1.4	3.6	9.4	27.3	72.8	205.9
White, male												
2004.....	10.2	6.3	0.5	0.2	0.3	0.6	1.6	4.8	11.7	32.2	87.6	196.5
2003.....	10.2	4.6	0.6	0.2	0.3	0.6	1.7	4.7	11.9	32.1	89.5	203.3
2002.....	10.3	6.2	0.5	0.3	0.3	0.7	1.6	4.7	11.7	34.1	91.5	202.9
2001.....	9.9	6.2	0.8	0.2	0.3	0.5	1.7	4.3	11.1	32.5	89.4	211.8
2000.....	9.4	5.8	0.6	0.2	0.2	0.6	1.7	4.4	10.9	30.8	84.6	209.8
1999.....	9.3	6.9	0.5	0.2	0.3	0.6	1.6	4.0	10.0	30.6	84.6	217.4
White, female												
2004.....	12.2	4.0	0.5	0.2	0.2	0.7	1.5	4.0	10.3	26.5	69.2	169.9
2003.....	12.8	4.5	*	0.2	0.3	0.6	1.7	3.8	10.5	26.2	73.8	185.8
2002.....	12.8	6.0	0.4	0.2	0.3	0.6	1.5	3.8	9.6	28.6	73.4	187.1
2001.....	12.3	4.6	0.6	0.1	0.2	0.5	1.2	3.7	9.7	26.6	69.0	188.3
2000.....	12.1	5.6	0.5	*	0.1	0.4	1.5	3.3	9.0	24.6	66.8	200.2
1999.....	11.9	4.0	0.5	0.2	0.2	0.5	1.2	3.2	9.0	24.5	65.1	201.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Septicemia (A40-A41)												
Black, both sexes												
2004.....	15.6	14.5	*	*	0.7	1.9	4.5	13.5	31.6	69.9	155.6	320.3
2003.....	16.3	18.6	0.9	*	0.6	1.9	4.9	13.7	32.2	73.0	157.5	372.0
2002.....	16.3	13.5	*	0.3	0.4	1.9	4.4	12.9	33.2	74.7	171.2	367.0
2001.....	15.8	19.3	0.9	*	0.7	1.8	4.6	13.4	32.1	71.8	160.2	364.8
2000.....	16.2	16.4	1.0	*	0.6	2.0	4.7	13.6	31.4	71.9	173.4	392.6
1999.....	16.1	17.7	*	*	0.6	1.5	4.9	12.9	30.3	74.4	168.3	412.3
Black, male												
2004.....	14.6	16.3	*	*	0.7	2.0	4.7	15.1	36.2	81.1	189.0	341.7
2003.....	15.3	21.1	*	*	*	2.1	5.0	15.3	37.8	85.7	189.0	404.5
2002.....	15.0	14.8	*	*	*	1.9	4.9	14.7	35.9	86.1	212.3	393.5
2001.....	14.9	20.1	*	*	0.8	1.8	4.7	14.9	36.9	85.6	195.1	424.4
2000.....	14.9	16.0	*	*	*	2.3	4.6	16.3	35.5	81.2	207.1	427.8
1999.....	15.3	19.8	*	*	*	1.6	5.8	15.1	37.1	90.3	201.7	453.5
Black, female												
2004.....	16.4	12.7	*	*	0.8	1.8	4.3	12.0	27.9	61.9	136.7	311.9
2003.....	17.2	16.1	*	*	0.7	1.6	4.9	12.3	27.8	63.9	139.7	359.3
2002.....	17.4	12.1	*	*	*	1.9	4.1	11.4	30.9	66.6	148.0	356.9
2001.....	16.7	18.6	*	*	*	1.8	4.6	12.1	28.3	62.0	140.6	342.7
2000.....	17.4	16.9	*	*	*	1.7	4.8	11.3	28.2	65.3	154.4	379.5
1999.....	16.9	15.4	*	*	*	1.5	4.2	11.1	24.8	63.2	149.6	396.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All Under 1-4 5-14 15-24 25-34 35-44 45-54 55-64 65-74 75-84 85 years
ages 1/ 1 year 2/ years years years years years years years years years and over

Measles (B05)

All races 3/, both sexes

Table with 12 columns (years 2004-1999) and 12 rows (ages 1-85+), showing asterisks for all data points.

All races, male

Table with 12 columns (years 2004-1999) and 12 rows (ages 1-85+), showing asterisks for all data points.

All races, female

Table with 12 columns (years 2004-1999) and 12 rows (ages 1-85+), showing asterisks for all data points.

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Viral hepatitis (B15-B19)

All races 3/, both sexes

2004.....	1.8	*	*	*	*	0.2	1.4	5.3	4.3	3.4	3.6	2.7
2003.....	1.9	*	*	*	*	0.2	1.5	5.5	3.9	3.8	4.2	2.4
2002.....	2.0	*	*	*	0.1	0.2	1.6	5.8	4.0	4.2	4.6	3.9
2001.....	2.0	*	*	*	0.1	0.2	1.8	5.3	4.0	4.6	4.9	3.8
2000.....	1.9	*	*	*	*	0.2	1.7	5.0	3.7	4.6	5.3	4.0
1999.....	1.7	*	*	*	*	0.2	1.6	4.1	3.3	4.8	5.3	4.7

All races, male

2004.....	2.4	*	*	*	*	0.2	1.8	8.0	6.1	4.0	3.6	3.0
2003.....	2.5	*	*	*	*	0.2	2.1	8.3	5.4	4.2	4.8	2.8
2002.....	2.7	*	*	*	*	0.3	2.3	8.9	5.5	4.5	5.3	5.0
2001.....	2.5	*	*	*	*	0.2	2.5	8.2	5.3	4.8	5.3	4.4
2000.....	2.4	*	*	*	*	0.2	2.4	7.7	5.0	4.8	6.4	5.7
1999.....	2.2	*	*	*	*	0.3	2.3	6.3	4.3	5.1	6.6	6.9

All races, female

2004.....	1.3	*	*	*	*	0.1	1.0	2.7	2.5	3.0	3.5	2.5
2003.....	1.3	*	*	*	*	0.1	0.9	2.9	2.5	3.5	3.8	2.3
2002.....	1.4	*	*	*	*	0.2	1.0	2.8	2.7	3.9	4.1	3.3
2001.....	1.4	*	*	*	*	0.2	1.1	2.5	2.8	4.4	4.6	3.5
2000.....	1.4	*	*	*	*	0.2	1.1	2.4	2.6	4.5	4.7	3.3
1999.....	1.3	*	*	*	*	0.2	0.9	2.0	2.4	4.5	4.5	3.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Viral hepatitis (B15-B19)

White, both sexes

2004.....	1.8	*	*	*	*	0.1	1.4	5.2	3.8	3.1	3.2	2.5
2003.....	1.8	*	*	*	*	0.2	1.5	5.4	3.3	3.4	3.9	2.3
2002.....	2.0	*	*	*	0.1	0.2	1.6	5.6	3.5	3.7	4.4	3.7
2001.....	1.9	*	*	*	*	0.2	1.8	5.0	3.6	4.1	4.7	3.6
2000.....	1.8	*	*	*	*	0.2	1.7	4.6	3.2	4.2	5.1	3.9
1999.....	1.7	*	*	*	*	0.2	1.5	3.8	2.9	4.3	5.1	4.5

White, male

2004.....	2.4	*	*	*	*	0.1	1.9	7.8	5.3	3.6	3.3	2.7
2003.....	2.4	*	*	*	*	0.2	2.1	8.2	4.7	3.8	4.4	2.8
2002.....	2.6	*	*	*	*	0.3	2.3	8.7	4.8	4.1	5.2	4.9
2001.....	2.5	*	*	*	*	0.2	2.5	7.8	4.8	4.5	5.4	4.5
2000.....	2.4	*	*	*	*	0.2	2.4	7.2	4.3	4.3	6.2	5.7
1999.....	2.2	*	*	*	*	0.3	2.3	5.7	3.9	4.8	6.5	6.5

White, female

2004.....	1.2	*	*	*	*	*	0.9	2.6	2.3	2.7	3.2	2.4
2003.....	1.2	*	*	*	*	*	0.9	2.7	2.0	3.0	3.5	2.1
2002.....	1.3	*	*	*	*	*	0.9	2.7	2.4	3.3	3.9	3.2
2001.....	1.3	*	*	*	*	0.1	1.0	2.2	2.4	3.7	4.2	3.2
2000.....	1.3	*	*	*	*	0.1	1.0	2.1	2.2	4.1	4.5	3.2
1999.....	1.2	*	*	*	*	0.2	0.8	1.9	2.0	3.9	4.1	3.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Viral hepatitis (B15-B19)

Black, both sexes

2004.....	2.2	*	*	*	*	*	1.4	7.5	8.7	4.6	5.0	*
2003.....	2.2	*	*	*	*	*	1.7	7.4	8.0	5.4	6.0	*
2002.....	2.3	*	*	*	*	0.4	1.9	8.0	7.1	6.1	5.2	*
2001.....	2.2	*	*	*	*	*	2.1	7.9	6.7	5.6	5.7	*
2000.....	2.2	*	*	*	*	0.4	2.0	7.9	5.9	5.7	6.2	*
1999.....	2.0	*	*	*	*	*	2.1	6.9	5.0	6.9	6.1	6.4

Black, male

2004.....	3.1	*	*	*	*	*	1.5	11.7	13.8	6.7	*	*
2003.....	3.0	*	*	*	*	*	2.4	11.5	11.8	6.1	6.2	*
2002.....	3.0	*	*	*	*	*	2.5	12.4	10.5	7.1	*	*
2001.....	2.7	*	*	*	*	*	2.9	11.8	8.9	4.9	*	*
2000.....	2.9	*	*	*	*	*	2.9	12.3	8.1	7.0	7.6	*
1999.....	2.6	*	*	*	*	*	3.0	11.6	6.2	5.9	*	*

Black, female

2004.....	1.5	*	*	*	*	*	1.4	3.8	4.5	3.2	5.2	*
2003.....	1.6	*	*	*	*	*	1.1	4.0	5.0	4.9	5.9	*
2002.....	1.6	*	*	*	*	*	1.4	4.3	4.3	5.4	5.4	*
2001.....	1.8	*	*	*	*	*	1.4	4.7	4.9	6.0	6.8	*
2000.....	1.5	*	*	*	*	*	1.3	4.2	4.1	4.7	5.5	*
1999.....	1.5	*	*	*	*	*	1.3	2.8	4.1	7.6	6.5	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Human immunodeficiency virus (HIV) disease (B20-B24)

All races 3/, both sexes

2004.....	4.4	*	*	0.1	0.5	3.7	10.9	10.6	5.4	2.4	0.8	*
2003.....	4.7	*	*	0.1	0.4	4.0	12.0	10.9	5.4	2.4	0.7	*
2002.....	4.9	*	*	0.1	0.4	4.6	12.7	11.2	5.1	2.2	0.8	*
2001.....	5.0	*	*	0.1	0.6	5.3	13.0	10.5	5.2	2.1	0.7	*
2000.....	5.1	*	*	0.1	0.5	6.1	13.1	11.0	5.1	2.2	0.7	*
1999.....	5.3	*	0.2	0.2	0.5	6.8	13.8	10.7	4.8	2.2	0.6	*

All races, male

2004.....	6.6	*	*	*	0.5	4.5	15.7	16.3	9.0	4.0	1.4	*
2003.....	7.1	*	*	*	0.4	5.1	17.5	17.2	9.1	4.0	1.5	*
2002.....	7.4	*	*	*	0.4	5.9	18.8	17.7	8.5	3.9	1.4	*
2001.....	7.6	*	*	0.1	0.5	7.1	19.5	16.8	8.6	3.5	1.5	*
2000.....	7.9	*	*	0.1	0.5	8.0	19.8	17.8	8.7	3.8	1.3	*
1999.....	8.2	*	*	0.2	0.5	9.5	21.0	17.5	8.3	3.8	1.0	*

All races, female

2004.....	2.4	*	*	*	0.4	2.8	6.2	5.2	2.0	1.0	0.5	*
2003.....	2.4	*	*	*	0.4	2.8	6.5	4.8	2.1	1.0	0.3	*
2002.....	2.5	*	*	*	0.4	3.3	6.7	4.8	1.9	0.8	0.3	*
2001.....	2.5	*	*	*	0.6	3.5	6.7	4.4	2.0	0.9	*	*
2000.....	2.5	*	*	0.1	0.4	4.2	6.5	4.4	1.8	0.8	0.3	*
1999.....	2.5	*	*	0.2	0.5	4.1	6.7	4.1	1.6	0.8	0.3	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease (B20-B24)

White, both sexes

2004.....	2.4	*	*	*	0.1	1.6	6.2	5.5	2.8	1.2	0.5	*
2003.....	2.5	*	*	*	0.2	2.0	6.8	5.8	2.9	1.2	0.4	*
2002.....	2.6	*	*	*	0.2	2.2	7.0	5.8	2.8	1.0	0.4	*
2001.....	2.7	*	*	*	0.2	2.6	7.3	5.5	2.8	1.0	0.3	*
2000.....	2.8	*	*	0.1	0.2	3.1	7.5	5.9	2.8	1.0	0.4	*
1999.....	3.0	*	*	0.1	0.2	3.7	8.0	5.7	2.5	1.1	0.3	*

White, male

2004.....	3.9	*	*	*	0.2	2.4	9.9	9.2	5.0	2.2	0.9	*
2003.....	4.2	*	*	*	0.2	3.0	11.0	10.0	5.0	2.1	0.9	*
2002.....	4.3	*	*	*	0.2	3.3	11.6	9.8	4.9	1.8	0.6	*
2001.....	4.5	*	*	*	0.2	3.9	12.1	9.5	4.8	1.9	0.7	*
2000.....	4.7	*	*	*	0.2	4.6	12.4	10.1	5.0	1.9	0.7	*
1999.....	5.0	*	*	*	0.3	5.8	13.2	9.9	4.7	2.0	0.7	*

White, female

2004.....	0.8	*	*	*	*	0.7	2.4	1.9	0.7	0.4	*	*
2003.....	0.9	*	*	*	0.1	0.9	2.5	1.7	0.9	0.4	*	*
2002.....	0.9	*	*	*	0.1	1.0	2.5	1.8	0.8	*	*	*
2001.....	0.9	*	*	*	0.2	1.2	2.4	1.5	0.8	0.3	*	*
2000.....	1.0	*	*	*	*	1.5	2.6	1.7	0.8	0.3	*	*
1999.....	1.0	*	*	*	0.2	1.6	2.7	1.6	0.6	0.3	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Black, both sexes

2004.....	18.8	*	*	*	2.2	17.1	44.6	50.0	29.7	13.6	4.5	*
2003.....	19.6	*	*	0.5	1.9	17.3	49.2	50.6	29.6	14.5	4.7	*
2002.....	20.8	*	*	*	2.0	20.6	52.8	53.9	26.1	14.4	5.6	*
2001.....	21.1	*	*	0.4	2.7	23.2	53.7	51.2	27.7	12.8	5.3	*
2000.....	21.4	*	*	0.3	2.1	26.3	53.5	53.4	26.0	13.5	4.8	*
1999.....	21.8	*	*	0.6	2.3	27.3	56.1	51.8	26.2	13.5	3.6	*

Black, male

2004.....	25.9	*	*	*	2.2	19.2	60.3	74.9	50.3	23.4	7.5	*
2003.....	27.7	*	*	*	1.9	20.8	68.0	77.8	50.7	25.8	9.0	*
2002.....	29.5	*	*	*	1.9	24.5	74.0	85.0	44.4	26.3	12.0	*
2001.....	30.1	*	*	*	2.3	29.8	75.9	80.7	47.1	21.7	11.5	*
2000.....	31.0	*	*	*	2.0	32.3	77.5	86.0	46.0	24.6	9.4	*
1999.....	31.9	*	*	0.7	1.9	36.0	81.9	85.0	45.5	24.7	6.5	*

Black, female

2004.....	12.4	*	*	*	2.2	15.1	30.6	28.6	13.1	6.5	*	*
2003.....	12.3	*	*	*	1.9	14.2	32.5	27.3	12.5	6.4	*	*
2002.....	12.8	*	*	*	2.1	17.1	34.0	27.2	11.3	6.0	*	*
2001.....	12.9	*	*	*	3.1	17.3	34.1	25.9	12.1	6.5	*	*
2000.....	12.8	*	*	*	2.2	20.8	32.2	25.3	10.0	5.8	*	*
1999.....	12.6	*	*	*	2.7	19.5	33.4	23.4	10.7	5.6	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All Under 1-4 5-14 15-24 25-34 35-44 45-54 55-64 65-74 75-84 85 years
ages 1/ 1 year 2/ years years years years years years years years years and over

Malaria (B50-B54)

All races 3/, both sexes

Table with 12 columns (years 2004-1999) and 12 rows (ages 1-85+). All cells contain an asterisk (*).

All races, male

Table with 12 columns (years 2004-1999) and 12 rows (ages 1-85+). All cells contain an asterisk (*).

All races, female

Table with 12 columns (years 2004-1999) and 12 rows (ages 1-85+). All cells contain an asterisk (*).

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All Under 1-4 5-14 15-24 25-34 35-44 45-54 55-64 65-74 75-84 85 years
ages 1/ 1 year 2/ years years years years years years years years years and over

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)

All races 3/, both sexes

2004.....	2.6	4.1	0.4	0.1	0.2	0.4	1.0	3.0	4.3	7.1	13.2	22.5
2003.....	2.6	4.3	0.5	0.1	0.2	0.4	1.0	2.9	4.1	7.3	13.3	22.5
2002.....	2.3	5.6	0.4	0.2	0.2	0.4	1.0	2.3	3.3	6.7	12.5	21.7
2001.....	2.0	4.7	0.4	0.2	0.2	0.4	0.8	1.7	3.0	6.2	10.6	20.1
2000.....	2.0	4.4	0.5	0.2	0.2	0.4	0.8	1.6	2.9	5.8	11.1	21.2
1999.....	2.6	5.3	0.5	0.2	0.2	0.6	1.2	2.3	3.8	7.8	14.2	27.0

All races, male

2004.....	2.8	4.8	0.4	0.1	0.3	0.4	1.2	4.1	5.4	8.4	15.1	29.6
2003.....	2.8	4.2	0.5	0.1	0.2	0.4	1.3	4.0	5.1	8.5	15.8	26.5
2002.....	2.6	6.1	0.5	0.2	0.2	0.5	1.3	3.2	4.2	8.0	15.4	26.8
2001.....	2.1	4.9	0.5	0.1	0.2	0.4	1.0	2.2	3.7	6.8	13.2	23.6
2000.....	2.1	4.8	0.5	0.2	0.3	0.4	1.0	1.9	3.3	7.2	13.6	27.6
1999.....	2.7	6.1	0.6	0.2	0.2	0.6	1.5	2.7	4.2	9.2	17.2	31.9

All races, female

2004.....	2.3	3.4	0.4	0.1	0.2	0.3	0.8	1.9	3.2	6.0	11.9	19.3
2003.....	2.3	4.5	0.5	0.1	0.2	0.3	0.7	1.9	3.1	6.3	11.7	20.8
2002.....	2.1	5.0	0.3	0.2	0.2	0.4	0.7	1.4	2.6	5.6	10.6	19.5
2001.....	1.9	4.6	0.4	0.2	0.2	0.3	0.6	1.2	2.4	5.8	8.9	18.7
2000.....	1.9	3.9	0.4	0.2	0.2	0.3	0.6	1.2	2.6	4.7	9.5	18.6
1999.....	2.5	4.5	0.4	0.2	0.2	0.6	0.9	1.9	3.4	6.7	12.3	25.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)

White, both sexes

2004.....	2.6	3.7	0.4	0.1	0.2	0.3	0.9	3.0	4.0	6.8	13.0	22.5
2003.....	2.6	3.5	0.4	0.1	0.2	0.3	0.9	2.8	3.8	7.3	13.0	23.0
2002.....	2.4	5.0	0.4	0.2	0.2	0.4	0.8	2.1	3.0	6.6	12.4	22.0
2001.....	2.0	3.8	0.4	0.2	0.2	0.3	0.7	1.6	2.8	6.0	10.5	20.1
2000.....	2.0	3.4	0.4	0.2	0.2	0.3	0.7	1.4	2.7	5.7	11.0	21.0
1999.....	2.5	4.5	0.5	0.2	0.2	0.4	1.0	1.9	3.4	7.5	13.7	26.5

White, male

2004.....	2.9	4.2	0.4	*	0.3	0.3	1.1	4.1	5.1	8.1	14.9	30.1
2003.....	2.8	3.2	0.4	*	0.2	0.4	1.2	3.8	4.8	8.3	15.5	27.2
2002.....	2.6	5.3	0.5	0.2	0.2	0.4	1.1	3.0	3.7	7.9	15.2	27.2
2001.....	2.1	3.9	0.4	0.1	0.2	0.4	0.9	2.0	3.4	6.6	13.1	23.8
2000.....	2.1	3.9	0.4	0.2	0.3	0.3	0.8	1.7	3.1	7.0	13.3	26.9
1999.....	2.6	5.6	0.6	0.2	0.2	0.5	1.3	2.3	3.8	8.8	16.3	30.6

White, female

2004.....	2.4	3.0	0.3	*	0.2	0.3	0.7	1.9	3.0	5.7	11.7	19.0
2003.....	2.4	3.9	0.5	0.1	0.1	0.3	0.6	1.8	2.9	6.4	11.3	21.2
2002.....	2.2	4.6	*	0.2	0.2	0.4	0.6	1.3	2.4	5.5	10.5	19.8
2001.....	1.9	3.8	*	0.2	0.2	0.2	0.5	1.1	2.2	5.6	8.7	18.5
2000.....	1.9	2.9	0.4	0.2	0.2	0.2	0.5	1.1	2.3	4.6	9.5	18.7
1999.....	2.5	3.4	*	0.2	0.2	0.4	0.6	1.5	3.0	6.4	12.0	24.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)

Black, both sexes

2004.....	2.5	6.7	*	*	0.3	0.8	1.6	3.4	6.2	8.7	16.5	24.3
2003.....	2.5	8.0	1.0	*	*	0.6	1.8	4.3	6.1	7.0	16.2	19.5
2002.....	2.5	8.7	*	*	0.4	0.9	2.2	3.9	5.8	7.2	13.8	21.7
2001.....	2.2	9.5	*	*	0.4	0.8	1.6	2.9	5.1	8.3	11.1	20.6
2000.....	2.3	9.9	*	*	0.4	0.9	1.6	3.0	4.9	7.8	13.3	24.1
1999.....	3.5	9.9	0.9	*	0.3	1.6	3.0	5.4	7.2	11.8	20.5	34.1

Black, male

2004.....	2.9	7.4	*	*	*	0.9	1.8	4.9	8.9	10.5	19.8	29.8
2003.....	2.9	8.9	*	*	*	0.8	2.1	5.7	8.0	10.7	18.6	22.9
2002.....	2.9	9.6	*	*	*	1.1	2.6	5.5	8.4	8.8	16.9	28.9
2001.....	2.4	10.8	*	*	*	*	2.2	3.9	6.3	10.0	12.4	*
2000.....	2.5	9.9	*	*	*	0.9	2.1	3.9	5.7	10.2	18.2	32.3
1999.....	3.6	10.1	*	*	*	1.8	3.1	6.1	7.8	14.0	28.9	45.7

Black, female

2004.....	2.2	*	*	*	*	*	1.4	2.1	4.0	7.4	14.6	22.1
2003.....	2.2	7.1	*	*	*	*	1.4	3.1	4.6	4.3	14.8	18.2
2002.....	2.1	7.9	*	*	*	*	1.8	2.5	3.6	6.1	12.0	18.9
2001.....	2.0	8.2	*	*	*	0.9	1.1	2.1	4.2	7.2	10.4	20.2
2000.....	2.1	9.9	*	*	*	1.0	1.2	2.2	4.2	6.1	10.6	21.0
1999.....	3.4	9.7	*	*	*	1.4	2.9	4.8	6.7	10.3	15.8	29.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over
Malignant neoplasms (C00-C97)												
All races 3/, both sexes												
2004.....	188.6	1.8	2.5	2.5	4.1	9.1	33.4	119.0	333.4	755.1	1,280.4	1,653.3
2003.....	191.5	1.9	2.5	2.6	4.0	9.4	35.0	122.2	343.0	770.3	1,302.5	1,698.2
2002.....	193.2	1.8	2.6	2.6	4.3	9.7	35.8	123.8	351.1	792.1	1,311.9	1,723.9
2001.....	194.4	1.6	2.7	2.5	4.3	10.1	36.8	126.5	356.5	802.8	1,315.8	1,765.6
2000.....	196.5	2.4	2.7	2.5	4.4	9.8	36.6	127.5	366.7	816.3	1,335.6	1,819.4
1999.....	197.0	1.8	2.7	2.5	4.5	10.0	37.1	127.6	374.6	827.1	1,331.5	1,805.8
All races, male												
2004.....	198.4	1.8	2.6	2.7	4.8	8.6	29.1	124.3	376.7	907.6	1,662.1	2,349.5
2003.....	201.3	1.7	2.8	2.8	4.6	8.9	30.8	127.4	386.8	931.7	1,695.4	2,413.8
2002.....	203.8	2.0	2.7	2.9	4.9	9.2	31.5	128.0	399.8	964.8	1,711.3	2,491.1
2001.....	205.3	1.5	2.9	2.5	5.0	9.3	32.6	130.3	405.2	984.6	1,727.1	2,613.6
2000.....	207.2	2.6	3.0	2.7	5.1	9.2	32.7	130.9	415.8	1,001.9	1,760.6	2,710.7
1999.....	208.9	1.9	2.9	2.7	5.2	9.7	33.3	131.7	428.4	1,019.4	1,767.0	2,722.2
All races, female												
2004.....	179.1	1.9	2.4	2.2	3.4	9.6	37.7	113.8	293.2	627.1	1,023.5	1,340.1
2003.....	182.0	2.1	2.1	2.4	3.4	9.9	39.1	117.1	302.3	635.3	1,040.1	1,381.9
2002.....	183.0	1.6	2.4	2.4	3.6	10.2	40.0	119.8	306.0	648.5	1,046.7	1,391.1
2001.....	183.9	1.8	2.5	2.4	3.5	10.9	41.0	122.7	311.5	652.2	1,045.4	1,410.7
2000.....	186.2	2.3	2.5	2.2	3.6	10.4	40.4	124.2	321.3	663.6	1,058.5	1,456.4
1999.....	185.6	1.7	2.5	2.3	3.7	10.3	40.9	123.7	325.0	669.5	1,048.5	1,436.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms (C00-C97)												
White, both sexes												
2004.....	200.7	2.0	2.5	2.5	4.0	8.8	32.0	113.3	326.9	753.3	1,284.3	1,659.7
2003.....	203.7	2.1	2.6	2.7	4.1	9.1	33.6	116.0	336.6	769.5	1,303.4	1,701.4
2002.....	205.5	1.8	2.5	2.7	4.4	9.5	34.5	118.5	343.3	790.1	1,311.0	1,725.7
2001.....	206.4	1.7	3.0	2.4	4.3	9.8	35.3	121.2	349.1	797.8	1,311.1	1,765.2
2000.....	208.6	2.5	2.8	2.6	4.3	9.7	34.5	121.8	359.4	811.0	1,330.3	1,815.7
1999.....	208.7	1.8	2.9	2.4	4.3	9.8	35.3	121.9	365.2	820.7	1,323.7	1,798.8
White, male												
2004.....	209.9	1.9	2.5	2.8	4.7	8.6	28.2	117.5	364.9	896.3	1,652.7	2,348.9
2003.....	213.1	1.8	2.9	2.9	4.7	8.9	29.9	119.9	375.6	922.7	1,683.6	2,412.1
2002.....	215.5	2.2	2.7	2.8	5.0	9.1	30.5	121.8	386.0	954.8	1,695.3	2,486.8
2001.....	216.4	1.5	3.1	2.4	5.1	9.3	31.3	123.6	392.1	969.4	1,704.6	2,597.6
2000.....	218.1	2.8	3.1	2.8	5.0	9.2	30.9	123.5	401.9	984.3	1,736.0	2,693.7
1999.....	219.5	2.0	3.2	2.7	5.0	9.5	31.9	123.5	412.7	1,002.3	1,741.8	2,696.3
White, female												
2004.....	191.7	2.1	2.5	2.2	3.3	9.0	35.8	109.2	290.8	630.8	1,033.1	1,348.9
2003.....	194.6	2.3	2.3	2.5	3.4	9.4	37.3	112.1	299.8	638.9	1,046.3	1,386.5
2002.....	195.8	1.4	2.3	2.5	3.7	9.9	38.5	115.3	303.1	650.4	1,053.1	1,395.1
2001.....	196.7	2.0	2.8	2.5	3.5	10.4	39.3	118.9	308.6	652.9	1,049.8	1,416.7
2000.....	199.4	2.3	2.4	2.3	3.6	10.1	38.2	120.1	319.7	665.6	1,063.4	1,459.1
1999.....	198.2	1.5	2.7	2.2	3.7	10.1	38.7	120.2	320.7	669.4	1,049.6	1,438.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms (C00-C97)												
Black, both sexes												
2004.....	161.9	*	2.5	2.5	4.8	12.0	46.8	177.6	458.5	915.9	1,446.0	1,774.0
2003.....	164.3	*	2.2	2.6	3.9	12.2	48.9	185.6	471.0	925.4	1,487.9	1,853.1
2002.....	165.9	*	2.9	2.5	4.0	12.3	50.0	181.7	485.6	962.6	1,520.7	1,885.7
2001.....	167.1	*	1.9	2.6	4.3	12.9	51.4	184.4	486.1	991.2	1,543.9	1,931.7
2000.....	169.3	*	2.8	2.2	5.1	11.9	54.0	192.6	496.0	1,005.3	1,566.3	1,998.9
1999.....	171.3	*	2.0	2.7	5.5	12.2	53.9	192.9	521.4	1,019.7	1,579.3	2,008.4
Black, male												
2004.....	176.2	*	2.8	2.5	5.4	10.0	38.4	197.0	569.2	1,209.7	2,087.2	2,748.8
2003.....	178.3	*	2.4	2.8	4.5	10.3	41.7	207.0	583.8	1,221.5	2,144.2	2,825.5
2002.....	181.5	*	2.6	3.1	4.5	11.2	43.0	197.3	610.3	1,274.7	2,223.0	2,976.1
2001.....	184.5	*	2.4	2.9	5.0	10.5	44.6	204.8	604.2	1,335.3	2,290.0	3,209.9
2000.....	188.5	*	2.7	2.5	6.1	10.1	48.4	214.2	626.4	1,363.8	2,351.8	3,264.8
1999.....	191.0	*	2.1	2.8	6.4	11.4	47.3	221.3	659.0	1,369.8	2,359.4	3,340.7
Black, female												
2004.....	148.9	*	2.2	2.6	4.2	13.9	54.2	160.9	369.4	706.2	1,083.6	1,387.7
2003.....	151.4	*	2.1	2.3	3.3	13.9	55.4	167.2	380.4	714.6	1,116.9	1,475.3
2002.....	151.7	*	3.2	2.0	3.6	13.3	56.2	168.2	385.4	741.1	1,123.1	1,468.0
2001.....	151.3	*	*	2.2	3.6	15.0	57.3	166.8	390.9	748.4	1,125.0	1,457.5
2000.....	151.8	*	2.9	1.9	4.1	13.5	58.9	173.9	391.0	753.1	1,124.0	1,527.7
1999.....	153.3	*	1.9	2.6	4.6	13.0	59.8	168.5	411.1	772.7	1,142.6	1,510.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

All races 3/, both sexes

2004.....	2.7	*	*	*	0.1	0.1	0.6	2.9	6.1	10.1	13.7	18.0
2003.....	2.7	*	*	*	0.1	0.1	0.6	2.8	6.5	10.2	13.1	18.6
2002.....	2.7	*	*	*	0.1	0.1	0.6	2.7	6.8	10.4	13.5	18.7
2001.....	2.7	*	*	*	0.1	0.1	0.6	2.7	6.9	10.4	14.2	18.5
2000.....	2.7	*	*	*	0.1	0.1	0.6	2.8	6.8	10.7	13.2	19.4
1999.....	2.7	*	*	*	0.1	0.1	0.6	2.9	6.8	10.3	14.3	19.3

All races, male

2004.....	3.7	*	*	*	*	0.2	0.8	4.6	10.0	15.6	20.3	24.9
2003.....	3.7	*	*	*	0.1	0.2	0.9	4.6	10.6	15.6	19.4	26.9
2002.....	3.7	*	*	*	*	0.2	0.8	4.4	10.9	15.9	19.6	26.9
2001.....	3.7	*	*	*	*	0.2	0.9	4.4	11.3	16.0	20.6	25.8
2000.....	3.6	*	*	*	*	0.1	1.0	4.5	10.9	16.0	19.2	27.1
1999.....	3.6	*	*	*	0.1	0.1	0.9	4.6	10.8	15.8	20.9	27.1

All races, female

2004.....	1.7	*	*	*	*	*	0.4	1.2	2.5	5.6	9.2	14.9
2003.....	1.7	*	*	*	*	0.1	0.4	1.0	2.8	5.7	9.0	15.0
2002.....	1.7	*	*	*	*	*	0.4	1.1	3.0	5.8	9.5	15.1
2001.....	1.8	*	*	*	*	0.1	0.4	1.2	2.8	5.8	10.0	15.4
2000.....	1.8	*	*	*	*	0.1	0.3	1.3	2.9	6.3	9.2	16.3
1999.....	1.8	*	*	*	*	*	0.4	1.3	3.1	5.8	9.9	16.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
White, both sexes												
2004.....	2.7	*	*	*	*	0.1	0.5	2.5	5.6	9.9	13.7	18.6
2003.....	2.7	*	*	*	*	0.1	0.5	2.4	6.2	9.8	12.9	19.0
2002.....	2.7	*	*	*	*	0.1	0.5	2.4	6.3	10.2	13.7	18.9
2001.....	2.7	*	*	*	0.1	0.1	0.5	2.4	6.3	10.0	14.4	19.0
2000.....	2.7	*	*	*	*	0.1	0.5	2.4	6.0	10.2	13.1	19.7
1999.....	2.7	*	*	*	*	0.1	0.5	2.3	6.1	9.9	14.1	19.8
White, male												
2004.....	3.6	*	*	*	*	0.2	0.6	3.9	9.1	14.8	19.9	24.8
2003.....	3.7	*	*	*	*	0.2	0.8	4.0	9.7	14.6	18.6	27.2
2002.....	3.7	*	*	*	*	0.2	0.7	3.8	9.9	15.3	19.7	27.0
2001.....	3.6	*	*	*	*	0.1	0.7	3.9	10.1	15.1	20.4	26.4
2000.....	3.5	*	*	*	*	0.1	0.7	3.7	9.6	14.9	18.7	27.6
1999.....	3.5	*	*	*	*	*	0.7	3.6	9.6	15.1	20.3	27.3
White, female												
2004.....	1.8	*	*	*	*	*	0.3	1.1	2.4	5.8	9.4	15.7
2003.....	1.8	*	*	*	*	*	0.2	0.9	2.8	5.7	9.1	15.4
2002.....	1.8	*	*	*	*	*	0.3	1.0	2.8	5.9	9.6	15.3
2001.....	1.9	*	*	*	*	*	0.3	1.0	2.7	5.8	10.3	15.9
2000.....	1.9	*	*	*	*	*	0.3	1.2	2.7	6.3	9.4	16.5
1999.....	1.9	*	*	*	*	*	0.3	1.0	2.8	5.6	10.1	16.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
Black, both sexes												
2004.....	3.0	*	*	*	*	*	1.3	6.0	11.1	13.8	15.3	12.4
2003.....	2.9	*	*	*	*	*	1.1	5.1	11.2	15.4	15.3	15.5
2002.....	2.8	*	*	*	*	*	1.1	5.5	12.1	12.9	11.8	16.0
2001.....	2.9	*	*	*	*	*	1.2	5.1	12.6	15.5	13.9	12.9
2000.....	3.0	*	*	*	*	*	1.1	6.0	13.5	15.5	15.1	15.3
1999.....	3.2	*	*	*	*	*	1.2	7.1	13.5	15.2	16.9	14.3
Black, male												
2004.....	4.7	*	*	*	*	*	1.9	10.4	19.8	26.6	26.4	26.8
2003.....	4.6	*	*	*	*	*	1.2	9.0	21.1	28.7	27.9	25.0
2002.....	4.3	*	*	*	*	*	1.6	9.5	21.9	23.0	18.4	26.8
2001.....	4.4	*	*	*	*	*	1.7	8.7	23.3	27.1	23.5	*
2000.....	4.8	*	*	*	*	*	1.7	10.5	24.3	28.2	28.8	*
1999.....	4.8	*	*	*	*	*	1.8	12.1	23.2	25.1	31.7	24.6
Black, female												
2004.....	1.4	*	*	*	*	*	0.8	2.3	4.2	4.6	9.0	*
2003.....	1.4	*	*	*	*	*	1.0	1.7	3.2	6.0	8.1	11.7
2002.....	1.4	*	*	*	*	*	0.7	2.0	4.2	5.8	8.1	11.9
2001.....	1.5	*	*	*	*	*	0.8	2.1	3.9	7.4	8.4	10.5
2000.....	1.4	*	*	*	*	*	*	2.0	4.8	6.5	7.3	14.2
1999.....	1.7	*	*	*	*	*	0.7	2.9	5.7	8.2	8.6	10.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of esophagus (C15)												
All races 3/, both sexes												
2004.....	4.4	*	*	*	*	0.1	0.7	3.4	10.3	19.2	26.7	25.8
2003.....	4.4	*	*	*	*	0.1	0.6	3.4	10.5	19.7	26.4	25.0
2002.....	4.4	*	*	*	*	*	0.7	3.5	10.4	20.2	26.6	24.6
2001.....	4.4	*	*	*	*	0.1	0.7	3.6	10.1	20.6	26.3	25.8
2000.....	4.3	*	*	*	*	0.1	0.7	3.6	10.8	20.0	25.6	25.9
1999.....	4.3	*	*	*	*	0.1	0.7	3.5	10.6	20.1	24.4	26.8
All races, male												
2004.....	7.0	*	*	*	*	0.1	1.1	5.9	17.9	33.4	48.8	48.5
2003.....	6.9	*	*	*	*	0.1	1.1	6.0	18.5	34.1	47.8	45.3
2002.....	6.9	*	*	*	*	*	1.1	6.0	18.2	35.6	48.3	43.4
2001.....	6.9	*	*	*	*	0.1	1.1	6.3	17.6	36.0	47.4	48.1
2000.....	6.7	*	*	*	*	0.2	1.2	6.3	18.6	34.1	45.8	47.8
1999.....	6.6	*	*	*	*	0.1	1.1	6.1	18.3	35.3	44.4	48.0
All races, female												
2004.....	2.0	*	*	*	*	*	0.2	1.0	3.2	7.3	11.9	15.7
2003.....	2.0	*	*	*	*	*	0.2	1.0	3.2	7.7	12.0	16.1
2002.....	2.0	*	*	*	*	*	0.2	1.0	3.2	7.4	12.2	16.5
2001.....	2.0	*	*	*	*	*	0.2	1.0	3.2	7.9	12.4	16.5
2000.....	2.1	*	*	*	*	*	0.1	1.0	3.5	8.3	12.4	17.1
1999.....	2.0	*	*	*	*	*	0.2	0.9	3.6	7.6	11.5	18.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of esophagus (C15)

White, both sexes

2004.....	4.7	*	*	*	*	0.1	0.7	3.3	10.0	19.2	27.1	26.2
2003.....	4.7	*	*	*	*	0.1	0.7	3.2	10.2	19.6	26.7	25.8
2002.....	4.6	*	*	*	*	*	0.7	3.4	9.9	19.8	26.7	25.1
2001.....	4.6	*	*	*	*	0.1	0.7	3.4	9.4	19.8	26.5	26.3
2000.....	4.5	*	*	*	*	0.1	0.7	3.3	10.0	19.5	25.3	26.6
1999.....	4.4	*	*	*	*	0.1	0.6	3.2	9.8	19.4	24.2	27.2

White, male

2004.....	7.5	*	*	*	*	0.1	1.2	5.8	17.6	33.5	49.7	49.2
2003.....	7.4	*	*	*	*	0.1	1.1	5.7	18.0	33.9	48.8	46.7
2002.....	7.3	*	*	*	*	*	1.2	6.0	17.5	35.2	48.8	44.6
2001.....	7.2	*	*	*	*	0.1	1.1	5.9	16.4	34.9	47.9	48.8
2000.....	7.0	*	*	*	*	0.2	1.2	5.9	17.5	33.6	45.6	49.3
1999.....	6.9	*	*	*	*	0.1	1.1	5.7	17.0	34.4	44.0	48.0

White, female

2004.....	2.0	*	*	*	*	*	0.2	0.8	2.8	6.9	11.7	15.9
2003.....	2.0	*	*	*	*	*	0.2	0.8	2.7	7.5	11.7	16.5
2002.....	2.0	*	*	*	*	*	0.2	0.8	2.8	6.8	11.9	16.7
2001.....	2.1	*	*	*	*	*	0.2	0.9	2.7	7.1	12.3	16.8
2000.....	2.1	*	*	*	*	*	0.1	0.8	2.9	7.8	11.9	17.3
1999.....	2.0	*	*	*	*	*	0.1	0.7	3.1	6.9	11.2	18.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of esophagus (C15)												
Black, both sexes												
2004.....	4.0	*	*	*	*	*	0.7	5.2	15.6	24.7	27.2	21.4
2003.....	4.2	*	*	*	*	*	0.8	6.0	16.9	25.6	27.9	21.3
2002.....	4.2	*	*	*	*	*	1.0	5.1	17.3	28.5	29.6	20.8
2001.....	4.5	*	*	*	*	*	0.9	5.9	18.5	32.3	29.5	23.6
2000.....	4.5	*	*	*	*	*	0.8	6.6	19.8	28.2	33.7	22.2
1999.....	4.6	*	*	*	*	*	0.9	6.5	20.0	29.9	31.1	23.9
Black, male												
2004.....	5.8	*	*	*	*	*	1.0	8.3	25.5	41.3	47.9	38.7
2003.....	6.2	*	*	*	*	*	1.3	10.2	28.7	44.3	46.0	38.6
2002.....	6.3	*	*	*	*	*	1.4	8.3	30.1	49.2	51.9	37.5
2001.....	7.0	*	*	*	*	*	1.5	10.2	32.5	55.3	54.7	46.5
2000.....	6.7	*	*	*	*	*	1.2	11.2	34.0	47.0	58.4	38.1
1999.....	7.0	*	*	*	*	*	1.7	11.3	34.8	50.6	56.9	51.6
Black, female												
2004.....	2.3	*	*	*	*	*	*	2.5	7.5	12.9	15.6	14.6
2003.....	2.3	*	*	*	*	*	*	2.5	7.4	12.2	17.7	14.6
2002.....	2.3	*	*	*	*	*	*	2.4	7.0	13.8	16.9	14.4
2001.....	2.3	*	*	*	*	*	*	2.2	7.3	16.1	15.4	15.2
2000.....	2.5	*	*	*	*	*	*	2.7	8.3	14.9	19.8	16.3
1999.....	2.3	*	*	*	*	*	*	2.4	8.2	15.2	16.7	13.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of stomach (C16)

All races 3/, both sexes

2004.....	4.0	*	*	*	*	0.3	1.1	2.7	6.5	14.9	26.3	42.2
2003.....	4.2	*	*	*	*	0.3	1.1	3.0	6.7	15.2	27.6	43.6
2002.....	4.2	*	*	*	0.1	0.3	1.0	2.8	6.9	15.4	28.7	47.6
2001.....	4.3	*	*	*	0.1	0.3	1.0	2.9	7.0	16.5	30.2	45.4
2000.....	4.5	*	*	*	*	0.3	1.1	2.9	7.4	16.6	31.0	52.7
1999.....	4.6	*	*	*	*	0.3	1.1	3.0	7.8	17.1	31.6	50.1

All races, male

2004.....	4.8	*	*	*	*	0.3	1.2	3.5	9.1	20.6	37.2	62.9
2003.....	4.9	*	*	*	*	0.3	1.2	3.8	9.5	21.9	38.4	60.0
2002.....	5.0	*	*	*	*	0.4	1.1	3.7	9.5	22.5	40.6	67.2
2001.....	5.1	*	*	*	*	0.4	1.1	3.8	10.2	23.2	43.1	62.6
2000.....	5.3	*	*	*	*	0.3	1.3	3.8	10.5	23.7	43.9	75.6
1999.....	5.5	*	*	*	*	0.4	1.3	4.2	11.0	24.2	46.5	71.3

All races, female

2004.....	3.3	*	*	*	*	0.3	0.9	2.0	4.1	10.1	19.0	32.9
2003.....	3.4	*	*	*	*	0.3	1.0	2.1	4.1	9.7	20.3	36.4
2002.....	3.5	*	*	*	0.1	0.3	0.9	1.9	4.5	9.5	20.8	39.1
2001.....	3.5	*	*	*	*	0.3	0.9	2.0	4.1	10.9	21.7	38.3
2000.....	3.7	*	*	*	*	0.3	0.9	2.1	4.6	10.8	22.6	43.4
1999.....	3.7	*	*	*	*	0.3	1.0	1.9	4.9	11.3	21.9	41.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of stomach (C16)												
White, both sexes												
2004.....	3.8	*	*	*	*	0.3	0.9	2.3	5.5	12.9	23.7	38.8
2003.....	3.9	*	*	*	*	0.3	0.9	2.4	5.9	13.5	24.6	38.7
2002.....	4.0	*	*	*	0.1	0.3	0.8	2.3	5.9	13.4	25.4	42.2
2001.....	4.1	*	*	*	*	0.3	0.8	2.4	6.1	14.5	26.9	41.1
2000.....	4.2	*	*	*	*	0.2	0.9	2.5	6.5	14.5	27.3	47.3
1999.....	4.3	*	*	*	*	0.3	0.9	2.5	6.6	15.2	28.1	44.8
White, male												
2004.....	4.5	*	*	*	*	0.3	1.1	3.0	7.8	17.5	33.2	58.4
2003.....	4.6	*	*	*	*	0.3	1.0	3.1	8.3	19.2	34.3	53.5
2002.....	4.8	*	*	*	*	0.4	0.9	3.1	8.3	19.7	36.6	60.5
2001.....	4.9	*	*	*	*	0.3	1.0	3.2	8.8	20.2	38.7	57.7
2000.....	5.0	*	*	*	*	0.3	1.0	3.2	9.3	20.9	39.0	68.9
1999.....	5.1	*	*	*	*	0.3	1.0	3.5	9.3	21.7	41.5	63.4
White, female												
2004.....	3.1	*	*	*	*	0.2	0.8	1.6	3.4	8.8	17.1	29.9
2003.....	3.2	*	*	*	*	0.3	0.8	1.7	3.6	8.7	17.9	32.1
2002.....	3.2	*	*	*	*	0.2	0.7	1.5	3.7	8.0	17.9	34.2
2001.....	3.3	*	*	*	*	0.2	0.7	1.7	3.5	9.7	19.0	34.1
2000.....	3.5	*	*	*	*	0.2	0.7	1.7	3.9	9.2	19.6	38.6
1999.....	3.4	*	*	*	*	0.2	0.8	1.5	4.0	9.8	19.3	37.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of stomach (C16)												
Black, both sexes												
2004.....	5.3	*	*	*	*	0.5	1.6	5.3	12.8	30.2	50.4	71.3
2003.....	5.4	*	*	*	*	0.4	1.8	5.7	12.7	27.1	54.3	85.8
2002.....	5.6	*	*	*	*	0.6	1.6	5.4	12.9	30.9	56.8	100.4
2001.....	5.6	*	*	*	*	0.5	2.1	5.4	12.9	30.5	58.4	88.1
2000.....	5.9	*	*	*	*	0.7	2.0	5.7	13.8	32.0	63.9	101.7
1999.....	6.1	*	*	*	*	0.7	2.2	5.9	15.9	31.1	64.6	102.4
Black, male												
2004.....	6.5	*	*	*	*	*	1.9	7.2	19.4	47.3	75.2	100.3
2003.....	6.5	*	*	*	*	*	2.0	8.3	19.6	44.0	79.5	106.3
2002.....	6.6	*	*	*	*	0.8	1.8	7.7	18.9	46.7	79.8	137.2
2001.....	6.7	*	*	*	*	*	2.3	7.9	19.7	48.2	87.1	104.4
2000.....	7.0	*	*	*	*	0.8	2.9	7.9	20.6	46.7	91.1	136.1
1999.....	7.3	*	*	*	*	*	2.7	8.5	23.1	46.0	100.5	147.6
Black, female												
2004.....	4.2	*	*	*	*	*	1.4	3.6	7.5	18.1	36.3	59.9
2003.....	4.3	*	*	*	*	*	1.6	3.5	7.1	15.0	40.0	77.8
2002.....	4.7	*	*	*	*	*	1.4	3.4	8.0	19.8	43.9	86.3
2001.....	4.5	*	*	*	*	*	1.9	3.2	7.5	18.0	42.3	82.1
2000.....	5.0	*	*	*	*	*	1.3	3.7	8.4	21.6	48.6	88.9
1999.....	5.0	*	*	*	*	*	1.8	3.7	10.1	20.6	44.4	85.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over
Malignant neoplasms of colon, rectum and anus (C18-C21)												
All races 3/, both sexes												
2004.....	18.3	*	*	*	0.1	0.7	3.0	10.3	28.2	64.7	128.9	225.6
2003.....	19.2	*	*	*	0.1	0.7	3.0	10.9	29.8	70.5	134.7	239.7
2002.....	19.7	*	*	*	0.1	0.7	3.2	11.4	31.7	73.3	137.4	241.4
2001.....	20.0	*	*	*	0.1	0.8	3.2	11.6	31.5	75.3	139.4	255.7
2000.....	20.4	*	*	*	0.1	0.7	3.0	11.2	32.7	76.7	146.0	270.4
1999.....	20.5	*	*	*	0.1	0.7	3.0	11.1	33.5	77.4	146.7	271.7
All races, male												
2004.....	18.6	*	*	*	0.2	0.7	3.2	11.9	34.2	80.6	156.9	254.7
2003.....	19.6	*	*	*	0.1	0.8	3.2	12.5	36.0	88.0	164.4	273.3
2002.....	20.1	*	*	*	0.2	0.8	3.6	13.0	39.4	91.3	167.4	275.1
2001.....	20.2	*	*	*	0.1	0.8	3.5	13.1	38.3	93.7	169.5	298.3
2000.....	20.6	*	*	*	0.1	0.8	3.4	12.8	39.6	95.4	179.5	315.6
1999.....	20.6	*	*	*	0.1	0.8	3.0	12.5	40.5	95.1	181.9	324.0
All races, female												
2004.....	18.0	*	*	*	0.1	0.6	2.9	8.7	22.7	51.4	110.0	212.5
2003.....	18.9	*	*	*	*	0.7	2.7	9.3	24.0	55.9	114.8	224.8
2002.....	19.2	*	*	*	*	0.7	2.8	9.8	24.5	58.3	117.5	226.8
2001.....	19.8	*	*	*	0.1	0.7	2.9	10.2	25.3	60.1	119.6	237.8
2000.....	20.2	*	*	*	*	0.6	2.7	9.8	26.4	61.2	124.2	251.9
1999.....	20.3	*	*	*	0.1	0.6	2.9	9.7	27.0	62.9	123.8	250.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of colon, rectum and anus (C18-C21)												
White, both sexes												
2004.....	19.2	*	*	*	0.1	0.7	2.8	9.6	26.9	62.8	127.4	225.1
2003.....	20.2	*	*	*	0.1	0.7	2.7	10.0	28.2	68.7	133.3	238.5
2002.....	20.7	*	*	*	0.1	0.7	2.9	10.7	30.1	71.8	135.2	240.7
2001.....	21.0	*	*	*	0.1	0.7	3.0	10.8	30.1	73.4	136.9	255.3
2000.....	21.5	*	*	*	0.1	0.6	2.7	10.4	31.2	75.1	144.0	269.2
1999.....	21.6	*	*	*	0.1	0.7	2.8	10.4	32.0	75.6	144.5	270.9
White, male												
2004.....	19.5	*	*	*	0.1	0.7	3.0	11.2	32.8	78.2	154.6	255.4
2003.....	20.5	*	*	*	0.1	0.7	3.0	11.5	34.3	86.2	162.8	272.9
2002.....	21.1	*	*	*	0.1	0.8	3.3	12.2	37.6	89.8	164.2	273.6
2001.....	21.2	*	*	*	*	0.7	3.2	12.3	36.8	91.8	167.0	297.3
2000.....	21.6	*	*	*	*	0.7	3.1	12.0	37.6	93.4	176.9	313.3
1999.....	21.8	*	*	*	*	0.7	2.9	11.9	38.8	93.7	179.3	324.5
White, female												
2004.....	18.9	*	*	*	*	0.6	2.6	7.9	21.3	49.6	108.9	211.4
2003.....	19.8	*	*	*	*	0.6	2.4	8.4	22.5	53.8	113.4	223.2
2002.....	20.3	*	*	*	*	0.7	2.6	9.1	23.1	56.4	115.7	226.4
2001.....	20.7	*	*	*	*	0.7	2.7	9.3	23.7	57.9	116.8	237.7
2000.....	21.4	*	*	*	*	0.5	2.4	9.0	25.2	59.7	122.3	251.3
1999.....	21.4	*	*	*	*	0.6	2.6	8.9	25.7	60.5	121.7	249.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4/2/ years	5-14/ years	15-24/ years	25-34/ years	35-44/ years	45-54/ years	55-64/ years	65-74/ years	75-84/ years	85 years and over
---	----------	----------------	--------------	-------------	--------------	--------------	--------------	--------------	--------------	--------------	--------------	-------------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

Black, both sexes

2004.....	17.2	*	*	*	*	1.0	4.4	16.6	44.8	94.6	165.1	260.8
2003.....	18.2	*	*	*	*	0.9	5.0	18.7	48.1	100.4	172.0	281.5
2002.....	18.2	*	*	*	*	0.7	4.9	18.6	49.5	99.2	183.6	274.7
2001.....	18.6	*	*	*	*	1.0	4.9	19.0	49.8	104.0	187.6	286.5
2000.....	18.8	*	*	*	*	1.0	5.1	19.5	50.1	104.7	190.1	310.0
1999.....	18.7	*	*	*	*	1.0	4.8	17.7	52.7	106.2	193.1	306.1

Black, male

2004.....	17.4	*	*	*	*	1.0	4.5	19.1	53.4	120.4	214.6	278.2
2003.....	18.3	*	*	*	*	1.0	5.2	22.0	58.9	125.8	213.0	317.0
2002.....	18.6	*	*	*	*	1.0	5.8	21.2	61.2	121.6	237.5	328.1
2001.....	18.4	*	*	*	*	1.2	5.7	20.8	58.4	128.1	228.4	347.2
2000.....	19.1	*	*	*	*	1.0	6.0	22.0	62.5	131.5	241.0	377.1
1999.....	18.6	*	*	*	*	1.2	4.7	20.1	65.8	126.6	244.4	350.4

Black, female

2004.....	17.0	*	*	*	*	0.9	4.3	14.4	37.8	76.2	137.0	254.0
2003.....	18.1	*	*	*	*	0.9	4.8	16.0	39.4	82.3	148.8	267.8
2002.....	17.8	*	*	*	*	*	4.1	16.3	40.2	83.3	153.0	254.2
2001.....	18.8	*	*	*	*	0.9	4.3	17.5	42.8	87.0	164.7	264.0
2000.....	18.5	*	*	*	*	1.0	4.3	17.2	40.2	85.8	161.4	285.0
1999.....	18.9	*	*	*	*	0.9	4.9	15.6	42.2	91.8	164.4	289.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)												
All races 3/, both sexes												
2004.....	5.2	*	0.1	*	0.1	0.2	0.9	5.6	10.7	19.9	31.3	32.4
2003.....	5.1	*	0.1	*	0.1	0.2	0.9	5.4	9.9	19.8	30.7	33.4
2002.....	4.9	*	0.2	*	0.1	0.3	0.9	4.9	9.6	19.6	29.3	35.0
2001.....	4.7	*	*	*	0.1	0.2	0.9	4.8	9.3	18.8	29.0	34.1
2000.....	4.6	*	0.1	0.0	0.1	0.2	1.0	4.5	9.3	18.7	28.8	32.7
1999.....	4.4	*	0.2	*	0.1	0.2	1.0	4.1	8.8	19.0	28.0	30.6
All races, male												
2004.....	6.8	*	*	*	0.1	0.3	1.3	9.2	16.6	28.3	42.7	41.5
2003.....	6.6	*	*	*	0.1	0.3	1.3	8.9	14.9	27.9	44.6	47.5
2002.....	6.4	*	*	*	*	0.3	1.4	8.2	14.5	28.5	41.6	46.2
2001.....	6.1	*	*	*	*	0.2	1.3	7.7	13.9	26.8	41.0	50.3
2000.....	5.9	*	*	*	0.1	0.3	1.3	7.2	13.8	27.0	42.1	44.7
1999.....	5.6	*	0.3	*	0.1	0.2	1.4	6.5	12.9	27.5	39.0	40.1
All races, female												
2004.....	3.7	*	*	*	*	0.1	0.4	2.2	5.3	12.8	23.6	28.3
2003.....	3.5	*	*	*	*	0.2	0.5	2.1	5.3	13.0	21.4	27.2
2002.....	3.4	*	*	*	*	0.2	0.5	1.8	5.1	12.2	21.1	30.2
2001.....	3.4	*	*	*	*	0.2	0.4	1.9	5.0	12.1	21.1	27.2
2000.....	3.3	*	*	*	*	0.1	0.6	1.9	5.1	11.8	20.1	27.8
1999.....	3.3	*	*	*	*	0.2	0.6	1.8	5.1	11.9	20.9	26.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)												
White, both sexes												
2004.....	5.1	*	*	*	0.1	0.1	0.7	4.8	9.4	18.3	30.0	31.2
2003.....	4.9	*	*	*	0.1	0.1	0.7	4.6	8.7	18.3	29.5	32.1
2002.....	4.8	*	0.2	*	*	0.2	0.8	4.1	8.5	18.3	28.4	33.8
2001.....	4.6	*	*	*	0.1	0.2	0.6	4.0	8.2	17.6	27.9	33.3
2000.....	4.5	*	*	*	0.1	0.1	0.7	3.7	8.1	17.3	27.8	32.0
1999.....	4.4	*	0.2	*	0.1	0.1	0.8	3.4	7.6	17.8	27.2	29.9
White, male												
2004.....	6.5	*	*	*	*	0.2	1.0	7.9	14.2	25.9	41.4	39.9
2003.....	6.4	*	*	*	*	0.1	1.0	7.4	13.1	26.0	43.0	45.6
2002.....	6.1	*	*	*	*	0.2	1.1	6.8	12.6	26.5	40.0	45.1
2001.....	5.9	*	*	*	*	0.1	0.9	6.4	12.2	25.4	39.6	49.7
2000.....	5.7	*	*	*	*	0.2	1.0	5.7	11.8	25.0	41.1	43.1
1999.....	5.5	*	*	*	*	0.1	1.1	5.2	11.1	25.8	38.0	40.6
White, female												
2004.....	3.7	*	*	*	*	*	0.4	1.9	4.9	11.8	22.3	27.3
2003.....	3.5	*	*	*	*	0.1	0.5	1.8	4.6	11.7	20.3	26.2
2002.....	3.5	*	*	*	*	0.2	0.4	1.5	4.5	11.4	20.6	29.0
2001.....	3.3	*	*	*	*	0.2	0.3	1.6	4.4	10.9	20.1	26.5
2000.....	3.3	*	*	*	*	*	0.4	1.7	4.6	10.9	19.0	27.4
1999.....	3.3	*	*	*	*	*	0.5	1.6	4.4	11.0	20.1	25.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)												
Black, both sexes												
2004.....	5.2	*	*	*	*	0.4	1.4	9.6	17.4	25.6	34.4	36.1
2003.....	4.9	*	*	*	*	0.5	1.4	9.8	16.3	22.9	31.3	41.7
2002.....	4.6	*	*	*	*	0.5	1.4	8.7	14.8	23.0	30.9	38.0
2001.....	4.5	*	*	*	*	*	1.7	8.7	14.8	21.7	32.4	35.0
2000.....	4.4	*	*	*	*	0.4	1.8	8.6	14.1	23.6	28.7	31.6
1999.....	4.2	*	*	*	*	0.4	1.6	7.2	13.2	24.2	29.4	34.4
Black, male												
2004.....	7.3	*	*	*	*	*	2.2	16.7	29.9	40.3	44.0	46.7
2003.....	6.8	*	*	*	*	*	2.2	17.1	25.9	33.5	44.3	63.6
2002.....	6.5	*	*	*	*	*	2.2	15.1	24.2	37.7	47.0	41.8
2001.....	5.9	*	*	*	*	*	2.6	14.8	23.6	30.0	43.0	42.0
2000.....	6.1	*	*	*	*	*	2.4	14.8	23.5	37.3	39.7	48.4
1999.....	5.5	*	*	*	*	*	2.4	12.8	20.0	36.5	41.2	30.5
Black, female												
2004.....	3.3	*	*	*	*	*	*	3.5	7.3	15.0	29.0	31.9
2003.....	3.2	*	*	*	*	*	*	3.5	8.6	15.3	23.9	33.2
2002.....	2.9	*	*	*	*	*	0.7	3.2	7.3	12.6	21.8	36.6
2001.....	3.2	*	*	*	*	*	0.8	3.5	7.7	15.9	26.4	32.4
2000.....	2.8	*	*	*	*	*	1.2	3.3	6.5	14.0	22.5	25.3
1999.....	3.0	*	*	*	*	*	0.9	2.4	7.8	15.6	22.7	35.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of pancreas (C25)												
All races 3/, both sexes												
2004.....	10.8	*	*	*	*	0.2	1.2	6.3	19.8	44.3	76.0	97.7
2003.....	10.6	*	*	*	*	0.2	1.2	6.2	19.1	44.2	75.4	95.2
2002.....	10.5	*	*	*	*	0.1	1.3	6.1	19.2	44.8	75.0	94.5
2001.....	10.5	*	*	*	*	0.1	1.3	6.1	19.2	44.5	74.1	100.1
2000.....	10.4	*	*	*	*	0.1	1.2	6.3	19.4	44.1	76.5	96.4
1999.....	10.4	*	*	*	*	0.1	1.3	6.4	19.8	45.0	73.1	99.7
All races, male												
2004.....	10.9	*	*	*	*	0.2	1.5	8.1	24.1	52.6	84.6	102.0
2003.....	10.5	*	*	*	*	0.2	1.5	7.7	23.1	50.9	85.4	95.6
2002.....	10.5	*	*	*	*	0.1	1.6	7.8	23.6	51.4	84.0	100.9
2001.....	10.3	*	*	*	*	0.1	1.4	7.6	23.6	50.8	84.0	105.1
2000.....	10.3	*	*	*	*	0.2	1.4	8.0	23.6	50.4	85.9	102.9
1999.....	10.4	*	*	*	*	0.2	1.6	8.3	23.9	52.4	81.8	107.5
All races, female												
2004.....	10.7	*	*	*	*	0.1	0.9	4.7	15.8	37.3	70.1	95.8
2003.....	10.6	*	*	*	*	0.1	1.0	4.8	15.3	38.6	68.8	95.0
2002.....	10.5	*	*	*	*	*	1.0	4.4	15.1	39.3	69.1	91.7
2001.....	10.6	*	*	*	*	0.1	1.1	4.7	15.2	39.4	67.6	98.1
2000.....	10.5	*	*	*	*	0.1	1.0	4.6	15.5	38.8	70.4	93.8
1999.....	10.5	*	*	*	*	0.1	1.0	4.7	16.0	39.0	67.4	96.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of pancreas (C25)												
White, both sexes												
2004.....	11.4	*	*	*	*	0.1	1.1	6.1	19.1	43.5	75.6	97.1
2003.....	11.2	*	*	*	*	0.2	1.2	5.9	18.4	43.7	74.8	94.4
2002.....	11.1	*	*	*	*	0.1	1.3	5.8	18.7	43.7	74.3	94.3
2001.....	11.1	*	*	*	*	0.1	1.3	5.9	19.0	43.8	73.3	99.0
2000.....	11.0	*	*	*	*	0.1	1.2	6.1	18.9	43.1	75.3	94.7
1999.....	11.0	*	*	*	*	0.1	1.3	6.2	19.1	44.2	71.7	98.6
White, male												
2004.....	11.5	*	*	*	*	0.2	1.4	7.7	23.4	51.6	85.1	102.9
2003.....	11.2	*	*	*	*	0.2	1.4	7.2	22.5	50.8	85.4	96.0
2002.....	11.1	*	*	*	*	*	1.6	7.5	23.0	50.7	84.2	101.6
2001.....	11.0	*	*	*	*	0.2	1.4	7.4	23.5	50.0	83.8	106.5
2000.....	10.9	*	*	*	*	0.2	1.4	7.8	23.0	49.8	85.2	102.4
1999.....	10.9	*	*	*	*	0.2	1.6	7.9	22.9	51.6	80.8	107.4
White, female												
2004.....	11.3	*	*	*	*	*	0.9	4.4	15.1	36.7	69.1	94.5
2003.....	11.2	*	*	*	*	0.1	1.0	4.5	14.6	37.6	67.7	93.8
2002.....	11.0	*	*	*	*	*	1.0	4.2	14.5	37.8	67.7	91.1
2001.....	11.2	*	*	*	*	*	1.1	4.5	14.7	38.5	66.3	95.9
2000.....	11.1	*	*	*	*	*	0.9	4.4	15.0	37.5	68.8	91.5
1999.....	11.0	*	*	*	*	*	0.9	4.6	15.5	37.9	65.8	95.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of pancreas (C25)												
Black, both sexes												
2004.....	9.5	*	*	*	*	*	1.6	9.3	28.4	57.9	88.0	113.6
2003.....	9.5	*	*	*	*	*	1.8	10.0	27.3	56.3	93.1	112.9
2002.....	9.3	*	*	*	*	*	1.8	8.7	27.6	60.7	89.9	106.6
2001.....	9.1	*	*	*	*	*	1.6	9.0	25.5	58.4	92.7	121.3
2000.....	9.3	*	*	*	*	*	1.7	9.0	27.5	59.1	97.8	119.8
1999.....	9.4	*	*	*	*	*	2.0	9.6	28.2	59.5	94.0	118.6
Black, male												
2004.....	9.4	*	*	*	*	*	2.0	11.4	36.1	72.3	89.6	100.3
2003.....	9.2	*	*	*	*	*	2.4	13.3	33.2	63.2	97.9	104.3
2002.....	9.0	*	*	*	*	*	2.1	11.4	34.1	68.4	96.1	100.8
2001.....	8.6	*	*	*	*	*	1.7	11.1	30.3	68.1	100.3	99.8
2000.....	9.0	*	*	*	*	*	2.0	11.8	34.2	65.8	106.9	109.6
1999.....	9.4	*	*	*	*	*	2.4	13.1	36.5	67.9	106.7	114.8
Black, female												
2004.....	9.6	*	*	*	*	*	1.2	7.5	22.2	47.7	87.0	118.9
2003.....	9.7	*	*	*	*	*	1.3	7.2	22.5	51.5	90.4	116.3
2002.....	9.5	*	*	*	*	*	1.4	6.4	22.5	55.3	86.4	108.8
2001.....	9.6	*	*	*	*	*	1.4	7.1	21.6	51.6	88.4	129.2
2000.....	9.7	*	*	*	*	*	1.4	6.6	22.1	54.4	92.7	123.6
1999.....	9.4	*	*	*	*	*	1.6	6.6	21.6	53.6	86.9	120.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of larynx (C32)												
All races 3/, both sexes												
2004.....	1.2	*	*	*	*	*	0.1	1.1	3.2	5.7	6.6	6.5
2003.....	1.3	*	*	*	*	*	0.2	1.0	3.4	6.2	6.9	6.5
2002.....	1.3	*	*	*	*	*	0.2	1.1	3.4	5.9	7.6	5.5
2001.....	1.3	*	*	*	*	*	0.2	1.1	3.6	6.4	7.2	6.7
2000.....	1.4	*	*	*	*	*	0.1	1.1	3.6	6.6	8.1	6.5
1999.....	1.4	*	*	*	*	*	0.2	1.1	3.8	6.7	7.3	7.0
All races, male												
2004.....	2.0	*	*	*	*	*	0.2	1.8	5.4	10.1	13.0	15.6
2003.....	2.1	*	*	*	*	*	0.3	1.8	5.8	10.8	13.7	15.3
2002.....	2.1	*	*	*	*	*	0.3	1.8	5.8	10.0	14.9	13.2
2001.....	2.2	*	*	*	*	*	0.3	1.8	6.1	11.1	14.2	18.3
2000.....	2.2	*	*	*	*	*	0.2	1.8	6.2	11.4	15.7	17.4
1999.....	2.2	*	*	*	*	*	0.3	1.9	6.4	11.4	14.1	17.8
All races, female												
2004.....	0.5	*	*	*	*	*	*	0.4	1.2	2.0	2.3	2.4
2003.....	0.5	*	*	*	*	*	*	0.3	1.2	2.4	2.4	2.7
2002.....	0.5	*	*	*	*	*	*	0.4	1.2	2.5	2.8	2.2
2001.....	0.5	*	*	*	*	*	0.1	0.4	1.3	2.5	2.6	1.9
2000.....	0.6	*	*	*	*	*	*	0.4	1.2	2.7	3.1	2.1
1999.....	0.6	*	*	*	*	*	*	0.3	1.4	2.9	2.8	2.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of larynx (C32)												
White, both sexes												
2004.....	1.3	*	*	*	*	*	0.1	0.9	3.0	5.4	6.4	6.3
2003.....	1.3	*	*	*	*	*	0.2	0.9	3.1	5.8	6.6	6.2
2002.....	1.3	*	*	*	*	*	0.1	0.9	3.1	5.7	7.3	5.6
2001.....	1.3	*	*	*	*	*	0.2	0.9	3.2	6.0	6.9	6.6
2000.....	1.3	*	*	*	*	*	0.1	0.9	3.1	6.1	7.7	6.6
1999.....	1.4	*	*	*	*	*	0.1	0.9	3.4	6.4	7.2	6.7
White, male												
2004.....	2.0	*	*	*	*	*	0.2	1.5	4.8	9.5	12.4	15.0
2003.....	2.1	*	*	*	*	*	0.3	1.5	5.2	9.8	12.9	14.5
2002.....	2.0	*	*	*	*	*	0.2	1.5	5.1	9.5	14.1	13.3
2001.....	2.1	*	*	*	*	*	0.3	1.4	5.4	10.3	13.3	18.3
2000.....	2.1	*	*	*	*	*	0.2	1.5	5.3	10.4	15.0	17.6
1999.....	2.1	*	*	*	*	*	0.2	1.6	5.7	10.7	13.8	16.8
White, female												
2004.....	0.5	*	*	*	*	*	*	0.4	1.2	1.9	2.3	2.5
2003.....	0.5	*	*	*	*	*	*	0.3	1.1	2.4	2.4	2.5
2002.....	0.6	*	*	*	*	*	*	0.3	1.2	2.5	2.6	2.3
2001.....	0.5	*	*	*	*	*	*	0.3	1.2	2.3	2.7	1.7
2000.....	0.6	*	*	*	*	*	*	0.4	1.1	2.6	2.9	2.1
1999.....	0.6	*	*	*	*	*	*	0.3	1.2	2.8	2.9	2.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of larynx (C32)												
Black, both sexes												
2004.....	1.7	*	*	*	*	*	*	2.3	6.6	10.3	11.2	8.2
2003.....	1.8	*	*	*	*	*	*	2.5	7.5	11.5	11.7	11.1
2002.....	1.8	*	*	*	*	*	0.5	2.6	7.5	10.3	13.7	*
2001.....	1.8	*	*	*	*	*	0.4	2.8	7.9	11.4	11.7	8.3
2000.....	2.0	*	*	*	*	*	*	2.9	9.0	12.8	14.1	6.9
1999.....	1.8	*	*	*	*	*	0.6	2.9	8.6	11.6	10.0	8.3
Black, male												
2004.....	2.9	*	*	*	*	*	*	4.2	12.6	20.4	24.8	23.8
2003.....	3.1	*	*	*	*	*	*	4.5	13.6	23.8	27.9	27.1
2002.....	3.0	*	*	*	*	*	0.7	4.5	14.6	19.3	29.3	*
2001.....	3.1	*	*	*	*	*	*	5.1	14.2	21.3	27.7	*
2000.....	3.3	*	*	*	*	*	*	5.1	17.2	24.3	28.5	*
1999.....	3.1	*	*	*	*	*	0.9	5.2	15.9	21.8	22.8	24.6
Black, female												
2004.....	0.5	*	*	*	*	*	*	*	1.7	3.2	3.5	*
2003.....	0.6	*	*	*	*	*	*	0.8	2.5	2.8	*	*
2002.....	0.7	*	*	*	*	*	*	1.1	1.8	4.0	4.9	*
2001.....	0.7	*	*	*	*	*	*	*	2.8	4.5	*	*
2000.....	0.7	*	*	*	*	*	*	1.0	2.4	4.6	6.0	*
1999.....	0.7	*	*	*	*	*	*	0.9	2.7	4.4	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)												
All races 3/, both sexes												
2004.....	53.8	*	*	*	0.1	0.4	5.5	29.9	106.9	265.5	373.6	297.5
2003.....	54.4	*	*	*	0.1	0.4	5.6	30.3	111.0	269.3	377.8	298.9
2002.....	54.7	*	*	*	0.1	0.4	6.0	30.3	115.3	275.0	377.6	297.2
2001.....	54.8	*	*	*	0.1	0.4	6.2	30.7	117.7	279.7	371.4	302.7
2000.....	55.3	*	*	*	0.1	0.5	6.1	31.6	122.4	284.2	370.8	302.1
1999.....	54.5	*	*	*	0.1	0.4	6.0	31.3	123.4	281.5	362.1	297.0
All races, male												
2004.....	62.0	*	*	*	*	0.4	5.8	35.8	131.7	337.6	517.3	466.8
2003.....	62.9	*	*	*	*	0.4	6.1	36.5	136.7	346.6	525.1	475.1
2002.....	63.7	*	*	*	*	0.4	6.2	36.6	144.0	355.9	527.9	482.2
2001.....	64.7	*	*	*	*	0.4	6.7	37.4	147.1	370.0	529.5	512.4
2000.....	65.5	*	*	*	*	0.5	6.9	38.5	154.0	377.9	532.2	521.2
1999.....	65.4	*	*	*	*	0.5	6.6	38.2	158.0	377.2	527.6	529.2
All races, female												
2004.....	45.9	*	*	*	*	0.3	5.2	24.2	83.9	205.0	277.0	221.3
2003.....	46.1	*	*	*	*	0.4	5.1	24.4	87.1	204.8	279.4	221.0
2002.....	46.0	*	*	*	*	0.4	5.8	24.3	88.8	207.7	277.8	217.0
2001.....	45.3	*	*	*	*	0.4	5.7	24.3	90.5	204.9	267.5	215.0
2000.....	45.4	*	*	*	*	0.5	5.3	25.0	93.3	206.9	265.6	212.8
1999.....	44.1	*	*	*	*	0.4	5.5	24.6	91.5	203.1	254.5	203.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)												
White, both sexes												
2004.....	58.1	*	*	*	*	0.3	5.3	28.7	106.3	270.4	378.6	302.0
2003.....	58.7	*	*	*	0.1	0.4	5.4	28.9	110.8	274.2	382.6	302.4
2002.....	59.1	*	*	*	*	0.4	5.8	29.1	114.5	280.0	382.9	300.9
2001.....	59.1	*	*	*	*	0.4	6.0	29.6	117.8	283.4	374.7	305.4
2000.....	59.5	*	*	*	*	0.5	5.7	30.2	122.5	287.0	374.7	304.5
1999.....	58.5	*	*	*	*	0.5	5.8	29.7	122.9	283.5	363.8	298.8
White, male												
2004.....	65.9	*	*	*	*	0.4	5.5	33.3	127.6	338.1	517.5	472.2
2003.....	66.9	*	*	*	*	0.4	5.7	34.1	133.1	347.1	524.6	478.5
2002.....	67.7	*	*	*	*	0.4	5.8	34.2	139.3	356.4	527.8	486.6
2001.....	68.6	*	*	*	*	0.4	6.3	34.9	143.8	368.1	526.4	513.9
2000.....	69.4	*	*	*	*	0.5	6.1	35.7	150.8	374.9	529.9	522.4
1999.....	69.1	*	*	*	*	0.5	6.1	35.2	153.9	373.4	523.5	531.5
White, female												
2004.....	50.4	*	*	*	*	0.3	5.2	24.1	86.1	212.5	283.8	225.3
2003.....	50.7	*	*	*	*	0.4	5.1	23.8	89.8	212.2	286.6	224.4
2002.....	50.7	*	*	*	*	0.4	5.8	24.2	91.0	215.1	285.8	220.2
2001.....	49.8	*	*	*	*	0.4	5.7	24.3	93.3	211.9	274.0	218.1
2000.....	49.9	*	*	*	*	0.5	5.2	24.8	96.1	213.2	272.7	215.9
1999.....	48.2	*	*	*	*	0.4	5.4	24.4	93.8	208.6	259.1	205.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Black, both sexes

2004.....	42.7	*	*	*	*	0.5	7.7	45.4	141.6	279.6	384.4	277.5
2003.....	42.9	*	*	*	*	*	7.7	47.4	144.0	288.6	383.1	280.9
2002.....	43.2	*	*	*	*	0.4	8.7	46.9	152.4	295.3	381.3	272.3
2001.....	43.1	*	*	*	*	*	8.9	46.5	147.0	306.5	385.4	280.7
2000.....	43.9	*	*	*	*	0.6	10.1	50.4	152.5	315.8	375.0	288.1
1999.....	44.1	*	*	*	*	0.4	9.3	50.6	158.1	313.1	387.9	290.5

Black, male

2004.....	54.3	*	*	*	*	*	8.8	62.5	206.9	410.4	611.1	481.8
2003.....	54.9	*	*	*	*	*	9.3	63.6	210.0	427.1	622.2	502.5
2002.....	56.0	*	*	*	*	*	10.6	64.6	223.6	444.6	626.2	484.6
2001.....	57.2	*	*	*	*	*	11.0	66.2	215.2	476.4	659.8	549.2
2000.....	58.3	*	*	*	*	*	13.4	70.7	223.5	488.8	642.5	562.8
1999.....	58.9	*	*	*	*	*	11.3	72.6	237.6	484.4	661.0	574.2

Black, female

2004.....	32.1	*	*	*	*	*	6.8	30.7	89.0	186.2	256.3	196.5
2003.....	32.0	*	*	*	*	*	6.2	33.4	91.0	190.0	248.0	194.8
2002.....	31.6	*	*	*	*	*	7.0	31.7	95.3	189.3	242.6	191.0
2001.....	30.2	*	*	*	*	*	7.1	29.5	92.0	186.7	231.3	181.0
2000.....	30.8	*	*	*	*	*	7.3	32.9	95.3	194.1	224.3	185.9
1999.....	30.7	*	*	*	*	*	7.6	31.9	94.4	192.3	234.9	184.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant melanoma of skin (C43)												
All races 3/, both sexes												
2004.....	2.7	*	*	*	0.1	0.5	1.2	2.8	5.3	9.3	14.7	17.7
2003.....	2.7	*	*	*	0.1	0.5	1.3	2.9	5.5	8.7	14.6	17.6
2002.....	2.6	*	*	*	0.1	0.5	1.3	2.8	5.1	8.8	13.7	17.9
2001.....	2.6	*	*	*	0.1	0.5	1.4	3.1	5.3	9.2	13.3	17.8
2000.....	2.6	*	*	*	0.1	0.5	1.4	3.2	5.4	8.9	13.1	18.9
1999.....	2.6	*	*	*	0.1	0.5	1.4	3.1	5.3	9.0	13.4	16.4
All races, male												
2004.....	3.5	*	*	*	*	0.6	1.4	3.7	7.6	13.7	23.7	31.4
2003.....	3.5	*	*	*	0.1	0.6	1.5	3.8	7.8	12.6	23.3	30.2
2002.....	3.3	*	*	*	0.1	0.6	1.6	3.8	7.2	13.0	21.2	29.2
2001.....	3.4	*	*	*	0.1	0.5	1.7	3.9	7.3	13.9	21.3	29.9
2000.....	3.3	*	*	*	0.1	0.5	1.6	4.2	7.4	12.9	20.8	31.8
1999.....	3.3	*	*	*	0.1	0.5	1.8	4.0	7.5	13.2	20.9	27.5
All races, female												
2004.....	1.9	*	*	*	0.1	0.4	1.0	2.0	3.2	5.5	8.7	11.5
2003.....	1.9	*	*	*	*	0.4	1.0	2.1	3.5	5.4	8.7	12.1
2002.....	1.9	*	*	*	0.1	0.4	1.1	1.9	3.2	5.4	8.8	12.9
2001.....	1.9	*	*	*	*	0.4	1.1	2.3	3.3	5.3	8.1	12.8
2000.....	2.0	*	*	*	0.1	0.4	1.2	2.2	3.5	5.7	8.2	13.7
1999.....	1.9	*	*	*	*	0.4	1.0	2.2	3.3	5.5	8.5	11.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant melanoma of skin (C43)												
White, both sexes												
2004.....	3.3	*	*	*	0.1	0.6	1.4	3.3	6.1	10.5	16.2	19.2
2003.....	3.2	*	*	*	0.1	0.6	1.5	3.4	6.3	9.8	16.0	19.1
2002.....	3.1	*	*	*	0.1	0.6	1.6	3.3	5.8	9.9	15.1	19.2
2001.....	3.2	*	*	*	0.1	0.6	1.7	3.6	6.1	10.3	14.6	19.3
2000.....	3.1	*	*	*	0.1	0.6	1.6	3.7	6.1	9.9	14.3	20.3
1999.....	3.1	*	*	*	0.1	0.6	1.7	3.6	6.1	10.0	14.5	17.6
White, male												
2004.....	4.3	*	*	*	*	0.7	1.7	4.3	8.7	15.6	26.0	34.2
2003.....	4.2	*	*	*	0.1	0.7	1.8	4.4	8.8	14.2	25.5	32.9
2002.....	4.0	*	*	*	*	0.7	1.9	4.4	8.1	14.4	23.3	31.7
2001.....	4.1	*	*	*	0.1	0.6	2.0	4.6	8.4	15.5	23.3	32.7
2000.....	4.0	*	*	*	0.1	0.7	1.9	4.8	8.5	14.3	22.5	34.7
1999.....	4.0	*	*	*	0.1	0.6	2.1	4.6	8.5	14.7	22.6	29.7
White, female												
2004.....	2.3	*	*	*	0.1	0.5	1.2	2.3	3.8	6.2	9.5	12.4
2003.....	2.3	*	*	*	*	0.5	1.2	2.4	4.0	6.1	9.5	12.9
2002.....	2.3	*	*	*	0.1	0.5	1.3	2.3	3.6	6.1	9.6	13.8
2001.....	2.3	*	*	*	*	0.5	1.3	2.7	3.8	5.9	8.8	13.7
2000.....	2.3	*	*	*	*	0.5	1.4	2.6	3.9	6.3	8.8	14.5
1999.....	2.3	*	*	*	*	0.5	1.2	2.6	3.9	6.1	9.2	12.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of breast (C50)												
All races 3/, both sexes												
2004.....	14.1	*	*	*	*	1.0	5.7	15.0	29.1	44.9	72.2	124.2
2003.....	14.4	*	*	*	*	1.0	6.1	15.6	29.6	45.5	74.9	132.5
2002.....	14.5	*	*	*	*	1.1	6.1	16.1	29.5	46.5	76.5	134.6
2001.....	14.7	*	*	*	*	1.2	6.3	16.8	30.2	47.6	76.8	134.2
2000.....	15.0	*	*	*	*	1.2	6.3	16.9	31.2	49.1	78.9	147.4
1999.....	14.9	*	*	*	*	1.1	6.1	16.9	31.1	49.5	78.9	144.4
All races, male												
2004.....	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.1	3.3
2003.....	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.0	3.9
2002.....	0.3	*	*	*	*	*	*	0.2	0.5	1.0	2.2	3.4
2001.....	0.3	*	*	*	*	*	*	0.2	0.6	1.4	2.3	3.5
2000.....	0.3	*	*	*	*	*	*	0.2	0.7	1.4	2.2	4.2
1999.....	0.3	*	*	*	*	*	*	0.2	0.6	1.2	2.0	4.5
All races, female												
2004.....	27.5	*	*	*	*	2.0	11.3	29.3	55.8	81.6	119.5	178.6
2003.....	28.2	*	*	*	*	2.1	12.2	30.4	56.6	82.6	123.7	189.4
2002.....	28.3	*	*	*	*	2.1	12.0	31.4	56.2	84.4	125.9	191.5
2001.....	28.6	*	*	*	*	2.4	12.4	32.8	57.5	85.8	125.8	188.9
2000.....	29.2	*	*	*	*	2.3	12.4	33.0	59.3	88.3	128.9	205.7
1999.....	28.9	*	*	*	*	2.2	12.0	32.9	59.2	88.9	128.9	200.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of breast (C50)												
White, both sexes												
2004.....	14.6	*	*	*	*	0.8	5.0	13.6	28.1	44.8	73.0	124.8
2003.....	15.0	*	*	*	*	0.9	5.5	14.4	28.4	45.1	75.1	132.5
2002.....	15.1	*	*	*	*	0.9	5.3	14.9	28.5	46.2	76.6	135.3
2001.....	15.2	*	*	*	*	1.0	5.6	15.7	29.1	47.0	77.1	134.8
2000.....	15.7	*	*	*	*	1.0	5.7	15.8	30.2	49.2	79.4	147.3
1999.....	15.5	*	*	*	*	0.9	5.4	15.7	29.9	49.0	79.0	146.0
White, male												
2004.....	0.3	*	*	*	*	*	*	0.1	0.4	1.0	2.1	3.0
2003.....	0.3	*	*	*	*	*	*	0.2	0.5	1.0	2.0	3.9
2002.....	0.3	*	*	*	*	*	*	0.2	0.5	0.9	2.1	3.3
2001.....	0.3	*	*	*	*	*	*	0.2	0.6	1.4	2.2	3.3
2000.....	0.3	*	*	*	*	*	*	0.2	0.7	1.4	2.2	3.9
1999.....	0.3	*	*	*	*	*	*	0.2	0.5	1.2	2.0	4.4
White, female												
2004.....	28.6	*	*	*	*	1.7	10.1	27.0	54.3	82.2	121.3	179.7
2003.....	29.3	*	*	*	*	1.8	11.1	28.4	54.9	82.6	124.6	189.4
2002.....	29.5	*	*	*	*	1.8	10.7	29.4	55.0	84.6	126.5	192.6
2001.....	29.8	*	*	*	*	2.0	11.3	31.0	56.0	85.6	126.9	189.9
2000.....	30.7	*	*	*	*	2.1	11.3	31.2	57.9	89.3	130.2	205.5
1999.....	30.2	*	*	*	*	1.9	10.8	30.9	57.5	88.9	129.5	202.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of breast (C50)												
Black, both sexes												
2004.....	14.5	*	*	*	*	2.1	10.9	26.0	43.7	56.3	82.3	143.5
2003.....	15.1	*	*	*	*	2.2	11.1	26.5	45.4	59.3	90.8	161.6
2002.....	14.9	*	*	*	*	2.3	11.7	27.0	43.1	60.1	94.0	152.6
2001.....	15.0	*	*	*	*	2.6	11.1	27.3	44.6	63.5	92.3	152.0
2000.....	14.8	*	*	*	*	2.3	11.2	27.8	45.4	58.6	90.5	176.4
1999.....	15.0	*	*	*	*	2.3	11.6	28.7	46.3	63.3	94.5	147.3
Black, male												
2004.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2004.....	27.5	*	*	*	*	4.0	20.5	48.0	78.2	95.1	127.4	196.9
2003.....	28.5	*	*	*	*	4.1	20.8	48.9	81.1	100.1	141.0	222.4
2002.....	28.2	*	*	*	*	4.3	22.0	49.8	76.6	101.1	145.0	209.1
2001.....	28.3	*	*	*	*	5.0	20.8	50.3	79.8	106.6	141.8	205.9
2000.....	27.9	*	*	*	*	4.2	20.9	51.5	80.9	98.6	139.8	238.7
1999.....	28.2	*	*	*	*	4.3	21.7	53.0	82.2	106.2	145.4	199.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of cervix uteri (C53)

All races 3/, both sexes

2004.....	1.3	*	*	*	*	0.5	1.4	2.0	2.8	3.3	3.7	5.1
2003.....	1.3	*	*	*	*	0.5	1.4	2.2	2.8	3.1	4.3	5.7
2002.....	1.4	*	*	*	*	0.5	1.5	2.2	2.7	3.4	4.1	6.2
2001.....	1.4	*	*	*	*	0.5	1.6	2.4	2.9	3.2	4.6	7.2
2000.....	1.5	*	*	*	*	0.6	1.5	2.4	3.1	3.7	4.8	7.3
1999.....	1.5	*	*	*	*	0.5	1.6	2.5	3.2	3.8	4.6	7.9

All races, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2004.....	2.6	*	*	*	*	1.1	2.8	4.0	5.4	6.0	6.2	7.3
2003.....	2.7	*	*	*	*	1.0	2.8	4.3	5.5	5.6	7.2	8.3
2002.....	2.7	*	*	*	*	1.1	3.0	4.4	5.2	6.3	6.9	8.9
2001.....	2.8	*	*	*	*	1.1	3.1	4.8	5.5	5.8	7.7	10.1
2000.....	2.9	*	*	*	*	1.2	3.1	4.8	5.9	6.8	8.0	10.3
1999.....	3.0	*	*	*	*	1.1	3.1	4.9	6.2	6.9	7.7	11.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of cervix uteri (C53)

White, both sexes

2004.....	1.2	*	*	*	*	0.5	1.3	1.8	2.6	2.7	3.3	4.1
2003.....	1.2	*	*	*	*	0.4	1.3	2.0	2.6	2.6	3.6	4.9
2002.....	1.3	*	*	*	*	0.5	1.4	2.0	2.4	2.9	3.6	5.1
2001.....	1.3	*	*	*	*	0.5	1.5	2.2	2.6	2.7	4.0	6.2
2000.....	1.4	*	*	*	*	0.5	1.4	2.2	2.7	3.1	4.3	6.3
1999.....	1.4	*	*	*	*	0.5	1.4	2.4	2.8	3.1	4.0	6.8

White, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

White, female

2004.....	2.4	*	*	*	*	1.0	2.6	3.7	5.0	5.1	5.5	5.9
2003.....	2.5	*	*	*	*	0.9	2.7	4.0	5.0	4.8	6.0	7.1
2002.....	2.5	*	*	*	*	1.0	2.8	4.0	4.7	5.3	6.0	7.3
2001.....	2.7	*	*	*	*	1.1	3.0	4.5	5.0	5.0	6.6	8.8
2000.....	2.7	*	*	*	*	1.1	2.8	4.3	5.2	5.7	7.1	8.8
1999.....	2.8	*	*	*	*	1.1	2.9	4.7	5.4	5.7	6.7	9.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of cervix uteri (C53)

Black, both sexes

2004	2.0	*	*	*	*	0.8	2.2	3.7	5.3	7.6	8.8	16.1
2003	2.1	*	*	*	*	0.9	2.2	3.6	5.5	7.4	11.3	14.6
2002	2.2	*	*	*	*	0.8	2.5	4.1	5.0	9.0	10.8	17.5
2001	2.1	*	*	*	*	0.8	2.4	3.8	5.2	7.8	11.9	19.0
2000	2.4	*	*	*	*	0.9	2.6	4.3	6.3	9.5	11.9	19.7
1999	2.3	*	*	*	*	0.7	2.5	3.4	7.1	10.7	11.3	21.7

Black, male

2004
2003
2002
2001
2000
1999

Black, female

2004	3.9	*	*	*	*	1.6	4.2	6.9	9.6	13.0	13.8	22.4
2003	4.0	*	*	*	*	1.7	4.2	6.8	10.0	12.6	17.7	20.3
2002	4.2	*	*	*	*	1.5	4.7	7.6	8.9	15.4	16.9	24.2
2001	4.0	*	*	*	*	1.5	4.5	7.0	9.3	13.3	18.5	26.1
2000	4.5	*	*	*	*	1.8	4.9	7.9	11.3	16.3	18.6	27.0
1999	4.4	*	*	*	*	1.2	4.7	6.4	12.8	18.3	17.6	29.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

All races 3/, both sexes

2004.....	2.4	*	*	*	*	0.0	0.4	1.3	4.4	10.1	15.5	22.3
2003.....	2.4	*	*	*	*	0.1	0.3	1.3	4.7	10.1	15.2	22.8
2002.....	2.4	*	*	*	*	0.1	0.3	1.4	4.5	10.0	15.4	24.6
2001.....	2.4	*	*	*	*	0.1	0.3	1.4	4.4	9.7	16.4	24.7
2000.....	2.3	*	*	*	*	0.1	0.3	1.3	4.6	9.7	15.8	24.7
1999.....	2.3	*	*	*	*	0.1	0.3	1.3	4.3	9.8	16.4	24.0

All races, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2004.....	4.7	*	*	*	*	0.1	0.8	2.6	8.6	18.5	25.9	32.3
2003.....	4.7	*	*	*	*	0.1	0.7	2.6	9.0	18.5	25.3	33.0
2002.....	4.7	*	*	*	*	0.1	0.6	2.8	8.7	18.4	25.6	35.2
2001.....	4.7	*	*	*	*	0.2	0.6	2.8	8.4	17.8	27.2	35.0
2000.....	4.6	*	*	*	*	0.1	0.6	2.6	8.9	17.7	26.2	34.8
1999.....	4.5	*	*	*	*	0.1	0.6	2.5	8.2	17.8	27.1	33.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year	2/	years	years	years	years	years	years	years	years	and over

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

White, both sexes

2004.....	2.4	*	*	*	*	*	0.4	1.3	4.1	8.8	14.9	22.4
2003.....	2.4	*	*	*	*	*	0.3	1.3	4.3	9.1	14.4	22.7
2002.....	2.4	*	*	*	*	*	0.3	1.3	4.1	8.9	14.4	23.8
2001.....	2.4	*	*	*	*	0.1	0.3	1.4	4.0	8.8	15.7	24.5
2000.....	2.3	*	*	*	*	0.1	0.3	1.3	4.3	8.5	14.8	24.4
1999.....	2.4	*	*	*	*	*	0.3	1.3	4.0	8.9	15.8	23.4

White, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

White, female

2004.....	4.7	*	*	*	*	*	0.7	2.5	7.9	16.4	25.1	32.6
2003.....	4.7	*	*	*	*	*	0.6	2.5	8.3	16.9	24.1	32.8
2002.....	4.7	*	*	*	*	*	0.6	2.6	8.0	16.5	24.1	34.2
2001.....	4.8	*	*	*	*	0.2	0.6	2.7	7.8	16.2	26.2	34.7
2000.....	4.6	*	*	*	*	0.1	0.6	2.6	8.3	15.7	24.6	34.3
1999.....	4.6	*	*	*	*	*	0.6	2.5	7.7	16.3	26.1	32.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Black, both sexes

2004.....	2.9	*	*	*	*	*	0.5	1.9	8.0	23.0	26.1	26.2
2003.....	2.9	*	*	*	*	*	0.5	1.8	9.0	21.0	27.8	30.0
2002.....	3.0	*	*	*	*	*	0.4	1.9	8.4	22.3	28.3	38.9
2001.....	2.8	*	*	*	*	*	0.4	2.1	7.7	21.2	28.4	32.2
2000.....	2.9	*	*	*	*	*	0.4	1.6	8.3	22.7	30.5	33.2
1999.....	2.7	*	*	*	*	*	0.5	1.4	7.9	21.3	27.3	35.4

Black, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Black, female

2004.....	5.5	*	*	*	*	*	1.0	3.6	14.4	39.4	40.9	36.6
2003.....	5.5	*	*	*	*	*	1.0	3.3	16.2	36.0	43.5	41.7
2002.....	5.7	*	*	*	*	*	0.7	3.6	15.2	38.1	44.3	53.8
2001.....	5.4	*	*	*	*	*	0.7	3.9	13.9	36.2	44.3	44.2
2000.....	5.5	*	*	*	*	*	0.7	3.0	14.9	38.6	47.7	45.5
1999.....	5.2	*	*	*	*	*	0.9	2.6	14.3	36.3	42.5	48.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of ovary (C56)												
All races 3/, both sexes												
2004.....	5.0	*	*	*	0.1	0.2	1.0	4.0	9.6	20.0	32.1	37.7
2003.....	5.0	*	*	*	0.1	0.2	1.0	4.0	10.4	19.5	32.6	38.4
2002.....	5.1	*	*	*	0.1	0.2	1.0	4.3	10.3	19.7	33.4	38.1
2001.....	5.1	*	*	*	0.1	0.3	1.0	4.1	10.3	20.0	33.2	39.6
2000.....	5.0	*	*	*	0.1	0.3	1.0	4.2	10.1	19.6	33.2	40.9
1999.....	4.9	*	*	*	0.1	0.2	1.1	4.2	10.2	19.6	31.7	37.6
All races, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races, female												
2004.....	9.9	*	*	*	0.1	0.5	2.1	7.8	18.5	36.8	53.7	54.6
2003.....	9.9	*	*	*	0.1	0.4	2.0	7.8	20.0	35.7	54.4	55.4
2002.....	10.0	*	*	*	0.2	0.5	2.1	8.4	19.9	36.1	55.6	54.6
2001.....	9.9	*	*	*	0.1	0.5	2.1	8.1	19.9	36.6	55.0	56.1
2000.....	9.8	*	*	*	0.1	0.5	2.1	8.3	19.4	35.7	54.8	57.6
1999.....	9.6	*	*	*	0.1	0.4	2.2	8.3	19.6	35.7	52.3	52.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of ovary (C56)												
White, both sexes												
2004.....	5.5	*	*	*	0.1	0.2	1.1	4.0	9.9	20.5	33.3	38.6
2003.....	5.6	*	*	*	0.1	0.2	1.0	4.1	10.8	20.0	33.6	39.1
2002.....	5.6	*	*	*	0.1	0.2	1.1	4.5	10.7	20.4	34.3	39.0
2001.....	5.6	*	*	*	*	0.3	1.1	4.4	10.5	20.6	34.2	40.4
2000.....	5.5	*	*	*	*	0.3	1.1	4.4	10.5	20.1	34.3	42.0
1999.....	5.4	*	*	*	0.1	0.2	1.1	4.4	10.6	20.1	32.4	38.7
White, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
White, female												
2004.....	10.9	*	*	*	0.1	0.5	2.1	8.0	19.3	38.0	56.0	56.0
2003.....	11.0	*	*	*	0.1	0.5	2.1	8.1	21.0	37.1	56.3	56.4
2002.....	11.1	*	*	*	0.2	0.5	2.2	8.9	20.8	37.7	57.4	55.9
2001.....	11.0	*	*	*	*	0.5	2.1	8.6	20.5	38.0	56.8	57.3
2000.....	10.9	*	*	*	*	0.5	2.2	8.7	20.4	37.0	56.8	59.0
1999.....	10.6	*	*	*	0.1	0.4	2.3	8.7	20.5	36.9	53.6	54.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of ovary (C56)												
Black, both sexes												
2004.....	3.3	*	*	*	*	*	0.9	3.9	8.6	20.1	27.3	32.7
2003.....	3.1	*	*	*	*	*	1.0	3.8	8.6	17.7	27.5	36.8
2002.....	3.1	*	*	*	*	*	1.0	3.4	9.1	17.1	28.8	34.7
2001.....	3.1	*	*	*	*	*	1.1	3.0	10.0	18.3	26.9	36.5
2000.....	2.9	*	*	*	*	*	1.0	3.3	8.4	18.1	26.6	34.4
1999.....	3.0	*	*	*	*	*	0.8	3.5	9.1	18.5	28.7	28.7
Black, male												
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
Black, female												
2004.....	6.2	*	*	*	*	*	1.7	7.2	15.6	34.5	42.8	45.7
2003.....	6.0	*	*	*	*	*	1.9	7.1	15.5	30.2	43.1	51.0
2002.....	5.9	*	*	*	*	*	1.9	6.4	16.4	29.3	45.2	48.1
2001.....	5.9	*	*	*	*	*	2.1	5.7	18.0	31.2	42.0	50.1
2000.....	5.6	*	*	*	*	*	2.0	6.2	15.2	30.8	41.6	47.2
1999.....	5.7	*	*	*	*	*	1.5	6.6	16.4	31.5	44.8	39.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over
Malignant neoplasms of kidney and renal pelvis (C64-C65)												
All races 3/, both sexes												
2004.....	4.2	*	*	0.1	*	0.1	0.6	3.0	8.7	16.7	26.8	32.7
2003.....	4.2	*	*	0.1	0.1	0.1	0.7	2.9	9.0	16.9	27.0	33.2
2002.....	4.2	*	*	0.1	0.0	0.1	0.7	3.3	8.8	17.7	26.3	32.5
2001.....	4.2	*	0.1	0.1	0.1	0.1	0.8	3.3	9.2	17.6	26.3	32.3
2000.....	4.2	*	*	0.1	0.1	0.1	0.7	3.3	9.0	17.1	26.6	33.8
1999.....	4.0	*	*	0.1	*	0.1	0.8	3.1	8.9	16.7	24.6	31.4
All races, male												
2004.....	5.3	*	*	*	*	0.2	0.8	4.3	12.8	23.9	38.2	47.0
2003.....	5.4	*	*	*	*	0.2	1.0	4.3	13.3	23.7	39.5	48.7
2002.....	5.4	*	*	*	*	0.2	1.0	4.8	13.0	25.1	37.9	50.1
2001.....	5.4	*	*	*	*	0.1	1.1	4.9	13.4	25.2	37.4	52.9
2000.....	5.3	*	*	*	*	0.1	1.0	4.8	13.1	24.2	38.2	55.1
1999.....	5.0	*	*	*	*	0.2	1.1	4.5	12.9	23.6	35.7	49.4
All races, female												
2004.....	3.1	*	*	*	*	*	0.4	1.7	5.0	10.6	19.1	26.3
2003.....	3.1	*	*	0.1	*	0.1	0.4	1.6	4.9	11.2	18.7	26.4
2002.....	3.1	*	*	*	*	*	0.4	1.8	4.9	11.6	18.7	24.9
2001.....	3.1	*	*	0.1	*	0.1	0.5	1.9	5.3	11.3	19.0	23.7
2000.....	3.1	*	*	*	*	*	0.5	1.9	5.3	11.3	18.9	25.1
1999.....	3.0	*	*	*	*	*	0.5	1.8	5.1	11.0	17.4	24.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of kidney and renal pelvis (C64-C65)												
White, both sexes												
2004.....	4.6	*	*	0.1	*	0.1	0.6	3.1	9.1	17.1	27.3	33.7
2003.....	4.6	*	*	0.1	*	0.1	0.7	3.0	9.1	17.5	27.7	33.8
2002.....	4.6	*	*	0.1	*	0.1	0.7	3.3	9.1	18.3	26.8	33.2
2001.....	4.6	*	*	0.1	*	0.1	0.7	3.4	9.4	18.2	26.7	32.8
2000.....	4.6	*	*	*	*	0.1	0.7	3.4	9.3	17.5	27.0	34.5
1999.....	4.3	*	*	0.1	*	0.1	0.7	3.1	9.1	17.1	24.9	31.5
White, male												
2004.....	5.8	*	*	*	*	*	0.8	4.5	13.2	24.4	39.0	48.6
2003.....	5.9	*	*	*	*	0.1	1.0	4.4	13.6	24.3	40.4	50.0
2002.....	5.9	*	*	*	*	*	1.0	4.9	13.3	25.9	38.6	51.9
2001.....	5.8	*	*	*	*	*	1.0	5.0	13.6	26.0	37.7	54.4
2000.....	5.7	*	*	*	*	*	1.0	4.9	13.4	24.4	38.7	56.6
1999.....	5.4	*	*	*	*	0.1	1.0	4.5	13.3	23.9	36.3	49.8
White, female												
2004.....	3.4	*	*	*	*	*	0.4	1.8	5.1	10.9	19.4	27.0
2003.....	3.4	*	*	*	*	*	0.4	1.7	4.9	11.6	19.1	26.6
2002.....	3.4	*	*	*	*	*	0.4	1.8	5.1	11.9	18.9	25.1
2001.....	3.4	*	*	*	*	*	0.4	1.8	5.4	11.6	19.4	23.8
2000.....	3.4	*	*	*	*	*	0.4	1.9	5.4	11.6	19.2	25.5
1999.....	3.2	*	*	*	*	*	0.4	1.8	5.3	11.3	17.5	24.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of kidney and renal pelvis (C64-C65)												
Black, both sexes												
2004.....	2.8	*	*	*	*	0.4	0.8	3.1	8.0	15.8	25.8	28.2
2003.....	2.9	*	*	*	*	*	1.1	3.2	9.5	14.6	23.8	33.8
2002.....	2.8	*	*	*	*	*	0.8	3.5	8.6	15.4	26.1	28.2
2001.....	3.0	*	*	*	*	*	1.2	3.7	9.5	15.0	25.6	29.8
2000.....	2.9	*	*	*	*	*	0.9	3.4	8.8	16.5	25.6	29.7
1999.....	2.9	*	*	*	*	0.4	1.1	3.8	8.6	16.1	23.5	31.2
Black, male												
2004.....	3.6	*	*	*	*	*	1.1	4.5	12.1	24.0	37.3	38.7
2003.....	3.7	*	*	*	*	*	1.6	4.9	13.8	22.2	37.0	43.8
2002.....	3.5	*	*	*	*	*	1.2	4.8	13.9	21.6	36.1	39.7
2001.....	3.8	*	*	*	*	*	1.5	5.4	14.9	22.3	40.3	46.5
2000.....	3.6	*	*	*	*	*	1.0	4.9	13.1	26.3	38.2	49.6
1999.....	3.6	*	*	*	*	*	1.6	5.8	13.2	23.5	32.0	49.2
Black, female												
2004.....	2.2	*	*	*	*	*	*	1.8	4.7	10.0	19.4	24.0
2003.....	2.2	*	*	*	*	*	0.7	1.8	6.0	9.2	16.4	30.0
2002.....	2.2	*	*	*	*	*	*	2.4	4.2	11.0	20.5	23.8
2001.....	2.2	*	*	*	*	*	1.0	2.2	5.2	9.9	17.4	23.6
2000.....	2.1	*	*	*	*	*	0.8	2.2	5.2	9.6	18.6	22.3
1999.....	2.2	*	*	*	*	*	*	2.1	5.0	10.9	18.8	24.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of bladder (C67)

All races 3/, both sexes

2004.....	4.4	*	*	*	*	*	0.3	1.3	5.0	14.9	37.4	67.9
2003.....	4.3	*	*	*	*	*	0.2	1.3	5.0	14.9	36.4	64.8
2002.....	4.4	*	*	*	*	*	0.3	1.3	4.9	15.9	37.0	66.6
2001.....	4.3	*	*	*	*	*	0.3	1.2	5.0	15.4	35.7	68.9
2000.....	4.3	*	*	*	*	*	0.2	1.2	5.0	15.5	35.7	69.6
1999.....	4.3	*	*	*	*	*	0.2	1.3	5.2	15.9	35.3	68.5

All races, male

2004.....	6.2	*	*	*	*	*	0.3	2.0	7.9	24.0	64.7	133.0
2003.....	6.0	*	*	*	*	*	0.3	2.0	7.8	23.9	63.1	125.1
2002.....	6.0	*	*	*	*	*	0.3	1.8	7.7	25.2	63.0	128.9
2001.....	6.0	*	*	*	*	*	0.3	1.8	7.9	25.2	62.5	134.8
2000.....	5.9	*	*	*	*	*	0.3	1.7	7.9	25.2	62.6	139.4
1999.....	5.9	*	*	*	*	*	0.3	2.0	8.2	25.7	62.5	134.8

All races, female

2004.....	2.7	*	*	*	*	*	0.2	0.6	2.3	7.3	19.0	38.6
2003.....	2.6	*	*	*	*	*	0.2	0.6	2.4	7.4	18.5	38.1
2002.....	2.8	*	*	*	*	*	0.2	0.7	2.4	8.2	19.7	39.6
2001.....	2.7	*	*	*	*	*	0.3	0.6	2.3	7.2	18.1	41.4
2000.....	2.7	*	*	*	*	*	0.2	0.6	2.4	7.5	18.1	41.2
1999.....	2.7	*	*	*	*	*	0.2	0.6	2.5	7.8	17.7	41.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of bladder (C67)

White, both sexes

2004.....	5.0	*	*	*	*	*	0.2	1.3	5.3	15.8	39.1	69.6
2003.....	4.8	*	*	*	*	*	0.2	1.3	5.3	15.7	37.5	66.6
2002.....	4.9	*	*	*	*	*	0.3	1.3	5.1	16.5	38.1	67.7
2001.....	4.8	*	*	*	*	*	0.3	1.2	5.2	16.1	36.7	70.8
2000.....	4.8	*	*	*	*	*	0.3	1.2	5.3	16.2	36.4	72.0
1999.....	4.8	*	*	*	*	*	0.2	1.3	5.4	16.3	36.3	69.8

White, male

2004.....	7.1	*	*	*	*	*	0.3	2.1	8.4	25.7	68.3	139.3
2003.....	6.8	*	*	*	*	*	0.3	2.0	8.3	25.3	65.7	131.0
2002.....	6.8	*	*	*	*	*	0.4	1.9	8.2	26.5	65.3	134.1
2001.....	6.8	*	*	*	*	*	0.3	1.9	8.3	26.6	65.0	141.1
2000.....	6.7	*	*	*	*	*	0.3	1.9	8.4	26.5	64.7	145.7
1999.....	6.7	*	*	*	*	*	0.3	1.9	8.5	26.9	64.8	140.7

White, female

2004.....	2.9	*	*	*	*	*	0.2	0.6	2.4	7.3	19.2	38.1
2003.....	2.9	*	*	*	*	*	0.1	0.5	2.4	7.6	18.4	38.1
2002.....	3.0	*	*	*	*	*	0.2	0.7	2.3	8.1	19.8	38.9
2001.....	2.9	*	*	*	*	*	0.3	0.6	2.2	7.2	17.8	41.3
2000.....	2.9	*	*	*	*	*	0.2	0.6	2.5	7.5	17.8	42.1
1999.....	2.9	*	*	*	*	*	0.2	0.6	2.4	7.6	17.7	41.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of bladder (C67)												
Black, both sexes												
2004.....	2.4	*	*	*	*	*	0.5	1.6	4.1	11.7	27.2	60.1
2003.....	2.4	*	*	*	*	*	0.4	1.6	3.9	12.0	32.3	53.4
2002.....	2.5	*	*	*	*	*	*	1.1	4.5	14.3	31.0	63.3
2001.....	2.4	*	*	*	*	*	0.4	1.3	4.9	11.9	31.2	55.0
2000.....	2.3	*	*	*	*	*	*	1.0	4.1	13.3	34.7	49.7
1999.....	2.5	*	*	*	*	*	*	1.5	5.6	15.0	29.1	60.3
Black, male												
2004.....	2.6	*	*	*	*	*	*	2.3	5.9	14.5	39.5	81.5
2003.....	2.8	*	*	*	*	*	*	2.2	5.4	17.5	48.8	80.3
2002.....	2.7	*	*	*	*	*	*	1.2	5.6	18.3	47.6	91.1
2001.....	2.5	*	*	*	*	*	*	1.7	6.4	16.1	43.0	78.3
2000.....	2.7	*	*	*	*	*	*	1.4	5.8	17.6	52.7	88.8
1999.....	2.9	*	*	*	*	*	*	2.6	7.4	19.1	46.1	86.7
Black, female												
2004.....	2.2	*	*	*	*	*	*	0.9	2.6	9.7	20.3	51.6
2003.....	2.1	*	*	*	*	*	*	1.2	2.8	8.1	23.0	42.9
2002.....	2.3	*	*	*	*	*	*	0.9	3.7	11.6	21.6	52.6
2001.....	2.2	*	*	*	*	*	*	0.8	3.8	8.9	24.6	46.3
2000.....	2.0	*	*	*	*	*	*	*	2.7	10.3	24.5	35.2
1999.....	2.2	*	*	*	*	*	*	*	4.2	12.1	19.6	50.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
All races 3/, both sexes												
2004.....	4.4	*	0.6	0.8	0.5	0.9	2.0	4.6	9.5	15.7	19.6	16.6
2003.....	4.4	*	0.7	0.8	0.6	1.0	2.4	4.8	9.7	15.5	19.5	15.2
2002.....	4.4	*	0.7	0.8	0.5	1.0	2.2	5.1	9.6	16.2	19.1	16.4
2001.....	4.4	*	0.7	0.7	0.6	1.1	2.3	5.0	9.7	15.4	20.4	16.4
2000.....	4.5	0.5	0.7	0.8	0.6	1.0	2.4	5.1	9.6	16.5	20.0	16.9
1999.....	4.6	*	0.7	0.8	0.6	1.1	2.4	5.1	10.3	16.9	20.4	16.0
All races, male												
2004.....	4.9	*	0.7	0.8	0.6	1.1	2.5	5.8	11.8	18.4	23.7	19.4
2003.....	5.0	*	0.7	0.8	0.6	1.1	2.9	6.2	11.4	18.6	24.6	20.0
2002.....	5.0	*	0.8	0.9	0.7	1.1	2.8	6.2	11.7	20.2	23.8	20.1
2001.....	5.0	*	0.7	0.7	0.6	1.3	2.8	6.1	12.1	19.2	25.8	20.1
2000.....	5.1	*	0.7	0.8	0.7	1.3	3.1	6.2	11.8	20.2	24.3	22.4
1999.....	5.1	*	0.7	0.7	0.7	1.4	2.8	6.2	12.4	20.8	25.5	20.7
All races, female												
2004.....	3.9	*	0.6	0.8	0.4	0.7	1.6	3.4	7.3	13.4	16.8	15.4
2003.....	3.9	*	0.6	0.8	0.5	0.9	2.0	3.5	8.1	12.8	16.1	13.1
2002.....	3.9	*	0.7	0.7	0.4	0.8	1.7	4.1	7.7	12.8	16.0	14.8
2001.....	3.9	*	0.6	0.8	0.5	0.8	1.7	3.9	7.6	12.3	16.8	14.8
2000.....	3.9	*	0.6	0.7	0.5	0.7	1.7	4.0	7.6	13.4	17.3	14.7
1999.....	4.1	*	0.8	0.8	0.5	0.8	1.9	4.1	8.4	13.7	17.1	14.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

White, both sexes

2004.....	4.9	*	0.7	0.8	0.6	1.0	2.3	5.0	10.3	17.0	20.8	17.3
2003.....	5.0	*	0.7	0.9	0.6	1.1	2.7	5.2	10.5	16.6	20.7	15.7
2002.....	5.0	*	0.7	0.8	0.5	1.1	2.5	5.6	10.4	17.4	20.1	17.1
2001.....	5.0	*	0.7	0.8	0.6	1.2	2.6	5.4	10.5	16.5	21.6	17.0
2000.....	5.1	*	0.7	0.8	0.6	1.1	2.6	5.6	10.4	17.8	21.0	17.5
1999.....	5.2	*	0.8	0.8	0.6	1.3	2.6	5.6	11.1	18.2	21.4	16.6

White, male

2004.....	5.5	*	0.7	0.8	0.7	1.1	2.7	6.2	12.8	20.0	24.8	20.6
2003.....	5.6	*	0.8	0.9	0.7	1.3	3.2	6.7	12.3	19.8	25.8	20.7
2002.....	5.7	*	0.7	0.9	0.7	1.2	3.1	6.8	12.4	21.6	24.8	21.2
2001.....	5.7	*	0.8	0.7	0.7	1.5	3.2	6.7	12.9	20.5	27.1	20.5
2000.....	5.7	*	0.7	0.9	0.7	1.4	3.4	6.8	12.8	21.7	25.5	23.5
1999.....	5.8	*	0.8	0.8	0.6	1.5	3.1	6.7	13.3	22.3	26.8	21.8

White, female

2004.....	4.4	*	0.6	0.7	0.4	0.8	1.8	3.7	8.0	14.4	18.1	15.8
2003.....	4.4	*	0.6	0.9	0.5	1.0	2.3	3.8	8.7	13.9	17.2	13.4
2002.....	4.4	*	0.6	0.8	0.4	1.0	1.9	4.5	8.5	13.8	16.9	15.4
2001.....	4.4	*	0.7	0.8	0.5	0.9	1.9	4.3	8.2	13.0	18.0	15.6
2000.....	4.4	*	0.6	0.7	0.5	0.9	1.8	4.5	8.2	14.5	18.0	15.0
1999.....	4.6	*	0.8	0.8	0.5	1.0	2.1	4.5	9.1	14.8	17.9	14.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
Black, both sexes												
2004.....	2.0	*	*	0.7	0.5	0.7	1.3	2.8	4.9	8.0	10.0	10.4
2003.....	2.1	*	*	0.7	0.5	0.7	1.4	3.1	5.5	8.9	10.5	10.5
2002.....	2.1	*	1.1	0.8	0.5	0.6	1.2	2.7	5.6	8.7	11.8	9.5
2001.....	2.0	*	*	0.6	0.5	0.5	1.1	3.0	5.8	9.2	9.9	11.7
2000.....	2.1	*	*	0.8	0.6	0.7	1.6	2.9	5.7	8.1	11.9	10.6
1999.....	2.1	*	*	0.8	0.6	0.7	1.3	3.2	6.0	8.2	11.8	9.9
Black, male												
2004.....	2.2	*	*	0.7	*	0.8	1.9	3.9	6.7	7.9	14.2	*
2003.....	2.5	*	*	0.8	0.7	*	2.1	4.0	6.7	12.0	15.2	*
2002.....	2.3	*	*	1.0	*	0.8	1.3	2.7	8.1	10.4	16.4	*
2001.....	2.2	*	*	0.7	*	*	1.2	3.6	7.9	10.1	12.4	*
2000.....	2.4	*	*	0.9	0.7	1.1	2.0	3.7	7.1	9.8	13.0	*
1999.....	2.3	*	*	0.7	0.8	0.9	1.8	4.4	7.2	10.0	12.6	*
Black, female												
2004.....	1.8	*	*	0.8	*	*	0.7	1.9	3.4	8.1	7.6	11.4
2003.....	1.8	*	*	0.6	*	*	0.9	2.2	4.6	6.7	7.8	11.7
2002.....	1.9	*	*	*	*	*	1.1	2.8	3.7	7.4	9.3	10.3
2001.....	1.8	*	*	*	*	*	1.0	2.4	4.2	8.5	8.6	9.7
2000.....	1.9	*	*	0.6	*	*	1.2	2.2	4.6	6.9	11.3	10.7
1999.....	1.9	*	*	0.9	*	*	1.0	2.1	5.0	6.9	11.4	9.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
All races 3/, both sexes												
2004.....	18.6	0.6	0.8	0.9	1.5	2.1	3.4	9.0	25.8	68.2	139.8	189.4
2003.....	19.1	0.8	0.9	1.0	1.5	2.2	3.9	9.3	26.8	70.3	143.3	198.0
2002.....	19.5	0.7	0.8	0.9	1.7	2.3	3.9	9.9	28.2	73.9	143.6	197.5
2001.....	19.8	0.7	1.0	0.9	1.8	2.5	4.1	10.4	29.5	74.6	145.2	199.9
2000.....	20.1	1.0	0.9	0.9	1.8	2.3	4.1	10.7	30.0	76.3	149.2	204.6
1999.....	20.2	0.8	0.9	0.9	1.9	2.4	4.2	10.8	31.8	77.8	148.0	199.4
All races, male												
2004.....	20.4	*	0.9	1.0	1.9	2.4	4.2	11.0	32.1	87.4	181.3	254.9
2003.....	21.0	*	1.0	1.1	1.8	2.6	4.7	11.4	33.9	90.4	184.8	268.7
2002.....	21.4	*	0.8	1.0	2.0	2.7	4.6	11.9	35.0	95.2	188.0	269.1
2001.....	21.5	*	1.0	1.0	2.1	2.9	5.1	12.7	36.3	94.7	187.0	275.3
2000.....	21.6	1.1	1.0	1.1	2.3	2.7	4.8	13.0	36.1	96.8	192.6	281.8
1999.....	21.7	*	1.0	1.0	2.2	2.8	5.1	13.2	38.8	97.3	189.0	279.0
All races, female												
2004.....	16.8	*	0.6	0.7	1.2	1.7	2.7	7.0	19.9	52.0	111.8	160.0
2003.....	17.3	1.1	0.8	0.8	1.2	1.7	3.1	7.2	20.1	53.4	115.6	166.8
2002.....	17.7	*	0.7	0.8	1.4	1.9	3.2	7.9	21.9	56.2	114.1	166.5
2001.....	18.1	*	1.0	0.7	1.4	2.1	3.2	8.1	23.1	57.9	117.7	168.4
2000.....	18.6	*	0.9	0.7	1.4	1.9	3.4	8.5	24.4	59.5	120.9	173.1
1999.....	18.7	*	0.8	0.7	1.5	2.0	3.4	8.5	25.3	61.9	121.3	167.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
White, both sexes												
2004.....	20.2	0.6	0.8	0.9	1.6	2.0	3.4	8.7	25.7	68.9	143.4	194.9
2003.....	20.9	0.9	0.9	1.0	1.5	2.1	3.8	9.0	27.0	71.5	146.9	204.5
2002.....	21.3	*	0.8	1.0	1.8	2.2	3.9	9.6	28.0	75.4	147.6	204.0
2001.....	21.5	0.7	1.1	0.9	1.8	2.4	4.0	10.2	29.2	75.8	148.7	205.9
2000.....	21.9	1.2	1.0	1.0	1.8	2.2	4.0	10.4	30.1	78.0	153.3	208.7
1999.....	21.9	0.8	1.0	0.9	1.8	2.3	4.2	10.7	31.5	79.3	151.3	204.1
White, male												
2004.....	22.2	*	0.8	1.1	1.9	2.4	4.1	10.7	32.2	88.2	186.3	264.0
2003.....	23.1	*	1.1	1.1	1.8	2.6	4.6	11.1	34.2	92.8	191.0	279.5
2002.....	23.4	*	0.9	1.0	2.1	2.6	4.5	11.6	34.8	97.3	193.7	279.4
2001.....	23.3	*	1.1	1.0	2.1	2.8	4.8	12.6	35.9	96.3	191.8	284.8
2000.....	23.6	*	1.0	1.1	2.2	2.6	4.8	12.6	36.1	98.7	198.2	290.2
1999.....	23.6	*	1.0	1.0	2.2	2.6	5.0	13.0	38.6	99.1	193.4	285.8
White, female												
2004.....	18.3	*	0.7	0.7	1.2	1.6	2.6	6.8	19.5	52.4	114.2	163.8
2003.....	18.8	1.4	0.8	0.8	1.2	1.6	3.0	7.0	20.1	53.4	117.2	171.3
2002.....	19.2	*	0.8	0.9	1.4	1.8	3.2	7.5	21.5	56.8	116.7	171.2
2001.....	19.7	*	1.1	0.8	1.4	2.0	3.2	7.9	22.8	58.5	120.1	172.8
2000.....	20.3	*	1.0	0.8	1.3	1.8	3.2	8.3	24.4	60.6	123.8	175.5
1999.....	20.4	*	0.9	0.7	1.5	2.0	3.3	8.4	24.9	62.8	123.7	171.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
Black, both sexes												
2004.....	13.6	*	*	0.7	1.6	3.0	4.5	12.6	32.4	75.1	123.4	154.0
2003.....	13.4	*	*	0.8	1.4	2.8	5.2	13.0	31.1	73.1	126.1	150.5
2002.....	13.6	*	*	0.8	1.5	2.9	4.8	13.6	35.9	76.0	122.3	152.3
2001.....	14.1	*	*	1.0	1.8	3.4	5.6	13.2	36.7	78.0	126.9	155.7
2000.....	13.9	*	0.8	0.6	2.2	2.9	5.3	14.7	35.8	75.0	121.0	175.5
1999.....	14.1	*	*	0.9	2.3	2.9	5.5	13.8	38.9	76.2	126.8	162.0
Black, male												
2004.....	14.4	*	*	0.8	1.9	3.3	5.2	15.8	39.7	98.5	150.6	193.7
2003.....	13.9	*	*	0.9	1.6	3.4	6.4	16.6	40.0	87.0	148.9	184.5
2002.....	14.6	*	*	1.0	1.4	3.6	6.2	16.1	44.4	97.8	152.6	202.6
2001.....	15.1	*	*	1.3	2.0	3.9	7.3	16.0	46.0	99.2	159.5	206.5
2000.....	14.7	*	*	0.8	2.7	3.5	5.7	17.9	44.5	96.0	155.0	224.9
1999.....	15.1	*	*	1.1	2.6	3.8	6.8	17.3	47.4	95.4	161.1	220.3
Black, female												
2004.....	12.8	*	*	0.7	1.3	2.7	3.9	9.8	26.6	58.3	108.1	138.2
2003.....	12.9	*	*	0.8	1.1	2.4	4.2	9.9	23.8	63.3	113.2	137.3
2002.....	12.8	*	*	0.6	1.5	2.2	3.5	11.4	29.1	60.5	105.1	133.1
2001.....	13.1	*	*	0.6	1.6	2.9	4.0	10.8	29.2	63.0	108.6	136.8
2000.....	13.1	*	*	*	1.8	2.4	5.0	12.0	28.8	60.2	101.8	157.1
1999.....	13.2	*	*	0.7	2.0	2.1	4.4	10.8	32.0	62.6	107.6	140.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hodgkin's disease (C81)												
All races 3/, both sexes												
2004.....	0.4	*	*	*	0.2	0.3	0.3	0.4	0.7	1.2	1.9	2.2
2003.....	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7	1.2	2.2	2.4
2002.....	0.5	*	*	*	0.2	0.4	0.4	0.4	0.6	1.3	1.9	2.7
2001.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.6	1.2	2.2	2.2
2000.....	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7	1.3	2.1	2.5
1999.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.7	1.5	2.1	2.4
All races, male												
2004.....	0.5	*	*	*	0.2	0.3	0.3	0.5	0.9	1.6	2.5	3.2
2003.....	0.5	*	*	*	0.1	0.4	0.4	0.5	0.9	1.5	2.6	2.6
2002.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.8	1.8	2.5	3.6
2001.....	0.5	*	*	*	0.2	0.4	0.5	0.6	0.7	1.6	2.6	2.6
2000.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.7	1.6	2.5	3.7
1999.....	0.6	*	*	*	0.3	0.4	0.5	0.5	1.0	1.9	2.5	3.8
All races, female												
2004.....	0.4	*	*	*	0.1	0.3	0.2	0.3	0.5	0.9	1.5	1.7
2003.....	0.4	*	*	*	0.2	0.4	0.3	0.3	0.5	0.9	1.9	2.3
2002.....	0.4	*	*	*	0.2	0.4	0.3	0.4	0.4	0.9	1.6	2.3
2001.....	0.4	*	*	*	0.2	0.4	0.3	0.4	0.4	0.8	2.0	2.0
2000.....	0.4	*	*	*	0.2	0.3	0.3	0.3	0.6	1.0	1.8	2.0
1999.....	0.4	*	*	*	0.2	0.3	0.3	0.4	0.5	1.2	1.9	1.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hodgkin's disease (C81)												
White, both sexes												
2004.....	0.5	*	*	*	0.2	0.3	0.3	0.4	0.7	1.3	2.0	2.3
2003.....	0.5	*	*	*	0.2	0.4	0.4	0.4	0.7	1.2	2.2	2.5
2002.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.6	1.3	2.1	2.9
2001.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.6	1.3	2.3	2.4
2000.....	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7	1.3	2.2	2.6
1999.....	0.6	*	*	*	0.2	0.4	0.4	0.5	0.8	1.6	2.3	2.4
White, male												
2004.....	0.5	*	*	*	0.2	0.3	0.3	0.5	1.0	1.7	2.6	3.5
2003.....	0.5	*	*	*	0.1	0.5	0.5	0.5	0.9	1.5	2.7	2.7
2002.....	0.6	*	*	*	0.2	0.4	0.4	0.5	0.8	1.9	2.6	3.9
2001.....	0.6	*	*	*	0.2	0.4	0.5	0.6	0.7	1.7	2.7	2.7
2000.....	0.5	*	*	*	0.2	0.4	0.4	0.6	0.7	1.7	2.7	3.9
1999.....	0.6	*	*	*	0.3	0.4	0.5	0.5	1.1	2.0	2.7	3.8
White, female												
2004.....	0.4	*	*	*	0.1	0.3	0.2	0.3	0.5	0.9	1.6	1.7
2003.....	0.5	*	*	*	0.2	0.4	0.3	0.3	0.6	1.0	2.0	2.4
2002.....	0.5	*	*	*	0.2	0.4	0.3	0.4	0.5	0.9	1.7	2.5
2001.....	0.4	*	*	*	0.2	0.4	0.3	0.4	0.4	0.9	2.0	2.2
2000.....	0.5	*	*	*	0.2	0.4	0.3	0.3	0.6	1.0	1.9	2.1
1999.....	0.5	*	*	*	0.2	0.3	0.3	0.4	0.5	1.3	2.1	1.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)												
All races 3/, both sexes												
2004.....	7.1	*	*	0.1	0.3	0.6	1.2	3.7	9.9	26.1	54.6	75.0
2003.....	7.4	*	*	0.1	0.3	0.6	1.4	3.8	10.8	26.8	56.1	78.8
2002.....	7.6	*	*	0.1	0.3	0.7	1.5	3.9	11.5	28.8	57.2	77.3
2001.....	7.8	*	*	0.1	0.4	0.8	1.7	4.4	12.2	29.3	58.6	79.6
2000.....	8.1	*	*	0.1	0.5	0.7	1.7	4.6	13.1	30.4	61.0	80.2
1999.....	8.2	*	*	0.1	0.4	0.7	1.8	4.8	13.3	31.7	61.8	77.7
All races, male												
2004.....	7.6	*	*	0.1	0.4	0.8	1.7	4.8	12.3	32.5	67.3	95.8
2003.....	8.0	*	*	0.1	0.4	0.8	1.8	4.8	13.8	33.9	69.1	103.6
2002.....	8.1	*	*	0.2	0.4	0.9	2.0	5.0	14.4	36.0	70.7	99.7
2001.....	8.4	*	*	0.1	0.5	0.9	2.2	5.8	15.0	36.5	71.3	106.6
2000.....	8.6	*	*	0.1	0.6	0.9	2.1	6.0	15.9	37.4	74.5	105.9
1999.....	8.6	*	*	0.1	0.5	0.9	2.3	6.2	16.2	38.3	74.7	104.4
All races, female												
2004.....	6.6	*	*	*	0.3	0.5	0.8	2.6	7.6	20.7	46.0	65.7
2003.....	6.8	*	*	*	0.3	0.4	1.0	2.8	8.0	20.9	47.4	67.8
2002.....	7.1	*	*	*	0.2	0.5	1.1	3.0	8.8	22.8	48.2	67.6
2001.....	7.3	*	*	*	0.3	0.6	1.1	3.1	9.7	23.3	50.2	68.3
2000.....	7.6	*	*	*	0.3	0.6	1.2	3.3	10.5	24.7	52.1	69.8
1999.....	7.7	*	*	*	0.3	0.6	1.2	3.4	10.6	26.4	53.4	66.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)												
White, both sexes												
2004.....	8.0	*	*	0.1	0.3	0.6	1.2	3.6	10.2	27.4	57.6	78.8
2003.....	8.3	*	*	0.1	0.3	0.6	1.4	3.8	11.2	28.2	59.1	83.3
2002.....	8.5	*	*	0.1	0.3	0.6	1.5	3.9	11.9	30.2	60.3	82.1
2001.....	8.8	*	*	0.1	0.3	0.7	1.6	4.5	12.6	30.7	61.6	83.4
2000.....	9.1	*	*	0.1	0.4	0.7	1.7	4.7	13.7	32.1	64.3	84.3
1999.....	9.2	*	*	0.1	0.4	0.7	1.7	4.9	13.8	33.5	64.9	81.2
White, male												
2004.....	8.5	*	*	0.1	0.4	0.8	1.7	4.7	12.6	33.9	70.8	101.2
2003.....	8.9	*	*	*	0.4	0.7	1.8	4.8	14.4	35.9	73.0	109.7
2002.....	9.1	*	*	0.2	0.5	0.8	1.9	5.0	14.8	37.7	74.4	105.6
2001.....	9.3	*	*	0.1	0.4	0.9	2.1	5.8	15.3	38.0	74.7	111.5
2000.....	9.6	*	*	0.1	0.6	0.8	2.2	6.0	16.5	39.2	78.4	111.8
1999.....	9.6	*	*	0.1	0.5	0.8	2.2	6.3	16.6	40.2	78.0	108.0
White, female												
2004.....	7.5	*	*	*	0.3	0.4	0.8	2.6	7.9	21.9	48.6	68.7
2003.....	7.7	*	*	*	0.3	0.4	1.0	2.8	8.3	21.7	49.7	71.6
2002.....	8.0	*	*	*	0.2	0.5	1.1	2.9	9.1	23.9	50.8	71.8
2001.....	8.2	*	*	*	0.3	0.5	1.1	3.1	10.1	24.5	52.9	71.6
2000.....	8.6	*	*	*	0.2	0.6	1.1	3.4	11.1	26.1	55.0	73.1
1999.....	8.8	*	*	*	0.3	0.5	1.2	3.5	11.1	27.9	56.3	70.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)												
Black, both sexes												
2004.....	3.7	*	*	*	0.5	1.1	1.5	4.8	9.0	18.5	27.9	38.6
2003.....	3.6	*	*	*	0.4	1.1	1.9	4.8	8.9	18.2	28.3	31.2
2002.....	3.9	*	*	*	0.4	1.2	2.0	4.7	10.8	21.2	30.2	30.6
2001.....	4.0	*	*	*	0.5	1.2	2.2	5.0	10.9	21.0	30.5	39.3
2000.....	3.7	*	*	*	0.6	1.0	2.1	5.0	10.5	19.3	26.8	36.3
1999.....	4.0	*	*	*	0.5	1.1	2.4	4.7	11.1	20.1	31.8	40.5
Black, male												
2004.....	4.0	*	*	*	0.6	1.4	1.9	6.5	11.8	23.4	32.9	41.7
2003.....	3.8	*	*	*	*	1.3	2.3	6.5	11.2	20.4	31.0	34.4
2002.....	4.3	*	*	*	*	1.5	2.9	5.5	13.7	27.8	35.3	42.9
2001.....	4.5	*	*	*	0.7	1.3	3.2	6.7	13.6	27.7	36.8	54.5
2000.....	4.2	*	*	*	*	1.2	2.6	6.9	14.5	24.9	31.2	43.8
1999.....	4.6	*	*	*	*	1.5	3.2	6.6	14.6	25.6	40.6	59.8
Black, female												
2004.....	3.3	*	*	*	*	0.8	1.2	3.3	6.8	14.9	25.2	37.4
2003.....	3.4	*	*	*	*	0.8	1.5	3.3	7.1	16.6	26.8	30.0
2002.....	3.5	*	*	*	*	0.8	1.3	4.0	8.5	16.5	27.3	25.9
2001.....	3.6	*	*	*	*	1.1	1.3	3.6	8.7	16.3	26.9	33.7
2000.....	3.3	*	*	*	*	0.9	1.6	3.3	7.3	15.4	24.4	33.5
1999.....	3.5	*	*	*	*	*	1.7	3.2	8.2	16.3	26.9	33.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leukemia (C91-C95)												
All races 3/, both sexes												
2004.....	7.3	0.6	0.8	0.8	1.0	1.0	1.6	3.3	9.2	25.4	52.4	78.8
2003.....	7.4	0.7	0.8	0.8	1.0	1.1	1.8	3.3	9.3	25.7	53.9	80.3
2002.....	7.5	0.5	0.7	0.8	1.2	1.2	1.7	3.6	9.8	26.4	53.1	80.5
2001.....	7.5	0.6	1.0	0.8	1.2	1.3	1.7	3.8	10.0	26.7	53.3	81.4
2000.....	7.6	1.0	0.9	0.8	1.2	1.2	1.8	3.7	9.9	27.2	54.5	83.6
1999.....	7.5	0.7	0.8	0.7	1.2	1.2	1.8	3.7	10.6	27.2	52.8	82.4
All races, male												
2004.....	8.3	*	0.9	0.9	1.3	1.2	1.8	3.8	11.9	34.4	73.2	110.7
2003.....	8.4	*	0.9	0.9	1.2	1.4	2.0	4.0	11.9	35.1	75.3	111.8
2002.....	8.5	*	0.7	0.8	1.4	1.4	1.9	4.2	12.2	36.3	74.9	114.5
2001.....	8.5	*	1.0	0.9	1.4	1.5	1.9	4.2	12.8	35.7	74.2	117.5
2000.....	8.5	1.1	0.9	0.9	1.5	1.4	1.9	4.3	12.5	37.2	75.7	119.8
1999.....	8.4	*	0.8	0.8	1.5	1.4	1.9	4.3	13.1	36.8	73.2	119.2
All races, female												
2004.....	6.3	*	0.6	0.7	0.8	0.9	1.4	2.8	6.8	17.8	38.4	64.5
2003.....	6.4	1.1	0.8	0.7	0.7	0.8	1.6	2.7	7.0	17.8	39.6	66.4
2002.....	6.5	*	0.6	0.7	1.0	1.0	1.5	3.1	7.6	18.1	38.7	65.7
2001.....	6.6	*	1.0	0.7	1.0	1.1	1.5	3.3	7.5	19.2	39.5	66.3
2000.....	6.7	*	0.9	0.6	0.9	1.0	1.7	3.2	7.6	19.0	40.6	68.8
1999.....	6.7	*	0.7	0.7	1.0	1.1	1.6	3.2	8.3	19.3	39.6	67.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leukemia (C91-C95)												
White, both sexes												
2004.....	8.0	*	0.7	0.8	1.1	1.0	1.6	3.3	9.4	26.0	54.2	81.6
2003.....	8.2	0.9	0.9	0.9	1.0	1.1	1.8	3.4	9.6	26.6	56.0	83.4
2002.....	8.2	*	0.7	0.8	1.2	1.2	1.8	3.6	9.9	27.5	54.9	83.0
2001.....	8.3	0.6	1.1	0.8	1.2	1.3	1.7	3.7	10.2	27.5	55.1	84.7
2000.....	8.3	1.2	1.0	0.8	1.2	1.1	1.8	3.7	10.0	28.3	56.6	86.0
1999.....	8.3	0.8	0.9	0.7	1.2	1.2	1.8	3.7	10.8	28.0	54.5	85.4
White, male												
2004.....	9.1	*	0.8	0.9	1.3	1.2	1.8	3.8	12.2	35.0	75.8	115.3
2003.....	9.4	*	1.0	1.0	1.2	1.3	1.9	4.1	12.2	36.4	78.7	116.9
2002.....	9.3	*	0.8	0.8	1.5	1.4	2.0	4.1	12.3	37.5	77.4	119.1
2001.....	9.3	*	1.1	0.8	1.5	1.5	1.9	4.3	13.0	36.7	76.8	123.1
2000.....	9.4	*	0.9	0.9	1.4	1.3	1.9	4.2	12.6	38.5	78.5	123.6
1999.....	9.2	*	0.9	0.8	1.5	1.3	1.9	4.2	13.3	37.9	75.5	124.1
White, female												
2004.....	6.9	*	0.7	0.7	0.8	0.9	1.4	2.8	6.8	18.4	39.5	66.4
2003.....	7.0	1.4	0.8	0.7	0.7	0.9	1.6	2.7	7.1	18.2	40.6	68.6
2002.....	7.1	*	0.6	0.8	1.0	0.9	1.6	3.1	7.6	18.9	39.8	67.3
2001.....	7.3	*	1.1	0.7	1.0	1.1	1.6	3.2	7.6	19.7	40.8	68.6
2000.....	7.4	*	1.0	0.7	0.9	0.9	1.6	3.2	7.7	19.8	42.1	70.8
1999.....	7.3	*	0.8	0.7	1.0	1.2	1.6	3.2	8.4	19.8	40.6	69.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leukemia (C91-C95)												
Black, both sexes												
2004.....	4.8	*	*	0.6	1.0	1.3	2.0	4.0	10.3	25.2	41.1	60.1
2003.....	4.6	*	*	0.7	0.9	1.2	2.3	3.7	9.6	23.1	41.6	59.2
2002.....	4.7	*	*	0.6	0.9	1.2	1.6	4.3	11.2	22.3	42.1	64.7
2001.....	4.7	*	*	0.9	1.1	1.4	2.1	4.2	9.8	24.9	41.6	57.1
2000.....	4.8	*	*	0.5	1.4	1.6	2.0	4.8	10.7	22.7	38.2	66.0
1999.....	4.7	*	*	0.8	1.4	1.2	1.9	4.3	10.6	24.8	41.2	55.8
Black, male												
2004.....	5.3	*	*	0.7	1.1	1.4	2.2	4.9	12.8	37.0	52.3	79.5
2003.....	5.0	*	*	0.7	1.1	1.6	2.8	4.5	12.3	31.1	53.3	77.2
2002.....	5.1	*	*	0.7	1.0	1.4	1.8	5.2	13.7	31.5	56.8	86.8
2001.....	5.2	*	*	1.1	1.2	1.7	2.5	4.5	13.0	34.0	55.0	77.2
2000.....	5.3	*	*	0.6	1.9	1.9	2.0	5.5	13.8	31.6	54.8	98.0
1999.....	5.2	*	*	1.0	1.6	1.5	2.0	5.5	13.4	32.9	54.7	77.3
Black, female												
2004.....	4.4	*	*	*	0.8	1.2	1.8	3.2	8.3	16.7	34.8	52.4
2003.....	4.2	*	*	0.7	0.6	0.8	1.9	3.0	7.4	17.4	34.9	52.3
2002.....	4.3	*	*	*	0.7	1.1	1.4	3.6	9.2	15.7	33.8	56.3
2001.....	4.3	*	*	0.6	1.0	1.1	1.7	3.9	7.3	18.4	34.0	49.7
2000.....	4.2	*	*	*	0.9	1.3	2.1	4.3	8.1	16.4	28.8	54.1
1999.....	4.3	*	*	0.6	1.3	1.0	1.7	3.3	8.3	19.0	33.6	47.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

All races 3/, both sexes

2004.....	3.7	*	*	*	*	*	0.3	1.6	5.9	15.4	30.7	33.1
2003.....	3.9	*	*	*	*	*	0.3	1.7	5.9	16.5	31.1	36.3
2002.....	4.0	*	*	*	*	*	0.3	1.8	6.3	17.4	31.2	36.7
2001.....	3.9	*	*	*	*	*	0.3	1.7	6.6	17.4	30.9	36.3
2000.....	3.9	*	*	*	*	*	0.3	1.9	6.3	17.3	31.5	37.8
1999.....	3.9	*	*	*	*	0.1	0.3	1.8	7.1	17.2	31.0	36.6

All races, male

2004.....	3.9	*	*	*	*	*	0.4	1.9	6.9	18.8	38.1	44.6
2003.....	4.1	*	*	*	*	*	0.4	2.1	7.4	19.8	37.7	50.4
2002.....	4.2	*	*	*	*	*	0.3	2.2	7.6	21.0	39.7	50.8
2001.....	4.1	*	*	*	*	*	0.4	2.1	7.8	20.8	38.6	48.0
2000.....	4.0	*	*	*	*	*	0.4	2.1	6.9	20.4	39.6	51.8
1999.....	4.1	*	*	*	*	*	0.4	2.1	8.5	20.1	38.3	51.3

All races, female

2004.....	3.5	*	*	*	*	*	0.3	1.3	5.0	12.5	25.7	28.0
2003.....	3.7	*	*	*	*	*	0.2	1.4	4.6	13.8	26.7	30.1
2002.....	3.7	*	*	*	*	*	0.3	1.5	5.1	14.3	25.5	30.6
2001.....	3.8	*	*	*	*	*	0.3	1.3	5.5	14.5	25.8	31.3
2000.....	3.8	*	*	*	*	*	0.3	1.6	5.7	14.7	26.3	32.2
1999.....	3.8	*	*	*	*	*	0.3	1.5	5.9	14.8	26.2	30.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Multiple myeloma and immunoproliferative neoplasms (C88,C90)												
White, both sexes												
2004.....	3.8	*	*	*	*	*	0.3	1.4	5.3	14.1	29.4	32.0
2003.....	3.9	*	*	*	*	*	0.3	1.4	5.4	15.4	29.5	35.0
2002.....	4.0	*	*	*	*	*	0.2	1.6	5.6	16.3	30.2	35.7
2001.....	3.9	*	*	*	*	*	0.3	1.5	5.8	16.3	29.5	35.0
2000.....	3.9	*	*	*	*	*	0.3	1.6	5.6	16.2	30.0	35.2
1999.....	3.9	*	*	*	*	*	0.3	1.6	6.1	16.1	29.4	34.8
White, male												
2004.....	4.0	*	*	*	*	*	0.3	1.6	6.4	17.6	36.8	43.5
2003.....	4.2	*	*	*	*	*	0.4	1.8	6.8	18.9	36.5	49.7
2002.....	4.3	*	*	*	*	*	0.3	2.0	6.9	20.1	39.0	50.3
2001.....	4.2	*	*	*	*	*	0.4	1.9	6.8	19.7	37.4	47.1
2000.....	4.1	*	*	*	*	*	0.3	1.8	6.3	19.3	38.2	50.2
1999.....	4.1	*	*	*	*	*	0.3	1.9	7.6	19.0	36.9	49.7
White, female												
2004.....	3.5	*	*	*	*	*	0.2	1.1	4.3	11.1	24.3	26.8
2003.....	3.6	*	*	*	*	*	0.2	1.1	4.1	12.4	24.8	28.5
2002.....	3.7	*	*	*	*	*	0.2	1.2	4.4	13.1	24.3	29.4
2001.....	3.7	*	*	*	*	*	0.2	1.1	4.8	13.4	24.3	29.9
2000.....	3.8	*	*	*	*	*	0.2	1.4	5.0	13.6	24.6	29.1
1999.....	3.8	*	*	*	*	*	0.2	1.2	4.8	13.7	24.5	28.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Multiple myeloma and immunoproliferative neoplasms (C88,C90)												
Black, both sexes												
2004.....	4.7	*	*	*	*	*	0.7	3.4	12.4	30.4	53.0	53.9
2003.....	4.8	*	*	*	*	*	0.7	4.2	11.8	30.8	55.3	59.8
2002.....	4.7	*	*	*	*	*	0.7	4.1	13.1	31.5	48.7	55.5
2001.....	4.9	*	*	*	*	*	0.9	3.6	15.1	31.3	53.1	58.3
2000.....	5.0	*	*	*	*	*	0.8	4.5	13.9	32.1	55.1	71.6
1999.....	5.0	*	*	*	*	*	0.9	4.3	16.6	30.2	53.1	63.1
Black, male												
2004.....	4.7	*	*	*	*	*	0.7	3.9	14.2	36.6	64.0	71.5
2003.....	4.7	*	*	*	*	*	0.9	5.2	15.3	33.6	63.5	73.0
2002.....	4.7	*	*	*	*	*	0.8	4.6	15.7	37.1	59.1	71.8
2001.....	4.9	*	*	*	*	*	1.0	4.3	18.4	36.6	65.6	71.5
2000.....	4.8	*	*	*	*	*	*	5.0	15.0	38.5	67.8	79.6
1999.....	4.9	*	*	*	*	*	1.1	4.4	18.6	35.1	64.6	80.9
Black, female												
2004.....	4.7	*	*	*	*	*	0.7	3.1	11.0	26.0	46.7	46.9
2003.....	4.9	*	*	*	*	*	*	3.3	9.0	28.7	50.7	54.7
2002.....	4.7	*	*	*	*	*	*	3.7	11.1	27.6	42.9	49.3
2001.....	4.9	*	*	*	*	*	0.8	3.0	12.5	27.6	46.1	53.5
2000.....	5.2	*	*	*	*	*	0.9	4.0	12.9	27.6	47.9	68.7
1999.....	5.1	*	*	*	*	*	0.7	4.2	15.0	26.8	46.6	56.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All races 3/, both sexes

2004.....	21.1	0.6	0.8	0.7	1.3	1.7	4.3	12.7	34.4	76.8	143.0	213.1
2003.....	21.5	*	0.7	0.7	1.3	1.7	4.2	13.8	35.2	77.9	146.7	224.0
2002.....	21.8	0.6	0.8	0.7	1.3	1.8	4.5	13.7	36.6	80.8	146.7	230.1
2001.....	22.0	*	0.8	0.7	1.3	1.8	4.6	14.1	37.3	82.0	148.7	234.5
2000.....	22.4	0.6	0.8	0.6	1.3	1.8	4.6	14.1	38.3	85.5	152.6	239.8
1999.....	23.7	*	0.8	0.7	1.3	1.9	5.1	15.0	41.5	91.4	160.8	250.4

All races, male

2004.....	21.9	*	0.7	0.8	1.7	2.1	4.6	14.4	40.6	92.5	177.9	260.6
2003.....	22.3	*	0.8	0.7	1.6	2.1	4.7	15.7	41.5	93.4	180.8	275.0
2002.....	22.7	*	0.8	0.7	1.6	2.3	5.0	15.9	43.2	98.1	180.0	290.8
2001.....	22.6	*	0.9	0.7	1.7	2.2	4.9	16.4	43.8	98.5	181.1	295.2
2000.....	23.0	*	1.0	0.7	1.6	2.1	5.1	16.0	44.9	103.2	187.8	300.7
1999.....	24.5	*	0.8	0.8	1.7	2.4	5.8	17.3	49.1	110.4	196.1	318.9

All races, female

2004.....	20.3	*	0.9	0.6	0.9	1.4	3.9	11.0	28.6	63.7	119.5	191.8
2003.....	20.8	*	0.5	0.6	0.9	1.3	3.8	11.9	29.3	64.8	123.9	201.4
2002.....	21.0	*	0.8	0.7	0.9	1.4	4.1	11.6	30.4	66.3	124.6	203.8
2001.....	21.4	*	0.6	0.6	0.9	1.5	4.2	11.7	31.3	68.4	127.4	209.1
2000.....	21.9	*	0.7	0.6	1.0	1.4	4.2	12.2	32.2	70.8	129.6	215.0
1999.....	23.1	*	0.7	0.7	1.0	1.4	4.5	12.8	34.5	75.8	137.9	222.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

White, both sexes

2004.....	22.6	*	0.8	0.7	1.3	1.8	4.1	12.0	33.6	77.0	144.8	217.6
2003.....	23.1	*	0.6	0.7	1.3	1.7	4.1	13.1	34.6	78.3	148.6	227.9
2002.....	23.3	0.7	0.7	0.7	1.3	1.8	4.4	13.1	35.8	80.7	147.8	233.0
2001.....	23.5	*	0.8	0.7	1.3	1.8	4.4	13.3	36.2	81.4	149.5	238.1
2000.....	23.9	*	0.8	0.7	1.3	1.8	4.3	13.3	37.5	85.3	153.3	243.7
1999.....	25.3	*	0.8	0.7	1.3	1.9	4.9	14.3	40.2	91.3	161.2	252.4

White, male

2004.....	23.4	*	0.7	0.8	1.6	2.2	4.4	13.5	39.3	92.1	180.7	266.8
2003.....	23.8	*	0.7	0.7	1.6	2.1	4.6	14.8	40.6	93.5	183.4	280.9
2002.....	24.2	*	0.8	0.7	1.6	2.3	4.7	15.3	42.0	98.4	181.6	296.5
2001.....	24.0	*	1.0	0.7	1.7	2.2	4.8	15.5	42.0	97.7	182.1	300.2
2000.....	24.4	*	1.0	0.7	1.6	2.3	4.7	15.1	43.2	103.1	188.2	307.0
1999.....	25.9	*	0.9	0.8	1.6	2.4	5.6	16.2	47.4	109.9	197.5	323.6

White, female

2004.....	21.8	*	0.8	0.5	0.9	1.3	3.8	10.6	28.2	64.0	120.3	195.4
2003.....	22.5	*	0.5	0.6	1.0	1.2	3.6	11.4	28.9	65.4	125.0	204.4
2002.....	22.5	*	0.7	0.7	1.0	1.4	4.1	11.1	29.9	65.6	125.1	205.5
2001.....	22.9	*	0.7	0.6	0.9	1.4	4.0	11.2	30.7	67.7	127.8	212.1
2000.....	23.5	*	0.6	0.6	1.0	1.3	3.9	11.6	32.1	70.3	130.3	218.0
1999.....	24.6	*	0.7	0.6	1.1	1.4	4.2	12.4	33.4	75.8	137.4	223.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Black, both sexes

2004.....	17.2	*	0.8	0.9	1.6	1.8	5.9	19.4	47.5	90.8	147.2	186.7
2003.....	17.3	*	0.9	0.8	1.1	2.3	5.8	20.7	48.2	88.2	150.1	206.3
2002.....	18.1	*	1.1	0.8	1.3	2.3	6.3	20.1	51.0	96.2	157.6	222.1
2001.....	18.3	*	*	0.7	1.3	2.3	6.0	21.7	52.9	100.8	156.9	217.1
2000.....	18.3	*	1.0	0.6	1.4	1.6	7.5	22.1	51.9	100.1	162.0	210.8
1999.....	19.5	*	*	0.7	1.4	2.0	7.5	22.9	59.6	104.7	170.9	242.6

Black, male

2004.....	18.4	*	*	0.8	1.9	1.9	6.5	24.1	60.9	116.4	182.6	222.5
2003.....	18.3	*	*	0.8	1.4	2.4	6.5	25.0	61.1	111.3	187.6	241.9
2002.....	18.8	*	*	0.8	1.8	2.6	7.4	24.0	63.8	116.7	195.6	267.0
2001.....	19.4	*	*	0.8	1.7	2.7	6.3	26.6	67.5	127.9	193.6	283.7
2000.....	19.4	*	*	0.7	1.9	1.8	8.5	26.2	68.1	122.6	210.4	253.7
1999.....	20.6	*	*	0.6	2.0	2.3	8.3	28.9	76.1	133.2	206.3	292.9

Black, female

2004.....	16.2	*	*	0.9	1.3	1.8	5.5	15.3	36.8	72.5	127.1	172.5
2003.....	16.5	*	*	0.7	0.8	2.2	5.1	17.0	37.8	71.8	128.8	192.4
2002.....	17.4	*	*	0.7	0.8	2.0	5.3	16.7	40.7	81.7	136.1	205.0
2001.....	17.3	*	*	0.6	0.8	1.9	5.7	17.6	41.2	81.7	136.3	192.4
2000.....	17.3	*	*	*	1.0	1.5	6.6	18.6	38.8	84.3	134.7	194.9
1999.....	18.5	*	*	0.8	0.9	1.6	6.7	17.8	46.4	84.5	151.1	223.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All races 3/, both sexes

2004.....	4.6	1.7	0.3	0.2	0.3	0.4	0.7	1.7	4.5	12.7	35.8	77.7
2003.....	4.7	1.5	0.3	0.2	0.2	0.4	0.8	1.7	4.4	12.8	36.4	80.5
2002.....	4.6	1.7	0.4	0.2	0.2	0.4	0.8	1.8	4.5	12.8	36.1	78.7
2001.....	4.8	1.3	0.4	0.3	0.3	0.4	0.8	1.8	5.0	14.1	37.0	82.7
2000.....	4.8	1.6	0.3	0.2	0.2	0.4	0.8	1.8	5.0	13.8	37.6	84.9
1999.....	4.7	1.5	0.4	0.2	0.3	0.4	0.8	1.8	4.8	14.6	36.6	82.9

All races, male

2004.....	4.7	1.7	0.3	0.2	0.3	0.5	0.8	1.9	5.4	15.7	45.6	98.9
2003.....	4.7	1.6	0.4	0.2	0.3	0.5	0.9	1.8	5.1	16.0	46.2	101.7
2002.....	4.6	1.5	0.4	0.2	0.2	0.4	0.9	1.9	5.3	15.8	45.9	101.1
2001.....	4.8	1.3	0.4	0.2	0.3	0.4	0.9	2.0	5.5	17.3	48.2	107.6
2000.....	4.8	1.6	0.4	0.2	0.3	0.4	0.8	2.1	5.6	17.0	47.5	108.9
1999.....	4.8	1.5	0.4	0.2	0.3	0.5	0.9	2.0	5.8	18.4	46.1	109.2

All races, female

2004.....	4.6	1.8	0.3	0.2	0.2	0.4	0.7	1.5	3.7	10.2	29.2	68.1
2003.....	4.7	1.5	0.3	0.2	0.2	0.4	0.6	1.5	3.8	10.2	29.9	71.1
2002.....	4.6	2.0	0.4	0.2	0.2	0.4	0.8	1.7	3.7	10.4	29.6	68.9
2001.....	4.8	1.3	0.4	0.3	0.3	0.4	0.8	1.6	4.4	11.3	29.7	72.3
2000.....	4.9	1.5	0.3	0.2	0.2	0.4	0.8	1.6	4.5	11.2	31.1	75.2
1999.....	4.7	1.5	0.4	0.3	0.2	0.4	0.6	1.6	3.9	11.5	30.4	72.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

White, both sexes												
2004.....	5.1	1.9	0.3	0.2	0.3	0.4	0.7	1.6	4.4	12.9	36.7	80.2
2003.....	5.2	1.5	0.3	0.2	0.2	0.5	0.7	1.6	4.4	13.1	37.7	83.0
2002.....	5.1	1.9	0.4	0.2	0.2	0.4	0.8	1.7	4.5	13.2	37.0	80.7
2001.....	5.3	1.2	0.4	0.2	0.2	0.4	0.8	1.9	5.0	14.3	38.0	83.9
2000.....	5.3	1.6	0.4	0.2	0.2	0.4	0.8	1.8	5.1	13.9	38.6	87.3
1999.....	5.2	1.4	0.4	0.3	0.2	0.4	0.7	1.7	4.7	14.8	37.4	85.3
White, male												
2004.....	5.1	1.8	0.3	0.2	0.3	0.5	0.7	1.8	5.2	16.1	46.8	103.2
2003.....	5.2	1.6	0.3	0.2	0.3	0.5	0.9	1.8	5.0	16.4	47.8	105.5
2002.....	5.1	1.7	0.3	0.2	0.2	0.4	0.9	1.9	5.3	16.3	47.0	104.2
2001.....	5.3	1.2	0.4	0.2	0.2	0.4	0.9	2.1	5.6	17.8	49.4	109.9
2000.....	5.2	1.6	0.4	0.2	0.3	0.4	0.8	2.0	5.6	17.0	49.1	112.6
1999.....	5.2	1.4	0.4	0.2	0.2	0.4	0.9	2.0	5.8	18.7	47.2	114.3
White, female												
2004.....	5.0	1.9	0.3	0.2	0.2	0.4	0.6	1.4	3.6	10.2	29.9	69.9
2003.....	5.1	1.5	*	0.2	0.2	0.4	0.6	1.5	3.9	10.3	30.9	73.0
2002.....	5.1	2.1	0.4	0.2	0.2	0.3	0.7	1.6	3.8	10.5	30.3	70.5
2001.....	5.2	*	0.4	0.3	0.2	0.4	0.7	1.6	4.4	11.4	30.4	73.0
2000.....	5.4	1.6	*	0.2	0.2	0.4	0.8	1.5	4.6	11.3	31.8	77.1
1999.....	5.1	1.4	0.4	0.3	0.2	0.4	0.6	1.5	3.7	11.5	30.9	73.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year	2/	years	years	years	years	years	years	years	years	and over

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Black, both sexes												
2004.....	3.1	*	*	0.3	*	0.7	1.2	2.3	6.2	12.8	31.5	57.0
2003.....	2.9	*	*	*	*	0.5	1.2	2.5	5.2	13.0	29.5	57.8
2002.....	3.0	*	*	*	*	0.6	1.2	2.5	5.5	12.6	31.3	63.6
2001.....	3.1	*	*	0.4	0.3	*	1.1	2.0	5.9	14.3	31.5	73.4
2000.....	3.1	*	*	*	*	0.6	1.1	2.7	5.5	15.0	31.0	64.1
1999.....	3.2	*	*	*	0.4	0.5	1.2	2.6	6.2	15.7	33.0	61.9
Black, male												
2004.....	3.0	*	*	*	*	*	1.2	2.5	7.7	15.4	39.3	61.6
2003.....	2.9	*	*	*	*	*	1.1	2.5	6.7	16.2	36.7	66.7
2002.....	2.8	*	*	*	*	*	1.0	2.2	7.1	14.1	38.7	79.3
2001.....	3.0	*	*	*	*	*	1.1	1.8	6.5	17.2	41.5	88.5
2000.....	3.0	*	*	*	*	*	1.2	2.8	7.0	19.2	36.0	76.1
1999.....	3.0	*	*	*	*	0.8	1.1	2.5	7.1	19.0	38.4	63.3
Black, female												
2004.....	3.2	*	*	*	*	0.8	1.1	2.1	5.0	11.0	27.1	55.1
2003.....	3.0	*	*	*	*	*	1.2	2.4	4.0	10.8	25.5	54.3
2002.....	3.3	*	*	*	*	0.8	1.4	2.7	4.2	11.5	27.1	57.5
2001.....	3.2	*	*	*	*	*	1.0	2.2	5.3	12.3	25.9	67.8
2000.....	3.2	*	*	*	*	0.8	1.0	2.6	4.4	12.0	28.1	59.7
1999.....	3.4	*	*	*	*	*	1.2	2.7	5.6	13.4	29.9	61.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Anemias (D50-D64)												
All races 3/, both sexes												
2004.....	1.6	*	0.2	0.1	0.3	0.4	0.5	0.6	1.1	2.6	8.6	38.1
2003.....	1.6	0.5	0.2	0.1	0.3	0.4	0.4	0.6	1.0	2.6	8.8	40.7
2002.....	1.6	*	0.1	0.1	0.3	0.5	0.4	0.6	1.0	2.8	9.1	40.3
2001.....	1.6	*	0.1	0.1	0.3	0.4	0.5	0.6	1.0	2.7	9.8	42.0
2000.....	1.6	*	0.2	0.1	0.2	0.4	0.4	0.6	1.1	2.8	9.5	42.4
1999.....	1.6	*	0.2	0.1	0.3	0.4	0.4	0.6	1.0	2.6	9.5	43.8
All races, male												
2004.....	1.3	*	*	0.1	0.3	0.4	0.5	0.5	1.2	2.7	8.9	36.5
2003.....	1.3	*	*	0.1	0.2	0.4	0.4	0.6	1.2	3.0	9.3	38.0
2002.....	1.3	*	*	0.1	0.3	0.5	0.4	0.6	1.0	2.9	8.9	38.3
2001.....	1.3	*	*	0.1	0.3	0.5	0.4	0.6	0.9	2.9	9.9	40.0
2000.....	1.2	*	*	0.1	0.2	0.3	0.3	0.6	1.1	2.8	9.5	40.1
1999.....	1.3	*	*	0.1	0.3	0.4	0.5	0.6	0.9	2.7	10.1	43.4
All races, female												
2004.....	1.8	*	*	0.1	0.3	0.4	0.4	0.6	1.0	2.5	8.4	38.8
2003.....	1.9	*	*	0.1	0.3	0.4	0.4	0.6	0.8	2.3	8.5	41.9
2002.....	1.9	*	*	0.1	0.2	0.4	0.4	0.7	1.0	2.6	9.2	41.2
2001.....	2.0	*	*	0.1	0.3	0.3	0.5	0.6	1.0	2.5	9.6	42.9
2000.....	2.0	*	*	0.1	0.2	0.4	0.5	0.6	1.0	2.8	9.4	43.3
1999.....	1.9	*	*	0.2	0.3	0.4	0.4	0.6	1.0	2.6	9.2	44.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Anemias (D50-D64)												
White, both sexes												
2004.....	1.5	*	*	0.1	0.1	0.1	0.2	0.3	0.7	2.4	8.5	37.7
2003.....	1.5	*	*	0.1	0.1	0.1	0.1	0.3	0.6	2.4	8.6	40.6
2002.....	1.5	*	*	0.1	0.1	0.1	0.1	0.3	0.7	2.5	8.7	39.7
2001.....	1.5	*	*	0.1	0.1	0.1	0.1	0.3	0.8	2.5	9.4	41.7
2000.....	1.5	*	*	0.1	0.1	0.1	0.1	0.3	0.8	2.5	9.3	42.0
1999.....	1.5	*	*	0.1	0.1	*	0.1	0.3	0.6	2.4	9.4	43.0
White, male												
2004.....	1.1	*	*	*	*	*	0.2	0.3	0.8	2.5	8.6	36.7
2003.....	1.1	*	*	*	*	0.1	0.1	0.3	0.7	2.6	9.1	37.9
2002.....	1.1	*	*	*	*	0.1	0.1	0.3	0.8	2.6	8.5	37.2
2001.....	1.1	*	*	*	*	0.1	0.1	0.2	0.7	2.6	9.8	40.4
2000.....	1.1	*	*	*	*	*	0.1	0.3	0.9	2.7	9.2	39.1
1999.....	1.1	*	*	*	*	*	0.1	0.3	0.5	2.5	9.9	43.4
White, female												
2004.....	1.8	*	*	*	*	*	0.1	0.2	0.7	2.2	8.4	38.2
2003.....	1.8	*	*	*	*	*	0.1	0.2	0.6	2.2	8.2	41.8
2002.....	1.8	*	*	*	*	*	*	0.3	0.7	2.4	8.8	40.8
2001.....	1.9	*	*	*	*	*	0.2	0.3	0.8	2.4	9.1	42.3
2000.....	1.9	*	*	*	*	*	0.1	0.3	0.8	2.4	9.3	43.2
1999.....	1.9	*	*	*	0.1	*	0.1	0.3	0.7	2.3	9.0	42.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Anemias (D50-D64)												
Black, both sexes												
2004.....	2.7	*	*	0.3	1.4	2.3	2.4	2.7	4.0	4.8	11.8	48.2
2003.....	2.7	*	*	0.4	1.3	2.1	2.3	3.0	4.2	5.6	14.5	49.3
2002.....	2.8	*	*	0.4	1.3	2.4	2.5	3.1	3.6	5.9	16.0	52.3
2001.....	2.7	*	*	0.4	1.2	2.2	2.7	3.1	3.0	5.5	16.0	48.5
2000.....	2.6	*	*	0.5	1.1	2.3	2.4	3.0	3.4	5.4	13.3	51.0
1999.....	2.8	*	0.8	0.5	1.4	2.6	2.5	3.1	4.2	5.0	13.4	52.9
Black, male												
2004.....	2.6	*	*	*	1.5	2.7	2.7	2.5	4.9	5.0	13.9	42.7
2003.....	2.7	*	*	*	1.3	2.2	2.6	3.3	5.4	7.9	14.9	49.0
2002.....	2.7	*	*	*	1.6	2.8	2.5	2.6	3.9	7.3	16.9	56.8
2001.....	2.5	*	*	*	1.4	2.9	2.4	3.6	3.1	6.7	15.0	42.0
2000.....	2.3	*	*	*	1.0	2.1	2.0	3.3	3.4	4.5	13.9	55.4
1999.....	2.7	*	*	*	1.7	2.8	2.7	3.3	5.1	5.4	12.9	44.5
Black, female												
2004.....	2.8	*	*	*	1.3	2.0	2.2	2.9	3.2	4.6	10.5	50.4
2003.....	2.8	*	*	*	1.2	2.0	2.1	2.8	3.2	3.9	14.2	49.4
2002.....	3.0	*	*	*	0.9	2.0	2.5	3.6	3.3	5.0	15.4	50.5
2001.....	2.8	*	*	*	1.1	1.6	2.9	2.6	2.8	4.6	16.5	50.9
2000.....	3.0	*	*	*	1.1	2.4	2.8	2.8	3.4	6.0	13.0	49.4
1999.....	3.0	*	*	*	1.2	2.3	2.3	3.0	3.6	4.7	13.6	56.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diabetes mellitus (E10-E14)												
All races 3/, both sexes												
2004.....	24.9	*	*	0.1	0.4	1.5	4.6	13.4	37.1	87.2	176.9	307.0
2003.....	25.5	*	*	0.1	0.4	1.6	4.6	13.9	38.5	90.8	181.1	317.5
2002.....	25.4	*	*	0.1	0.4	1.6	4.8	13.7	37.7	91.4	182.8	320.6
2001.....	25.1	*	*	0.1	0.4	1.5	4.3	13.6	37.8	91.4	181.4	321.8
2000.....	24.6	*	*	0.1	0.4	1.6	4.3	13.1	37.8	90.7	179.5	319.7
1999.....	24.5	*	*	0.1	0.4	1.4	4.3	12.9	38.3	91.8	178.0	317.2
All races, male												
2004.....	24.4	*	*	*	0.4	1.7	5.7	16.5	44.0	100.0	202.5	333.9
2003.....	24.8	*	*	*	0.5	1.9	5.7	16.7	45.0	104.0	207.2	340.7
2002.....	24.2	*	*	0.1	0.5	1.9	5.9	16.4	44.5	101.4	202.6	348.5
2001.....	23.5	*	*	0.1	0.4	1.7	5.3	16.4	43.1	101.7	198.9	343.1
2000.....	22.9	*	*	*	0.5	1.9	5.1	15.3	42.4	99.1	199.6	345.1
1999.....	22.8	*	*	*	0.4	1.6	5.1	15.6	42.6	99.8	198.2	345.7
All races, female												
2004.....	25.4	*	*	0.1	0.4	1.3	3.5	10.4	30.7	76.4	159.6	294.9
2003.....	26.2	*	*	*	0.3	1.4	3.5	11.2	32.4	79.8	163.6	307.2
2002.....	26.5	*	*	*	0.4	1.3	3.7	11.1	31.4	83.1	169.7	308.4
2001.....	26.6	*	*	*	0.3	1.3	3.4	11.0	32.9	82.8	169.9	312.9
2000.....	26.3	*	*	*	0.4	1.2	3.4	11.1	33.6	83.7	166.3	309.4
1999.....	26.2	*	*	*	0.4	1.3	3.5	10.4	34.3	85.2	164.9	305.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diabetes mellitus (E10-E14)												
White, both sexes												
2004.....	24.4	*	*	*	0.3	1.3	4.0	11.6	32.4	78.1	164.2	289.4
2003.....	25.0	*	*	0.1	0.3	1.4	4.0	11.8	33.5	81.4	168.4	300.8
2002.....	24.9	*	*	0.1	0.3	1.4	4.3	11.7	32.7	81.5	170.4	303.7
2001.....	24.6	*	*	0.1	0.3	1.2	3.8	11.7	32.7	81.9	168.9	306.0
2000.....	24.1	*	*	*	0.3	1.3	3.8	11.1	33.1	81.2	166.0	303.5
1999.....	23.9	*	*	0.1	0.4	1.3	3.7	10.9	32.4	81.8	165.1	298.9
White, male												
2004.....	24.3	*	*	*	0.3	1.4	5.0	14.7	39.2	91.6	193.3	323.0
2003.....	24.8	*	*	*	0.4	1.5	5.1	14.7	40.2	95.8	198.6	333.4
2002.....	24.2	*	*	*	0.4	1.6	5.3	14.3	39.6	93.3	194.7	344.3
2001.....	23.5	*	*	*	0.3	1.3	4.7	14.3	38.4	94.6	189.2	337.3
2000.....	22.9	*	*	*	0.4	1.6	4.7	13.3	38.0	91.8	190.5	337.4
1999.....	22.7	*	*	*	0.4	1.4	4.5	13.5	37.3	92.3	189.9	333.4
White, female												
2004.....	24.5	*	*	*	0.3	1.1	2.9	8.6	25.9	66.5	144.4	274.3
2003.....	25.2	*	*	*	0.3	1.2	2.9	9.0	27.2	69.2	148.0	286.3
2002.....	25.6	*	*	*	0.3	1.1	3.2	9.2	26.2	71.5	154.0	286.1
2001.....	25.7	*	*	*	0.2	1.1	2.9	9.2	27.4	71.2	155.5	293.0
2000.....	25.3	*	*	*	0.3	1.0	3.0	9.0	28.5	72.4	149.8	289.7
1999.....	25.0	*	*	*	0.3	1.2	3.0	8.4	27.8	73.1	148.9	285.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diabetes mellitus (E10-E14)												
Black, both sexes												
2004.....	33.2	*	*	*	0.9	3.3	9.5	28.1	80.5	179.3	332.8	532.4
2003.....	33.8	*	*	*	0.7	3.8	9.5	30.1	85.7	184.9	336.1	527.5
2002.....	33.6	*	*	*	1.0	3.6	9.6	30.1	84.2	188.1	336.9	537.8
2001.....	33.1	*	*	*	1.0	3.7	8.8	29.6	85.0	188.4	335.2	529.7
2000.....	32.8	*	*	*	0.9	3.5	8.0	30.2	82.5	188.4	350.6	524.9
1999.....	33.0	*	*	*	0.8	3.0	8.9	30.3	90.2	190.8	333.5	529.0
Black, male												
2004.....	30.2	*	*	*	1.0	3.9	11.8	33.1	91.6	198.4	340.8	505.7
2003.....	29.8	*	*	*	0.8	4.7	11.5	33.7	96.0	197.2	337.3	453.5
2002.....	29.0	*	*	*	1.2	4.4	11.7	34.8	94.7	190.1	329.9	417.0
2001.....	28.5	*	*	*	1.0	4.8	10.2	34.5	92.2	187.7	341.9	450.5
2000.....	27.4	*	*	*	1.0	3.8	9.4	33.4	88.2	184.9	337.0	452.1
1999.....	27.7	*	*	*	*	3.3	10.3	34.7	93.8	186.5	314.8	506.2
Black, female												
2004.....	36.0	*	*	*	0.9	2.7	7.6	23.8	71.6	165.7	328.2	543.0
2003.....	37.4	*	*	*	*	3.0	7.8	27.0	77.4	176.1	335.4	556.2
2002.....	37.8	*	*	*	0.7	2.9	7.8	26.0	75.8	186.7	340.9	584.1
2001.....	37.2	*	*	*	1.0	2.8	7.6	25.4	79.1	188.9	331.5	559.1
2000.....	37.8	*	*	*	0.9	3.1	6.8	27.5	77.8	190.9	358.3	552.0
1999.....	37.8	*	*	*	0.9	2.6	7.6	26.6	87.4	193.9	343.9	537.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Nutritional deficiencies (E40-E64)

All races 3/, both sexes

2004.....	1.0	*	*	*	*	*	0.1	0.3	0.6	1.6	6.9	30.3
2003.....	1.1	*	*	*	*	*	0.1	0.2	0.6	1.9	7.7	35.5
2002.....	1.3	*	*	*	*	*	0.1	0.3	0.6	2.1	8.9	41.5
2001.....	1.3	*	*	*	*	*	0.1	0.3	0.6	2.0	8.7	42.9
2000.....	1.5	*	*	*	*	*	0.1	0.3	0.7	2.5	10.4	50.0
1999.....	1.5	*	*	*	*	*	0.1	0.3	0.7	2.3	10.2	53.9

All races, male

2004.....	0.8	*	*	*	*	*	*	0.2	0.6	1.7	6.8	27.4
2003.....	0.8	*	*	*	*	*	*	0.2	0.6	2.0	7.8	30.6
2002.....	0.9	*	*	*	*	*	0.1	0.2	0.7	2.2	8.5	37.0
2001.....	0.9	*	*	*	*	*	0.1	0.3	0.7	2.1	8.5	35.2
2000.....	1.0	*	*	*	*	*	*	0.2	0.8	2.7	9.7	43.7
1999.....	1.1	*	*	*	*	*	0.1	0.3	0.9	2.7	10.3	48.4

All races, female

2004.....	1.3	*	*	*	*	*	0.1	0.3	0.5	1.5	7.0	31.7
2003.....	1.5	*	*	*	*	*	*	0.3	0.6	1.7	7.7	37.6
2002.....	1.7	*	*	*	*	*	0.1	0.3	0.5	2.0	9.1	43.5
2001.....	1.7	*	*	*	*	*	0.1	0.3	0.5	2.0	8.9	46.1
2000.....	2.0	*	*	*	*	*	0.2	0.3	0.6	2.4	10.9	52.6
1999.....	2.0	*	*	*	*	*	0.2	0.3	0.6	2.1	10.2	56.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4/2/ years	5-14/ years	15-24/ years	25-34/ years	35-44/ years	45-54/ years	55-64/ years	65-74/ years	75-84/ years	85 years and over
Nutritional deficiencies (E40-E64)												
White, both sexes												
2004.....	1.1	*	*	*	*	*	0.1	0.2	0.5	1.5	6.6	29.2
2003.....	1.2	*	*	*	*	*	0.1	0.2	0.5	1.7	7.4	34.2
2002.....	1.4	*	*	*	*	*	0.1	0.2	0.5	1.9	8.6	40.2
2001.....	1.4	*	*	*	*	*	0.1	0.2	0.5	1.9	8.2	42.2
2000.....	1.6	*	*	*	*	*	0.1	0.2	0.6	2.3	10.0	48.8
1999.....	1.6	*	*	*	*	*	0.1	0.2	0.6	2.0	9.5	52.8
White, male												
2004.....	0.7	*	*	*	*	*	*	0.2	0.5	1.5	6.4	26.3
2003.....	0.8	*	*	*	*	*	*	0.2	0.5	1.8	7.1	29.6
2002.....	1.0	*	*	*	*	*	0.1	0.2	0.5	2.1	8.1	36.3
2001.....	0.9	*	*	*	*	*	*	0.2	0.5	1.8	8.0	33.8
2000.....	1.0	*	*	*	*	*	*	0.2	0.6	2.3	9.1	42.8
1999.....	1.1	*	*	*	*	*	0.1	0.3	0.6	2.1	9.3	46.5
White, female												
2004.....	1.4	*	*	*	*	*	*	0.3	0.5	1.4	6.8	30.4
2003.....	1.6	*	*	*	*	*	*	0.2	0.6	1.7	7.5	36.3
2002.....	1.8	*	*	*	*	*	*	0.3	0.5	1.8	8.9	41.9
2001.....	1.8	*	*	*	*	*	*	0.3	0.6	2.0	8.3	45.7
2000.....	2.1	*	*	*	*	*	0.2	0.3	0.6	2.3	10.5	51.3
1999.....	2.1	*	*	*	*	*	0.2	0.2	0.5	1.9	9.6	55.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nutritional deficiencies (E40-E64)												
Black, both sexes												
2004.....	1.1	*	*	*	*	*	*	0.6	1.2	3.3	10.6	46.8
2003.....	1.2	*	*	*	*	*	*	0.7	1.4	3.2	13.2	55.1
2002.....	1.3	*	*	*	*	*	*	0.4	1.7	4.2	14.5	60.6
2001.....	1.3	*	*	*	*	*	*	0.7	1.2	3.7	15.9	54.7
2000.....	1.5	*	*	*	*	*	*	0.6	1.7	4.7	16.1	66.9
1999.....	1.6	*	*	*	*	*	*	0.7	2.0	5.5	19.4	70.8
Black, male												
2004.....	0.9	*	*	*	*	*	*	*	1.6	4.4	12.2	45.7
2003.....	1.0	*	*	*	*	*	*	*	1.9	4.4	17.8	48.0
2002.....	1.0	*	*	*	*	*	*	*	2.8	4.4	16.4	47.2
2001.....	1.2	*	*	*	*	*	*	1.2	1.9	5.5	16.5	57.9
2000.....	1.2	*	*	*	*	*	*	*	2.8	6.4	17.9	63.4
1999.....	1.5	*	*	*	*	*	*	*	2.8	8.1	22.8	75.0
Black, female												
2004.....	1.2	*	*	*	*	*	*	*	*	2.6	9.6	47.3
2003.....	1.4	*	*	*	*	*	*	*	*	2.3	10.7	57.9
2002.....	1.6	*	*	*	*	*	*	*	*	4.1	13.5	65.7
2001.....	1.4	*	*	*	*	*	*	*	*	2.4	15.5	53.5
2000.....	1.7	*	*	*	*	*	*	*	*	3.5	15.2	68.3
1999.....	1.8	*	*	*	*	*	*	*	*	3.6	17.6	69.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malnutrition (E40-E46)												
All races 3/, both sexes												
2004.....	1.0	*	*	*	*	*	0.1	0.3	0.5	1.6	6.7	28.3
2003.....	1.1	*	*	*	*	*	0.1	0.2	0.6	1.7	7.4	33.4
2002.....	1.2	*	*	*	*	*	0.1	0.2	0.6	1.9	8.4	38.4
2001.....	1.2	*	*	*	*	*	0.1	0.3	0.6	1.9	8.2	40.2
2000.....	1.4	*	*	*	*	*	0.1	0.2	0.7	2.3	9.7	46.3
1999.....	1.4	*	*	*	*	*	0.1	0.3	0.6	2.2	9.5	50.0
All races, male												
2004.....	0.7	*	*	*	*	*	*	0.2	0.6	1.7	6.6	25.3
2003.....	0.8	*	*	*	*	*	*	0.2	0.6	1.9	7.4	28.2
2002.....	0.9	*	*	*	*	*	0.1	0.2	0.7	2.1	8.0	35.1
2001.....	0.8	*	*	*	*	*	*	0.2	0.6	1.9	8.0	33.6
2000.....	0.9	*	*	*	*	*	*	0.2	0.7	2.5	8.9	40.7
1999.....	1.0	*	*	*	*	*	0.1	0.3	0.8	2.5	9.4	44.4
All races, female												
2004.....	1.2	*	*	*	*	*	0.1	0.3	0.5	1.5	6.7	29.6
2003.....	1.4	*	*	*	*	*	*	0.2	0.6	1.7	7.4	35.6
2002.....	1.6	*	*	*	*	*	0.1	0.2	0.5	1.8	8.6	39.8
2001.....	1.6	*	*	*	*	*	0.1	0.3	0.5	1.9	8.2	43.0
2000.....	1.8	*	*	*	*	*	0.2	0.3	0.6	2.1	10.2	48.6
1999.....	1.8	*	*	*	*	*	0.1	0.3	0.5	2.0	9.5	52.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malnutrition (E40-E46)												
White, both sexes												
2004.....	1.0	*	*	*	*	*	0.1	0.2	0.5	1.4	6.3	26.9
2003.....	1.1	*	*	*	*	*	0.1	0.2	0.5	1.6	7.0	32.1
2002.....	1.3	*	*	*	*	*	0.1	0.2	0.5	1.8	8.0	37.0
2001.....	1.3	*	*	*	*	*	0.1	0.2	0.5	1.8	7.6	39.5
2000.....	1.5	*	*	*	*	*	0.1	0.2	0.6	2.1	9.3	45.2
1999.....	1.5	*	*	*	*	*	0.1	0.2	0.5	1.9	8.8	48.9
White, male												
2004.....	0.7	*	*	*	*	*	*	0.2	0.5	1.5	6.1	24.1
2003.....	0.8	*	*	*	*	*	*	0.2	0.4	1.7	6.7	27.2
2002.....	0.9	*	*	*	*	*	0.1	0.2	0.5	1.9	7.5	34.3
2001.....	0.8	*	*	*	*	*	*	0.1	0.5	1.7	7.4	32.2
2000.....	1.0	*	*	*	*	*	*	0.2	0.6	2.2	8.4	39.7
1999.....	1.0	*	*	*	*	*	*	0.2	0.6	2.0	8.5	42.2
White, female												
2004.....	1.3	*	*	*	*	*	*	0.3	0.5	1.4	6.5	28.2
2003.....	1.5	*	*	*	*	*	*	0.2	0.6	1.6	7.3	34.2
2002.....	1.6	*	*	*	*	*	*	0.3	0.5	1.7	8.4	38.2
2001.....	1.7	*	*	*	*	*	*	0.3	0.5	1.9	7.7	42.6
2000.....	1.9	*	*	*	*	*	0.1	0.2	0.6	2.0	9.9	47.4
1999.....	2.0	*	*	*	*	*	0.2	0.2	0.4	1.8	9.0	51.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malnutrition (E40-E46)												
Black, both sexes												
2004.....	1.0	*	*	*	*	*	*	0.6	1.1	3.2	10.4	46.0
2003.....	1.2	*	*	*	*	*	*	0.7	1.4	3.0	13.0	53.4
2002.....	1.3	*	*	*	*	*	*	*	1.7	3.6	14.0	57.6
2001.....	1.2	*	*	*	*	*	*	0.6	1.1	3.4	14.8	51.6
2000.....	1.4	*	*	*	*	*	*	0.5	1.5	4.7	14.9	63.8
1999.....	1.5	*	*	*	*	*	*	0.6	1.8	5.4	17.9	66.0
Black, male												
2004.....	0.9	*	*	*	*	*	*	*	1.5	4.3	12.0	43.7
2003.....	1.0	*	*	*	*	*	*	*	1.9	4.2	17.2	45.9
2002.....	1.0	*	*	*	*	*	*	*	2.6	3.9	15.2	46.1
2001.....	1.1	*	*	*	*	*	*	1.0	*	4.9	15.9	55.6
2000.....	1.2	*	*	*	*	*	*	*	2.4	6.3	17.3	61.1
1999.....	1.4	*	*	*	*	*	*	*	2.5	7.8	20.3	73.8
Black, female												
2004.....	1.2	*	*	*	*	*	*	*	*	2.5	9.4	46.9
2003.....	1.3	*	*	*	*	*	*	*	*	2.2	10.7	56.3
2002.....	1.5	*	*	*	*	*	*	*	*	3.4	13.3	62.0
2001.....	1.3	*	*	*	*	*	*	*	*	2.2	14.2	50.1
2000.....	1.6	*	*	*	*	*	*	*	*	3.5	13.6	64.8
1999.....	1.6	*	*	*	*	*	*	*	*	3.6	16.5	63.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other nutritional deficiencies (E50-E64)

All races 3/, both sexes

2004.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.1
2003.....	0.1	*	*	*	*	*	*	*	*	0.1	0.3	2.1
2002.....	0.1	*	*	*	*	*	*	*	*	0.2	0.5	3.1
2001.....	0.1	*	*	*	*	*	*	*	*	0.1	0.6	2.7
2000.....	0.1	*	*	*	*	*	*	*	*	0.2	0.7	3.7
1999.....	0.1	*	*	*	*	*	*	*	*	0.1	0.7	3.9

All races, male

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	2.1
2003.....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.4
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.6	1.9
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.5	1.6
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.8	3.0
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.9	3.9

All races, female

2004.....	0.1	*	*	*	*	*	*	*	*	*	*	2.1
2003.....	0.1	*	*	*	*	*	*	*	*	*	*	2.0
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.5	3.6
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.6	3.1
2000.....	0.1	*	*	*	*	*	*	*	*	0.2	0.7	4.0
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.6	3.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other nutritional deficiencies (E50-E64)

White, both sexes

2004.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
2003.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.5	3.1
2001.....	0.1	*	*	*	*	*	*	*	*	0.1	0.6	2.7
2000.....	0.1	*	*	*	*	*	*	*	*	0.2	0.7	3.7
1999.....	0.1	*	*	*	*	*	*	*	*	0.1	0.7	3.9

White, male

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	2.2
2003.....	0.1	*	*	*	*	*	*	*	*	*	*	2.4
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.5	2.0
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.5	*
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.7	3.2
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.8	4.3

White, female

2004.....	0.1	*	*	*	*	*	*	*	*	*	*	2.2
2003.....	0.1	*	*	*	*	*	*	*	*	*	*	2.1
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.5	3.6
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.6	3.1
2000.....	0.2	*	*	*	*	*	*	*	*	0.3	0.6	3.9
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.6	3.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Meningitis (G00,G03)

All races 3/, both sexes

2004.....	0.2	1.6	0.2	0.1	0.1	0.1	0.2	0.3	0.3	0.5	0.7	0.9
2003.....	0.3	1.9	0.2	*	0.1	0.1	0.2	0.3	0.4	0.5	0.7	1.1
2002.....	0.2	1.8	*	0.1	0.1	0.1	0.1	0.3	0.3	0.6	0.7	1.2
2001.....	0.3	1.9	0.2	0.1	0.1	0.1	0.2	0.3	0.4	0.5	0.8	0.9
2000.....	0.3	2.4	0.2	0.1	*	0.1	0.2	0.3	0.4	0.6	1.0	1.2
1999.....	0.3	3.1	0.3	0.1	0.1	0.1	0.2	0.3	0.4	0.6	1.0	1.5

All races, male

2004.....	0.2	1.5	*	*	*	0.1	0.2	0.4	0.3	0.4	0.7	*
2003.....	0.3	1.9	*	*	0.1	0.1	0.2	0.4	0.5	0.5	0.8	*
2002.....	0.3	2.0	*	*	*	*	0.2	0.4	0.4	0.7	0.8	*
2001.....	0.3	1.9	*	*	0.1	0.1	0.2	0.3	0.5	0.6	0.9	*
2000.....	0.3	2.5	0.3	*	*	0.1	0.2	0.3	0.5	0.7	1.3	*
1999.....	0.3	3.1	0.3	*	0.1	0.1	0.2	0.4	0.4	0.7	1.0	1.7

All races, female

2004.....	0.2	1.7	*	*	*	*	0.1	0.2	0.3	0.6	0.6	0.7
2003.....	0.2	2.0	*	*	*	*	0.1	0.2	0.3	0.5	0.6	1.0
2002.....	0.2	1.7	*	*	*	*	*	0.2	0.3	0.5	0.6	1.1
2001.....	0.2	2.0	*	*	*	*	0.1	0.2	0.4	0.4	0.6	0.8
2000.....	0.2	2.3	*	*	*	0.1	0.1	0.2	0.3	0.5	0.8	1.0
1999.....	0.3	3.0	*	*	*	0.1	0.1	0.2	0.4	0.5	1.0	1.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Meningitis (G00,G03)

White, both sexes

2004.....	0.2	1.4	*	*	*	*	0.1	0.2	0.3	0.4	0.6	0.8
2003.....	0.2	1.7	*	*	0.1	0.1	0.1	0.2	0.3	0.5	0.7	1.0
2002.....	0.2	1.7	*	0.1	*	0.1	0.1	0.2	0.3	0.6	0.6	1.2
2001.....	0.2	1.5	*	*	0.1	0.1	0.1	0.2	0.4	0.5	0.7	0.8
2000.....	0.2	1.8	0.2	0.1	*	0.1	0.1	0.2	0.3	0.5	0.9	1.0
1999.....	0.3	2.3	0.3	*	0.1	0.1	0.1	0.2	0.3	0.6	0.9	1.3

White, male

2004.....	0.2	1.5	*	*	*	*	0.2	0.3	0.3	0.3	0.7	*
2003.....	0.3	2.0	*	*	*	*	0.1	0.3	0.4	0.5	0.8	*
2002.....	0.2	1.8	*	*	*	*	0.1	0.3	0.3	0.7	0.8	*
2001.....	0.2	1.4	*	*	*	*	0.1	0.3	0.4	0.6	0.9	*
2000.....	0.3	2.0	*	*	*	*	0.1	0.3	0.4	0.7	1.2	*
1999.....	0.3	3.0	0.3	*	*	*	0.1	0.3	0.4	0.6	0.9	*

White, female

2004.....	0.2	1.4	*	*	*	*	*	0.1	0.3	0.5	0.5	*
2003.....	0.2	1.3	*	*	*	*	*	0.2	0.2	0.5	0.6	0.9
2002.....	0.2	1.5	*	*	*	*	*	0.2	0.2	0.4	0.5	1.2
2001.....	0.2	1.6	*	*	*	*	*	0.2	0.3	0.4	0.6	0.8
2000.....	0.2	1.6	*	*	*	*	*	0.2	0.2	0.4	0.8	1.0
1999.....	0.3	1.5	*	*	*	0.1	0.1	0.2	0.3	0.5	0.9	1.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Parkinson's disease (G20-G21)												
All races 3/, both sexes												
2004.....	6.1	*	*	*	*	*	*	0.2	1.2	12.0	67.5	135.8
2003.....	6.2	*	*	*	*	*	*	0.2	1.3	12.7	67.8	138.2
2002.....	5.9	*	*	*	*	*	*	0.1	1.2	12.2	63.9	135.2
2001.....	5.8	*	*	*	*	*	*	0.1	1.2	11.7	64.6	134.2
2000.....	5.6	*	*	*	*	*	*	0.1	1.1	11.5	61.9	131.9
1999.....	5.2	*	*	*	*	*	*	0.1	1.0	11.0	58.2	124.4
All races, male												
2004.....	7.2	*	*	*	*	*	*	0.2	1.6	17.3	101.7	220.0
2003.....	7.1	*	*	*	*	*	*	0.2	1.6	18.2	100.7	222.4
2002.....	6.8	*	*	*	*	*	*	0.2	1.5	17.9	94.7	220.1
2001.....	6.7	*	*	*	*	*	*	0.2	1.8	16.8	97.0	224.4
2000.....	6.4	*	*	*	*	*	*	0.2	1.5	16.1	92.8	222.3
1999.....	6.0	*	*	*	*	*	*	0.2	1.4	15.7	89.5	205.6
All races, female												
2004.....	5.1	*	*	*	*	*	*	*	0.8	7.5	44.4	97.9
2003.....	5.3	*	*	*	*	*	*	0.1	0.9	8.1	45.8	101.0
2002.....	5.0	*	*	*	*	*	*	*	0.9	7.4	43.5	98.4
2001.....	4.9	*	*	*	*	*	*	*	0.7	7.5	43.3	96.4
2000.....	4.8	*	*	*	*	*	*	0.1	0.8	7.6	41.7	95.0
1999.....	4.5	*	*	*	*	*	*	*	0.7	7.1	37.8	91.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Parkinson's disease (G20-G21)

White, both sexes

2004.....	7.2	*	*	*	*	*	*	0.2	1.3	13.0	71.9	143.1
2003.....	7.2	*	*	*	*	*	*	0.2	1.3	13.6	72.2	146.0
2002.....	6.9	*	*	*	*	*	*	0.2	1.3	13.0	68.2	143.3
2001.....	6.8	*	*	*	*	*	*	0.1	1.2	12.6	68.8	141.8
2000.....	6.5	*	*	*	*	*	*	0.1	1.2	12.3	65.6	139.5
1999.....	6.1	*	*	*	*	*	*	0.1	1.1	11.7	61.5	131.2

White, male

2004.....	8.3	*	*	*	*	*	*	0.2	1.7	18.6	107.9	231.3
2003.....	8.2	*	*	*	*	*	*	0.2	1.6	19.3	106.6	233.3
2002.....	7.9	*	*	*	*	*	*	0.2	1.6	19.0	100.5	232.7
2001.....	7.8	*	*	*	*	*	*	0.2	1.9	17.9	103.0	237.6
2000.....	7.4	*	*	*	*	*	*	0.2	1.6	17.2	97.5	235.3
1999.....	7.0	*	*	*	*	*	*	0.2	1.5	16.8	94.4	216.7

White, female

2004.....	6.0	*	*	*	*	*	*	*	0.9	8.2	47.4	103.3
2003.....	6.2	*	*	*	*	*	*	0.2	1.0	8.7	48.8	107.4
2002.....	5.9	*	*	*	*	*	*	*	0.9	7.9	46.5	104.4
2001.....	5.8	*	*	*	*	*	*	*	0.7	8.1	46.2	101.7
2000.....	5.6	*	*	*	*	*	*	*	0.9	8.2	44.7	100.6
1999.....	5.2	*	*	*	*	*	*	*	0.6	7.5	40.0	96.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Parkinson's disease (G20-G21)

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Black, both sexes												
2004.....	1.6	*	*	*	*	*	*	*	0.7	5.4	29.2	61.2
2003.....	1.6	*	*	*	*	*	*	*	1.2	7.0	29.0	52.8
2002.....	1.5	*	*	*	*	*	*	*	*	6.8	25.4	51.7
2001.....	1.5	*	*	*	*	*	*	*	0.8	5.5	25.2	59.0
2000.....	1.5	*	*	*	*	*	*	*	*	5.7	27.4	51.6
1999.....	1.4	*	*	*	*	*	*	*	0.9	5.9	24.4	52.0
Black, male												
2004.....	1.9	*	*	*	*	*	*	*	*	8.5	43.4	106.3
2003.....	1.9	*	*	*	*	*	*	*	1.7	10.3	45.4	97.0
2002.....	1.7	*	*	*	*	*	*	*	*	11.1	39.6	89.0
2001.....	1.7	*	*	*	*	*	*	*	*	8.5	37.4	99.8
2000.....	1.8	*	*	*	*	*	*	*	*	8.8	47.8	86.5
1999.....	1.6	*	*	*	*	*	*	*	*	8.7	37.8	86.7
Black, female												
2004.....	1.4	*	*	*	*	*	*	*	*	3.2	21.1	43.3
2003.....	1.4	*	*	*	*	*	*	*	*	4.7	19.8	35.6
2002.....	1.2	*	*	*	*	*	*	*	*	3.6	17.4	37.4
2001.....	1.3	*	*	*	*	*	*	*	*	3.3	18.3	43.8
2000.....	1.2	*	*	*	*	*	*	*	*	3.5	15.9	38.6
1999.....	1.2	*	*	*	*	*	*	*	*	3.9	16.9	39.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alzheimer's disease (G30)												
All races 3/, both sexes												
2004.....	22.5	*	*	*	*	*	*	0.2	1.9	19.7	168.7	818.8
2003.....	21.8	*	*	*	*	*	*	0.2	2.0	20.9	164.4	802.4
2002.....	20.4	*	*	*	*	*	*	0.1	1.9	19.7	158.1	752.3
2001.....	18.9	*	*	*	*	*	*	0.2	2.1	18.7	147.5	710.3
2000.....	17.6	*	*	*	*	*	*	0.2	2.0	18.7	139.6	667.7
1999.....	16.0	*	*	*	*	*	*	0.2	1.9	17.4	129.5	601.3
All races, male												
2004.....	13.1	*	*	*	*	*	*	0.2	1.8	18.5	149.0	619.6
2003.....	12.8	*	*	*	*	*	*	0.2	1.8	20.2	147.6	606.5
2002.....	12.0	*	*	*	*	*	*	0.1	1.8	18.6	142.2	572.0
2001.....	11.3	*	*	*	*	*	*	0.2	1.9	17.7	136.3	556.1
2000.....	10.5	*	*	*	*	*	*	0.2	1.8	18.7	127.6	522.1
1999.....	9.8	*	*	*	*	*	*	0.2	1.9	17.3	122.6	485.7
All races, female												
2004.....	31.5	*	*	*	*	*	*	0.3	1.9	20.8	181.9	908.4
2003.....	30.5	*	*	*	*	*	*	0.2	2.1	21.5	175.7	889.0
2002.....	28.5	*	*	*	*	*	*	0.1	2.0	20.6	168.7	830.5
2001.....	26.3	*	*	*	*	*	*	0.2	2.3	19.5	154.8	774.9
2000.....	24.5	*	*	*	*	*	*	0.2	2.2	18.6	147.4	727.0
1999.....	21.9	*	*	*	*	*	*	0.2	1.8	17.5	134.0	648.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alzheimer's disease (G30)												
White, both sexes												
2004.....	25.6	*	*	*	*	*	*	0.3	2.0	20.5	174.1	846.8
2003.....	25.0	*	*	*	*	*	*	0.2	2.2	21.7	170.7	834.0
2002.....	23.5	*	*	*	*	*	*	0.1	2.1	20.7	164.8	781.8
2001.....	21.7	*	*	*	*	*	*	0.2	2.3	19.6	152.6	738.1
2000.....	20.2	*	*	*	*	*	*	0.2	2.2	19.5	144.8	695.3
1999.....	18.3	*	*	*	*	*	*	0.2	2.0	18.1	134.5	627.9
White, male												
2004.....	14.9	*	*	*	*	*	*	0.2	1.9	19.0	153.7	640.9
2003.....	14.6	*	*	*	*	*	*	0.2	1.9	20.8	152.4	630.3
2002.....	13.7	*	*	*	*	*	*	0.1	1.9	19.3	147.3	594.8
2001.....	12.8	*	*	*	*	*	*	0.2	2.0	18.5	140.5	577.2
2000.....	11.9	*	*	*	*	*	*	0.2	1.9	19.5	132.5	542.1
1999.....	11.2	*	*	*	*	*	*	0.2	2.1	17.9	127.4	506.9
White, female												
2004.....	36.1	*	*	*	*	*	*	0.3	2.1	21.7	188.0	939.6
2003.....	35.2	*	*	*	*	*	*	0.2	2.4	22.5	183.1	924.3
2002.....	33.0	*	*	*	*	*	*	0.1	2.2	21.8	176.5	863.0
2001.....	30.3	*	*	*	*	*	*	0.2	2.6	20.6	160.6	805.4
2000.....	28.2	*	*	*	*	*	*	0.2	2.4	19.5	152.9	757.6
1999.....	25.3	*	*	*	*	*	*	0.2	2.0	18.3	139.1	676.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alzheimer's disease (G30)												
Black, both sexes												
2004.....	11.0	*	*	*	*	*	*	*	1.3	19.3	153.1	657.9
2003.....	9.7	*	*	*	*	*	*	*	1.0	19.2	138.1	583.8
2002.....	8.9	*	*	*	*	*	*	*	1.2	16.8	123.1	543.7
2001.....	8.4	*	*	*	*	*	*	*	1.1	15.2	124.7	506.7
2000.....	7.5	*	*	*	*	*	*	*	1.2	15.8	109.9	445.1
1999.....	6.5	*	*	*	*	*	*	*	1.2	15.1	98.2	372.4
Black, male												
2004.....	6.2	*	*	*	*	*	*	*	*	19.1	136.7	497.7
2003.....	6.0	*	*	*	*	*	*	*	*	20.3	136.2	459.8
2002.....	5.3	*	*	*	*	*	*	*	*	18.0	122.5	412.8
2001.....	5.1	*	*	*	*	*	*	*	*	16.1	121.5	414.1
2000.....	4.5	*	*	*	*	*	*	*	*	16.5	97.5	371.3
1999.....	4.0	*	*	*	*	*	*	*	*	16.5	89.8	317.5
Black, female												
2004.....	15.3	*	*	*	*	*	*	*	1.4	19.4	162.4	721.4
2003.....	13.2	*	*	*	*	*	*	*	*	18.4	139.2	631.9
2002.....	12.0	*	*	*	*	*	*	*	*	15.9	123.4	593.9
2001.....	11.3	*	*	*	*	*	*	*	*	14.6	126.4	541.0
2000.....	10.2	*	*	*	*	*	*	*	*	15.2	116.9	472.6
1999.....	8.7	*	*	*	*	*	*	*	*	14.2	102.9	392.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Major cardiovascular diseases (I00-I78)												
All races 3/, both sexes												
2004.....	293.3	13.8	1.5	0.8	3.2	10.0	36.6	110.9	268.2	697.7	2,035.6	6,616.6
2003.....	310.3	13.9	1.5	0.8	3.5	10.3	38.2	113.4	284.9	748.3	2,170.8	7,138.0
2002.....	318.6	15.9	1.4	0.9	3.2	10.0	37.8	114.3	294.8	788.4	2,257.0	7,389.6
2001.....	323.9	14.9	1.9	0.9	3.2	10.2	37.1	113.4	301.2	810.4	2,322.2	7,673.7
2000.....	332.9	16.7	1.5	0.9	3.3	9.5	36.8	115.7	319.2	849.3	2,396.2	8,030.3
1999.....	340.6	17.0	1.5	0.9	3.4	9.6	37.7	116.2	327.3	887.3	2,474.9	8,197.1
All races, male												
2004.....	281.2	14.7	1.5	0.8	4.0	12.8	49.0	156.7	371.5	903.9	2,462.3	6,818.1
2003.....	296.3	15.2	1.4	1.0	4.2	12.9	51.1	160.9	393.3	976.3	2,636.1	7,329.3
2002.....	303.3	16.6	1.5	0.9	4.1	12.7	50.8	162.3	407.0	1,027.8	2,731.1	7,638.8
2001.....	306.2	15.3	1.9	0.9	3.9	12.8	49.9	160.4	414.0	1,057.6	2,823.3	7,936.0
2000.....	315.8	17.5	1.6	1.0	3.9	11.8	49.5	165.1	441.6	1,114.1	2,926.6	8,448.1
1999.....	323.0	17.9	1.6	1.0	4.0	12.1	51.2	165.8	452.0	1,168.4	3,024.0	8,603.2
All races, female												
2004.....	304.9	12.9	1.5	0.8	2.3	7.1	24.2	66.6	172.2	524.5	1,748.4	6,526.0
2003.....	323.9	12.5	1.5	0.7	2.8	7.5	25.3	67.5	184.4	557.8	1,860.0	7,053.5
2002.....	333.3	15.1	1.3	0.8	2.2	7.2	24.9	68.0	191.0	589.1	1,942.2	7,281.5
2001.....	340.9	14.5	1.8	0.9	2.4	7.6	24.3	68.1	196.9	605.6	1,992.9	7,563.9
2000.....	349.5	15.9	1.4	0.8	2.8	7.2	24.3	68.0	206.3	631.2	2,050.4	7,860.1
1999.....	357.5	16.0	1.4	0.8	2.7	7.2	24.4	68.5	212.5	657.0	2,118.2	8,033.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Major cardiovascular diseases (I00-I78)												
White, both sexes												
2004.....	311.7	11.9	1.2	0.7	2.7	8.2	32.1	97.1	244.7	665.2	2,016.2	6,704.8
2003.....	330.0	11.0	1.2	0.7	2.9	8.5	32.9	99.0	259.8	712.2	2,151.4	7,233.1
2002.....	339.4	13.4	1.3	0.8	2.7	8.1	33.0	100.6	269.9	755.1	2,234.8	7,482.7
2001.....	344.8	12.2	1.5	0.8	2.7	8.5	31.7	99.5	275.3	774.1	2,298.9	7,770.6
2000.....	354.7	14.6	1.3	0.8	2.8	7.9	31.6	102.0	292.7	814.2	2,370.8	8,133.9
1999.....	362.7	14.6	1.3	0.8	2.9	8.1	32.1	102.7	300.0	850.9	2,449.9	8,299.2
White, male												
2004.....	296.7	12.7	1.3	0.7	3.5	10.7	44.4	141.3	343.3	870.1	2,452.6	6,940.8
2003.....	312.6	11.5	1.1	0.9	3.5	10.9	45.6	144.6	364.8	938.3	2,623.0	7,466.8
2002.....	320.9	14.3	1.5	0.8	3.5	10.5	45.7	147.1	378.3	994.8	2,721.2	7,766.6
2001.....	323.5	13.2	1.6	0.8	3.3	10.9	44.2	145.1	385.3	1,020.8	2,809.7	8,060.7
2000.....	334.2	15.3	1.6	0.9	3.3	10.1	44.1	150.2	411.8	1,080.7	2,912.4	8,585.0
1999.....	341.2	15.3	1.5	0.9	3.4	10.4	45.1	150.5	420.7	1,133.1	3,012.8	8,717.5
White, female												
2004.....	326.5	11.0	1.1	0.7	1.9	5.5	19.5	53.6	151.4	489.8	1,718.6	6,598.4
2003.....	347.0	10.4	1.4	0.6	2.3	5.9	20.0	53.9	160.6	519.6	1,832.5	7,129.6
2002.....	357.5	12.5	1.1	0.7	1.8	5.6	20.0	54.7	167.7	551.7	1,908.4	7,359.3
2001.....	365.5	11.1	1.5	0.8	2.0	6.0	19.0	54.6	171.9	565.8	1,959.8	7,649.2
2000.....	374.8	13.9	1.1	0.6	2.2	5.5	19.0	54.6	180.9	590.8	2,014.4	7,950.6
1999.....	383.6	13.8	1.2	0.7	2.4	5.7	19.0	55.7	186.8	615.7	2,081.0	8,131.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Major cardiovascular diseases (I00-I78)												
Black, both sexes												
2004.....	260.3	21.9	2.9	1.5	5.9	22.7	74.3	231.0	525.5	1,122.3	2,572.7	6,368.6
2003.....	273.1	26.2	2.8	1.4	6.9	22.9	78.8	238.8	556.2	1,207.1	2,713.5	6,862.0
2002.....	276.1	27.6	2.0	1.4	5.9	23.3	76.0	237.4	568.3	1,226.2	2,824.0	7,081.7
2001.....	280.4	28.6	3.5	1.3	6.6	22.6	78.3	238.2	578.0	1,276.2	2,891.8	7,316.7
2000.....	285.0	27.8	2.5	1.4	6.8	21.5	77.4	241.3	600.3	1,316.2	2,968.8	7,518.9
1999.....	289.4	29.2	2.4	1.4	6.8	20.7	81.6	241.9	614.6	1,368.1	3,024.7	7,582.5
Black, male												
2004.....	257.2	23.7	2.8	1.4	7.6	27.7	92.6	305.7	717.0	1,432.4	3,033.4	6,198.0
2003.....	268.6	31.2	3.3	1.5	8.5	28.0	98.1	317.1	742.4	1,542.1	3,265.7	6,547.7
2002.....	268.5	28.2	1.9	1.5	7.2	28.9	94.7	313.1	762.5	1,556.1	3,287.6	6,929.0
2001.....	272.4	28.1	3.6	1.2	8.0	27.9	97.4	312.5	761.1	1,620.3	3,435.3	7,363.9
2000.....	275.6	29.1	2.1	1.4	7.5	25.4	95.8	317.5	796.8	1,639.9	3,501.8	7,687.4
1999.....	282.5	30.6	2.5	1.5	8.4	25.4	102.7	323.2	822.6	1,723.6	3,526.5	7,921.1
Black, female												
2004.....	263.2	20.1	3.0	1.5	4.1	18.1	58.0	166.9	371.5	901.1	2,312.3	6,436.3
2003.....	277.2	21.0	2.4	1.3	5.3	18.1	61.7	171.5	406.5	968.7	2,401.2	6,984.1
2002.....	283.0	26.9	2.1	1.3	4.5	18.1	59.5	172.5	412.2	992.0	2,561.6	7,140.2
2001.....	287.8	29.0	3.4	1.4	5.1	17.7	61.4	174.3	430.5	1,033.4	2,586.6	7,299.2
2000.....	293.4	26.5	2.9	1.4	6.1	17.9	61.1	175.6	441.9	1,088.3	2,668.7	7,456.2
1999.....	295.7	27.9	2.4	1.2	5.1	16.4	63.0	172.3	447.7	1,117.3	2,743.8	7,456.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13,I20-I51)												
All races 3/, both sexes												
2004.....	222.2	10.3	1.2	0.6	2.5	7.9	29.3	90.2	218.8	541.6	1,506.3	4,895.9
2003.....	235.6	11.0	1.2	0.6	2.7	8.2	30.7	92.5	233.2	585.0	1,611.1	5,278.4
2002.....	241.7	12.4	1.1	0.6	2.5	7.9	30.5	93.7	241.5	615.9	1,677.2	5,446.8
2001.....	245.8	11.9	1.5	0.7	2.5	8.0	29.6	92.9	246.9	635.1	1,725.7	5,664.2
2000.....	252.6	13.0	1.2	0.7	2.6	7.4	29.2	94.2	261.2	665.6	1,780.3	5,926.1
1999.....	259.9	13.8	1.2	0.7	2.8	7.6	30.2	95.7	269.9	701.7	1,849.9	6,063.0
All races, male												
2004.....	222.8	10.9	1.1	0.6	3.2	10.5	40.9	132.3	312.8	723.8	1,893.6	5,239.3
2003.....	235.0	12.1	1.1	0.7	3.4	10.5	42.8	136.2	331.7	785.3	2,030.3	5,621.5
2002.....	240.7	12.9	1.1	0.7	3.3	10.5	43.1	138.4	343.4	827.1	2,110.1	5,823.5
2001.....	242.5	11.8	1.5	0.7	3.2	10.3	41.7	136.6	349.8	851.3	2,177.3	6,040.5
2000.....	249.8	13.3	1.4	0.8	3.2	9.6	41.4	140.2	371.7	898.3	2,248.1	6,430.0
1999.....	257.0	13.9	1.3	0.8	3.4	9.8	43.0	142.0	383.0	950.6	2,336.4	6,561.6
All races, female												
2004.....	221.6	9.7	1.2	0.6	1.7	5.2	17.7	49.6	131.5	388.6	1,245.6	4,741.5
2003.....	236.2	9.8	1.3	0.5	2.1	5.7	18.6	50.2	141.9	417.5	1,331.1	5,126.7
2002.....	242.7	11.8	1.0	0.6	1.7	5.2	18.0	50.6	147.2	440.1	1,389.7	5,283.3
2001.....	249.0	12.0	1.4	0.7	1.8	5.6	17.6	50.7	151.8	455.9	1,428.9	5,506.8
2000.....	255.3	12.5	1.0	0.5	2.1	5.2	17.2	49.8	159.3	474.0	1,475.1	5,720.9
1999.....	262.6	13.7	1.1	0.6	2.1	5.4	17.5	51.0	165.6	497.9	1,533.9	5,862.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over
Diseases of heart (I00-I09,I11,I13,I20-I51)												
White, both sexes												
2004.....	237.4	9.0	0.9	0.5	2.1	6.5	26.3	81.2	203.4	521.2	1,500.0	4,977.6
2003.....	251.7	8.8	1.0	0.5	2.3	6.7	27.2	82.9	216.5	561.8	1,603.6	5,363.5
2002.....	258.5	10.6	0.9	0.6	2.1	6.5	27.3	84.8	224.5	595.1	1,666.9	5,529.1
2001.....	262.8	9.9	1.2	0.6	2.0	6.6	25.9	83.9	229.6	612.2	1,714.3	5,752.1
2000.....	270.2	11.5	1.1	0.6	2.1	6.1	25.8	85.4	243.2	642.5	1,767.8	6,020.8
1999.....	277.7	11.8	1.0	0.6	2.3	6.4	26.3	86.8	251.1	677.7	1,837.5	6,152.7
White, male												
2004.....	236.5	9.6	1.0	0.5	2.7	8.8	37.8	122.2	294.4	703.2	1,897.1	5,348.4
2003.....	249.5	9.4	0.8	0.6	2.8	9.0	39.1	125.3	313.2	761.1	2,030.1	5,747.2
2002.....	256.0	11.3	1.0	0.6	2.8	8.8	39.7	128.6	324.0	807.8	2,112.0	5,939.8
2001.....	257.8	10.6	1.2	0.6	2.6	8.7	37.8	127.0	330.8	829.1	2,175.8	6,157.2
2000.....	265.8	11.9	1.3	0.7	2.6	8.1	37.8	130.7	351.8	877.8	2,247.0	6,560.8
1999.....	272.9	12.0	1.2	0.7	2.7	8.4	38.6	132.1	361.4	928.4	2,338.8	6,669.0
White, female												
2004.....	238.3	8.4	0.9	0.5	1.4	4.0	14.5	40.7	117.2	365.4	1,229.1	4,810.4
2003.....	253.8	8.1	1.1	0.4	1.7	4.4	15.1	41.1	125.2	392.0	1,315.2	5,193.6
2002.....	261.0	9.9	0.8	0.5	1.4	4.0	14.8	41.7	130.6	414.7	1,368.2	5,350.6
2001.....	267.7	9.2	1.1	0.6	1.5	4.4	13.9	41.5	134.3	429.0	1,407.9	5,582.5
2000.....	274.5	11.1	0.8	0.4	1.6	4.0	13.7	40.9	141.3	445.2	1,452.4	5,801.4
1999.....	282.4	11.6	0.9	0.5	1.8	4.3	13.8	42.2	147.8	468.7	1,509.0	5,945.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13,I20-I51)												
Black, both sexes												
2004.....	192.3	15.3	2.4	1.2	4.9	17.7	56.5	176.1	402.7	839.8	1,844.1	4,563.8
2003.....	202.8	20.2	2.2	1.1	5.6	18.1	59.5	181.4	427.8	913.0	1,962.1	4,941.2
2002.....	205.6	21.3	1.6	1.0	4.9	18.4	57.8	180.9	442.9	924.5	2,048.8	5,115.2
2001.....	208.8	21.6	3.0	1.1	5.4	17.9	59.5	179.9	447.6	965.7	2,108.4	5,275.7
2000.....	211.8	19.8	2.0	1.1	5.6	17.2	57.4	181.5	466.0	1,002.6	2,159.7	5,389.5
1999.....	217.2	23.8	2.0	1.1	5.8	16.7	62.1	185.4	481.3	1,048.6	2,211.1	5,487.4
Black, male												
2004.....	196.7	16.3	2.4	1.1	6.4	22.5	72.8	240.0	560.2	1,096.6	2,235.5	4,637.3
2003.....	206.0	24.1	2.5	1.2	7.3	22.2	77.1	248.1	580.9	1,195.5	2,426.6	4,850.3
2002.....	206.3	22.1	*	1.0	6.2	23.9	75.6	246.0	605.3	1,192.7	2,449.6	5,125.7
2001.....	209.0	19.5	3.2	1.0	6.9	22.8	76.7	242.6	602.2	1,245.8	2,569.3	5,459.9
2000.....	211.1	19.8	1.9	1.2	6.6	21.2	74.3	247.2	631.2	1,268.8	2,597.6	5,633.5
1999.....	218.5	23.4	2.1	1.3	7.4	21.0	81.9	255.0	660.0	1,346.4	2,623.5	5,858.8
Black, female												
2004.....	188.3	14.2	2.3	1.3	3.4	13.3	42.0	121.2	276.0	656.5	1,622.9	4,534.7
2003.....	200.0	16.1	2.0	1.0	3.9	14.4	43.9	124.1	304.7	712.0	1,699.6	4,976.5
2002.....	205.0	20.6	1.8	1.0	3.7	13.5	42.1	124.9	312.3	734.0	1,821.9	5,111.2
2001.....	208.6	23.9	2.7	1.1	3.8	13.5	44.2	125.9	323.1	768.0	1,849.6	5,207.3
2000.....	212.6	19.8	2.0	1.0	4.7	13.6	42.4	125.0	332.8	815.2	1,913.1	5,298.7
1999.....	216.0	24.2	1.9	1.0	4.3	12.8	44.6	125.8	337.9	838.6	1,980.3	5,348.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
All races 3/, both sexes												
2004.....	1.1	*	*	*	*	0.1	0.2	0.5	1.1	3.3	8.8	17.1
2003.....	1.2	*	*	*	*	0.1	0.2	0.5	1.3	3.8	9.6	20.5
2002.....	1.2	*	*	*	*	0.1	0.2	0.6	1.6	3.8	9.6	17.8
2001.....	1.2	*	*	*	*	0.1	0.3	0.6	1.4	4.1	9.4	17.7
2000.....	1.3	*	*	*	*	0.1	0.3	0.7	1.7	4.5	9.5	16.8
1999.....	1.3	*	*	*	*	0.1	0.3	0.7	1.7	4.6	10.0	18.0
All races, male												
2004.....	0.7	*	*	*	*	*	0.2	0.4	0.8	2.4	7.3	11.5
2003.....	0.8	*	*	*	*	*	0.2	0.4	1.1	2.9	7.5	16.2
2002.....	0.8	*	*	*	*	*	0.2	0.5	1.2	3.0	6.7	11.8
2001.....	0.7	*	*	*	*	*	0.2	0.5	1.1	2.8	6.6	12.8
2000.....	0.8	*	*	*	*	*	0.3	0.6	1.4	2.9	6.6	12.2
1999.....	0.8	*	*	*	*	*	0.2	0.5	1.2	3.5	6.8	10.3
All races, female												
2004.....	1.5	*	*	*	*	*	0.3	0.6	1.4	4.1	9.9	19.5
2003.....	1.7	*	*	*	*	*	0.2	0.7	1.5	4.5	11.0	22.3
2002.....	1.7	*	*	*	*	*	0.2	0.7	2.0	4.6	11.6	20.4
2001.....	1.7	*	*	*	*	*	0.3	0.7	1.7	5.1	11.3	19.8
2000.....	1.8	*	*	*	*	0.1	0.4	0.8	2.0	5.8	11.4	18.7
1999.....	1.9	*	*	*	*	0.1	0.3	0.8	2.2	5.6	12.2	21.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
White, both sexes												
2004.....	1.2	*	*	*	*	*	0.2	0.4	1.0	3.4	9.3	18.2
2003.....	1.4	*	*	*	*	*	0.2	0.4	1.2	3.8	10.2	21.8
2002.....	1.4	*	*	*	*	0.1	0.2	0.5	1.5	4.0	10.1	18.9
2001.....	1.3	*	*	*	*	*	0.2	0.5	1.4	4.1	10.0	18.7
2000.....	1.4	*	*	*	*	0.1	0.3	0.6	1.7	4.6	10.1	17.6
1999.....	1.4	*	*	*	*	0.1	0.2	0.6	1.7	4.8	10.5	18.9
White, male												
2004.....	0.8	*	*	*	*	*	0.1	0.4	0.7	2.5	7.8	12.4
2003.....	0.9	*	*	*	*	*	0.1	0.3	0.9	3.0	8.1	17.3
2002.....	0.8	*	*	*	*	*	0.2	0.4	1.1	3.0	7.1	12.6
2001.....	0.8	*	*	*	*	*	0.2	0.4	1.1	2.8	6.9	13.3
2000.....	0.9	*	*	*	*	*	0.2	0.6	1.5	2.9	7.0	12.7
1999.....	0.8	*	*	*	*	*	0.2	0.5	1.1	3.7	7.0	11.1
White, female												
2004.....	1.7	*	*	*	*	*	0.2	0.5	1.4	4.2	10.4	20.8
2003.....	1.8	*	*	*	*	*	0.2	0.5	1.4	4.6	11.6	23.8
2002.....	1.9	*	*	*	*	*	0.2	0.6	1.9	4.8	12.1	21.6
2001.....	1.8	*	*	*	*	*	0.2	0.6	1.6	5.2	12.0	21.0
2000.....	1.9	*	*	*	*	*	0.3	0.7	1.9	6.0	12.2	19.6
1999.....	2.0	*	*	*	*	*	0.3	0.7	2.2	5.7	12.8	22.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year	2/	years	years	years	years	years	years	years	years	and over

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Black, both sexes

2004.....	0.6	*	*	*	*	*	0.6	1.0	1.6	2.7	4.0	5.9
2003.....	0.8	*	*	*	*	*	0.5	1.3	2.6	3.9	4.1	7.6
2002.....	0.7	*	*	*	*	*	0.3	0.9	2.4	2.9	4.4	*
2001.....	0.8	*	*	*	*	*	0.7	1.3	2.1	4.1	4.2	6.8
2000.....	0.7	*	*	*	*	*	0.6	1.0	2.0	3.5	3.9	8.4
1999.....	0.8	*	*	*	*	*	0.6	1.2	2.2	3.4	4.4	8.3

Black, male

2004.....	0.5	*	*	*	*	*	*	0.9	*	*	*	*
2003.....	0.6	*	*	*	*	*	*	0.9	2.7	3.0	*	*
2002.....	0.5	*	*	*	*	*	*	*	1.9	3.0	*	*
2001.....	0.5	*	*	*	*	*	*	1.0	*	*	*	*
2000.....	0.5	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.5	*	*	*	*	*	*	*	*	*	*	*

Black, female

2004.....	0.8	*	*	*	*	*	*	1.1	1.7	3.3	5.0	*
2003.....	1.0	*	*	*	*	*	*	1.7	2.5	4.5	5.1	8.1
2002.....	0.8	*	*	*	*	*	*	1.1	2.8	2.8	5.0	*
2001.....	1.1	*	*	*	*	*	0.9	1.7	2.6	5.3	5.3	*
2000.....	1.0	*	*	*	*	*	0.8	1.3	2.7	4.3	4.8	*
1999.....	1.0	*	*	*	*	*	0.7	1.4	2.6	4.1	4.5	11.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart disease (I11)												
All races 3/, both sexes												
2004.....	9.7	*	*	*	0.1	0.8	3.0	7.8	13.4	21.7	49.3	192.8
2003.....	9.7	*	*	*	0.1	0.8	2.9	7.9	13.1	22.1	51.0	195.0
2002.....	9.2	*	*	*	0.1	0.7	2.7	7.2	12.5	21.3	49.6	187.0
2001.....	8.7	*	*	*	0.1	0.6	2.4	6.3	10.9	19.8	48.7	189.5
2000.....	8.4	*	*	*	0.1	0.5	2.2	5.6	11.0	19.6	48.2	193.1
1999.....	8.1	*	*	*	0.1	0.5	2.1	5.3	10.7	19.1	47.7	184.9
All races, male												
2004.....	8.9	*	*	*	0.1	1.1	4.1	10.7	18.5	27.3	48.0	145.4
2003.....	8.8	*	*	*	0.2	1.0	4.0	10.8	17.2	27.1	51.0	142.2
2002.....	8.2	*	*	*	0.1	1.1	3.8	9.8	16.9	25.2	49.4	134.6
2001.....	7.5	*	*	*	*	0.8	3.3	8.6	14.4	23.6	48.4	134.4
2000.....	7.2	*	*	*	0.1	0.7	2.9	7.8	14.5	23.7	46.5	139.1
1999.....	6.8	*	*	*	0.1	0.7	2.9	7.0	13.7	21.9	46.8	133.6
All races, female												
2004.....	10.5	*	*	*	*	0.4	1.9	5.1	8.6	16.9	50.2	214.1
2003.....	10.7	*	*	*	*	0.5	1.9	5.2	9.3	17.8	51.0	218.3
2002.....	10.2	*	*	*	*	0.4	1.6	4.6	8.4	18.1	49.8	209.7
2001.....	9.8	*	*	*	*	0.4	1.6	4.1	7.8	16.6	48.8	212.6
2000.....	9.7	*	*	*	*	0.4	1.5	3.5	7.8	16.2	49.4	215.1
1999.....	9.4	*	*	*	*	0.3	1.3	3.6	7.9	16.8	48.3	205.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart disease (I11)												
White, both sexes												
2004.....	8.9	*	*	*	0.1	0.5	2.1	5.6	10.0	17.6	43.2	183.7
2003.....	8.8	*	*	*	0.1	0.5	2.0	5.4	9.8	17.3	45.1	185.3
2002.....	8.3	*	*	*	*	0.4	1.9	5.1	9.1	16.7	42.9	177.3
2001.....	7.8	*	*	*	*	0.4	1.6	4.2	7.9	15.2	42.6	179.9
2000.....	7.6	*	*	*	*	0.3	1.5	3.7	7.6	15.0	41.5	181.8
1999.....	7.3	*	*	*	*	0.3	1.3	3.4	7.5	14.4	40.9	175.3
White, male												
2004.....	7.8	*	*	*	*	0.7	3.1	8.1	13.9	22.3	41.9	136.4
2003.....	7.7	*	*	*	0.1	0.7	3.1	7.9	13.4	21.3	44.8	131.9
2002.....	7.1	*	*	*	*	0.7	2.8	7.4	12.7	19.9	42.6	124.8
2001.....	6.4	*	*	*	*	0.6	2.4	6.1	10.9	18.8	41.9	123.7
2000.....	6.1	*	*	*	*	0.4	2.1	5.5	10.4	18.6	39.6	129.3
1999.....	5.8	*	*	*	*	0.6	2.0	4.9	10.0	16.6	39.7	123.7
White, female												
2004.....	9.9	*	*	*	*	0.2	1.1	3.1	6.3	13.5	44.0	205.0
2003.....	10.0	*	*	*	*	0.2	1.0	2.9	6.4	13.9	45.2	209.0
2002.....	9.5	*	*	*	*	0.2	0.9	2.8	5.8	14.1	43.1	200.1
2001.....	9.2	*	*	*	*	0.2	0.8	2.4	5.1	12.3	43.0	203.5
2000.....	9.0	*	*	*	*	0.2	0.9	1.9	5.0	11.9	42.9	203.1
1999.....	8.7	*	*	*	*	*	0.7	2.0	5.1	12.5	41.7	196.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart disease (I11)												
Black, both sexes												
2004.....	17.6	*	*	*	0.3	2.7	9.2	26.2	45.1	64.6	126.8	314.7
2003.....	17.9	*	*	*	0.4	2.5	9.4	27.9	44.7	69.3	126.3	332.3
2002.....	17.2	*	*	*	*	2.8	8.6	24.3	44.4	68.0	133.0	320.4
2001.....	16.0	*	*	*	*	2.1	8.3	23.0	39.2	65.6	124.4	326.1
2000.....	16.1	*	*	*	*	2.1	7.4	21.2	42.2	66.8	131.9	340.0
1999.....	15.5	*	*	*	0.3	1.8	7.9	20.9	39.6	67.2	131.9	305.1
Black, male												
2004.....	18.4	*	*	*	*	3.5	11.7	33.6	64.6	86.6	131.7	267.2
2003.....	18.1	*	*	*	*	3.0	11.2	35.0	57.1	92.3	137.1	287.8
2002.....	17.5	*	*	*	*	3.9	11.6	31.3	60.8	83.6	141.1	267.0
2001.....	15.9	*	*	*	*	2.6	10.3	30.2	49.8	79.1	137.1	284.8
2000.....	15.6	*	*	*	*	2.6	9.2	27.8	54.9	81.4	139.3	273.3
1999.....	15.2	*	*	*	*	2.1	10.1	26.3	50.6	80.4	142.3	263.6
Black, female												
2004.....	16.9	*	*	*	*	1.9	7.0	19.8	29.4	48.9	124.0	333.5
2003.....	17.8	*	*	*	*	2.0	7.9	21.8	34.7	52.9	120.2	349.6
2002.....	16.9	*	*	*	*	1.7	6.0	18.3	31.2	56.8	128.5	340.9
2001.....	16.2	*	*	*	*	1.7	6.5	16.8	30.6	56.1	117.3	341.4
2000.....	16.5	*	*	*	*	1.7	5.8	15.5	31.9	56.6	127.7	364.9
1999.....	15.9	*	*	*	*	1.5	5.9	16.2	30.8	57.9	126.0	320.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart and renal disease (I13)												
All races 3/, both sexes												
2004.....	1.0	*	*	*	*	0.1	0.2	0.5	0.9	2.1	6.6	24.8
2003.....	1.1	*	*	*	*	0.1	0.2	0.5	1.0	2.2	7.0	25.6
2002.....	1.0	*	*	*	*	0.1	0.2	0.4	1.0	2.2	6.8	23.5
2001.....	1.0	*	*	*	*	0.1	0.2	0.4	1.0	2.3	6.8	23.3
2000.....	1.0	*	*	*	*	0.1	0.2	0.4	0.9	2.3	7.0	24.3
1999.....	1.2	*	*	*	*	0.1	0.3	0.4	1.1	2.6	8.9	29.1
All races, male												
2004.....	0.9	*	*	*	*	*	0.3	0.6	1.1	2.2	7.2	24.3
2003.....	0.9	*	*	*	*	*	0.3	0.6	1.2	2.3	7.4	26.7
2002.....	0.9	*	*	*	*	*	0.2	0.5	1.2	2.4	7.0	23.1
2001.....	0.8	*	*	*	*	*	0.3	0.5	1.0	2.5	6.8	24.2
2000.....	0.8	*	*	*	*	*	0.2	0.5	0.9	2.5	7.2	27.8
1999.....	1.0	*	*	*	*	*	0.3	0.6	1.3	2.7	9.6	31.3
All races, female												
2004.....	1.2	*	*	*	*	*	0.1	0.4	0.7	1.9	6.2	25.0
2003.....	1.2	*	*	*	*	*	0.1	0.3	0.7	2.1	6.8	25.2
2002.....	1.2	*	*	*	*	*	0.1	0.3	0.8	2.0	6.7	23.7
2001.....	1.1	*	*	*	*	*	0.2	0.2	0.9	2.1	6.8	22.9
2000.....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.1	6.9	22.9
1999.....	1.4	*	*	*	*	*	0.2	0.3	0.9	2.4	8.5	28.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart and renal disease (I13)												
White, both sexes												
2004.....	0.9	*	*	*	*	*	0.1	0.2	0.5	1.3	5.5	23.0
2003.....	0.9	*	*	*	*	*	0.1	0.2	0.5	1.4	6.0	23.4
2002.....	0.8	*	*	*	*	*	0.1	0.1	0.5	1.4	5.7	21.0
2001.....	0.8	*	*	*	*	*	0.1	0.2	0.5	1.5	5.7	21.0
2000.....	0.8	*	*	*	*	*	0.1	0.2	0.4	1.5	5.6	22.0
1999.....	1.0	*	*	*	*	*	0.1	0.2	0.5	1.5	7.3	26.1
White, male												
2004.....	0.7	*	*	*	*	*	*	0.2	0.6	1.5	6.1	22.9
2003.....	0.7	*	*	*	*	*	*	0.2	0.6	1.6	6.6	24.3
2002.....	0.7	*	*	*	*	*	*	0.1	0.6	1.6	6.0	20.6
2001.....	0.7	*	*	*	*	*	*	0.2	0.5	1.6	5.9	22.7
2000.....	0.7	*	*	*	*	*	0.1	0.2	0.5	1.7	6.2	24.4
1999.....	0.8	*	*	*	*	*	*	0.2	0.6	1.7	8.2	28.8
White, female												
2004.....	1.0	*	*	*	*	*	*	0.1	0.4	1.2	5.1	23.0
2003.....	1.0	*	*	*	*	*	*	*	0.4	1.3	5.5	22.9
2002.....	1.0	*	*	*	*	*	*	*	0.4	1.2	5.6	21.1
2001.....	1.0	*	*	*	*	*	*	0.1	0.5	1.4	5.5	20.3
2000.....	1.0	*	*	*	*	*	*	0.1	0.4	1.4	5.2	21.1
1999.....	1.1	*	*	*	*	*	*	0.1	0.5	1.4	6.7	25.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart and renal disease (I13)												
Black, both sexes												
2004.....	2.3	*	*	*	*	*	1.2	2.8	4.9	8.5	19.6	48.2
2003.....	2.4	*	*	*	*	0.4	1.2	3.0	5.4	9.0	19.6	53.1
2002.....	2.4	*	*	*	*	*	0.9	2.7	6.1	9.9	19.6	56.1
2001.....	2.2	*	*	*	*	0.4	1.2	2.1	5.0	9.9	20.1	50.1
2000.....	2.2	*	*	*	*	*	0.8	2.2	4.6	9.9	22.8	50.4
1999.....	2.8	*	*	*	*	*	1.5	2.4	6.0	12.5	26.9	65.4
Black, male												
2004.....	2.2	*	*	*	*	*	1.9	3.5	6.1	8.7	23.1	41.7
2003.....	2.3	*	*	*	*	*	1.7	4.0	7.1	8.9	18.9	50.0
2002.....	2.3	*	*	*	*	*	1.0	3.8	7.5	10.1	19.2	61.1
2001.....	2.1	*	*	*	*	*	1.6	3.2	5.9	11.9	18.5	44.3
2000.....	2.0	*	*	*	*	*	1.0	2.8	5.3	11.3	20.9	63.4
1999.....	2.5	*	*	*	*	*	1.8	3.5	7.6	13.5	25.8	59.8
Black, female												
2004.....	2.4	*	*	*	*	*	*	2.3	4.0	8.3	17.6	50.8
2003.....	2.5	*	*	*	*	*	0.8	2.1	4.0	9.1	19.9	54.3
2002.....	2.5	*	*	*	*	*	0.9	1.8	4.9	9.7	19.8	54.2
2001.....	2.3	*	*	*	*	*	0.8	1.1	4.2	8.6	21.0	52.2
2000.....	2.4	*	*	*	*	*	*	1.6	4.1	9.0	23.9	45.5
1999.....	3.0	*	*	*	*	*	1.2	1.5	4.8	11.7	27.5	67.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Ischemic heart diseases (I20-I25)

All races 3/, both sexes

2004.....	153.7	*	*	*	0.3	2.7	16.2	59.9	158.2	395.9	1,072.6	3,287.4
2003.....	165.1	0.5	*	0.1	0.4	2.7	17.0	62.2	169.8	430.2	1,160.7	3,610.2
2002.....	171.4	0.6	*	0.0	0.3	2.5	17.1	63.8	177.7	458.9	1,220.2	3,775.0
2001.....	176.3	0.6	*	0.0	0.4	2.8	16.7	64.0	184.0	477.6	1,271.3	3,958.2
2000.....	183.1	0.8	*	*	0.3	2.3	16.7	66.1	196.7	505.1	1,324.2	4,182.5
1999.....	189.8	0.7	*	0.1	0.4	2.5	17.2	67.7	203.5	536.6	1,384.8	4,318.1

All races, male

2004.....	161.6	*	*	*	0.4	4.0	24.4	92.8	234.8	549.1	1,410.3	3,680.3
2003.....	172.2	*	*	*	0.5	4.0	25.7	97.2	251.5	598.8	1,524.4	4,005.2
2002.....	178.4	*	*	*	0.4	3.7	26.1	99.4	263.3	640.1	1,598.9	4,205.3
2001.....	181.7	*	*	*	0.6	4.1	25.5	99.5	271.6	663.7	1,663.5	4,395.8
2000.....	188.7	*	*	*	0.4	3.5	25.8	103.5	290.2	705.2	1,736.8	4,701.7
1999.....	195.4	*	*	*	0.5	3.7	26.6	106.2	299.3	749.8	1,813.7	4,837.5

All races, female

2004.....	146.1	*	*	*	0.1	1.4	8.0	28.0	87.1	267.3	845.3	3,110.6
2003.....	158.1	*	*	*	0.3	1.4	8.4	28.3	94.0	289.2	917.7	3,435.5
2002.....	164.7	*	*	*	0.2	1.3	8.2	29.5	98.5	308.1	968.9	3,588.3
2001.....	171.2	*	*	*	0.2	1.5	8.1	29.7	103.1	323.4	1,013.6	3,775.1
2000.....	177.6	*	*	*	0.2	1.2	7.6	29.9	110.4	340.3	1,055.2	3,971.0
1999.....	184.5	*	*	*	0.2	1.4	7.9	30.7	115.1	362.0	1,106.3	4,108.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ischemic heart diseases (I20-I25)												
White, both sexes												
2004.....	166.1	*	*	*	0.2	2.5	15.7	56.8	151.3	385.6	1,073.4	3,343.9
2003.....	178.3	*	*	*	0.3	2.5	16.4	58.8	162.0	418.9	1,159.0	3,671.7
2002.....	185.4	*	*	*	0.3	2.3	16.5	60.7	169.5	448.8	1,217.8	3,836.4
2001.....	190.5	*	*	*	0.3	2.6	15.9	60.7	175.1	465.8	1,268.5	4,022.7
2000.....	197.9	0.8	*	*	0.3	2.1	15.8	62.8	187.3	493.4	1,320.9	4,257.7
1999.....	205.1	*	*	*	0.3	2.3	16.2	64.4	194.3	524.8	1,383.7	4,389.2
White, male												
2004.....	174.3	*	*	*	0.3	3.7	24.1	89.6	227.1	539.9	1,422.1	3,768.8
2003.....	185.5	*	*	*	0.5	3.8	25.1	93.5	243.5	587.9	1,529.5	4,106.1
2002.....	192.5	*	*	*	0.4	3.5	25.6	96.3	254.3	632.9	1,608.0	4,299.7
2001.....	195.8	*	*	*	0.5	3.9	24.7	96.3	262.3	653.8	1,672.4	4,494.0
2000.....	203.6	*	*	*	0.4	3.3	25.1	100.5	280.3	697.3	1,744.8	4,813.9
1999.....	210.5	*	*	*	0.4	3.4	25.5	102.8	289.0	741.3	1,827.8	4,933.9
White, female												
2004.....	158.1	*	*	*	*	1.2	7.2	24.4	79.5	253.5	835.6	3,152.3
2003.....	171.4	*	*	*	0.2	1.2	7.5	24.6	85.0	274.9	908.5	3,479.3
2002.....	178.4	*	*	*	0.2	1.1	7.3	25.5	89.5	292.7	956.0	3,635.2
2001.....	185.3	*	*	*	0.2	1.3	7.0	25.6	93.1	307.2	1,000.3	3,825.5
2000.....	192.4	*	*	*	0.2	0.9	6.5	25.6	100.0	322.5	1,042.0	4,031.8
1999.....	200.0	*	*	*	0.1	1.3	6.8	26.7	105.5	344.3	1,092.6	4,170.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ischemic heart diseases (I20-I25)												
Black, both sexes												
2004.....	121.0	*	*	*	0.7	4.7	23.2	96.0	255.0	568.6	1,243.5	3,014.3
2003.....	129.2	*	*	*	0.7	4.3	24.7	99.3	273.1	615.2	1,363.0	3,310.1
2002.....	132.9	*	*	*	0.5	3.9	25.1	101.6	288.0	636.1	1,422.7	3,462.9
2001.....	136.9	*	*	*	0.9	4.6	25.6	102.3	297.9	673.0	1,479.0	3,615.1
2000.....	140.1	*	*	*	0.4	4.3	25.6	105.6	314.6	702.2	1,523.2	3,684.7
1999.....	143.7	*	*	*	0.8	3.9	27.5	108.5	320.1	735.6	1,552.7	3,805.8
Black, male												
2004.....	125.2	*	*	*	1.1	6.5	31.9	138.4	362.4	761.9	1,537.6	3,100.5
2003.....	133.4	*	*	*	0.9	6.2	35.0	144.5	381.8	827.3	1,744.8	3,285.3
2002.....	135.3	*	*	*	*	5.1	35.2	144.3	407.3	840.8	1,748.8	3,556.2
2001.....	138.9	*	*	*	1.2	6.6	36.0	145.7	412.8	894.0	1,825.4	3,775.0
2000.....	141.5	*	*	*	*	5.9	35.6	149.4	439.2	907.9	1,877.0	3,885.2
1999.....	146.0	*	*	*	0.9	5.7	39.5	156.9	452.9	962.3	1,870.8	4,080.1
Black, female												
2004.....	117.1	*	*	*	*	3.0	15.5	59.6	168.6	430.7	1,077.3	2,980.1
2003.....	125.3	*	*	*	*	2.5	15.6	60.5	185.6	464.3	1,147.2	3,319.7
2002.....	130.8	*	*	*	*	2.8	16.3	64.9	192.2	490.8	1,238.1	3,427.2
2001.....	135.2	*	*	*	*	2.8	16.4	65.0	205.3	517.1	1,284.5	3,555.7
2000.....	138.8	*	*	*	*	2.8	16.7	68.0	214.1	557.4	1,324.0	3,610.0
1999.....	141.7	*	*	*	0.7	2.3	16.9	67.0	213.6	575.8	1,374.6	3,703.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute myocardial infarction (I21-I22)												
All races 3/, both sexes												
2004.....	53.4	*	*	*	0.1	1.1	6.4	24.2	64.5	153.4	370.3	992.9
2003.....	58.7	*	*	*	0.2	1.0	7.0	25.5	70.9	168.6	412.7	1,118.8
2002.....	62.3	*	*	*	0.2	1.0	7.0	26.5	74.9	186.1	444.4	1,191.3
2001.....	64.9	*	*	*	0.2	1.2	7.1	27.6	78.4	193.2	471.2	1,265.4
2000.....	68.5	0.5	*	*	0.1	0.9	6.9	28.9	86.6	209.1	503.3	1,338.5
1999.....	71.5	*	*	*	0.2	1.1	7.2	29.6	90.0	225.0	529.3	1,379.3
All races, male												
2004.....	57.4	*	*	*	0.2	1.6	9.6	37.3	95.3	209.5	477.0	1,120.5
2003.....	62.6	*	*	*	0.2	1.5	10.4	39.8	104.1	231.0	530.6	1,263.5
2002.....	66.2	*	*	*	0.2	1.4	10.5	40.6	109.2	256.2	569.2	1,366.4
2001.....	68.6	*	*	*	0.2	1.7	10.6	42.4	114.0	264.7	604.1	1,461.3
2000.....	72.7	*	*	*	0.1	1.3	10.6	44.7	125.7	286.9	648.3	1,568.3
1999.....	76.1	*	*	*	0.2	1.5	11.0	46.3	131.8	310.0	684.4	1,622.1
All races, female												
2004.....	49.6	*	*	*	*	0.6	3.3	11.6	35.9	106.3	298.5	935.5
2003.....	54.8	*	*	*	0.1	0.6	3.6	11.7	40.0	116.5	333.9	1,054.8
2002.....	58.4	*	*	*	0.1	0.5	3.6	12.9	43.2	127.7	361.5	1,115.3
2001.....	61.3	*	*	*	*	0.7	3.5	13.3	45.4	134.0	383.9	1,183.5
2000.....	64.6	*	*	*	0.1	0.5	3.3	13.6	50.5	145.1	408.7	1,244.8
1999.....	67.0	*	*	*	0.1	0.6	3.4	13.5	51.5	155.4	428.6	1,281.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute myocardial infarction (I21-I22)												
White, both sexes												
2004.....	57.8	*	*	*	0.1	1.1	6.4	23.6	63.5	151.2	372.1	1,004.1
2003.....	63.4	*	*	*	0.2	1.0	6.8	24.7	69.5	166.8	412.7	1,132.6
2002.....	67.3	*	*	*	0.1	1.0	6.9	25.9	73.5	184.2	443.3	1,201.2
2001.....	70.1	*	*	*	0.1	1.1	6.8	26.9	76.8	190.8	471.1	1,277.8
2000.....	74.1	*	*	*	0.1	0.9	6.8	28.2	84.9	206.7	502.6	1,353.5
1999.....	77.4	*	*	*	0.1	1.0	7.0	29.1	88.6	223.3	530.2	1,393.2
White, male												
2004.....	62.3	*	*	*	0.1	1.6	9.7	37.0	95.1	208.6	484.2	1,143.7
2003.....	67.7	*	*	*	0.2	1.4	10.3	39.2	103.6	230.7	532.6	1,289.3
2002.....	71.8	*	*	*	0.2	1.4	10.5	40.4	108.9	257.3	572.0	1,386.1
2001.....	74.4	*	*	*	0.2	1.6	10.6	42.4	113.7	264.7	609.1	1,481.1
2000.....	78.8	*	*	*	0.2	1.3	10.6	44.7	125.0	287.2	652.5	1,592.9
1999.....	82.7	*	*	*	0.2	1.4	10.9	46.4	131.3	311.1	692.3	1,644.0
White, female												
2004.....	53.3	*	*	*	*	0.5	3.0	10.4	33.5	102.1	295.6	941.1
2003.....	59.2	*	*	*	*	0.5	3.2	10.4	37.2	112.3	331.5	1,063.2
2002.....	62.8	*	*	*	*	0.5	3.3	11.5	40.1	122.2	357.0	1,120.8
2001.....	66.0	*	*	*	*	0.7	3.1	11.7	42.0	128.4	379.6	1,192.7
2000.....	69.6	*	*	*	*	0.4	2.9	11.9	47.3	139.3	404.1	1,256.2
1999.....	72.3	*	*	*	*	0.6	3.1	12.1	48.6	150.1	423.9	1,292.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute myocardial infarction (I21-I22)												
Black, both sexes												
2004.....	42.0	*	*	*	0.3	1.7	8.8	35.4	91.1	204.7	420.0	979.9
2003.....	45.7	*	*	*	*	1.6	10.1	37.5	98.6	220.0	479.9	1,094.9
2002.....	48.5	*	*	*	*	1.6	9.7	37.9	106.9	238.8	517.3	1,201.2
2001.....	50.0	*	*	*	0.3	1.8	10.0	39.2	109.8	252.6	537.8	1,239.9
2000.....	52.2	*	*	*	*	1.7	9.9	41.1	120.1	268.7	571.1	1,275.7
1999.....	53.3	*	*	*	*	1.6	10.3	40.9	120.9	279.5	584.6	1,318.8
Black, male												
2004.....	42.4	*	*	*	*	2.0	11.6	50.3	123.5	265.8	486.1	1,009.3
2003.....	46.6	*	*	*	*	2.3	13.8	53.0	133.3	283.0	600.2	1,113.5
2002.....	48.3	*	*	*	*	2.1	13.3	52.0	141.6	300.1	623.4	1,280.1
2001.....	49.2	*	*	*	*	2.5	13.2	51.9	143.8	317.1	638.0	1,382.0
2000.....	52.0	*	*	*	*	2.2	13.4	54.9	160.3	335.8	687.3	1,418.5
1999.....	53.4	*	*	*	*	2.3	14.2	56.6	165.5	353.6	685.6	1,498.7
Black, female												
2004.....	41.6	*	*	*	*	1.4	6.2	22.5	65.0	161.1	382.7	968.3
2003.....	44.9	*	*	*	*	1.0	6.7	24.1	70.7	175.2	411.9	1,087.7
2002.....	48.8	*	*	*	*	1.2	6.5	25.9	79.1	195.3	457.2	1,171.0
2001.....	50.6	*	*	*	*	1.2	7.1	28.2	82.4	207.0	481.5	1,187.2
2000.....	52.5	*	*	*	*	1.3	6.8	29.1	87.7	221.6	505.6	1,222.5
1999.....	53.2	*	*	*	*	0.9	6.9	27.4	85.1	227.4	528.1	1,251.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute ischemic heart diseases (I24)												
All races 3/, both sexes												
2004.....	0.8	*	*	*	*	*	0.2	0.5	1.2	2.2	5.2	14.8
2003.....	1.1	*	*	*	*	*	0.2	0.7	1.4	3.2	6.5	20.0
2002.....	1.2	*	*	*	*	*	0.2	0.9	1.7	3.5	7.1	21.1
2001.....	1.2	*	*	*	*	*	0.2	0.8	1.8	3.5	6.9	21.5
2000.....	1.2	*	*	*	*	*	0.2	0.7	2.0	3.5	7.3	21.9
1999.....	1.2	*	*	*	*	*	0.2	0.7	1.9	3.7	7.0	21.9
All races, male												
2004.....	0.9	*	*	*	*	*	0.3	0.7	1.8	3.0	6.3	16.3
2003.....	1.1	*	*	*	*	*	0.3	1.0	2.1	4.3	8.0	19.5
2002.....	1.2	*	*	*	*	*	0.2	1.3	2.5	4.7	8.7	21.4
2001.....	1.2	*	*	*	*	*	0.3	1.2	2.6	4.7	8.5	20.7
2000.....	1.3	*	*	*	*	*	0.3	1.2	2.9	4.8	9.1	20.7
1999.....	1.2	*	*	*	*	*	0.3	1.0	2.8	4.9	8.8	22.3
All races, female												
2004.....	0.8	*	*	*	*	*	0.1	0.3	0.7	1.4	4.4	14.1
2003.....	1.0	*	*	*	*	*	0.1	0.4	0.9	2.2	5.5	20.2
2002.....	1.1	*	*	*	*	*	0.1	0.5	1.0	2.4	5.9	21.0
2001.....	1.1	*	*	*	*	*	0.1	0.4	1.1	2.5	5.9	21.8
2000.....	1.1	*	*	*	*	*	0.1	0.3	1.1	2.4	6.1	22.4
1999.....	1.1	*	*	*	*	*	*	0.4	1.1	2.6	5.8	21.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute ischemic heart diseases (I24)												
White, both sexes												
2004.....	0.8	*	*	*	*	*	0.2	0.4	1.1	2.0	4.9	15.0
2003.....	1.1	*	*	*	*	*	0.2	0.6	1.3	3.0	6.3	20.2
2002.....	1.2	*	*	*	*	*	0.2	0.8	1.6	3.3	7.0	21.4
2001.....	1.2	*	*	*	*	*	0.2	0.7	1.6	3.3	6.8	21.5
2000.....	1.2	*	*	*	*	*	0.2	0.7	1.8	3.3	7.1	22.1
1999.....	1.2	*	*	*	*	*	0.1	0.6	1.7	3.5	6.8	22.4
White, male												
2004.....	0.9	*	*	*	*	*	0.3	0.6	1.6	2.6	6.1	16.2
2003.....	1.1	*	*	*	*	*	0.3	0.9	1.8	4.1	7.6	19.7
2002.....	1.3	*	*	*	*	*	0.2	1.1	2.3	4.4	8.6	21.9
2001.....	1.2	*	*	*	*	*	0.2	1.1	2.4	4.4	8.5	20.7
2000.....	1.3	*	*	*	*	*	0.3	1.1	2.6	4.4	8.9	21.0
1999.....	1.2	*	*	*	*	*	0.2	0.8	2.5	4.7	8.6	22.9
White, female												
2004.....	0.8	*	*	*	*	*	*	0.2	0.6	1.4	4.2	14.4
2003.....	1.1	*	*	*	*	*	*	0.3	0.8	2.0	5.5	20.5
2002.....	1.2	*	*	*	*	*	*	0.4	0.9	2.3	5.9	21.1
2001.....	1.2	*	*	*	*	*	*	0.3	0.9	2.4	5.7	21.7
2000.....	1.2	*	*	*	*	*	*	0.3	0.9	2.3	5.9	22.6
1999.....	1.2	*	*	*	*	*	*	0.3	1.1	2.5	5.6	22.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute ischemic heart diseases (I24)												
Black, both sexes												
2004.....	1.0	*	*	*	*	*	*	1.4	3.0	4.1	8.8	15.2
2003.....	1.1	*	*	*	*	*	0.5	1.3	3.0	5.5	9.9	19.8
2002.....	1.2	*	*	*	*	*	0.3	1.7	3.0	5.9	9.3	21.1
2001.....	1.2	*	*	*	*	*	0.5	1.5	3.9	5.7	8.5	24.0
2000.....	1.2	*	*	*	*	*	0.4	1.5	4.2	5.6	10.8	21.9
1999.....	1.2	*	*	*	*	*	0.5	1.8	3.6	5.7	10.7	17.2
Black, male												
2004.....	1.3	*	*	*	*	*	*	1.9	4.6	7.1	11.7	22.8
2003.....	1.4	*	*	*	*	*	*	1.9	4.8	7.6	16.1	24.0
2002.....	1.4	*	*	*	*	*	*	2.7	4.2	9.1	11.8	21.4
2001.....	1.4	*	*	*	*	*	*	2.4	5.5	8.5	10.3	23.8
2000.....	1.5	*	*	*	*	*	*	2.4	6.1	8.6	13.9	*
1999.....	1.4	*	*	*	*	*	0.7	2.6	6.1	8.2	12.6	*
Black, female												
2004.....	0.8	*	*	*	*	*	*	0.9	1.7	2.0	7.2	12.2
2003.....	0.9	*	*	*	*	*	*	*	1.5	4.0	6.4	18.2
2002.....	1.0	*	*	*	*	*	*	0.8	2.0	3.6	8.0	20.9
2001.....	1.1	*	*	*	*	*	*	*	2.6	3.7	7.4	24.0
2000.....	1.0	*	*	*	*	*	*	*	2.7	3.5	9.0	22.3
1999.....	1.0	*	*	*	*	*	*	1.1	1.5	3.8	9.6	16.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)												
All races 3/, both sexes												
2004.....	99.5	*	*	*	0.2	1.6	9.6	35.1	92.5	240.3	697.1	2,279.7
2003.....	105.3	*	*	*	0.2	1.7	9.8	35.9	97.5	258.4	741.5	2,471.4
2002.....	108.0	*	*	*	0.2	1.4	9.9	36.5	101.0	269.4	768.8	2,562.5
2001.....	110.3	*	*	*	0.2	1.6	9.5	35.6	103.8	280.9	793.1	2,671.3
2000.....	113.3	*	*	*	0.2	1.4	9.5	36.4	108.1	292.4	813.6	2,822.1
1999.....	117.2	*	*	*	0.2	1.4	9.8	37.5	111.5	308.0	848.5	2,917.0
All races, male												
2004.....	103.3	*	*	*	0.2	2.4	14.5	54.8	137.7	336.5	926.9	2,543.6
2003.....	108.5	*	*	*	0.3	2.5	14.9	56.4	145.4	363.5	985.8	2,722.2
2002.....	110.9	*	*	*	0.2	2.2	15.4	57.5	151.5	379.3	1,021.0	2,817.4
2001.....	111.9	*	*	*	0.3	2.4	14.6	55.9	154.9	394.3	1,050.9	2,913.8
2000.....	114.8	*	*	*	0.2	2.1	14.8	57.7	161.6	413.5	1,079.4	3,112.7
1999.....	118.0	*	*	*	0.3	2.1	15.3	59.0	164.7	434.9	1,120.5	3,193.1
All races, female												
2004.....	95.7	*	*	*	0.7	4.6	16.2	50.5	159.6	542.4	2,161.0	
2003.....	102.3	*	*	*	0.1	0.8	4.7	16.2	53.1	170.5	578.3	2,360.5
2002.....	105.2	*	*	*	0.7	4.4	16.2	54.3	178.0	601.4	2,451.9	
2001.....	108.8	*	*	*	0.1	0.7	4.4	16.0	56.7	186.9	623.7	2,569.8
2000.....	111.9	*	*	*	0.6	4.3	16.0	58.8	192.8	640.3	2,703.8	
1999.....	116.4	*	*	*	0.1	0.7	4.3	16.7	62.6	204.0	671.8	2,805.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)												
White, both sexes												
2004.....	107.5	*	*	*	0.1	1.4	9.2	32.8	86.7	232.4	696.4	2,324.8
2003.....	113.8	*	*	*	0.2	1.5	9.4	33.5	91.3	249.1	740.0	2,518.8
2002.....	116.9	*	*	*	0.2	1.4	9.4	34.0	94.4	261.3	767.5	2,613.9
2001.....	119.1	*	*	*	0.2	1.4	8.9	33.1	96.7	271.7	790.5	2,723.5
2000.....	122.6	*	*	*	0.2	1.2	8.9	33.9	100.6	283.4	811.2	2,882.1
1999.....	126.5	*	*	*	0.1	1.3	9.0	34.7	104.0	298.0	846.7	2,973.6
White, male												
2004.....	111.1	*	*	*	0.2	2.2	14.2	52.1	130.5	328.7	931.8	2,608.9
2003.....	116.6	*	*	*	0.2	2.3	14.4	53.4	138.1	353.1	989.3	2,797.1
2002.....	119.4	*	*	*	0.2	2.1	14.9	54.8	143.1	371.2	1,027.4	2,891.6
2001.....	120.1	*	*	*	0.3	2.2	13.9	52.9	146.2	384.6	1,054.9	2,992.2
2000.....	123.5	*	*	*	0.2	1.9	14.2	54.8	152.7	405.6	1,083.4	3,200.0
1999.....	126.5	*	*	*	0.2	2.0	14.4	55.5	155.2	425.5	1,126.9	3,267.1
White, female												
2004.....	103.9	*	*	*	*	0.6	4.1	13.8	45.4	149.9	535.9	2,196.8
2003.....	111.0	*	*	*	*	0.7	4.2	13.9	47.0	160.6	571.4	2,395.5
2002.....	114.4	*	*	*	*	0.6	3.9	13.6	48.5	168.1	593.1	2,493.2
2001.....	118.1	*	*	*	*	0.6	3.9	13.6	50.2	176.4	615.0	2,611.0
2000.....	121.6	*	*	*	*	0.5	3.5	13.4	51.8	180.9	632.0	2,753.0
1999.....	126.5	*	*	*	*	0.6	3.6	14.3	55.9	191.7	663.1	2,855.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/	1-4 2/	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)												
Black, both sexes												
2004.....	78.0	*	*	*	0.4	2.9	14.2	59.2	161.0	359.8	814.7	2,019.1
2003.....	82.3	*	*	*	0.4	2.6	14.2	60.6	171.5	389.7	873.2	2,195.3
2002.....	83.2	*	*	*	*	2.2	15.1	61.9	178.1	391.4	896.1	2,240.6
2001.....	85.8	*	*	*	0.5	2.7	15.1	61.6	184.2	414.8	932.7	2,351.2
2000.....	86.6	*	*	*	*	2.5	15.3	63.1	190.2	427.8	941.3	2,387.1
1999.....	89.3	*	*	*	0.5	2.3	16.8	65.8	195.7	450.4	957.4	2,469.8
Black, male												
2004.....	81.5	*	*	*	*	4.3	19.8	86.2	234.3	489.0	1,039.8	2,068.3
2003.....	85.5	*	*	*	*	3.9	20.6	89.6	243.7	536.7	1,128.5	2,147.8
2002.....	85.7	*	*	*	*	2.9	21.7	89.5	261.4	531.5	1,113.6	2,254.6
2001.....	88.2	*	*	*	0.7	4.2	22.2	91.3	263.6	568.3	1,177.1	2,369.1
2000.....	88.0	*	*	*	*	3.6	21.6	92.1	272.7	563.5	1,175.7	2,446.0
1999.....	91.1	*	*	*	*	3.3	24.6	97.7	281.3	600.5	1,172.6	2,561.5
Black, female												
2004.....	74.7	*	*	*	*	1.5	9.1	36.1	102.0	267.6	687.4	1,999.6
2003.....	79.4	*	*	*	*	1.4	8.5	35.7	113.4	285.0	728.9	2,213.8
2002.....	81.0	*	*	*	*	1.5	9.4	38.2	111.1	291.9	772.9	2,235.2
2001.....	83.5	*	*	*	*	1.5	8.9	36.1	120.3	306.4	795.5	2,344.5
2000.....	85.3	*	*	*	*	1.5	9.7	38.1	123.7	332.3	809.4	2,365.2
1999.....	87.6	*	*	*	*	1.3	9.9	38.5	127.0	344.6	836.9	2,435.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)												
All races 3/, both sexes												
2004.....	21.7	*	*	*	*	0.5	3.3	13.6	32.0	58.7	129.0	400.4
2003.....	23.1	*	*	*	*	0.5	3.4	13.9	32.7	62.7	142.5	442.9
2002.....	23.6	*	*	*	*	0.5	3.4	13.4	33.1	64.7	146.2	474.5
2001.....	23.9	*	*	*	0.1	0.5	3.2	12.9	33.3	65.6	152.7	493.6
2000.....	24.7	*	*	*	*	0.4	3.0	13.2	34.0	68.3	158.1	534.6
1999.....	25.8	*	*	*	*	0.4	3.3	13.0	35.3	72.7	166.2	563.4
All races, male												
2004.....	23.9	*	*	*	*	0.8	5.2	22.0	49.0	83.6	164.9	414.0
2003.....	25.2	*	*	*	*	0.9	5.2	22.7	49.9	88.9	181.1	462.1
2002.....	25.4	*	*	*	*	0.7	5.4	21.5	51.2	91.8	185.3	491.8
2001.....	25.3	*	*	*	*	0.7	5.0	21.0	51.4	92.4	193.9	502.0
2000.....	26.0	*	*	*	*	0.6	4.9	21.5	52.6	97.2	199.3	550.8
1999.....	26.8	*	*	*	*	0.6	5.3	20.9	53.7	103.0	207.0	578.6
All races, female												
2004.....	19.5	*	*	*	*	0.2	1.4	5.5	16.1	37.7	104.8	394.3
2003.....	21.1	*	*	*	*	0.2	1.5	5.5	16.7	40.8	116.7	434.4
2002.....	21.9	*	*	*	*	0.2	1.3	5.5	16.2	42.1	120.2	467.1
2001.....	22.5	*	*	*	*	0.2	1.4	5.1	16.7	43.3	125.5	490.1
2000.....	23.5	*	*	*	*	0.1	1.2	5.2	16.8	44.6	131.3	528.0
1999.....	24.8	*	*	*	*	0.2	1.3	5.3	18.3	47.9	139.8	557.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)												
White, both sexes												
2004.....	22.1	*	*	*	*	0.4	3.2	12.3	28.7	53.2	122.2	395.6
2003.....	23.6	*	*	*	*	0.5	3.2	12.6	29.0	56.5	135.5	439.3
2002.....	24.1	*	*	*	*	0.4	3.1	12.1	29.4	58.1	138.4	470.6
2001.....	24.3	*	*	*	*	0.4	3.0	11.6	28.8	58.3	144.5	489.7
2000.....	25.1	*	*	*	*	0.3	2.8	11.8	29.4	61.2	149.2	529.5
1999.....	26.1	*	*	*	*	0.3	3.0	11.3	30.4	64.8	157.0	557.5
White, male												
2004.....	24.1	*	*	*	*	0.7	5.1	20.4	44.1	76.6	156.3	408.8
2003.....	25.3	*	*	*	*	0.8	5.0	20.8	44.8	80.0	171.9	458.7
2002.....	25.5	*	*	*	*	0.7	5.1	19.9	45.9	82.9	174.7	488.0
2001.....	25.3	*	*	*	*	0.6	4.7	19.2	45.1	82.7	184.2	498.6
2000.....	26.0	*	*	*	*	0.5	4.6	19.6	46.3	87.8	187.5	545.6
1999.....	26.6	*	*	*	*	0.5	4.9	18.6	46.9	92.5	195.2	571.2
White, female												
2004.....	20.1	*	*	*	*	0.2	1.3	4.3	14.1	33.1	99.0	389.7
2003.....	21.9	*	*	*	*	0.2	1.4	4.5	14.1	36.5	110.8	430.7
2002.....	22.7	*	*	*	*	0.1	1.1	4.5	13.8	37.0	114.0	463.0
2001.....	23.3	*	*	*	*	0.2	1.2	4.1	13.4	37.8	118.2	486.0
2000.....	24.3	*	*	*	*	*	1.0	4.2	13.5	39.0	123.9	522.9
1999.....	25.6	*	*	*	*	0.2	1.0	4.2	15.0	41.6	131.9	552.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/	1-4 2/	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)												
Black, both sexes												
2004.....	25.0	*	*	*	*	0.9	4.8	25.7	66.7	120.0	226.9	505.1
2003.....	26.0	*	*	*	*	0.9	4.9	25.7	69.7	128.1	241.6	543.2
2002.....	27.0	*	*	*	*	0.8	5.8	25.5	71.4	136.3	259.0	575.5
2001.....	27.7	*	*	*	*	0.7	5.6	25.3	77.5	143.4	266.5	594.5
2000.....	28.5	*	*	*	*	0.7	5.1	26.9	78.2	145.5	278.4	646.6
1999.....	29.9	*	*	*	*	0.7	6.3	27.5	81.4	158.6	291.2	668.6
Black, male												
2004.....	28.9	*	*	*	*	1.5	6.9	38.1	104.6	172.0	306.2	531.5
2003.....	30.5	*	*	*	*	1.5	7.7	40.4	106.0	190.3	331.4	583.9
2002.....	31.1	*	*	*	*	1.1	8.6	38.7	111.4	197.2	357.2	599.3
2001.....	31.5	*	*	*	*	0.9	8.5	39.2	117.4	208.2	351.7	609.3
2000.....	32.3	*	*	*	*	1.0	7.8	42.0	120.0	207.3	369.7	701.2
1999.....	33.8	*	*	*	*	1.0	9.6	42.8	123.3	229.3	380.0	715.9
Black, female												
2004.....	21.5	*	*	*	*	*	3.0	15.1	36.2	83.0	182.0	494.6
2003.....	21.9	*	*	*	*	*	2.6	13.1	40.5	83.9	190.9	527.4
2002.....	23.3	*	*	*	*	*	3.3	14.2	39.2	93.0	203.5	566.4
2001.....	24.3	*	*	*	*	*	3.1	13.3	45.4	97.7	218.6	589.0
2000.....	25.0	*	*	*	*	*	2.8	14.0	44.6	102.0	227.1	626.3
1999.....	26.4	*	*	*	*	*	3.4	14.4	47.8	108.8	241.5	650.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
All races 3/, both sexes												
2004.....	77.8	*	*	*	0.1	1.1	6.2	21.5	60.5	181.7	568.1	1,879.3
2003.....	82.2	*	*	*	0.2	1.1	6.5	22.0	64.9	195.7	599.0	2,028.5
2002.....	84.4	*	*	*	0.1	1.0	6.5	23.1	68.0	204.7	622.6	2,088.0
2001.....	86.4	*	*	*	0.2	1.1	6.2	22.7	70.5	215.3	640.5	2,177.7
2000.....	88.6	*	*	*	0.1	1.0	6.5	23.2	74.1	224.1	655.5	2,287.5
1999.....	91.4	*	*	*	0.2	1.0	6.5	24.5	76.3	235.2	682.3	2,353.6
All races, male												
2004.....	79.4	*	*	*	0.2	1.7	9.3	32.8	88.7	252.9	762.0	2,129.6
2003.....	83.3	*	*	*	0.2	1.6	9.7	33.7	95.5	274.6	804.7	2,260.1
2002.....	85.5	*	*	*	0.2	1.5	10.0	35.9	100.3	287.5	835.7	2,325.6
2001.....	86.5	*	*	*	0.2	1.8	9.5	34.9	103.5	301.9	856.9	2,411.8
2000.....	88.8	*	*	*	0.2	1.5	9.9	36.2	109.0	316.3	880.1	2,561.9
1999.....	91.2	*	*	*	0.2	1.5	10.0	38.1	111.0	331.9	913.6	2,614.4
All races, female												
2004.....	76.2	*	*	*	*	0.5	3.2	10.6	34.3	121.9	437.6	1,766.7
2003.....	81.1	*	*	*	0.1	0.6	3.2	10.8	36.4	129.7	461.6	1,926.2
2002.....	83.3	*	*	*	*	0.5	3.1	10.7	38.0	135.9	481.1	1,984.9
2001.....	86.2	*	*	*	0.1	0.5	3.0	11.0	40.0	143.6	498.2	2,079.8
2000.....	88.4	*	*	*	*	0.5	3.0	10.7	42.0	148.2	509.0	2,175.8
1999.....	91.6	*	*	*	0.1	0.5	3.0	11.4	44.3	156.1	532.1	2,248.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/	1-4 2/	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
White, both sexes												
2004.....	85.4	*	*	*	0.1	1.0	6.0	20.5	58.1	179.1	574.2	1,929.2
2003.....	90.2	*	*	*	0.1	1.0	6.2	20.9	62.2	192.6	604.6	2,079.5
2002.....	92.8	*	*	*	0.1	0.9	6.3	21.9	65.1	203.2	629.1	2,143.3
2001.....	94.8	*	*	*	0.1	1.0	5.9	21.5	67.9	213.4	646.0	2,233.8
2000.....	97.4	*	*	*	0.1	0.9	6.1	22.1	71.2	222.2	662.0	2,352.7
1999.....	100.4	*	*	*	0.1	1.0	6.1	23.4	73.5	233.2	689.7	2,416.1
White, male												
2004.....	87.0	*	*	*	0.2	1.5	9.1	31.6	86.4	252.0	775.6	2,200.2
2003.....	91.3	*	*	*	0.2	1.5	9.4	32.5	93.3	273.1	817.4	2,338.4
2002.....	93.9	*	*	*	0.1	1.4	9.8	34.9	97.3	288.3	852.7	2,403.6
2001.....	94.8	*	*	*	0.2	1.6	9.2	33.7	101.1	301.9	870.7	2,493.6
2000.....	97.5	*	*	*	0.1	1.4	9.6	35.2	106.4	317.8	895.9	2,654.4
1999.....	99.9	*	*	*	0.2	1.4	9.5	36.9	108.3	333.0	931.7	2,695.9
White, female												
2004.....	83.8	*	*	*	*	0.5	2.9	9.4	31.3	116.7	436.9	1,807.0
2003.....	89.1	*	*	*	*	0.5	2.9	9.4	32.9	124.0	460.7	1,964.9
2002.....	91.7	*	*	*	*	0.4	2.8	9.1	34.7	131.1	479.1	2,030.2
2001.....	94.9	*	*	*	*	0.4	2.6	9.6	36.7	138.6	496.8	2,125.0
2000.....	97.3	*	*	*	*	0.4	2.5	9.3	38.3	141.9	508.1	2,230.1
1999.....	100.8	*	*	*	*	0.5	2.6	10.1	40.9	150.1	531.1	2,303.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Black, both sexes												
2004.....	52.9	*	*	*	0.3	2.0	9.3	33.5	94.3	239.8	587.8	1,514.0
2003.....	56.3	*	*	*	0.3	1.8	9.3	34.9	101.8	261.5	631.6	1,652.1
2002.....	56.2	*	*	*	*	1.4	9.4	36.4	106.7	255.1	637.0	1,665.1
2001.....	58.0	*	*	*	0.4	2.0	9.5	36.3	106.7	271.4	666.3	1,756.7
2000.....	58.1	*	*	*	*	1.8	10.2	36.1	112.0	282.3	662.9	1,740.5
1999.....	59.3	*	*	*	0.3	1.5	10.4	38.3	114.2	291.8	666.2	1,801.2
Black, male												
2004.....	52.6	*	*	*	*	2.8	12.9	48.1	129.7	317.0	733.6	1,536.8
2003.....	55.0	*	*	*	*	2.4	13.0	49.1	137.6	346.5	797.1	1,563.9
2002.....	54.5	*	*	*	*	1.8	13.1	50.8	150.1	334.3	756.5	1,655.3
2001.....	56.7	*	*	*	*	3.2	13.7	52.1	146.2	360.1	825.4	1,759.8
2000.....	55.8	*	*	*	*	2.6	13.9	50.2	152.7	356.1	806.0	1,744.8
1999.....	57.3	*	*	*	*	2.3	15.0	54.9	157.9	371.2	792.6	1,845.5
Black, female												
2004.....	53.2	*	*	*	*	1.2	6.2	21.0	65.8	184.6	505.4	1,505.0
2003.....	57.5	*	*	*	*	1.2	6.0	22.6	72.9	201.1	538.0	1,686.4
2002.....	57.7	*	*	*	*	1.1	6.1	24.0	71.9	198.9	569.4	1,668.8
2001.....	59.2	*	*	*	*	1.0	5.9	22.8	75.0	208.7	576.9	1,755.5
2000.....	60.3	*	*	*	*	1.1	6.9	24.1	79.2	230.3	582.3	1,738.9
1999.....	61.2	*	*	*	*	0.8	6.4	24.1	79.1	235.8	595.4	1,784.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other heart diseases (I26-I51)

All races 3/, both sexes

2004.....	56.6	10.0	1.1	0.5	2.1	4.3	9.7	21.5	45.2	118.6	368.9	1,373.9
2003.....	58.5	10.4	1.1	0.6	2.2	4.5	10.3	21.4	48.0	126.8	382.8	1,427.2
2002.....	58.8	11.7	1.0	0.6	2.1	4.5	10.3	21.7	48.7	129.6	390.9	1,443.5
2001.....	58.6	11.1	1.4	0.6	2.0	4.4	9.9	21.6	49.5	131.4	389.5	1,475.5
2000.....	58.8	12.0	1.1	0.6	2.2	4.4	9.9	21.5	50.9	134.2	391.3	1,509.4
1999.....	59.4	12.9	1.2	0.6	2.3	4.4	10.3	21.6	52.9	138.8	398.5	1,512.9

All races, male

2004.....	50.6	10.5	1.1	0.5	2.6	5.3	12.0	27.7	57.7	142.8	420.8	1,377.7
2003.....	52.3	11.4	1.0	0.7	2.6	5.3	12.7	27.2	60.7	154.2	440.0	1,431.3
2002.....	52.4	12.1	1.0	0.6	2.7	5.6	12.8	28.2	60.8	156.5	448.2	1,448.6
2001.....	51.8	10.8	1.5	0.6	2.4	5.2	12.4	27.6	61.7	158.7	452.0	1,473.4
2000.....	52.2	12.2	1.2	0.7	2.6	5.2	12.2	27.8	64.7	164.0	451.0	1,549.1
1999.....	53.1	13.1	1.3	0.7	2.7	5.3	12.9	27.7	67.5	172.7	459.7	1,548.9

All races, female

2004.....	62.4	9.4	1.1	0.6	1.5	3.3	7.4	15.5	33.6	98.4	334.0	1,372.1
2003.....	64.5	9.3	1.2	0.5	1.7	3.7	7.9	15.7	36.3	103.8	344.5	1,425.4
2002.....	65.0	11.4	1.0	0.5	1.4	3.4	7.8	15.5	37.5	107.2	352.8	1,441.3
2001.....	65.2	11.5	1.3	0.6	1.5	3.6	7.5	15.9	38.3	108.7	348.4	1,476.4
2000.....	65.1	11.8	0.9	0.5	1.8	3.5	7.5	15.3	38.2	109.6	352.3	1,493.2
1999.....	65.5	12.7	1.0	0.6	1.8	3.6	7.7	15.6	39.4	111.1	358.7	1,498.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other heart diseases (I26-I51)

White, both sexes

2004.....	60.4	8.7	0.9	0.5	1.8	3.5	8.1	18.2	40.6	113.4	368.6	1,408.8
2003.....	62.2	8.2	0.9	0.5	1.8	3.6	8.6	18.1	43.1	120.3	383.4	1,461.4
2002.....	62.6	10.0	0.9	0.5	1.7	3.6	8.7	18.4	43.9	124.2	390.3	1,475.5
2001.....	62.3	9.3	1.1	0.5	1.6	3.6	8.1	18.4	44.6	125.6	387.6	1,509.6
2000.....	62.5	10.6	1.0	0.5	1.7	3.6	8.1	18.2	46.1	128.1	389.6	1,541.6
1999.....	62.9	11.0	1.0	0.5	1.9	3.6	8.4	18.1	47.2	132.2	395.1	1,543.3

White, male

2004.....	53.0	9.1	0.9	0.5	2.2	4.3	10.3	23.9	52.1	137.0	419.2	1,407.9
2003.....	54.7	8.8	0.8	0.6	2.2	4.4	10.8	23.3	54.7	147.2	441.1	1,467.6
2002.....	54.9	10.5	1.0	0.5	2.3	4.5	11.1	24.3	55.3	150.4	448.3	1,482.1
2001.....	54.1	9.6	1.2	0.5	2.0	4.2	10.4	24.0	55.9	152.2	448.7	1,503.4
2000.....	54.6	10.8	1.2	0.6	2.1	4.3	10.3	23.9	59.2	157.4	449.5	1,580.5
1999.....	55.1	11.4	1.1	0.6	2.2	4.3	10.9	23.7	60.6	165.1	456.1	1,571.5

White, female

2004.....	67.6	8.3	0.8	0.5	1.3	2.6	5.9	12.7	29.8	93.1	334.0	1,409.2
2003.....	69.6	7.6	1.0	0.4	1.5	2.8	6.3	13.0	32.1	97.3	344.4	1,458.6
2002.....	70.2	9.6	0.8	0.5	1.1	2.6	6.4	12.6	33.1	101.9	351.4	1,472.6
2001.....	70.4	8.9	1.1	0.5	1.3	2.9	5.7	12.8	34.0	103.1	347.1	1,512.2
2000.....	70.2	10.4	0.7	0.4	1.4	2.8	6.0	12.5	33.9	103.4	350.1	1,525.8
1999.....	70.5	10.7	0.8	0.5	1.6	2.8	6.0	12.6	34.5	104.7	355.1	1,531.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other heart diseases (I26-I51)												
Black, both sexes												
2004.....	50.8	14.8	2.3	1.1	3.7	10.0	22.2	50.0	96.1	195.5	450.2	1,180.7
2003.....	52.5	19.6	2.0	1.0	4.4	10.9	23.6	49.9	102.1	215.7	449.2	1,238.1
2002.....	52.5	20.2	1.5	0.8	4.1	11.3	22.8	51.4	102.0	207.6	469.1	1,270.4
2001.....	52.8	20.1	2.9	1.0	4.2	10.5	23.7	51.1	103.5	213.0	480.6	1,277.7
2000.....	52.7	19.0	1.9	1.1	4.7	10.3	23.0	51.6	102.6	220.2	477.9	1,306.0
1999.....	54.4	22.3	2.0	1.0	4.5	10.5	24.7	52.4	113.3	229.9	495.2	1,302.8
Black, male												
2004.....	50.4	15.7	2.3	1.0	4.7	12.2	26.7	63.6	125.6	237.6	540.9	1,224.9
2003.....	51.5	23.2	2.2	1.1	5.7	12.4	28.6	63.8	132.3	264.0	523.5	1,220.9
2002.....	50.8	20.9	*	0.9	5.1	14.2	27.4	65.9	127.9	255.2	537.3	1,238.3
2001.....	51.7	17.7	3.2	0.9	5.3	12.8	28.3	62.5	132.1	258.5	585.9	1,349.1
2000.....	51.5	18.9	1.9	1.2	5.4	12.2	28.1	66.5	130.5	266.0	558.0	1,402.3
1999.....	54.3	21.8	2.1	1.1	5.8	12.7	30.1	67.3	147.3	287.8	580.2	1,454.1
Black, female												
2004.....	51.1	13.9	2.3	1.1	2.7	8.0	18.3	38.4	72.3	165.4	398.9	1,163.2
2003.....	53.4	15.8	1.8	0.8	3.2	9.4	19.1	38.0	77.8	181.3	407.1	1,244.8
2002.....	54.0	19.4	1.8	0.8	3.1	8.7	18.7	39.0	81.1	173.8	430.6	1,282.7
2001.....	53.8	22.7	2.5	1.0	3.0	8.5	19.6	41.3	80.5	180.9	421.5	1,251.2
2000.....	53.9	19.2	2.0	1.0	4.1	8.7	18.6	38.7	80.1	187.9	432.8	1,270.2
1999.....	54.5	22.8	1.8	0.8	3.3	8.5	19.9	39.7	86.1	189.1	447.7	1,246.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and subacute endocarditis (I33)												
All races 3/, both sexes												
2004.....	0.4	*	*	*	*	0.1	0.2	0.5	0.6	1.3	2.5	3.2
2003.....	0.4	*	*	*	*	0.1	0.3	0.4	0.7	1.3	2.5	3.0
2002.....	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.3	2.4	2.6
2001.....	0.4	*	*	*	*	0.1	0.3	0.5	0.6	1.3	2.5	2.5
2000.....	0.4	*	*	*	*	0.1	0.3	0.5	0.7	1.2	2.4	2.8
1999.....	0.4	*	*	*	*	0.1	0.3	0.4	0.7	1.3	2.4	2.9
All races, male												
2004.....	0.5	*	*	*	*	0.1	0.2	0.6	0.7	1.4	3.1	4.8
2003.....	0.5	*	*	*	*	*	0.3	0.6	0.9	1.4	3.2	4.1
2002.....	0.5	*	*	*	*	0.2	0.3	0.6	0.7	1.6	3.0	3.5
2001.....	0.5	*	*	*	*	*	0.3	0.7	0.8	1.6	3.5	3.8
2000.....	0.5	*	*	*	*	0.1	0.3	0.6	0.8	1.5	3.5	4.2
1999.....	0.5	*	*	*	*	0.1	0.3	0.6	0.9	1.4	3.2	4.7
All races, female												
2004.....	0.4	*	*	*	*	*	0.1	0.3	0.4	1.1	2.0	2.5
2003.....	0.4	*	*	*	*	*	0.2	0.3	0.6	1.1	2.0	2.5
2002.....	0.3	*	*	*	*	*	0.1	0.3	0.6	1.1	2.0	2.2
2001.....	0.4	*	*	*	*	*	0.2	0.3	0.5	1.1	1.8	2.0
2000.....	0.4	*	*	*	*	*	0.2	0.3	0.5	1.0	1.8	2.3
1999.....	0.3	*	*	*	*	*	0.2	0.2	0.5	1.2	1.8	2.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and subacute endocarditis (I33)												
White, both sexes												
2004.....	0.4	*	*	*	*	0.1	0.2	0.3	0.5	1.2	2.4	3.2
2003.....	0.4	*	*	*	*	0.1	0.2	0.3	0.6	1.2	2.5	3.1
2002.....	0.4	*	*	*	*	0.1	0.2	0.3	0.5	1.2	2.5	2.7
2001.....	0.4	*	*	*	*	0.1	0.2	0.4	0.5	1.2	2.4	2.6
2000.....	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.2	2.4	2.7
1999.....	0.4	*	*	*	*	0.1	0.2	0.3	0.6	1.2	2.4	3.0
White, male												
2004.....	0.5	*	*	*	*	*	0.3	0.4	0.7	1.4	3.1	4.9
2003.....	0.5	*	*	*	*	*	0.3	0.5	0.8	1.3	3.3	4.4
2002.....	0.5	*	*	*	*	0.2	0.2	0.5	0.6	1.5	3.2	3.6
2001.....	0.5	*	*	*	*	*	0.2	0.5	0.7	1.3	3.5	3.9
2000.....	0.5	*	*	*	*	*	0.3	0.5	0.8	1.4	3.5	4.2
1999.....	0.5	*	*	*	*	*	0.2	0.4	0.7	1.4	3.4	5.1
White, female												
2004.....	0.3	*	*	*	*	*	*	0.2	0.4	1.1	1.9	2.4
2003.....	0.4	*	*	*	*	*	0.2	0.2	0.5	1.0	1.9	2.6
2002.....	0.3	*	*	*	*	*	0.1	0.2	0.5	1.0	2.0	2.3
2001.....	0.3	*	*	*	*	*	0.2	0.2	0.4	1.1	1.7	2.1
2000.....	0.3	*	*	*	*	*	0.2	0.2	0.4	1.0	1.7	2.1
1999.....	0.3	*	*	*	*	*	0.1	0.2	0.4	1.1	1.8	2.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute and subacute endocarditis (I33)

Black, both sexes

2004.....	0.5	*	*	*	*	*	*	1.4	1.2	2.1	2.7	*
2003.....	0.6	*	*	*	*	*	0.6	1.3	1.9	2.2	3.0	*
2002.....	0.5	*	*	*	*	*	0.4	1.2	1.6	2.3	*	*
2001.....	0.6	*	*	*	*	*	0.7	1.3	1.6	2.6	4.1	*
2000.....	0.6	*	*	*	*	*	0.5	1.3	1.6	1.9	2.8	*
1999.....	0.6	*	*	*	*	*	0.7	1.2	2.0	2.6	*	*

Black, male

2004.....	0.6	*	*	*	*	*	*	1.8	*	*	*	*
2003.....	0.7	*	*	*	*	*	*	2.0	2.2	*	*	*
2002.....	0.6	*	*	*	*	*	*	1.5	1.9	2.9	*	*
2001.....	0.8	*	*	*	*	*	0.7	1.9	2.1	3.8	*	*
2000.....	0.6	*	*	*	*	*	0.8	1.8	*	*	*	*
1999.....	0.7	*	*	*	*	*	1.0	1.9	2.5	*	*	*

Black, female

2004.....	0.5	*	*	*	*	*	*	1.0	*	2.0	*	*
2003.....	0.5	*	*	*	*	*	*	*	1.7	2.0	*	*
2002.....	0.5	*	*	*	*	*	*	1.0	*	*	*	*
2001.....	0.5	*	*	*	*	*	0.7	*	*	*	3.3	*
2000.....	0.5	*	*	*	*	*	*	1.0	1.6	*	*	*
1999.....	0.5	*	*	*	*	*	*	*	1.6	2.8	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
All races 3/, both sexes												
2004.....	0.3	0.5	0.2	*	0.1	0.1	0.2	0.3	0.4	0.6	1.2	1.8
2003.....	0.3	0.8	0.1	0.1	0.1	0.1	0.2	0.3	0.4	0.8	1.2	1.6
2002.....	0.3	0.6	0.1	0.1	0.1	0.1	0.2	0.3	0.4	0.8	1.2	2.1
2001.....	0.3	0.8	0.2	0.1	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.9
2000.....	0.3	0.5	0.2	0.0	0.1	0.1	0.2	0.2	0.4	0.7	1.2	1.8
1999.....	0.3	0.7	0.2	0.1	0.1	0.1	0.3	0.3	0.5	0.8	1.3	2.0
All races, male												
2004.....	0.3	*	*	*	*	*	0.2	0.3	0.5	0.7	1.3	2.3
2003.....	0.3	*	*	*	0.2	0.1	0.2	0.4	0.4	1.0	1.3	1.5
2002.....	0.3	*	*	*	0.1	0.2	0.2	0.4	0.5	0.9	1.2	2.4
2001.....	0.3	*	*	*	0.1	0.2	0.2	0.3	0.5	0.7	1.2	1.8
2000.....	0.3	*	*	*	0.1	0.1	0.3	0.2	0.4	0.8	1.5	2.1
1999.....	0.3	*	*	*	0.1	0.2	0.3	0.3	0.6	0.8	1.4	2.5
All races, female												
2004.....	0.3	*	*	*	*	0.1	0.1	0.2	0.3	0.5	1.2	1.6
2003.....	0.3	*	*	*	0.1	*	0.2	0.2	0.3	0.7	1.1	1.7
2002.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.7	1.3	2.0
2001.....	0.3	*	*	*	*	*	0.2	0.3	0.3	0.6	1.2	2.0
2000.....	0.3	*	*	*	*	0.1	0.1	0.2	0.3	0.6	1.0	1.7
1999.....	0.3	*	*	*	*	*	0.2	0.3	0.4	0.8	1.3	1.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

White, both sexes

2004.....	0.2	*	*	*	*	0.1	0.1	0.2	0.4	0.6	1.2	1.8
2003.....	0.3	*	*	*	0.1	0.1	0.2	0.2	0.4	0.8	1.2	1.7
2002.....	0.3	*	*	*	0.1	0.1	0.2	0.2	0.4	0.8	1.1	2.2
2001.....	0.3	*	*	*	0.1	0.1	0.2	0.2	0.4	0.7	1.2	1.9
2000.....	0.3	*	*	*	0.1	0.1	0.2	0.2	0.3	0.7	1.2	1.8
1999.....	0.3	*	*	*	0.1	0.1	0.2	0.3	0.5	0.8	1.3	2.0

White, male

2004.....	0.3	*	*	*	*	*	0.1	0.3	0.5	0.7	1.3	2.2
2003.....	0.3	*	*	*	0.1	*	0.2	0.3	0.4	1.1	1.3	1.7
2002.....	0.3	*	*	*	*	0.1	0.2	0.3	0.5	0.9	1.1	2.5
2001.....	0.3	*	*	*	*	0.2	0.2	0.3	0.5	0.7	1.2	2.0
2000.....	0.3	*	*	*	*	*	0.2	0.2	0.4	0.7	1.5	2.1
1999.....	0.3	*	*	*	*	0.1	0.3	0.3	0.6	0.8	1.4	2.5

White, female

2004.....	0.2	*	*	*	*	*	*	0.2	0.3	0.6	1.1	1.6
2003.....	0.3	*	*	*	*	*	*	0.2	0.3	0.7	1.1	1.7
2002.....	0.3	*	*	*	*	*	*	0.2	0.3	0.7	1.2	2.0
2001.....	0.3	*	*	*	*	*	0.2	0.2	0.3	0.6	1.3	1.9
2000.....	0.2	*	*	*	*	*	*	0.2	0.3	0.6	1.0	1.6
1999.....	0.3	*	*	*	*	*	0.2	0.3	0.4	0.7	1.2	1.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Heart failure (I50)												
All races 3/, both sexes												
2004.....	19.5	0.7	*	*	0.1	0.2	0.7	2.5	8.4	32.2	131.2	621.2
2003.....	19.8	0.9	*	*	0.1	0.2	0.7	2.4	8.8	33.5	133.2	641.1
2002.....	19.6	0.6	*	*	0.1	0.3	0.6	2.5	8.8	33.8	132.8	645.0
2001.....	20.0	0.6	*	*	0.1	0.2	0.7	2.4	9.4	34.3	136.3	674.6
2000.....	19.8	0.8	*	0.1	0.1	0.2	0.6	2.2	9.1	34.8	136.1	684.9
1999.....	19.7	*	*	*	0.1	0.2	0.6	2.1	8.9	35.4	137.2	681.9
All races, male												
2004.....	15.4	*	*	*	*	0.3	0.8	3.0	10.1	37.5	147.0	605.5
2003.....	15.7	1.4	*	*	0.1	0.3	0.9	2.9	10.8	39.7	148.6	629.8
2002.....	15.3	*	*	*	*	0.3	0.8	3.1	10.5	39.2	149.0	621.4
2001.....	15.5	*	*	*	*	0.3	0.8	3.1	11.1	40.3	153.8	646.4
2000.....	15.3	1.1	*	*	0.1	0.2	0.7	2.7	10.9	41.1	153.7	672.2
1999.....	15.2	*	*	*	0.1	0.2	0.7	2.7	10.8	42.5	152.0	667.0
All races, female												
2004.....	23.4	*	*	*	*	0.2	0.6	2.0	6.8	27.8	120.5	628.2
2003.....	23.7	*	*	*	*	0.2	0.5	1.9	7.0	28.3	122.9	646.1
2002.....	23.7	*	*	*	*	0.2	0.5	1.8	7.2	29.2	122.0	655.3
2001.....	24.3	*	*	*	*	0.2	0.6	1.8	7.8	29.3	124.7	686.4
2000.....	24.1	*	*	*	*	0.1	0.5	1.6	7.5	29.6	124.6	690.0
1999.....	24.0	*	*	*	*	0.2	0.5	1.6	7.1	29.6	127.6	687.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Heart failure (I50)												
White, both sexes												
2004.....	21.5	0.6	*	*	0.1	0.2	0.5	1.9	7.4	31.4	132.4	641.5
2003.....	21.8	0.6	*	*	0.1	0.2	0.5	1.9	7.8	32.2	134.3	660.4
2002.....	21.6	*	*	*	*	0.2	0.5	1.9	8.0	32.5	133.3	664.5
2001.....	22.1	*	*	*	*	0.2	0.5	2.0	8.5	33.0	137.0	695.1
2000.....	21.8	*	*	*	*	0.1	0.4	1.8	8.2	33.4	136.7	703.7
1999.....	21.6	*	*	*	*	0.1	0.5	1.7	8.0	34.1	136.7	699.2
White, male												
2004.....	16.8	*	*	*	*	0.2	0.7	2.3	8.8	36.8	148.2	625.5
2003.....	17.1	*	*	*	*	0.2	0.6	2.3	9.6	38.6	150.4	650.8
2002.....	16.7	*	*	*	*	0.2	0.6	2.5	9.5	37.7	150.3	641.9
2001.....	16.9	*	*	*	*	0.2	0.6	2.6	10.3	39.0	154.6	666.5
2000.....	16.7	*	*	*	*	0.1	0.6	2.2	10.0	40.0	154.6	690.0
1999.....	16.4	*	*	*	*	*	0.6	2.1	9.8	41.3	152.1	680.0
White, female												
2004.....	26.1	*	*	*	*	0.1	0.4	1.5	6.0	26.8	121.7	648.7
2003.....	26.3	*	*	*	*	0.1	0.3	1.4	6.1	26.8	123.4	664.7
2002.....	26.4	*	*	*	*	0.1	0.3	1.4	6.5	28.1	121.8	674.3
2001.....	27.1	*	*	*	*	0.2	0.4	1.4	6.9	28.0	125.3	707.1
2000.....	26.7	*	*	*	*	*	0.3	1.3	6.6	27.9	124.9	709.3
1999.....	26.6	*	*	*	*	*	0.3	1.2	6.3	28.0	126.6	707.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Heart failure (I50)												
Black, both sexes												
2004.....	13.5	*	*	*	*	0.7	1.9	7.5	19.4	48.2	150.4	506.5
2003.....	14.0	*	*	*	*	0.7	2.3	6.8	20.6	54.3	150.4	531.8
2002.....	13.7	*	*	*	*	0.8	1.8	7.2	18.6	53.2	156.9	528.0
2001.....	13.8	*	*	*	*	0.6	1.9	6.4	19.6	54.5	157.3	543.8
2000.....	13.6	*	*	*	*	0.4	1.6	5.7	18.6	55.4	154.4	558.4
1999.....	13.8	*	*	*	*	0.5	1.6	6.4	18.9	56.4	163.4	552.9
Black, male												
2004.....	11.8	*	*	*	*	0.9	2.2	9.3	25.0	55.7	176.2	503.7
2003.....	11.8	*	*	*	*	0.8	2.7	8.3	26.3	63.2	163.9	513.0
2002.....	11.5	*	*	*	*	1.0	2.2	9.0	22.9	64.7	174.4	479.2
2001.....	11.6	*	*	*	*	0.9	2.2	7.4	22.2	65.3	183.3	541.2
2000.....	11.2	*	*	*	*	*	1.7	7.1	21.4	61.9	173.5	581.2
1999.....	11.8	*	*	*	*	*	1.9	8.4	22.9	65.2	179.2	603.5
Black, female												
2004.....	15.1	*	*	*	*	*	1.7	5.9	15.0	42.8	135.8	507.6
2003.....	15.9	*	*	*	*	*	1.9	5.6	16.0	48.0	142.7	539.2
2002.....	15.8	*	*	*	*	*	1.5	5.6	15.1	45.1	147.0	546.7
2001.....	15.8	*	*	*	*	*	1.7	5.5	17.6	46.9	142.6	544.8
2000.....	15.7	*	*	*	*	*	1.5	4.5	16.4	50.9	143.7	549.9
1999.....	15.7	*	*	*	*	*	1.4	4.7	15.7	50.2	154.6	534.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
All races 3/, both sexes												
2004.....	36.5	8.7	0.9	0.5	1.9	3.9	8.7	18.3	35.8	84.5	234.1	747.7
2003.....	38.0	8.6	0.9	0.5	1.9	4.1	9.1	18.2	38.1	91.2	246.0	781.5
2002.....	38.5	10.5	0.8	0.5	1.9	4.0	9.3	18.6	38.8	93.7	254.4	793.7
2001.....	37.9	9.7	1.1	0.5	1.8	4.0	8.8	18.5	39.1	95.1	249.5	796.5
2000.....	38.3	10.6	0.9	0.5	2.0	4.0	8.8	18.6	40.8	97.4	251.6	820.0
1999.....	39.0	11.9	0.9	0.5	2.1	4.1	9.2	18.7	42.8	101.3	257.6	826.1
All races, male												
2004.....	34.5	9.2	0.9	0.5	2.4	4.8	10.7	23.8	46.3	103.1	269.4	765.2
2003.....	35.8	9.2	0.9	0.6	2.3	4.8	11.3	23.3	48.5	112.1	287.0	795.9
2002.....	36.3	10.9	0.8	0.5	2.5	5.0	11.5	24.1	49.0	114.8	294.9	821.3
2001.....	35.6	9.1	1.2	0.5	2.2	4.7	11.1	23.6	49.4	116.1	293.4	821.3
2000.....	36.1	10.5	1.0	0.6	2.3	4.8	10.9	24.3	52.6	120.7	292.4	870.7
1999.....	37.1	12.1	1.0	0.6	2.5	4.8	11.6	24.2	55.3	127.9	303.1	874.7
All races, female												
2004.....	38.5	8.2	0.8	0.5	1.3	2.9	6.6	13.0	26.0	68.9	210.3	739.9
2003.....	40.1	7.9	1.0	0.4	1.5	3.4	7.0	13.3	28.4	73.7	218.6	775.1
2002.....	40.6	10.0	0.8	0.4	1.3	3.1	7.1	13.2	29.3	76.2	227.6	781.7
2001.....	40.2	10.4	1.0	0.5	1.3	3.2	6.5	13.5	29.6	77.7	220.7	786.1
2000.....	40.4	10.7	0.7	0.4	1.6	3.2	6.8	13.1	29.9	78.3	225.0	799.3
1999.....	40.9	11.6	0.8	0.5	1.7	3.3	6.9	13.5	31.3	79.6	228.0	806.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
White, both sexes												
2004.....	38.2	7.7	0.7	0.4	1.6	3.1	7.3	15.8	32.4	80.1	232.6	762.4
2003.....	39.8	7.1	0.7	0.4	1.6	3.3	7.7	15.6	34.3	86.0	245.4	796.1
2002.....	40.4	9.1	0.7	0.4	1.6	3.2	7.9	15.9	34.9	89.7	253.4	806.1
2001.....	39.6	8.2	0.9	0.4	1.5	3.2	7.2	15.7	35.2	90.7	246.9	809.9
2000.....	40.0	9.6	0.8	0.4	1.6	3.3	7.3	15.8	37.0	92.8	249.3	833.4
1999.....	40.6	10.3	0.8	0.5	1.8	3.3	7.6	15.9	38.1	96.1	254.7	839.0
White, male												
2004.....	35.5	8.1	0.7	0.4	2.0	3.9	9.2	20.9	42.1	98.2	266.7	775.3
2003.....	36.9	7.4	0.7	0.5	1.9	4.0	9.7	20.2	43.9	106.3	286.1	810.7
2002.....	37.5	9.5	0.8	0.5	2.1	4.0	10.0	21.0	44.6	110.3	293.7	834.2
2001.....	36.5	8.2	0.9	0.5	1.8	3.8	9.4	20.6	44.6	111.2	289.4	831.0
2000.....	37.1	9.4	0.9	0.5	1.9	4.0	9.2	20.9	48.0	115.3	290.0	884.2
1999.....	37.9	10.7	1.0	0.5	2.0	4.0	9.8	20.9	49.4	121.6	299.2	883.9
White, female												
2004.....	40.9	7.3	0.6	0.4	1.1	2.3	5.3	10.7	23.1	64.6	209.3	756.5
2003.....	42.6	6.8	0.8	0.3	1.3	2.6	5.7	11.1	25.2	68.8	217.9	789.7
2002.....	43.2	8.7	0.6	0.4	1.0	2.3	5.8	10.9	25.8	72.2	226.4	793.9
2001.....	42.6	8.2	0.8	0.4	1.1	2.6	5.0	11.0	26.4	73.3	218.8	801.1
2000.....	42.8	9.7	0.7	0.3	1.2	2.5	5.4	10.8	26.6	73.9	222.6	812.7
1999.....	43.2	10.0	0.7	0.4	1.5	2.6	5.4	11.0	27.4	74.9	225.5	821.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
Black, both sexes												
2004.....	36.3	12.4	1.8	0.9	3.4	9.0	19.6	40.7	74.9	144.2	294.9	668.3
2003.....	37.5	15.0	1.6	0.8	3.9	9.7	20.3	41.3	79.2	158.2	294.5	703.7
2002.....	37.8	17.2	1.1	0.7	3.8	10.1	20.3	42.4	81.0	150.9	308.2	738.5
2001.....	38.0	17.2	2.3	0.9	3.7	9.6	20.7	42.8	81.5	155.1	318.1	730.2
2000.....	38.2	15.6	1.4	0.9	4.3	9.4	20.5	44.0	81.7	161.8	319.6	742.0
1999.....	39.6	20.2	1.4	0.8	4.2	9.6	21.8	44.3	91.7	169.8	328.2	746.1
Black, male												
2004.....	37.5	13.3	1.8	0.8	4.4	10.8	23.7	51.7	98.6	177.9	359.4	715.3
2003.....	38.6	17.8	1.8	0.9	5.0	10.9	24.8	52.9	103.3	197.6	355.1	707.9
2002.....	38.3	18.9	*	0.8	4.7	12.7	24.3	54.5	102.0	186.7	359.2	753.7
2001.....	38.9	15.0	2.6	0.8	4.8	11.4	25.1	52.4	106.8	188.2	395.5	805.6
2000.....	39.2	14.7	1.6	1.0	4.9	11.2	25.0	57.2	106.9	200.2	379.7	816.5
1999.....	41.5	19.8	*	1.1	5.4	11.5	26.6	56.4	121.0	218.8	398.1	849.5
Black, female												
2004.....	35.1	11.4	1.7	1.0	2.4	7.3	15.9	31.2	55.8	120.2	258.4	649.7
2003.....	36.5	12.1	*	0.7	2.7	8.5	16.2	31.4	59.7	130.2	260.2	702.0
2002.....	37.4	15.4	*	*	2.9	7.7	16.7	32.0	64.1	125.5	279.3	732.8
2001.....	37.2	19.5	2.0	0.9	2.6	7.9	16.7	34.5	61.1	131.7	274.6	702.2
2000.....	37.3	16.5	*	0.8	3.7	7.8	16.5	32.7	61.4	134.8	285.7	714.3
1999.....	37.9	20.5	*	*	3.0	7.9	17.5	34.0	68.2	135.1	289.0	707.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

All races 3/, both sexes

2004.....	7.9	*	*	*	0.1	0.3	0.8	2.7	6.3	17.1	52.6	198.5
2003.....	7.5	*	*	*	0.1	0.2	0.8	2.5	6.3	16.9	51.7	188.9
2002.....	7.0	*	*	*	0.1	0.2	0.8	2.3	5.7	16.0	48.2	180.4
2001.....	6.8	*	*	*	0.1	0.3	0.7	2.4	5.8	15.5	47.7	171.9
2000.....	6.4	*	*	*	*	0.2	0.8	2.3	5.9	15.1	45.5	162.9
1999.....	6.1	*	*	*	*	0.2	0.7	2.2	5.5	15.2	43.6	152.1

All races, male

2004.....	6.1	*	*	*	*	0.3	1.0	3.5	7.4	18.2	53.3	166.0
2003.....	5.7	*	*	*	*	0.3	1.0	3.3	7.4	18.4	48.9	153.5
2002.....	5.4	*	*	*	*	0.2	1.0	2.8	6.6	17.1	47.3	153.4
2001.....	5.2	*	*	*	*	0.3	0.9	3.0	6.5	16.6	45.8	150.8
2000.....	5.0	*	*	*	*	0.2	1.0	3.0	7.0	15.8	45.2	142.4
1999.....	4.7	*	*	*	*	0.2	0.9	2.7	6.4	16.5	42.9	132.4

All races, female

2004.....	9.5	*	*	*	*	0.2	0.6	2.0	5.3	16.1	52.1	213.2
2003.....	9.3	*	*	*	*	0.2	0.7	1.8	5.3	15.7	53.5	204.6
2002.....	8.6	*	*	*	*	0.2	0.6	1.8	4.9	15.1	48.8	192.1
2001.....	8.3	*	*	*	*	0.2	0.6	1.8	5.2	14.5	48.9	180.8
2000.....	7.8	*	*	*	*	0.2	0.6	1.7	4.9	14.6	45.7	171.3
1999.....	7.4	*	*	*	*	0.2	0.5	1.7	4.7	14.1	43.9	160.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

White, both sexes

2004.....	7.4	*	*	*	*	0.1	0.5	1.7	4.4	13.5	46.7	188.1
2003.....	7.2	*	*	*	*	0.1	0.5	1.7	4.4	13.0	46.2	181.0
2002.....	6.6	*	*	*	*	0.1	0.4	1.5	4.1	12.8	42.6	170.4
2001.....	6.3	*	*	*	*	0.1	0.4	1.4	4.0	12.1	42.3	163.3
2000.....	6.0	*	*	*	*	0.1	0.4	1.4	4.1	11.8	40.4	153.2
1999.....	5.6	*	*	*	*	0.1	0.4	1.3	3.7	12.1	37.9	142.0

White, male

2004.....	5.6	*	*	*	*	0.2	0.6	2.3	5.3	14.6	47.1	159.4
2003.....	5.2	*	*	*	*	0.1	0.7	2.2	5.2	14.5	43.8	145.3
2002.....	4.9	*	*	*	*	0.1	0.5	1.9	4.9	13.8	41.5	145.2
2001.....	4.7	*	*	*	*	0.2	0.5	1.8	4.5	13.2	40.4	142.2
2000.....	4.5	*	*	*	*	0.1	0.6	1.9	5.0	12.6	40.3	132.5
1999.....	4.2	*	*	*	*	0.2	0.5	1.7	4.5	13.4	37.3	121.0

White, female

2004.....	9.2	*	*	*	*	*	0.3	1.2	3.4	12.5	46.5	201.1
2003.....	9.1	*	*	*	*	0.1	0.4	1.2	3.6	11.7	47.8	196.8
2002.....	8.3	*	*	*	*	*	0.3	1.0	3.3	11.9	43.4	181.3
2001.....	8.0	*	*	*	*	*	0.3	1.0	3.5	11.3	43.6	172.2
2000.....	7.4	*	*	*	*	*	0.3	0.9	3.2	11.2	40.4	161.6
1999.....	7.0	*	*	*	*	*	0.2	1.0	3.0	11.0	38.3	150.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year	2/	years	years	years	years	years	years	years	years	and over

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

Black, both sexes

2004.....	12.3	*	*	*	*	1.0	3.1	10.2	23.9	51.9	122.6	335.6
2003.....	11.8	*	*	*	*	1.0	3.1	9.3	24.1	54.4	119.5	301.1
2002.....	11.1	*	*	*	*	0.8	3.3	9.0	21.0	47.6	118.0	308.8
2001.....	10.9	*	*	*	*	1.0	3.1	9.9	22.0	47.4	112.6	288.0
2000.....	10.6	*	*	*	*	0.6	3.1	9.7	22.9	48.7	107.1	283.4
1999.....	10.3	*	*	*	*	0.7	3.1	9.1	21.5	46.7	111.7	269.4

Black, male

2004.....	10.7	*	*	*	*	1.1	3.9	12.9	28.3	56.8	137.0	263.3
2003.....	10.4	*	*	*	*	1.3	3.4	12.7	29.3	60.5	119.6	274.2
2002.....	9.6	*	*	*	*	0.9	4.1	11.0	23.5	52.0	127.4	274.5
2001.....	9.7	*	*	*	*	1.1	3.8	12.9	25.5	54.4	118.3	275.7
2000.....	9.1	*	*	*	*	*	3.7	11.9	27.8	50.5	109.0	269.9
1999.....	9.1	*	*	*	*	*	3.7	11.6	24.6	51.3	118.7	266.0

Black, female

2004.....	13.8	*	*	*	*	0.9	2.4	7.8	20.3	48.4	114.5	364.2
2003.....	13.0	*	*	*	*	*	2.7	6.4	20.0	50.1	119.4	311.5
2002.....	12.5	*	*	*	*	0.7	2.5	7.2	19.0	44.5	112.7	322.0
2001.....	11.9	*	*	*	*	0.9	2.4	7.4	19.2	42.5	109.4	292.6
2000.....	12.0	*	*	*	*	*	2.6	7.7	18.9	47.5	106.0	288.5
1999.....	11.4	*	*	*	*	0.7	2.6	7.0	19.0	43.4	107.7	270.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebrovascular diseases (I60-I69)												
All races 3/, both sexes												
2004.....	51.1	3.1	0.3	0.2	0.5	1.4	5.4	14.9	34.3	107.8	386.2	1,245.9
2003.....	54.2	2.5	0.3	0.2	0.5	1.5	5.5	15.0	35.6	112.9	410.7	1,370.1
2002.....	56.4	2.9	0.3	0.2	0.4	1.4	5.4	15.1	37.2	120.3	431.0	1,445.9
2001.....	57.4	2.7	0.4	0.2	0.5	1.5	5.5	15.1	38.0	123.4	443.9	1,500.2
2000.....	59.6	3.3	0.3	0.2	0.5	1.5	5.8	16.0	41.0	128.6	461.3	1,589.2
1999.....	60.0	2.7	0.3	0.2	0.5	1.4	5.7	15.2	40.6	130.8	469.8	1,614.8
All races, male												
2004.....	40.7	3.4	0.3	0.2	0.5	1.4	5.6	16.7	39.5	121.1	402.9	1,118.1
2003.....	42.9	2.8	0.3	0.2	0.5	1.6	5.8	16.7	40.8	127.8	431.4	1,236.0
2002.....	44.2	3.2	0.4	0.2	0.5	1.4	5.3	16.7	42.7	135.0	445.9	1,317.9
2001.....	45.2	3.1	0.3	0.2	0.5	1.6	5.7	16.7	43.4	140.4	467.3	1,380.2
2000.....	46.9	3.8	*	0.2	0.5	1.5	5.8	17.5	47.2	145.0	490.8	1,484.3
1999.....	47.1	3.3	0.3	0.2	0.5	1.5	5.8	16.7	46.5	147.3	500.4	1,512.8
All races, female												
2004.....	61.2	2.8	*	0.2	0.5	1.4	5.1	13.1	29.5	96.6	374.9	1,303.4
2003.....	65.1	2.2	0.3	0.1	0.5	1.4	5.3	13.4	30.9	100.5	396.8	1,429.4
2002.....	68.2	2.5	0.3	0.2	0.3	1.4	5.5	13.6	32.1	108.1	421.2	1,501.5
2001.....	69.2	2.3	0.4	0.2	0.5	1.5	5.4	13.6	32.9	109.3	428.6	1,550.4
2000.....	71.8	2.7	0.4	0.2	0.5	1.5	5.7	14.5	35.3	115.1	442.1	1,632.0
1999.....	72.3	2.1	0.3	0.2	0.5	1.4	5.6	13.8	35.1	117.2	449.8	1,656.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebrovascular diseases (I60-I69)												
White, both sexes												
2004.....	53.7	2.5	0.2	0.2	0.5	1.2	4.4	11.4	28.6	99.0	377.7	1,257.8
2003.....	57.0	2.0	0.2	0.2	0.5	1.2	4.2	11.4	29.4	103.7	403.0	1,382.2
2002.....	59.5	2.3	0.3	0.2	0.4	1.2	4.2	11.6	31.4	110.7	423.5	1,459.5
2001.....	60.5	1.9	0.3	0.2	0.4	1.3	4.4	11.4	31.7	113.0	436.4	1,511.2
2000.....	62.8	2.5	0.3	0.2	0.5	1.3	4.5	12.4	34.8	119.4	452.4	1,601.6
1999.....	63.3	2.3	0.3	0.2	0.4	1.3	4.4	11.7	34.0	121.1	461.6	1,631.6
White, male												
2004.....	41.8	2.8	*	0.2	0.5	1.2	4.7	12.8	32.4	110.8	393.7	1,129.3
2003.....	44.2	2.0	*	0.2	0.5	1.3	4.4	12.9	33.3	117.3	422.4	1,247.0
2002.....	45.7	2.6	0.4	0.2	0.5	1.2	4.2	12.9	35.6	123.8	437.5	1,327.4
2001.....	46.6	2.2	*	0.2	0.5	1.4	4.5	12.7	36.1	128.5	458.8	1,386.2
2000.....	48.4	3.0	*	0.2	0.5	1.4	4.5	13.6	39.7	133.8	480.0	1,490.7
1999.....	48.7	2.8	*	0.2	0.5	1.3	4.6	12.8	38.9	136.0	490.3	1,522.7
White, female												
2004.....	65.3	2.2	*	0.2	0.4	1.1	4.1	10.1	25.1	89.0	366.8	1,315.7
2003.....	69.5	2.0	*	0.1	0.4	1.2	4.0	10.0	25.8	92.1	389.9	1,442.1
2002.....	73.0	2.0	*	0.2	0.3	1.2	4.3	10.4	27.4	99.5	414.1	1,516.9
2001.....	74.0	1.6	0.4	0.2	0.4	1.3	4.2	10.2	27.6	99.9	421.6	1,563.5
2000.....	76.9	2.0	*	0.2	0.4	1.3	4.4	11.2	30.2	107.3	434.2	1,646.7
1999.....	77.6	1.9	*	0.1	0.4	1.2	4.3	10.7	29.4	108.6	442.8	1,675.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebrovascular diseases (I60-I69)												
Black, both sexes												
2004.....	46.9	6.0	*	*	0.8	3.1	12.3	39.0	83.9	195.3	511.7	1,211.2
2003.....	49.3	4.9	*	*	0.9	3.1	13.8	41.0	89.8	201.1	531.1	1,348.7
2002.....	50.0	5.5	*	0.4	0.6	3.1	12.8	40.7	88.0	215.1	552.3	1,383.9
2001.....	51.1	6.6	*	*	0.8	2.9	13.2	42.6	91.1	222.5	560.3	1,452.7
2000.....	52.5	7.5	*	*	0.7	2.9	14.5	43.4	93.8	222.9	588.8	1,530.0
1999.....	52.2	4.6	*	*	0.7	2.7	14.3	41.5	96.3	231.9	590.5	1,492.5
Black, male												
2004.....	41.5	6.8	*	*	1.0	2.9	12.8	44.8	107.4	235.2	551.0	1,061.0
2003.....	43.2	5.9	*	*	0.8	3.3	14.3	46.9	112.1	237.4	588.9	1,180.3
2002.....	43.5	5.8	*	*	0.7	2.9	12.2	46.5	110.3	262.9	587.8	1,252.2
2001.....	44.6	8.1	*	*	0.7	3.1	13.2	48.8	111.9	269.2	613.9	1,349.1
2000.....	46.1	8.6	*	*	*	2.7	14.7	49.5	115.4	268.5	659.2	1,458.8
1999.....	45.9	*	*	*	*	2.8	14.7	48.6	117.9	274.7	650.6	1,448.3
Black, female												
2004.....	51.9	*	*	*	*	3.3	11.8	33.9	65.0	166.8	489.5	1,270.7
2003.....	54.8	*	*	*	1.1	2.8	13.3	36.0	71.8	175.3	498.3	1,414.2
2002.....	55.8	*	*	*	*	3.3	13.4	35.7	70.1	181.2	532.2	1,434.3
2001.....	56.9	*	*	*	1.0	2.8	13.1	37.3	74.4	189.5	530.3	1,491.2
2000.....	58.3	*	*	*	0.8	3.1	14.4	38.1	76.4	190.9	549.2	1,556.5
1999.....	57.9	*	*	*	0.8	2.7	13.9	35.3	78.9	201.7	556.8	1,509.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Atherosclerosis (I70)

All races 3/, both sexes

2004.....	4.0	*	*	*	*	*	0.1	0.4	1.6	6.1	26.3	137.0
2003.....	4.5	*	*	*	*	*	0.1	0.5	1.7	6.6	29.8	154.6
2002.....	4.8	*	*	*	*	*	0.1	0.5	1.8	7.1	31.9	168.1
2001.....	4.9	*	*	*	*	*	0.1	0.5	1.9	7.3	33.1	179.1
2000.....	5.1	*	*	*	*	*	0.1	0.5	1.9	7.7	34.6	188.8
1999.....	5.4	*	*	*	*	*	0.1	0.5	2.0	8.0	35.9	202.0

All races, male

2004.....	3.1	*	*	*	*	*	0.1	0.5	2.0	7.3	29.7	128.2
2003.....	3.4	*	*	*	*	*	0.2	0.7	2.2	7.8	33.6	138.6
2002.....	3.7	*	*	*	*	*	0.1	0.6	2.4	8.8	35.5	156.6
2001.....	3.7	*	*	*	*	*	0.1	0.6	2.4	9.0	35.9	167.2
2000.....	3.9	*	*	*	*	*	0.2	0.6	2.5	9.6	39.7	176.5
1999.....	4.0	*	*	*	*	*	0.1	0.7	2.7	9.6	40.0	189.0

All races, female

2004.....	4.9	*	*	*	*	*	*	0.2	1.2	5.1	24.0	141.0
2003.....	5.5	*	*	*	*	*	*	0.3	1.2	5.6	27.3	161.7
2002.....	5.9	*	*	*	*	*	0.1	0.3	1.3	5.7	29.5	173.1
2001.....	6.2	*	*	*	*	*	*	0.3	1.4	5.9	31.2	184.1
2000.....	6.3	*	*	*	*	*	*	0.3	1.2	6.2	31.3	193.8
1999.....	6.7	*	*	*	*	*	*	0.3	1.4	6.7	33.2	207.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerosis (I70)												
White, both sexes												
2004.....	4.6	*	*	*	*	*	0.1	0.4	1.6	6.1	27.1	142.0
2003.....	5.0	*	*	*	*	*	0.1	0.4	1.6	6.5	30.3	159.5
2002.....	5.4	*	*	*	*	*	0.1	0.4	1.7	7.1	32.1	174.7
2001.....	5.6	*	*	*	*	*	0.1	0.5	1.8	7.3	33.8	185.9
2000.....	5.7	*	*	*	*	*	0.1	0.4	1.8	7.7	35.1	195.2
1999.....	6.0	*	*	*	*	*	0.1	0.5	2.0	8.0	36.4	207.7
White, male												
2004.....	3.5	*	*	*	*	*	*	0.5	1.9	7.4	30.4	134.4
2003.....	3.7	*	*	*	*	*	0.2	0.6	2.1	7.5	33.7	143.3
2002.....	4.0	*	*	*	*	*	0.1	0.5	2.3	8.8	35.6	163.2
2001.....	4.1	*	*	*	*	*	0.1	0.6	2.3	9.0	36.4	173.9
2000.....	4.3	*	*	*	*	*	0.2	0.5	2.5	9.7	40.3	181.4
1999.....	4.4	*	*	*	*	*	0.1	0.7	2.6	9.5	40.2	193.7
White, female												
2004.....	5.6	*	*	*	*	*	*	0.2	1.2	5.0	24.9	145.4
2003.....	6.3	*	*	*	*	*	*	0.3	1.1	5.6	28.0	166.7
2002.....	6.7	*	*	*	*	*	*	0.3	1.3	5.7	29.8	179.7
2001.....	7.0	*	*	*	*	*	*	0.3	1.3	5.9	32.1	190.9
2000.....	7.1	*	*	*	*	*	*	0.2	1.1	6.1	31.7	200.8
1999.....	7.6	*	*	*	*	*	*	0.3	1.4	6.8	34.0	213.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerosis (I70)												
Black, both sexes												
2004.....	2.2	*	*	*	*	*	*	0.5	2.3	7.4	23.9	102.9
2003.....	2.6	*	*	*	*	*	*	0.8	2.6	9.2	30.7	121.1
2002.....	2.7	*	*	*	*	*	*	1.0	2.9	8.0	36.5	117.0
2001.....	2.6	*	*	*	*	*	*	0.7	3.3	8.9	30.7	125.6
2000.....	2.9	*	*	*	*	*	*	1.1	3.2	8.9	35.2	138.3
1999.....	3.0	*	*	*	*	*	*	0.7	3.2	9.2	34.9	154.3
Black, male												
2004.....	1.7	*	*	*	*	*	*	*	3.0	8.6	30.1	78.5
2003.....	2.3	*	*	*	*	*	*	1.2	3.6	12.4	40.3	104.3
2002.....	2.3	*	*	*	*	*	*	1.4	4.2	10.1	44.2	110.4
2001.....	2.1	*	*	*	*	*	*	1.0	3.8	10.9	37.1	113.5
2000.....	2.4	*	*	*	*	*	*	1.4	3.9	11.0	40.3	147.6
1999.....	2.5	*	*	*	*	*	*	*	4.4	11.8	44.0	164.0
Black, female												
2004.....	2.6	*	*	*	*	*	*	*	1.7	6.5	20.5	112.6
2003.....	2.9	*	*	*	*	*	*	*	1.8	7.0	25.2	127.6
2002.....	3.1	*	*	*	*	*	*	*	1.9	6.5	32.2	119.5
2001.....	3.1	*	*	*	*	*	*	*	2.9	7.5	27.1	130.1
2000.....	3.3	*	*	*	*	*	*	*	2.7	7.5	32.4	134.8
1999.....	3.4	*	*	*	*	*	*	*	2.3	7.4	29.8	150.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Other diseases of circulatory system (I71-I78)

All races 3/, both sexes

2004.....	8.1	*	*	*	0.1	0.4	1.0	2.7	7.2	25.1	64.3	139.3
2003.....	8.5	*	*	*	0.2	0.4	1.0	2.9	8.1	26.9	67.6	146.0
2002.....	8.6	*	*	*	0.2	0.4	1.0	2.7	8.6	29.1	68.7	148.4
2001.....	8.9	*	*	*	0.2	0.4	1.1	2.6	8.7	29.2	71.8	158.2
2000.....	9.3	*	*	*	0.2	0.4	1.0	2.8	9.2	32.2	74.6	163.2
1999.....	9.2	*	*	*	0.1	0.4	1.1	2.7	9.3	31.5	75.8	165.2

All races, male

2004.....	8.5	*	*	*	0.2	0.6	1.4	3.8	9.9	33.4	82.8	166.5
2003.....	9.2	*	*	*	0.2	0.5	1.4	4.1	11.3	37.0	91.8	179.7
2002.....	9.4	*	*	*	0.2	0.5	1.3	3.7	11.9	39.8	92.3	187.4
2001.....	9.6	*	*	*	0.2	0.6	1.5	3.6	12.0	40.3	97.0	197.3
2000.....	10.2	*	*	*	0.2	0.5	1.2	3.8	13.1	45.4	102.9	214.9
1999.....	10.1	*	*	*	0.2	0.5	1.3	3.7	13.3	44.4	104.2	207.5

All races, female

2004.....	7.6	*	*	*	*	0.3	0.7	1.6	4.7	18.1	51.8	127.0
2003.....	7.8	*	*	*	0.1	0.2	0.7	1.8	5.2	18.4	51.4	131.1
2002.....	7.9	*	*	*	0.1	0.3	0.7	1.7	5.5	20.1	53.0	131.5
2001.....	8.2	*	*	*	0.1	0.3	0.7	1.7	5.6	19.9	55.3	141.8
2000.....	8.3	*	*	*	0.1	0.3	0.8	1.8	5.6	21.4	56.2	142.2
1999.....	8.4	*	*	*	*	0.2	0.8	1.7	5.6	21.0	57.3	148.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Other diseases of circulatory system (I71-I78)

White, both sexes

2004.....	8.7	*	*	*	0.1	0.4	0.9	2.4	6.8	25.4	64.7	139.3
2003.....	9.2	*	*	*	0.2	0.3	0.9	2.5	7.9	27.3	68.3	146.9
2002.....	9.3	*	*	*	0.2	0.3	0.9	2.3	8.2	29.4	69.7	148.9
2001.....	9.6	*	*	*	0.1	0.4	0.9	2.3	8.3	29.5	72.1	158.2
2000.....	10.0	*	*	*	0.1	0.3	0.8	2.4	8.8	32.7	75.1	163.0
1999.....	10.0	*	*	*	0.1	0.3	1.0	2.4	9.1	32.1	76.4	165.1

White, male

2004.....	9.2	*	*	*	0.2	0.5	1.2	3.4	9.3	34.1	84.2	169.3
2003.....	10.0	*	*	*	0.2	0.4	1.2	3.6	11.0	37.9	93.1	184.1
2002.....	10.2	*	*	*	0.2	0.4	1.2	3.2	11.4	40.6	94.6	190.9
2001.....	10.4	*	*	*	0.2	0.6	1.2	3.1	11.6	41.1	98.3	201.3
2000.....	11.1	*	*	*	0.2	0.5	1.0	3.4	12.8	46.7	104.7	219.6
1999.....	11.0	*	*	*	0.2	0.5	1.2	3.3	13.4	45.8	106.2	211.1

White, female

2004.....	8.2	*	*	*	*	0.2	0.6	1.4	4.4	17.9	51.3	125.7
2003.....	8.3	*	*	*	*	0.2	0.5	1.5	4.9	18.3	51.5	130.5
2002.....	8.5	*	*	*	*	0.3	0.6	1.4	5.2	19.9	52.9	130.7
2001.....	8.8	*	*	*	*	0.3	0.6	1.5	5.2	19.7	54.7	140.1
2000.....	8.9	*	*	*	*	0.2	0.6	1.4	5.1	21.0	55.6	140.1
1999.....	9.1	*	*	*	*	0.2	0.7	1.5	5.2	20.6	56.9	146.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of circulatory system (I71-I78)

Black, both sexes

2004.....	6.6	*	*	*	*	0.8	2.2	5.3	12.7	28.0	70.4	155.1
2003.....	6.6	*	*	*	*	0.7	2.3	6.2	11.8	29.3	70.1	150.0
2002.....	6.7	*	*	*	*	0.9	2.0	5.9	13.5	31.1	68.3	156.8
2001.....	7.1	*	*	*	*	0.7	2.4	5.1	13.9	31.7	79.8	174.7
2000.....	7.1	*	*	*	*	0.7	2.2	5.5	14.4	33.0	78.0	177.7
1999.....	6.7	*	*	*	*	0.5	2.0	5.2	12.3	31.7	76.6	178.9

Black, male

2004.....	6.7	*	*	*	*	1.2	3.0	7.5	18.1	35.2	79.9	158.0
2003.....	6.7	*	*	*	*	1.1	3.0	8.2	16.4	36.3	90.3	138.7
2002.....	6.7	*	*	*	*	1.2	2.6	8.1	19.3	38.4	78.6	166.2
2001.....	6.9	*	*	*	*	0.9	3.4	7.3	17.8	40.0	96.8	165.7
2000.....	6.9	*	*	*	*	0.9	2.8	7.5	18.4	41.1	95.7	177.6
1999.....	6.5	*	*	*	*	0.9	2.3	7.2	15.7	39.4	89.8	184.0

Black, female

2004.....	6.6	*	*	*	*	*	1.6	3.4	8.4	22.8	65.0	154.0
2003.....	6.5	*	*	*	*	*	1.7	4.5	8.2	24.2	58.7	154.3
2002.....	6.6	*	*	*	*	*	1.4	4.0	8.9	25.8	62.5	153.2
2001.....	7.2	*	*	*	*	*	1.6	3.3	10.8	26.0	70.2	178.1
2000.....	7.2	*	*	*	*	*	1.5	3.9	11.1	27.3	68.0	177.7
1999.....	7.0	*	*	*	*	*	1.7	3.5	9.6	26.2	69.2	177.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Aortic aneurysm and dissection (I71)												
All races 3/, both sexes												
2004.....	4.7	*	*	*	0.1	0.3	0.7	1.8	4.5	15.9	39.6	64.8
2003.....	5.1	*	*	*	0.1	0.3	0.7	2.0	5.3	17.8	42.5	70.0
2002.....	5.1	*	*	*	0.1	0.3	0.7	1.8	5.5	19.1	43.3	69.4
2001.....	5.3	*	*	*	0.1	0.3	0.7	1.7	5.5	19.2	45.7	77.5
2000.....	5.6	*	*	*	0.1	0.3	0.7	1.8	6.2	21.8	47.2	79.2
1999.....	5.7	*	*	*	0.1	0.2	0.7	1.7	6.0	21.7	48.6	80.3
All races, male												
2004.....	5.6	*	*	*	0.2	0.5	1.1	2.7	6.9	23.1	54.8	95.0
2003.....	6.3	*	*	*	0.1	0.4	1.1	3.1	8.2	26.0	61.9	105.5
2002.....	6.4	*	*	*	0.2	0.4	1.0	2.7	8.5	28.5	62.7	106.6
2001.....	6.5	*	*	*	0.2	0.4	1.2	2.6	8.5	28.7	66.4	114.8
2000.....	7.0	*	*	*	0.1	0.4	1.0	2.7	9.5	32.9	70.6	127.3
1999.....	7.0	*	*	*	0.1	0.4	1.0	2.7	9.5	32.7	72.7	125.3
All races, female												
2004.....	3.8	*	*	*	*	0.2	0.3	0.8	2.3	9.8	29.3	51.2
2003.....	3.9	*	*	*	*	0.1	0.3	1.0	2.6	11.0	29.5	54.2
2002.....	4.0	*	*	*	*	0.1	0.3	0.8	2.7	11.2	30.5	53.3
2001.....	4.2	*	*	*	*	0.1	0.3	0.8	2.8	11.3	32.1	61.9
2000.....	4.3	*	*	*	*	0.1	0.4	0.9	3.0	12.7	32.0	59.6
1999.....	4.4	*	*	*	*	*	0.4	0.8	2.8	12.7	32.9	62.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Aortic aneurysm and dissection (I71)												
White, both sexes												
2004.....	5.1	*	*	*	0.1	0.3	0.6	1.5	4.3	16.5	40.6	66.8
2003.....	5.6	*	*	*	0.1	0.2	0.6	1.7	5.2	18.6	44.0	72.4
2002.....	5.7	*	*	*	0.1	0.2	0.6	1.5	5.4	19.9	45.0	71.2
2001.....	5.9	*	*	*	0.1	0.3	0.6	1.5	5.4	19.9	46.9	79.7
2000.....	6.2	*	*	*	0.1	0.2	0.6	1.6	6.0	22.7	48.5	80.8
1999.....	6.3	*	*	*	0.1	0.2	0.6	1.5	6.1	22.6	50.1	82.2
White, male												
2004.....	6.1	*	*	*	0.2	0.4	0.9	2.4	6.5	24.1	56.6	98.7
2003.....	6.9	*	*	*	0.1	0.3	1.0	2.7	8.2	27.3	64.3	110.0
2002.....	7.0	*	*	*	0.1	0.3	0.9	2.3	8.4	29.8	65.5	111.8
2001.....	7.2	*	*	*	0.1	0.5	1.0	2.3	8.4	29.9	68.5	120.3
2000.....	7.8	*	*	*	*	0.3	0.8	2.5	9.5	34.5	73.0	132.8
1999.....	7.8	*	*	*	*	0.4	0.9	2.4	9.8	34.3	75.8	129.8
White, female												
2004.....	4.1	*	*	*	*	0.2	0.3	0.7	2.1	10.1	29.8	52.4
2003.....	4.3	*	*	*	*	0.2	0.2	0.8	2.4	11.1	30.3	55.7
2002.....	4.3	*	*	*	*	*	0.3	0.7	2.6	11.4	31.2	53.5
2001.....	4.6	*	*	*	*	*	0.2	0.6	2.6	11.5	32.5	62.6
2000.....	4.6	*	*	*	*	0.1	0.3	0.7	2.8	12.8	32.3	59.6
1999.....	4.8	*	*	*	*	*	0.4	0.7	2.7	12.8	33.3	63.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Aortic aneurysm and dissection (I71)												
Black, both sexes												
2004.....	3.2	*	*	*	*	0.6	1.6	3.6	7.6	13.0	32.0	43.7
2003.....	3.2	*	*	*	*	0.5	1.6	4.2	7.1	14.5	30.0	46.4
2002.....	3.2	*	*	*	*	0.6	1.4	4.2	7.5	14.6	29.9	50.5
2001.....	3.4	*	*	*	*	0.4	1.6	3.5	7.6	14.6	37.6	55.3
2000.....	3.5	*	*	*	*	0.5	1.6	3.2	8.2	17.0	36.3	62.3
1999.....	3.3	*	*	*	*	0.4	1.4	3.7	6.5	16.7	33.8	60.3
Black, male												
2004.....	3.8	*	*	*	*	1.0	2.5	5.5	11.8	18.2	36.7	56.6
2003.....	3.8	*	*	*	*	1.0	2.4	6.4	10.8	18.0	39.2	51.1
2002.....	3.8	*	*	*	*	1.0	2.2	6.6	12.1	19.0	35.0	49.3
2001.....	3.8	*	*	*	*	*	2.6	5.3	11.2	20.0	47.4	48.8
2000.....	3.9	*	*	*	*	*	2.4	4.7	11.5	22.1	46.9	69.2
1999.....	3.6	*	*	*	*	*	1.9	5.8	9.1	21.9	38.4	76.2
Black, female												
2004.....	2.6	*	*	*	*	*	0.8	2.0	4.2	9.4	29.3	38.6
2003.....	2.7	*	*	*	*	*	1.0	2.4	4.2	11.9	24.7	44.6
2002.....	2.7	*	*	*	*	*	*	2.2	3.8	11.5	27.0	50.9
2001.....	3.0	*	*	*	*	*	0.7	1.9	4.6	10.8	32.1	57.7
2000.....	3.2	*	*	*	*	*	0.9	2.0	5.5	13.4	30.3	59.7
1999.....	3.0	*	*	*	*	*	0.9	2.0	4.4	13.0	31.2	54.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/	1-4 2/	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)												
All races 3/, both sexes												
2004.....	3.4	*	*	*	*	0.1	0.3	1.0	2.7	9.2	24.7	74.5
2003.....	3.4	*	*	*	0.1	0.1	0.3	0.9	2.8	9.1	25.0	76.0
2002.....	3.5	*	*	*	0.1	0.1	0.3	0.9	3.1	10.0	25.3	79.0
2001.....	3.5	*	*	*	*	0.1	0.4	0.9	3.2	10.0	26.1	80.7
2000.....	3.6	*	*	*	0.1	0.1	0.3	1.0	3.1	10.4	27.4	84.0
1999.....	3.6	*	*	*	*	0.1	0.3	0.9	3.3	9.8	27.2	84.9
All races, male												
2004.....	2.9	*	*	*	*	0.1	0.3	1.1	3.0	10.3	28.0	71.6
2003.....	3.0	*	*	*	*	0.1	0.3	1.0	3.1	11.0	29.9	74.2
2002.....	3.0	*	*	*	*	*	0.3	1.0	3.4	11.3	29.6	80.8
2001.....	3.1	*	*	*	*	0.1	0.3	0.9	3.5	11.6	30.6	82.5
2000.....	3.2	*	*	*	*	0.1	0.3	1.1	3.6	12.5	32.3	87.6
1999.....	3.1	*	*	*	*	0.1	0.3	1.0	3.8	11.7	31.5	82.2
All races, female												
2004.....	3.9	*	*	*	*	0.1	0.4	0.8	2.4	8.3	22.5	75.8
2003.....	3.8	*	*	*	*	*	0.3	0.9	2.6	7.5	21.8	76.9
2002.....	4.0	*	*	*	*	0.2	0.3	0.8	2.8	8.9	22.5	78.2
2001.....	4.0	*	*	*	*	0.1	0.4	0.9	2.9	8.6	23.1	79.9
2000.....	4.0	*	*	*	*	0.1	0.4	0.9	2.6	8.7	24.2	82.6
1999.....	4.1	*	*	*	*	0.1	0.4	0.9	2.8	8.3	24.5	86.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)												
White, both sexes												
2004.....	3.5	*	*	*	*	0.1	0.3	0.9	2.5	8.8	24.0	72.5
2003.....	3.6	*	*	*	*	0.1	0.3	0.8	2.7	8.7	24.2	74.6
2002.....	3.7	*	*	*	*	0.1	0.3	0.8	2.8	9.6	24.7	77.8
2001.....	3.7	*	*	*	*	0.1	0.3	0.9	2.9	9.5	25.2	78.5
2000.....	3.8	*	*	*	*	0.1	0.3	0.8	2.8	10.0	26.6	82.3
1999.....	3.8	*	*	*	*	0.1	0.3	0.9	3.0	9.5	26.3	82.9
White, male												
2004.....	3.0	*	*	*	*	*	0.3	1.0	2.7	10.1	27.7	70.7
2003.....	3.1	*	*	*	*	*	0.2	0.9	2.9	10.6	28.8	74.1
2002.....	3.2	*	*	*	*	*	0.3	0.9	3.1	10.8	29.1	79.1
2001.....	3.2	*	*	*	*	*	0.2	0.8	3.2	11.2	29.8	80.9
2000.....	3.4	*	*	*	*	0.1	0.2	0.9	3.3	12.2	31.6	86.9
1999.....	3.2	*	*	*	*	*	0.3	0.9	3.6	11.4	30.4	81.3
White, female												
2004.....	4.0	*	*	*	*	*	0.3	0.7	2.3	7.8	21.6	73.3
2003.....	4.0	*	*	*	*	*	0.3	0.7	2.5	7.2	21.2	74.8
2002.....	4.2	*	*	*	*	0.1	0.3	0.7	2.6	8.5	21.7	77.2
2001.....	4.2	*	*	*	*	0.1	0.4	0.9	2.6	8.2	22.1	77.5
2000.....	4.2	*	*	*	*	*	0.3	0.7	2.3	8.2	23.3	80.4
1999.....	4.3	*	*	*	*	*	0.3	0.8	2.5	7.9	23.6	83.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Black, both sexes												
2004.....	3.4	*	*	*	*	*	0.6	1.7	5.2	14.9	38.4	111.4
2003.....	3.4	*	*	*	*	*	0.7	2.0	4.7	14.8	40.1	103.6
2002.....	3.5	*	*	*	*	*	0.6	1.7	6.1	16.5	38.5	106.3
2001.....	3.7	*	*	*	*	*	0.9	1.7	6.4	17.1	42.2	119.4
2000.....	3.6	*	*	*	*	*	0.6	2.3	6.2	16.0	41.7	115.4
1999.....	3.5	*	*	*	*	*	0.6	1.5	5.8	15.0	42.8	118.6
Black, male												
2004.....	2.9	*	*	*	*	*	*	2.0	6.3	17.0	43.2	101.3
2003.....	2.9	*	*	*	*	*	*	1.8	5.6	18.3	51.1	87.6
2002.....	3.0	*	*	*	*	*	*	1.5	7.2	19.4	43.6	116.9
2001.....	3.2	*	*	*	*	*	0.9	2.0	6.5	20.0	49.4	116.9
2000.....	3.1	*	*	*	*	*	*	2.8	7.0	19.0	48.7	108.4
1999.....	2.9	*	*	*	*	*	*	1.4	6.6	17.5	51.3	107.8
Black, female												
2004.....	3.9	*	*	*	*	*	0.8	1.4	4.2	13.5	35.7	115.4
2003.....	3.8	*	*	*	*	*	0.7	2.1	4.0	12.3	34.0	109.8
2002.....	3.9	*	*	*	*	*	0.8	1.9	5.1	14.4	35.6	102.3
2001.....	4.2	*	*	*	*	*	0.8	1.4	6.2	15.1	38.2	120.4
2000.....	4.1	*	*	*	*	*	0.7	1.8	5.5	13.9	37.7	118.0
1999.....	4.0	*	*	*	*	*	0.8	1.6	5.2	13.3	38.0	122.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Other disorders of circulatory system (I80-I99)

All races 3/, both sexes

2004.....	1.6	0.7	*	*	0.1	0.3	0.7	1.3	2.1	3.6	9.0	23.3
2003.....	1.6	0.9	*	*	0.1	0.3	0.8	1.4	2.0	4.1	9.0	24.0
2002.....	1.6	0.7	*	*	0.1	0.3	0.7	1.3	2.2	3.8	9.8	24.4
2001.....	1.6	0.5	*	*	0.1	0.3	0.6	1.2	2.2	4.6	9.4	26.1
2000.....	1.6	0.7	*	*	0.1	0.3	0.7	1.4	2.1	4.3	9.4	25.6
1999.....	1.4	0.6	*	*	0.1	0.2	0.6	1.0	1.9	3.8	9.1	23.4

All races, male

2004.....	1.4	*	*	*	*	0.4	0.7	1.6	2.5	3.9	9.7	19.6
2003.....	1.4	1.0	*	*	0.1	0.3	0.8	1.6	2.4	4.2	8.5	19.1
2002.....	1.4	*	*	*	0.1	0.3	0.7	1.4	2.4	3.9	9.5	20.5
2001.....	1.4	*	*	*	*	0.3	0.7	1.4	2.5	4.9	9.8	22.0
2000.....	1.4	*	*	*	*	0.3	0.8	1.6	2.3	4.7	9.7	22.2
1999.....	1.3	*	*	*	*	0.2	0.7	1.1	2.0	4.2	9.6	20.8

All races, female

2004.....	1.7	*	*	*	*	0.3	0.7	1.1	1.8	3.4	8.5	25.0
2003.....	1.9	*	*	*	0.1	0.4	0.8	1.2	1.7	4.0	9.4	26.2
2002.....	1.9	*	*	*	*	0.4	0.7	1.2	2.1	3.8	10.1	26.2
2001.....	1.9	*	*	*	0.1	0.4	0.6	1.0	2.0	4.4	9.2	27.8
2000.....	1.8	*	*	*	0.1	0.4	0.6	1.1	2.0	4.0	9.2	26.9
1999.....	1.6	*	*	*	*	0.2	0.5	1.0	1.8	3.4	8.8	24.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4/2/ years	5-14/ years	15-24/ years	25-34/ years	35-44/ years	45-54/ years	55-64/ years	65-74/ years	75-84/ years	85 years and over
---	----------	----------------	--------------	-------------	--------------	--------------	--------------	--------------	--------------	--------------	--------------	-------------------

Other disorders of circulatory system (I80-I99)

White, both sexes

2004.....	1.6	0.7	*	*	0.1	0.3	0.6	1.2	2.0	3.4	8.8	23.2
2003.....	1.6	0.9	*	*	0.1	0.3	0.6	1.2	1.9	3.8	8.6	23.6
2002.....	1.6	0.6	*	*	0.1	0.3	0.6	1.2	2.0	3.6	9.5	24.3
2001.....	1.7	*	*	*	0.1	0.3	0.5	1.0	2.0	4.3	9.1	26.0
2000.....	1.6	0.7	*	*	0.1	0.3	0.6	1.2	1.9	4.0	9.1	25.5
1999.....	1.5	*	*	*	*	0.2	0.5	0.9	1.8	3.5	8.9	23.2

White, male

2004.....	1.4	*	*	*	*	0.3	0.6	1.4	2.4	3.7	9.4	19.7
2003.....	1.3	*	*	*	*	0.2	0.7	1.4	2.2	3.9	7.9	18.5
2002.....	1.3	*	*	*	*	0.3	0.6	1.3	2.2	3.6	9.0	19.9
2001.....	1.4	*	*	*	*	0.2	0.5	1.3	2.2	4.6	9.5	21.7
2000.....	1.4	*	*	*	*	0.2	0.8	1.5	2.0	4.4	9.5	22.0
1999.....	1.2	*	*	*	*	0.2	0.6	0.9	1.9	3.9	9.2	20.7

White, female

2004.....	1.8	*	*	*	*	0.3	0.5	0.9	1.6	3.2	8.4	24.8
2003.....	1.9	*	*	*	*	0.3	0.5	1.1	1.5	3.8	9.1	25.8
2002.....	1.9	*	*	*	*	0.3	0.6	1.1	1.9	3.6	9.8	26.3
2001.....	1.9	*	*	*	*	0.3	0.5	0.8	1.8	4.1	8.9	27.9
2000.....	1.9	*	*	*	*	0.3	0.5	1.0	1.8	3.6	8.9	26.9
1999.....	1.7	*	*	*	*	0.2	0.4	0.9	1.7	3.2	8.7	24.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other disorders of circulatory system (I80-I99)

Black, both sexes

2004.....	2.1	*	*	*	*	0.9	1.9	2.9	4.2	6.6	13.6	28.8
2003.....	2.2	*	*	*	*	0.9	1.9	2.6	4.3	7.5	15.4	33.6
2002.....	2.1	*	*	*	*	1.0	1.6	2.6	4.9	6.8	16.6	30.0
2001.....	2.0	*	*	*	*	0.8	1.5	2.2	4.9	7.8	14.4	30.1
2000.....	2.1	*	*	*	*	0.9	1.5	2.8	4.5	8.9	14.4	30.0
1999.....	1.7	*	*	*	*	0.5	1.4	2.4	2.9	6.6	13.8	28.4

Black, male

2004.....	1.9	*	*	*	*	1.1	2.0	3.4	4.6	7.1	15.0	24.8
2003.....	2.0	*	*	*	*	0.9	1.3	3.4	5.2	8.7	17.5	32.3
2002.....	2.0	*	*	*	*	1.0	1.8	3.0	5.6	7.7	18.4	32.2
2001.....	1.9	*	*	*	*	*	1.7	2.4	6.4	9.0	15.9	29.5
2000.....	1.9	*	*	*	*	0.9	1.8	3.4	5.1	9.1	13.3	30.0
1999.....	1.6	*	*	*	*	*	1.8	2.8	2.6	8.2	16.9	*

Black, female

2004.....	2.2	*	*	*	*	0.8	1.9	2.5	3.9	6.3	12.7	30.3
2003.....	2.3	*	*	*	*	0.9	2.5	1.9	3.6	6.7	14.2	34.0
2002.....	2.2	*	*	*	*	0.9	1.5	2.3	4.4	6.1	15.6	29.2
2001.....	2.1	*	*	*	*	0.9	1.4	2.1	3.6	7.1	13.6	30.3
2000.....	2.2	*	*	*	*	0.9	1.3	2.3	3.9	8.8	15.0	30.0
1999.....	1.8	*	*	*	*	*	1.1	2.0	3.1	5.5	12.0	30.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza and pneumonia (J10-J18)												
All races 3/, both sexes												
2004.....	20.3	6.7	0.7	0.2	0.4	0.8	2.0	4.6	10.8	34.6	139.3	582.6
2003.....	22.4	8.0	1.0	0.4	0.5	0.9	2.2	5.2	11.2	37.3	151.1	666.1
2002.....	22.8	6.5	0.7	0.2	0.4	0.9	2.2	4.8	11.2	37.5	156.9	696.6
2001.....	21.8	7.4	0.7	0.2	0.5	0.9	2.2	4.6	10.7	36.3	148.5	685.6
2000.....	23.2	7.6	0.7	0.2	0.5	0.9	2.4	4.7	11.9	39.1	160.3	744.1
1999.....	22.8	8.4	0.8	0.2	0.5	0.8	2.4	4.6	11.0	37.2	157.0	751.8
All races, male												
2004.....	18.6	8.0	0.6	0.2	0.6	1.0	2.2	5.6	13.0	42.8	172.7	677.2
2003.....	20.1	9.4	1.1	0.3	0.6	1.1	2.5	6.4	13.2	46.5	186.8	757.8
2002.....	20.4	7.8	0.8	0.2	0.5	0.9	2.4	5.8	13.6	46.4	194.7	805.8
2001.....	19.6	8.1	0.7	0.2	0.5	1.0	2.5	5.7	13.1	45.0	186.4	808.4
2000.....	20.8	8.4	0.7	0.2	0.5	1.0	2.8	5.8	14.4	48.3	199.8	892.0
1999.....	20.3	8.8	0.8	0.3	0.5	1.0	2.6	5.9	12.8	46.9	196.5	894.7
All races, female												
2004.....	22.0	5.4	0.9	0.2	0.3	0.6	1.8	3.6	8.8	27.7	116.8	540.0
2003.....	24.6	6.6	0.9	0.4	0.5	0.7	1.9	4.1	9.4	29.5	127.2	625.6
2002.....	25.1	5.1	0.6	0.3	0.4	0.8	1.9	3.8	9.0	30.1	131.8	649.2
2001.....	23.9	6.7	0.7	0.2	0.4	0.8	1.9	3.5	8.5	29.1	123.7	634.2
2000.....	25.6	6.8	0.7	0.2	0.5	0.8	1.9	3.7	9.5	31.5	134.6	683.9
1999.....	25.3	8.0	0.9	0.2	0.5	0.7	2.1	3.4	9.4	29.3	131.3	694.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza and pneumonia (J10-J18)												
White, both sexes												
2004.....	22.0	5.1	0.6	0.2	0.4	0.7	1.8	4.1	10.1	33.4	138.5	590.7
2003.....	24.4	6.7	0.9	0.3	0.5	0.8	2.0	4.7	10.3	35.9	151.5	679.3
2002.....	24.9	5.2	0.5	0.2	0.4	0.8	1.9	4.4	10.4	36.7	156.9	709.1
2001.....	23.6	5.3	0.7	0.2	0.4	0.7	1.8	4.0	9.9	35.3	147.2	695.0
2000.....	25.2	5.8	0.6	0.2	0.4	0.8	2.1	4.1	10.9	37.1	160.0	758.3
1999.....	24.8	6.2	0.6	0.2	0.4	0.7	1.9	4.0	10.1	36.1	157.8	764.3
White, male												
2004.....	19.8	6.2	0.5	0.2	0.6	0.9	1.9	4.9	12.1	40.6	170.2	685.5
2003.....	21.4	8.0	1.1	0.3	0.6	1.0	2.2	5.6	11.6	44.0	184.9	767.1
2002.....	21.9	6.6	0.6	0.2	0.4	0.8	2.1	5.2	12.3	45.2	193.1	815.9
2001.....	20.7	6.2	0.7	0.2	0.5	0.8	2.1	4.9	11.8	43.0	182.6	814.4
2000.....	22.0	6.4	0.5	0.2	0.5	0.9	2.5	4.8	12.9	45.1	197.8	906.1
1999.....	21.5	6.7	0.5	0.3	0.4	0.8	2.1	4.9	11.5	44.7	196.1	905.7
White, female												
2004.....	24.2	3.9	0.7	0.2	0.3	0.6	1.6	3.2	8.3	27.3	116.9	548.0
2003.....	27.3	5.4	0.8	0.4	0.5	0.6	1.8	3.7	9.0	29.0	129.0	640.4
2002.....	27.8	3.7	0.5	0.3	0.3	0.7	1.7	3.6	8.5	29.6	132.7	662.6
2001.....	26.4	4.4	0.6	0.2	0.3	0.7	1.6	3.2	8.1	28.9	123.8	645.0
2000.....	28.2	5.1	0.7	0.2	0.4	0.8	1.7	3.4	8.9	30.3	135.0	698.3
1999.....	27.9	5.6	0.7	0.2	0.4	0.6	1.8	3.0	8.7	28.9	132.6	707.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza and pneumonia (J10-J18)												
Black, both sexes												
2004.....	14.8	14.1	1.3	0.3	0.6	1.2	4.0	9.1	19.0	49.8	157.2	505.1
2003.....	15.4	14.1	1.1	0.4	0.7	1.9	4.4	10.4	20.6	54.2	155.9	525.4
2002.....	15.6	12.9	1.6	*	0.6	1.5	4.3	8.7	21.2	50.3	166.2	575.2
2001.....	15.5	16.3	1.0	*	0.7	1.6	4.8	9.7	18.9	48.3	169.0	581.3
2000.....	16.4	16.7	1.2	*	0.7	1.7	4.5	10.0	22.0	60.3	171.1	600.9
1999.....	16.2	19.0	1.7	*	0.8	2.0	5.5	10.6	20.9	54.1	157.4	634.8
Black, male												
2004.....	15.1	16.3	*	*	*	1.7	4.8	11.8	24.9	72.5	219.4	598.0
2003.....	16.0	16.1	*	*	0.8	2.3	5.3	13.2	29.1	78.2	218.6	641.2
2002.....	15.4	13.9	1.7	*	0.6	1.7	5.1	11.2	29.0	65.9	228.9	689.4
2001.....	15.9	15.9	*	*	0.8	2.1	5.7	13.3	26.3	69.4	242.2	714.8
2000.....	16.7	17.9	1.7	*	0.7	2.3	5.6	13.9	30.0	83.5	234.1	755.4
1999.....	16.4	18.9	2.0	*	0.8	2.2	6.6	15.1	27.2	76.9	213.7	803.8
Black, female												
2004.....	14.5	11.7	1.8	*	0.7	0.8	3.3	6.8	14.3	33.6	122.1	468.2
2003.....	14.9	12.1	*	*	*	1.5	3.6	7.9	13.8	37.2	120.4	480.4
2002.....	15.7	11.8	*	*	*	1.3	3.7	6.5	15.0	39.1	130.7	531.5
2001.....	15.2	16.7	*	*	0.7	1.3	3.9	6.7	13.0	33.4	127.9	531.7
2000.....	16.0	15.5	*	*	0.8	1.2	3.4	6.7	15.6	43.9	135.7	543.4
1999.....	16.1	19.1	*	*	0.9	1.7	4.4	6.8	15.8	38.0	125.8	571.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza (J10-J11)												
All races 3/, both sexes												
2004.....	0.4	*	*	*	*	*	*	0.1	0.2	0.5	2.2	11.4
2003.....	0.6	0.8	0.4	0.1	0.1	*	0.1	0.1	0.3	1.0	3.6	17.0
2002.....	0.3	*	*	*	*	*	*	*	0.1	0.3	1.4	8.8
2001.....	0.1	*	*	*	*	*	*	*	0.1	0.1	0.4	2.2
2000.....	0.6	*	*	*	*	*	0.0	0.1	0.4	1.1	4.0	20.2
1999.....	0.6	*	*	*	*	*	*	0.1	0.3	0.9	3.7	21.2
All races, male												
2004.....	0.3	*	*	*	*	*	*	*	0.2	0.7	2.4	11.1
2003.....	0.5	*	0.4	*	*	*	*	0.2	0.3	1.1	3.7	17.0
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.6	8.9
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.2
2000.....	0.5	*	*	*	*	*	*	0.1	0.4	1.3	4.0	20.7
1999.....	0.4	*	*	*	*	*	*	*	0.3	0.9	4.2	19.9
All races, female												
2004.....	0.5	*	*	*	*	*	*	*	0.2	0.4	2.1	11.5
2003.....	0.7	*	0.3	0.1	*	*	*	0.1	0.3	0.9	3.5	17.1
2002.....	0.3	*	*	*	*	*	*	*	*	0.2	1.3	8.7
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.5	2.2
2000.....	0.8	*	*	*	*	*	*	0.1	0.4	0.9	4.0	19.9
1999.....	0.7	*	*	*	*	*	*	*	0.3	0.9	3.4	21.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza (J10-J11)												
White, both sexes												
2004.....	0.4	*	*	*	*	*	*	0.1	0.2	0.6	2.3	12.3
2003.....	0.7	0.7	0.3	0.1	0.1	*	0.1	0.1	0.3	1.1	3.8	18.2
2002.....	0.3	*	*	*	*	*	*	*	0.1	0.3	1.5	9.4
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.5	2.3
2000.....	0.7	*	*	*	*	*	*	0.1	0.4	1.1	4.2	21.4
1999.....	0.7	*	*	*	*	*	*	0.1	0.3	0.9	3.9	22.8
White, male												
2004.....	0.3	*	*	*	*	*	*	*	0.2	0.7	2.4	12.1
2003.....	0.5	*	0.4	*	*	*	*	0.2	0.3	1.1	3.9	18.3
2002.....	0.2	*	*	*	*	*	*	*	*	0.4	1.7	9.7
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.3
2000.....	0.5	*	*	*	*	*	*	*	0.4	1.3	4.2	21.6
1999.....	0.5	*	*	*	*	*	*	*	0.4	0.9	4.4	21.7
White, female												
2004.....	0.5	*	*	*	*	*	*	*	0.2	0.4	2.3	12.4
2003.....	0.8	*	*	*	*	*	*	0.1	0.3	1.0	3.7	18.1
2002.....	0.4	*	*	*	*	*	*	*	*	*	1.4	9.3
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.5	2.3
2000.....	0.9	*	*	*	*	*	*	0.1	0.4	0.9	4.2	21.3
1999.....	0.9	*	*	*	*	*	*	*	0.3	0.9	3.5	23.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia (J12-J18)												
All races 3/, both sexes												
2004.....	19.9	6.3	0.6	0.2	0.4	0.7	2.0	4.5	10.7	34.0	137.0	571.2
2003.....	21.8	7.3	0.7	0.2	0.5	0.9	2.2	5.1	10.9	36.3	147.5	649.1
2002.....	22.5	6.3	0.7	0.2	0.4	0.9	2.1	4.7	11.1	37.2	155.5	687.8
2001.....	21.7	7.2	0.7	0.2	0.4	0.8	2.2	4.6	10.6	36.2	148.1	683.4
2000.....	22.6	7.4	0.6	0.2	0.4	0.9	2.3	4.6	11.5	38.0	156.4	723.9
1999.....	22.2	8.1	0.8	0.2	0.4	0.8	2.3	4.6	10.7	36.4	153.3	730.6
All races, male												
2004.....	18.3	7.5	0.5	0.2	0.5	0.9	2.2	5.5	12.8	42.1	170.3	666.1
2003.....	19.6	8.5	0.7	0.2	0.5	1.1	2.5	6.2	12.9	45.4	183.2	740.9
2002.....	20.2	7.6	0.8	0.2	0.4	0.9	2.4	5.7	13.4	46.0	193.1	796.9
2001.....	19.5	7.8	0.7	0.2	0.5	0.9	2.5	5.7	13.0	44.9	186.0	806.2
2000.....	20.3	8.1	0.6	0.2	0.5	1.0	2.7	5.6	14.0	47.1	195.8	871.3
1999.....	19.8	8.5	0.7	0.2	0.4	0.9	2.6	5.8	12.5	46.0	192.3	874.7
All races, female												
2004.....	21.5	5.0	0.8	0.2	0.3	0.5	1.8	3.5	8.6	27.3	114.6	528.5
2003.....	23.9	6.0	0.6	0.3	0.4	0.7	1.9	4.0	9.1	28.6	123.7	608.5
2002.....	24.8	5.0	0.6	0.2	0.3	0.8	1.9	3.8	9.0	29.8	130.6	640.5
2001.....	23.8	6.7	0.7	0.2	0.4	0.7	1.9	3.5	8.4	29.0	123.2	632.0
2000.....	24.8	6.6	0.6	0.2	0.4	0.8	1.9	3.6	9.1	30.6	130.6	663.9
1999.....	24.6	7.7	0.8	0.2	0.4	0.7	2.0	3.4	9.1	28.4	128.0	672.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia (J12-J18)												
White, both sexes												
2004.....	21.6	4.7	0.5	0.1	0.4	0.7	1.7	4.0	9.9	32.8	136.2	578.4
2003.....	23.7	6.0	0.6	0.2	0.5	0.8	1.9	4.5	10.0	34.8	147.7	661.1
2002.....	24.6	5.0	0.5	0.2	0.4	0.8	1.9	4.3	10.2	36.4	155.4	699.6
2001.....	23.5	5.1	0.6	0.2	0.4	0.7	1.8	4.0	9.8	35.2	146.8	692.7
2000.....	24.4	5.6	0.5	0.2	0.4	0.8	2.0	4.0	10.4	36.0	155.8	737.0
1999.....	24.1	5.9	0.6	0.2	0.4	0.7	1.9	3.9	9.8	35.2	153.9	741.6
White, male												
2004.....	19.5	5.8	0.4	0.2	0.6	0.9	1.9	4.9	11.9	39.8	167.8	673.4
2003.....	20.9	7.1	0.7	0.2	0.5	1.0	2.2	5.5	11.3	42.8	181.0	748.8
2002.....	21.7	6.4	0.5	0.1	0.4	0.8	2.1	5.1	12.1	44.8	191.4	806.2
2001.....	20.6	5.8	0.6	0.2	0.5	0.8	2.1	4.8	11.7	42.9	182.2	812.1
2000.....	21.5	6.1	0.4	0.2	0.5	0.8	2.4	4.7	12.5	43.8	193.6	884.4
1999.....	21.0	6.7	0.4	0.2	0.4	0.8	2.1	4.8	11.1	43.8	191.7	884.0
White, female												
2004.....	23.6	3.5	0.6	0.1	0.3	0.5	1.6	3.1	8.1	26.8	114.6	535.6
2003.....	26.5	4.8	0.5	0.2	0.4	0.6	1.7	3.6	8.7	28.0	125.2	622.3
2002.....	27.4	3.6	0.5	0.2	0.3	0.7	1.7	3.5	8.5	29.3	131.3	653.3
2001.....	26.2	4.4	0.6	0.2	0.3	0.7	1.6	3.1	8.0	28.7	123.3	642.8
2000.....	27.3	5.0	0.6	0.2	0.3	0.7	1.7	3.2	8.5	29.4	130.8	677.1
1999.....	27.1	5.2	0.7	0.2	0.3	0.5	1.7	3.0	8.5	27.9	129.1	684.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia (J12-J18)												
Black, both sexes												
2004.....	14.6	13.5	1.2	*	0.6	1.2	4.0	9.0	18.9	49.4	155.6	502.5
2003.....	15.1	13.0	0.9	*	0.6	1.8	4.4	10.1	20.3	53.7	154.5	519.6
2002.....	15.5	12.7	1.5	*	0.6	1.5	4.3	8.7	21.1	49.9	165.8	572.8
2001.....	15.5	16.3	1.0	*	0.7	1.6	4.7	9.7	18.8	48.2	168.6	579.7
2000.....	16.1	16.4	1.2	*	0.7	1.7	4.4	9.9	21.8	59.3	169.7	591.5
1999.....	16.1	18.3	1.7	*	0.8	1.9	5.4	10.5	20.7	53.3	155.7	629.1
Black, male												
2004.....	14.9	15.1	*	*	*	1.6	4.8	11.7	24.8	71.9	216.6	597.1
2003.....	15.8	15.2	*	*	0.7	2.2	5.2	12.9	28.7	77.7	217.5	638.1
2002.....	15.3	13.7	1.6	*	*	1.7	5.1	11.2	28.7	65.5	228.4	687.2
2001.....	15.8	15.9	*	*	0.7	2.0	5.7	13.3	26.1	69.1	242.2	712.6
2000.....	16.5	17.9	1.7	*	0.7	2.2	5.5	13.6	29.7	82.8	232.5	742.7
1999.....	16.3	17.9	1.9	*	0.8	2.2	6.6	15.0	27.1	76.0	211.8	800.3
Black, female												
2004.....	14.4	11.7	1.7	*	0.6	0.8	3.2	6.7	14.1	33.3	121.2	465.0
2003.....	14.6	10.8	*	*	*	1.4	3.6	7.7	13.6	36.6	119.0	473.5
2002.....	15.6	11.8	*	*	*	1.3	3.7	6.5	14.9	38.8	130.4	529.0
2001.....	15.1	16.7	*	*	0.7	1.3	3.9	6.7	13.0	33.4	127.2	530.5
2000.....	15.8	14.9	*	*	0.8	1.2	3.4	6.6	15.4	42.7	134.3	535.3
1999.....	15.9	18.8	*	*	0.9	1.7	4.3	6.8	15.5	37.2	124.3	565.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other acute lower respiratory infections (J20-J22)

All races 3/, both sexes												
2004.....	0.1	1.5	*	*	*	*	*	*	*	0.1	0.4	2.6
2003.....	0.1	1.3	0.2	*	*	*	*	*	0.1	0.2	0.6	3.5
2002.....	0.1	1.3	0.1	*	*	*	*	*	*	0.2	0.5	3.3
2001.....	0.1	1.2	*	*	*	*	*	*	*	0.2	0.5	3.3
2000.....	0.2	1.1	*	*	*	*	*	*	*	0.2	0.7	4.5
1999.....	0.2	1.7	0.2	*	*	*	*	*	0.1	0.2	0.9	6.0
All races, male												
2004.....	0.1	1.6	*	*	*	*	*	*	*	*	*	2.5
2003.....	0.1	1.7	*	*	*	*	*	*	*	*	0.7	2.4
2002.....	0.1	1.9	*	*	*	*	*	*	*	*	0.5	2.8
2001.....	0.1	1.3	*	*	*	*	*	*	*	*	0.4	2.3
2000.....	0.1	1.3	*	*	*	*	*	*	*	0.3	0.7	3.8
1999.....	0.1	1.9	*	*	*	*	*	*	*	*	1.1	4.0
All races, female												
2004.....	0.1	1.3	*	*	*	*	*	*	*	*	0.5	2.7
2003.....	0.2	*	*	*	*	*	*	*	*	*	0.6	3.9
2002.....	0.2	*	*	*	*	*	*	*	*	0.2	0.5	3.5
2001.....	0.2	1.2	*	*	*	*	*	*	*	*	0.5	3.7
2000.....	0.2	*	*	*	*	*	*	*	*	0.2	0.6	4.8
1999.....	0.2	1.5	*	*	*	*	*	*	*	*	0.8	6.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other acute lower respiratory infections (J20-J22)

White, both sexes												
2004.....	0.1	1.4	*	*	*	*	*	*	*	*	0.4	2.8
2003.....	0.2	1.3	*	*	*	*	*	*	*	0.2	0.7	3.7
2002.....	0.1	1.1	0.2	*	*	*	*	*	*	0.2	0.5	3.5
2001.....	0.1	1.0	*	*	*	*	*	*	*	0.1	0.5	3.5
2000.....	0.2	0.8	*	*	*	*	*	*	*	0.3	0.7	4.7
1999.....	0.2	1.2	*	*	*	*	*	*	*	0.2	0.9	6.3
White, male												
2004.....	0.1	1.5	*	*	*	*	*	*	*	*	*	2.6
2003.....	0.1	1.7	*	*	*	*	*	*	*	*	0.7	2.7
2002.....	0.1	1.6	*	*	*	*	*	*	*	*	0.5	2.9
2001.....	0.1	*	*	*	*	*	*	*	*	*	*	2.4
2000.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	3.8
1999.....	0.1	*	*	*	*	*	*	*	*	*	1.1	4.0
White, female												
2004.....	0.1	*	*	*	*	*	*	*	*	*	0.5	2.8
2003.....	0.2	*	*	*	*	*	*	*	*	*	0.6	4.2
2002.....	0.2	*	*	*	*	*	*	*	*	*	0.5	3.7
2001.....	0.2	*	*	*	*	*	*	*	*	*	0.5	3.9
2000.....	0.2	*	*	*	*	*	*	*	*	0.2	0.7	5.1
1999.....	0.3	*	*	*	*	*	*	*	*	*	0.8	7.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute bronchitis and bronchiolitis (J20-J21)

All races 3/, both sexes

2004.....	0.1	1.4	*	*	*	*	*	*	*	*	0.2	1.4
2003.....	0.1	1.3	0.2	*	*	*	*	*	*	0.1	0.3	2.1
2002.....	0.1	1.2	0.1	*	*	*	*	*	*	0.1	0.3	2.0
2001.....	0.1	1.2	*	*	*	*	*	*	*	0.1	0.3	2.1
2000.....	0.1	1.1	*	*	*	*	*	*	*	0.2	0.4	2.6
1999.....	0.1	1.6	0.2	*	*	*	*	*	*	0.2	0.6	3.6

All races, male

2004.....	0.1	1.6	*	*	*	*	*	*	*	*	*	1.8
2003.....	0.1	1.7	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	1.8	*	*	*	*	*	*	*	*	*	1.7
2001.....	0.1	1.2	*	*	*	*	*	*	*	*	*	*
2000.....	0.1	1.2	*	*	*	*	*	*	*	*	0.6	2.4
1999.....	0.1	1.9	*	*	*	*	*	*	*	*	0.7	2.6

All races, female

2004.....	0.1	1.3	*	*	*	*	*	*	*	*	0.3	1.3
2003.....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.5
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
2001.....	0.1	1.2	*	*	*	*	*	*	*	*	0.4	2.4
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.7
1999.....	0.2	1.4	*	*	*	*	*	*	*	*	0.5	4.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute bronchitis and bronchiolitis (J20-J21)

White, both sexes

2004.....	0.1	1.3	*	*	*	*	*	*	*	*	0.2	1.5
2003.....	0.1	1.3	*	*	*	*	*	*	*	*	0.4	2.3
2002.....	0.1	1.0	0.2	*	*	*	*	*	*	0.1	0.3	2.1
2001.....	0.1	1.0	*	*	*	*	*	*	*	*	0.3	2.1
2000.....	0.1	0.8	*	*	*	*	*	*	*	0.2	0.4	2.7
1999.....	0.1	1.2	*	*	*	*	*	*	*	0.1	0.6	3.8

White, male

2004.....	0.1	1.4	*	*	*	*	*	*	*	*	*	1.8
2003.....	0.1	1.6	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	1.5	*	*	*	*	*	*	*	*	*	1.7
2001.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.6	2.4
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.7	2.6

White, female

2004.....	0.1	*	*	*	*	*	*	*	*	*	0.3	1.3
2003.....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.7
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	2.3
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.5
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.9
1999.....	0.2	*	*	*	*	*	*	*	*	*	0.5	4.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Unspecified acute lower respiratory infection (J22)

All races 3/, both sexes

2004.....	0.0	*	*	*	*	*	*	*	*	*	0.2	1.2
2003.....	0.0	*	*	*	*	*	*	*	*	*	0.3	1.3
2002.....	0.0	*	*	*	*	*	*	*	*	*	0.2	1.3
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	1.2
2000.....	0.0	*	*	*	*	*	*	*	*	*	0.2	1.9
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.4

All races, male

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*	*	*	*

All races, female

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	1.4
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	1.4
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	1.3
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	1.3
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.1
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Unspecified acute lower respiratory infection (J22)

White, both sexes

2004.....	0.0	*	*	*	*	*	*	*	*	*	0.2	1.3
2003.....	0.0	*	*	*	*	*	*	*	*	*	0.3	1.4
2002.....	0.0	*	*	*	*	*	*	*	*	*	0.2	1.4
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	1.3
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.0
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.5

White, male

2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*	*	*	*

White, female

2004.....	0.1	*	*	*	*	*	*	*	*	*	*	1.5
2003.....	0.1	*	*	*	*	*	*	*	*	*	*	1.5
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	1.4
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	1.5
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Chronic lower respiratory diseases (J40-J47)

All races 3/, both sexes

2004.....	41.5	0.9	0.3	0.3	0.4	0.6	2.0	8.4	40.4	153.8	366.7	601.7
2003.....	43.5	0.8	0.3	0.3	0.5	0.7	2.1	8.7	43.3	163.2	383.0	635.1
2002.....	43.3	1.0	0.4	0.3	0.5	0.8	2.2	8.7	42.4	163.0	386.7	637.6
2001.....	43.2	1.0	0.3	0.3	0.4	0.7	2.2	8.5	44.1	167.9	379.8	644.7
2000.....	43.4	0.9	0.3	0.3	0.5	0.7	2.1	8.6	44.2	169.4	386.1	648.6
1999.....	44.5	0.9	0.4	0.3	0.5	0.8	2.0	8.5	47.5	177.2	397.8	646.0

All races, male

2004.....	40.6	1.2	0.4	0.4	0.5	0.7	2.0	8.8	43.1	172.1	445.6	811.1
2003.....	42.4	1.1	0.5	0.4	0.5	0.8	1.9	9.1	46.5	183.6	464.9	865.9
2002.....	42.9	1.1	0.6	0.4	0.6	0.8	2.2	9.1	45.2	184.8	480.8	894.8
2001.....	42.7	1.1	0.4	0.3	0.5	0.7	2.0	8.8	46.9	191.3	475.1	916.9
2000.....	43.5	1.2	0.4	0.4	0.6	0.8	1.9	9.0	47.8	195.2	488.5	967.9
1999.....	45.6	*	0.4	0.4	0.6	0.8	1.8	8.6	52.3	210.7	513.2	996.7

All races, female

2004.....	42.5	*	*	0.2	0.3	0.5	2.1	8.1	38.0	138.4	313.6	507.4
2003.....	44.4	*	*	0.2	0.4	0.6	2.3	8.2	40.3	146.0	328.3	533.0
2002.....	43.7	*	0.3	0.3	0.4	0.7	2.3	8.2	39.8	144.9	324.1	526.0
2001.....	43.7	*	*	0.2	0.4	0.7	2.3	8.1	41.5	148.5	317.3	530.8
2000.....	43.2	*	0.3	0.3	0.4	0.7	2.2	8.3	41.0	148.2	319.2	518.5
1999.....	43.4	*	0.3	0.2	0.5	0.9	2.1	8.4	43.1	149.8	322.9	504.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Chronic lower respiratory diseases (J40-J47)

White, both sexes

2004.....	47.4	0.7	0.2	0.2	0.3	0.5	1.8	8.2	42.3	164.7	387.0	630.1
2003.....	49.5	0.7	0.2	0.2	0.3	0.5	1.9	8.5	45.2	173.8	403.0	664.5
2002.....	49.2	0.8	0.2	0.2	0.3	0.6	1.9	8.4	44.2	173.2	405.6	666.2
2001.....	49.0	0.7	0.2	0.2	0.3	0.5	1.8	8.3	46.0	177.7	398.6	671.8
2000.....	49.0	0.8	0.2	0.2	0.3	0.5	1.7	8.2	45.8	178.7	404.4	675.6
1999.....	50.2	*	0.3	0.2	0.4	0.5	1.6	8.1	49.3	186.2	414.3	672.9

White, male

2004.....	45.5	*	*	0.2	0.3	0.5	1.7	8.5	43.8	180.6	462.4	837.7
2003.....	47.4	1.3	*	0.2	0.4	0.5	1.7	8.9	47.6	191.6	478.5	894.4
2002.....	47.8	*	0.3	0.2	0.4	0.6	1.8	8.8	46.0	192.3	495.2	923.4
2001.....	47.6	*	*	0.2	0.3	0.5	1.7	8.6	48.0	198.3	489.4	943.6
2000.....	48.3	*	*	0.2	0.4	0.5	1.6	8.4	48.6	201.4	503.6	997.4
1999.....	50.6	*	*	0.3	0.4	0.5	1.5	8.1	53.1	217.3	525.4	1,029.4

White, female

2004.....	49.3	*	*	0.1	0.2	0.4	1.9	7.9	40.8	151.1	335.5	536.5
2003.....	51.5	*	*	0.1	0.3	0.6	2.1	8.1	42.9	158.6	352.0	562.8
2002.....	50.5	*	*	0.2	0.3	0.6	2.0	8.1	42.4	157.0	345.4	554.5
2001.....	50.3	*	*	0.1	0.3	0.5	1.9	8.0	44.1	160.4	338.3	557.9
2000.....	49.7	*	*	0.2	0.3	0.5	1.8	7.9	43.2	159.6	339.1	544.8
1999.....	49.8	*	*	0.2	0.4	0.6	1.8	8.1	45.6	160.3	341.5	529.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Chronic lower respiratory diseases (J40-J47)

Black, both sexes

2004.....	19.2	*	0.8	1.0	1.2	1.8	3.9	12.6	38.1	102.1	222.8	333.9
2003.....	20.2	*	1.0	0.9	1.3	1.9	4.3	12.5	40.5	113.2	235.9	353.0
2002.....	20.7	*	1.2	1.1	1.3	1.9	5.1	12.9	40.7	113.5	248.7	368.2
2001.....	20.4	*	0.9	0.8	1.3	2.4	4.8	12.5	40.7	118.9	234.1	375.8
2000.....	20.8	*	0.8	1.0	1.4	2.1	4.8	14.2	43.9	121.7	234.7	373.8
1999.....	21.9	*	*	1.0	1.3	2.6	4.6	13.8	45.8	128.5	262.6	376.2

Black, male

2004.....	22.0	*	*	1.3	1.5	2.5	4.0	13.9	50.1	140.4	336.3	566.3
2003.....	23.3	*	1.6	1.2	1.5	2.8	4.0	13.3	50.5	155.1	382.2	601.6
2002.....	24.1	*	1.6	1.2	1.9	2.2	5.7	14.4	52.3	158.0	392.2	645.4
2001.....	23.6	*	*	0.9	1.6	2.7	4.7	13.3	49.8	168.0	380.8	671.7
2000.....	24.3	*	*	1.2	1.7	2.8	4.8	15.0	54.6	176.9	370.3	693.1
1999.....	26.2	*	*	1.4	1.8	2.7	4.7	15.3	59.3	184.6	434.4	701.9

Black, female

2004.....	16.6	*	*	0.6	1.0	1.2	3.9	11.5	28.5	74.9	158.6	241.8
2003.....	17.3	*	*	0.6	1.1	1.1	4.6	11.9	32.4	83.3	153.2	256.4
2002.....	17.7	*	*	1.0	0.6	1.6	4.6	11.6	31.5	82.0	167.4	262.0
2001.....	17.5	*	*	0.7	1.0	2.1	4.9	11.7	33.3	84.3	151.7	266.1
2000.....	17.6	*	*	0.8	1.0	1.5	4.7	13.4	35.3	82.9	158.4	255.0
1999.....	18.0	*	*	0.6	0.9	2.6	4.6	12.6	34.9	88.9	166.4	254.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Bronchitis, chronic and unspecified (J40-J42)

All races 3/, both sexes

2004.....	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.5	6.5
2003.....	0.3	0.5	*	*	*	*	*	0.1	0.2	0.6	1.9	7.3
2002.....	0.3	0.6	*	*	*	*	*	0.1	0.2	0.7	2.2	8.1
2001.....	0.3	0.5	*	*	*	*	0.0	0.1	0.2	0.7	2.1	9.1
2000.....	0.4	*	*	*	*	*	0.0	0.1	0.3	0.9	2.7	11.8
1999.....	0.4	0.5	*	*	*	*	*	0.1	0.4	0.9	2.8	10.7

All races, male

2004.....	0.2	*	*	*	*	*	*	0.1	0.3	0.6	1.6	7.0
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	2.2	7.0
2002.....	0.3	*	*	*	*	*	*	0.1	0.2	0.9	2.5	8.8
2001.....	0.3	*	*	*	*	*	*	0.1	0.3	0.9	2.3	9.2
2000.....	0.4	*	*	*	*	*	*	*	0.3	1.0	3.6	12.9
1999.....	0.4	*	*	*	*	*	*	*	0.4	1.1	3.3	11.9

All races, female

2004.....	0.3	*	*	*	*	*	*	*	0.2	0.4	1.5	6.3
2003.....	0.3	*	*	*	*	*	*	*	0.1	0.6	1.7	7.5
2002.....	0.4	*	*	*	*	*	*	*	0.2	0.6	2.1	7.8
2001.....	0.4	*	*	*	*	*	*	0.1	0.2	0.6	1.9	9.1
2000.....	0.5	*	*	*	*	*	*	0.1	0.3	0.8	2.1	11.4
1999.....	0.5	*	*	*	*	*	*	*	0.4	0.8	2.5	10.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bronchitis, chronic and unspecified (J40-J42)												
White, both sexes												
2004.....	0.3	*	*	*	*	*	*	0.1	0.2	0.6	1.6	6.7
2003.....	0.3	*	*	*	*	*	*	0.1	0.2	0.6	2.0	7.5
2002.....	0.4	*	*	*	*	*	*	0.1	0.2	0.8	2.3	8.6
2001.....	0.4	*	*	*	*	*	*	0.1	0.3	0.7	2.1	9.6
2000.....	0.5	*	*	*	*	*	*	0.1	0.3	0.9	2.8	12.7
1999.....	0.5	*	*	*	*	*	*	0.1	0.4	1.0	2.8	11.2
White, male												
2004.....	0.3	*	*	*	*	*	*	*	0.2	0.7	1.7	7.1
2003.....	0.3	*	*	*	*	*	*	*	0.2	0.5	2.2	7.3
2002.....	0.3	*	*	*	*	*	*	0.1	0.2	0.9	2.6	9.3
2001.....	0.3	*	*	*	*	*	*	*	0.3	0.9	2.3	9.4
2000.....	0.4	*	*	*	*	*	*	*	0.3	1.0	3.7	13.8
1999.....	0.4	*	*	*	*	*	*	*	0.5	1.2	3.2	12.4
White, female												
2004.....	0.3	*	*	*	*	*	*	*	0.2	0.5	1.6	6.5
2003.....	0.4	*	*	*	*	*	*	*	*	0.6	1.8	7.6
2002.....	0.4	*	*	*	*	*	*	*	0.2	0.7	2.2	8.3
2001.....	0.4	*	*	*	*	*	*	*	0.2	0.6	2.0	9.7
2000.....	0.5	*	*	*	*	*	*	*	0.3	0.8	2.2	12.2
1999.....	0.5	*	*	*	*	*	*	*	0.4	0.8	2.5	10.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Emphysema (J43)												
All races 3/, both sexes												
2004.....	4.6	*	*	*	*	*	0.2	1.0	5.4	20.5	41.0	49.9
2003.....	5.1	*	*	*	*	*	0.2	1.1	6.3	22.2	46.0	54.6
2002.....	5.4	*	*	*	*	*	0.2	1.1	6.4	24.0	48.0	59.3
2001.....	5.7	*	*	*	*	*	0.2	1.1	7.2	26.2	50.4	62.1
2000.....	5.9	*	*	*	*	*	0.2	1.3	7.4	27.8	52.7	63.3
1999.....	6.4	*	*	*	*	*	0.2	1.3	8.5	30.7	56.9	62.1
All races, male												
2004.....	4.9	*	*	*	*	*	0.2	1.2	6.2	24.3	52.2	74.4
2003.....	5.3	*	*	*	*	*	0.2	1.4	6.9	25.6	58.1	79.2
2002.....	5.7	*	*	*	*	*	0.3	1.4	7.2	28.1	63.5	90.0
2001.....	6.0	*	*	*	*	*	0.2	1.4	7.9	30.8	65.3	101.2
2000.....	6.3	*	*	*	*	*	0.2	1.5	8.1	33.0	69.1	104.2
1999.....	6.9	*	*	*	*	*	0.2	1.5	9.5	37.8	76.4	105.4
All races, female												
2004.....	4.4	*	*	*	*	*	0.1	0.8	4.7	17.4	33.5	38.9
2003.....	4.9	*	*	*	*	*	0.2	0.8	5.7	19.2	37.9	43.7
2002.....	5.0	*	*	*	*	*	0.1	0.9	5.6	20.6	37.6	46.0
2001.....	5.4	*	*	*	*	*	0.1	0.9	6.6	22.5	40.6	45.8
2000.....	5.6	*	*	*	*	*	0.2	1.0	6.8	23.6	41.9	46.7
1999.....	5.8	*	*	*	*	*	0.2	1.1	7.5	24.9	44.2	44.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Emphysema (J43)												
White, both sexes												
2004.....	5.4	*	*	*	*	*	0.2	1.0	5.8	22.3	43.7	52.5
2003.....	5.9	*	*	*	*	*	0.2	1.1	6.7	23.9	48.5	57.0
2002.....	6.2	*	*	*	*	*	0.2	1.2	6.9	25.7	50.7	62.3
2001.....	6.6	*	*	*	*	*	0.2	1.2	7.7	28.2	53.5	65.1
2000.....	6.8	*	*	*	*	*	0.2	1.2	7.9	29.7	55.6	65.8
1999.....	7.3	*	*	*	*	*	0.2	1.3	9.1	32.9	59.8	64.7
White, male												
2004.....	5.5	*	*	*	*	*	0.2	1.1	6.4	25.8	54.5	77.1
2003.....	5.9	*	*	*	*	*	0.2	1.4	7.2	27.1	59.8	80.4
2002.....	6.4	*	*	*	*	*	0.3	1.3	7.5	29.3	65.8	93.0
2001.....	6.8	*	*	*	*	*	0.2	1.4	8.2	32.3	68.1	105.1
2000.....	7.0	*	*	*	*	*	0.2	1.4	8.4	34.3	72.1	105.5
1999.....	7.8	*	*	*	*	*	0.2	1.4	9.9	39.4	79.0	109.2
White, female												
2004.....	5.2	*	*	*	*	*	0.1	0.8	5.2	19.3	36.3	41.4
2003.....	5.8	*	*	*	*	*	0.2	0.9	6.2	21.2	40.8	46.6
2002.....	5.9	*	*	*	*	*	*	1.0	6.2	22.7	40.5	48.9
2001.....	6.3	*	*	*	*	*	0.1	0.9	7.2	24.7	43.7	48.4
2000.....	6.5	*	*	*	*	*	0.2	1.0	7.4	25.9	44.7	49.7
1999.....	6.8	*	*	*	*	*	0.2	1.1	8.3	27.4	47.2	46.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Emphysema (J43)												
Black, both sexes												
2004.....	1.9	*	*	*	*	*	*	1.3	4.5	10.7	22.1	27.6
2003.....	2.1	*	*	*	*	*	*	1.2	5.1	12.9	27.3	30.9
2002.....	2.3	*	*	*	*	*	0.4	1.2	4.4	15.9	28.0	32.1
2001.....	2.2	*	*	*	*	*	*	1.3	5.6	15.3	23.6	35.3
2000.....	2.5	*	*	*	*	*	*	2.0	6.0	17.0	26.9	37.2
1999.....	2.6	*	*	*	*	*	0.4	1.8	6.1	17.1	31.3	37.6
Black, male												
2004.....	2.5	*	*	*	*	*	*	2.0	6.9	17.0	36.5	56.6
2003.....	2.7	*	*	*	*	*	*	1.7	7.3	18.2	46.0	70.9
2002.....	3.0	*	*	*	*	*	*	1.8	6.5	24.8	48.8	64.3
2001.....	2.7	*	*	*	*	*	*	1.6	8.0	22.6	39.7	68.1
2000.....	3.2	*	*	*	*	*	*	2.9	8.1	27.4	41.5	87.6
1999.....	3.5	*	*	*	*	*	*	2.9	8.9	29.0	54.7	72.6
Black, female												
2004.....	1.3	*	*	*	*	*	*	*	2.6	6.2	14.0	16.1
2003.....	1.6	*	*	*	*	*	*	*	3.4	9.2	16.7	15.4
2002.....	1.6	*	*	*	*	*	*	*	2.7	9.5	16.2	19.7
2001.....	1.7	*	*	*	*	*	*	1.1	3.6	10.2	14.5	23.2
2000.....	1.8	*	*	*	*	*	*	1.3	4.4	9.7	18.8	18.5
1999.....	1.7	*	*	*	*	*	*	1.0	3.9	8.7	18.2	24.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Asthma (J45-J46)												
All races 3/, both sexes												
2004.....	1.3	*	0.2	0.3	0.4	0.5	0.9	1.4	1.8	2.7	5.2	14.1
2003.....	1.4	*	0.2	0.3	0.4	0.6	0.9	1.5	2.0	2.9	5.8	14.2
2002.....	1.5	*	0.3	0.3	0.4	0.6	1.1	1.5	2.0	3.2	6.4	14.7
2001.....	1.5	*	0.2	0.2	0.4	0.6	1.0	1.5	2.2	3.5	6.4	16.1
2000.....	1.6	*	0.2	0.3	0.4	0.6	1.0	1.6	2.2	4.0	6.9	16.7
1999.....	1.7	*	0.2	0.3	0.5	0.7	1.0	1.7	2.5	4.2	7.6	17.3
All races, male												
2004.....	1.0	*	*	0.3	0.5	0.6	0.7	1.0	1.3	2.1	3.5	11.1
2003.....	1.0	*	0.4	0.3	0.4	0.6	0.7	1.2	1.6	2.1	4.5	10.7
2002.....	1.1	*	0.4	0.3	0.5	0.6	0.9	1.2	1.6	2.4	5.0	10.9
2001.....	1.1	*	0.3	0.3	0.4	0.6	0.9	1.1	1.5	2.7	4.5	12.1
2000.....	1.2	*	0.3	0.4	0.5	0.7	0.9	1.3	1.7	3.0	5.0	12.9
1999.....	1.2	*	*	0.4	0.5	0.6	0.8	1.1	1.8	3.2	5.7	12.7
All races, female												
2004.....	1.6	*	*	0.2	0.3	0.4	1.0	1.6	2.2	3.2	6.3	15.4
2003.....	1.8	*	*	0.2	0.3	0.5	1.1	1.9	2.4	3.6	6.7	15.8
2002.....	1.8	*	*	0.3	0.3	0.5	1.2	1.8	2.4	3.9	7.3	16.3
2001.....	1.9	*	*	0.2	0.3	0.6	1.1	1.9	2.8	4.1	7.6	17.8
2000.....	2.0	*	*	0.3	0.3	0.5	1.2	2.0	2.7	4.8	8.1	18.2
1999.....	2.1	*	*	0.2	0.4	0.7	1.2	2.2	3.0	5.0	8.8	19.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Asthma (J45-J46)												
White, both sexes												
2004.....	1.1	*	*	0.1	0.2	0.3	0.6	0.9	1.4	2.4	4.7	13.7
2003.....	1.2	*	*	0.2	0.2	0.4	0.7	1.1	1.6	2.4	5.4	14.3
2002.....	1.3	*	*	0.1	0.3	0.5	0.7	1.1	1.5	2.7	6.1	14.7
2001.....	1.3	*	*	0.1	0.2	0.4	0.7	1.1	1.6	3.0	6.0	15.7
2000.....	1.4	*	*	0.2	0.3	0.4	0.7	1.1	1.7	3.4	6.5	15.9
1999.....	1.5	*	*	0.2	0.3	0.4	0.7	1.2	1.9	3.5	7.2	16.6
White, male												
2004.....	0.7	*	*	0.1	0.3	0.3	0.5	0.7	0.9	1.8	2.8	10.3
2003.....	0.8	*	*	0.2	0.3	0.4	0.5	0.8	1.2	1.7	4.0	10.6
2002.....	0.9	*	*	0.2	0.3	0.5	0.5	0.8	1.2	1.9	4.6	10.7
2001.....	0.8	*	*	0.2	0.2	0.3	0.6	0.8	1.1	2.2	4.0	11.3
2000.....	0.9	*	*	0.2	0.3	0.4	0.6	1.0	1.2	2.5	4.7	11.4
1999.....	0.9	*	*	0.2	0.4	0.4	0.5	0.8	1.4	2.5	5.3	10.8
White, female												
2004.....	1.5	*	*	0.1	0.2	0.3	0.8	1.2	1.8	3.0	6.0	15.2
2003.....	1.6	*	*	0.1	0.2	0.5	0.8	1.4	2.0	3.1	6.3	15.9
2002.....	1.7	*	*	*	0.2	0.5	0.9	1.3	1.9	3.3	7.2	16.4
2001.....	1.7	*	*	*	0.2	0.4	0.8	1.5	2.1	3.7	7.3	17.6
2000.....	1.8	*	*	0.2	0.2	0.4	0.8	1.3	2.1	4.1	7.7	17.7
1999.....	2.0	*	*	0.1	0.3	0.4	0.9	1.7	2.4	4.4	8.5	18.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Asthma (J45-J46)												
Black, both sexes												
2004.....	2.6	*	*	0.8	1.1	1.6	2.5	4.5	5.6	5.7	9.1	15.5
2003.....	2.7	*	0.9	0.9	1.2	1.6	2.7	4.8	5.1	7.0	9.1	12.0
2002.....	2.9	*	0.9	1.0	1.2	1.6	3.4	4.8	6.4	7.8	8.0	11.9
2001.....	3.0	*	0.8	0.8	1.1	2.1	3.3	4.6	6.8	7.7	10.6	16.3
2000.....	3.2	*	*	1.0	1.3	1.9	3.1	5.4	6.9	9.2	10.4	22.2
1999.....	3.2	*	*	1.0	1.2	2.4	2.9	5.2	6.8	9.6	10.4	20.4
Black, male												
2004.....	2.5	*	*	1.1	1.4	2.3	2.3	3.9	5.2	5.5	8.6	*
2003.....	2.4	*	1.5	1.2	1.4	2.4	2.3	3.9	4.2	5.3	8.7	*
2002.....	2.8	*	*	1.1	1.8	2.0	3.6	3.9	5.7	7.3	6.9	*
2001.....	2.6	*	*	0.9	1.4	2.4	3.0	4.1	4.6	6.8	8.8	*
2000.....	2.8	*	*	1.2	1.5	2.5	2.9	3.8	6.1	7.9	6.4	25.4
1999.....	2.8	*	*	1.4	1.6	2.6	2.8	3.8	5.4	9.1	8.3	*
Black, female												
2004.....	2.7	*	*	*	0.9	1.1	2.7	5.0	6.0	5.7	9.4	16.9
2003.....	2.9	*	*	*	1.0	0.9	3.1	5.6	5.8	8.2	9.2	13.8
2002.....	3.0	*	*	1.0	*	1.2	3.3	5.6	7.0	8.2	8.6	12.3
2001.....	3.3	*	*	0.7	0.9	1.9	3.5	5.0	8.5	8.3	11.6	16.8
2000.....	3.5	*	*	0.8	1.0	1.4	3.3	6.8	7.5	10.2	12.6	21.0
1999.....	3.5	*	*	0.6	0.9	2.1	3.1	6.4	7.9	10.0	11.5	21.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over
Other chronic lower respiratory diseases (J44,J47)												
All races 3/, both sexes												
2004.....	35.3	*	*	*	*	0.1	0.9	6.0	33.0	130.0	319.0	531.1
2003.....	36.6	*	*	*	0.1	0.1	1.0	5.9	34.8	137.5	329.2	558.9
2002.....	36.1	*	*	*	*	0.1	0.9	5.9	33.8	135.1	330.1	555.5
2001.....	35.7	*	*	*	0.1	0.1	0.9	5.7	34.5	137.5	321.0	557.4
2000.....	35.4	*	*	*	*	0.1	0.8	5.6	34.3	136.7	323.8	556.7
1999.....	36.0	*	*	*	*	0.1	0.7	5.5	36.2	141.4	330.6	555.8
All races, male												
2004.....	34.5	*	*	*	*	0.1	1.0	6.5	35.2	145.0	388.4	718.6
2003.....	35.9	*	*	*	*	0.1	1.0	6.4	37.8	155.4	400.1	769.0
2002.....	35.7	*	*	*	*	0.1	1.0	6.4	36.2	153.4	409.8	785.1
2001.....	35.3	*	*	*	*	0.1	0.9	6.2	37.2	157.0	403.0	794.5
2000.....	35.6	*	*	*	*	*	0.8	6.1	37.7	158.3	410.9	838.0
1999.....	37.1	*	*	*	*	*	0.8	5.9	40.5	168.6	427.9	866.7
All races, female												
2004.....	36.2	*	*	*	*	*	0.9	5.5	30.8	117.3	272.2	446.8
2003.....	37.4	*	*	*	*	*	1.0	5.5	32.1	122.6	281.9	466.0
2002.....	36.5	*	*	*	*	0.1	0.9	5.4	31.5	119.8	277.2	455.8
2001.....	36.0	*	*	*	*	0.1	1.0	5.2	32.0	121.3	267.1	458.2
2000.....	35.2	*	*	*	*	*	0.8	5.2	31.2	119.0	267.1	442.2
1999.....	35.0	*	*	*	*	0.2	0.7	5.0	32.2	119.1	267.4	430.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)												
White, both sexes												
2004.....	40.6	*	*	*	*	0.1	1.0	6.2	34.9	139.4	336.9	557.2
2003.....	42.1	*	*	*	*	0.1	1.0	6.2	36.7	146.9	347.2	585.8
2002.....	41.3	*	*	*	*	0.1	1.0	6.1	35.6	144.0	346.4	580.7
2001.....	40.8	*	*	*	*	0.1	0.9	5.8	36.5	145.9	337.0	581.3
2000.....	40.4	*	*	*	*	*	0.8	5.7	36.0	144.7	339.5	581.2
1999.....	41.0	*	*	*	*	0.1	0.7	5.6	37.8	148.8	344.5	580.3
White, male												
2004.....	39.0	*	*	*	*	0.1	1.0	6.6	36.3	152.3	403.4	743.2
2003.....	40.4	*	*	*	*	*	0.9	6.6	39.0	162.2	412.5	796.0
2002.....	40.2	*	*	*	*	*	1.0	6.5	37.1	160.2	422.1	810.4
2001.....	39.7	*	*	*	*	0.1	0.9	6.3	38.5	162.9	415.0	817.8
2000.....	39.9	*	*	*	*	*	0.8	6.0	38.7	163.6	423.0	866.7
1999.....	41.5	*	*	*	*	*	0.7	5.9	41.4	174.2	438.0	897.0
White, female												
2004.....	42.2	*	*	*	*	*	0.9	5.8	33.6	128.3	291.6	473.4
2003.....	43.7	*	*	*	*	*	1.0	5.8	34.5	133.8	303.0	492.7
2002.....	42.5	*	*	*	*	*	0.9	5.7	34.1	130.3	295.6	480.9
2001.....	41.8	*	*	*	*	*	1.0	5.4	34.6	131.5	285.3	482.3
2000.....	40.9	*	*	*	*	*	0.8	5.5	33.4	128.9	284.4	465.2
1999.....	40.4	*	*	*	*	0.1	0.7	5.2	34.5	127.7	283.2	453.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)												
Black, both sexes												
2004.....	14.5	*	*	*	*	*	1.1	6.7	27.7	85.5	190.7	285.9
2003.....	15.2	*	*	*	*	*	1.2	6.3	29.9	92.9	198.3	304.6
2002.....	15.4	*	*	*	*	*	1.2	6.7	29.7	89.5	211.3	320.1
2001.....	15.1	*	*	*	*	*	1.3	6.4	28.1	95.0	198.7	320.9
2000.....	14.9	*	*	*	*	*	1.3	6.6	30.9	94.5	195.2	310.6
1999.....	15.9	*	*	*	*	*	1.2	6.4	32.5	100.8	218.0	313.7
Black, male												
2004.....	16.8	*	*	*	*	*	1.2	7.7	37.8	117.4	290.6	492.7
2003.....	18.0	*	*	*	*	*	1.3	7.4	38.7	131.3	325.5	520.3
2002.....	18.2	*	*	*	*	*	1.4	8.3	39.9	125.3	335.1	565.0
2001.....	18.1	*	*	*	*	*	1.4	7.4	37.0	137.4	330.5	583.2
2000.....	18.2	*	*	*	*	*	1.3	8.3	40.1	140.8	320.4	576.6
1999.....	19.6	*	*	*	*	*	1.5	8.2	44.6	145.4	366.5	607.0
Black, female												
2004.....	12.4	*	*	*	*	*	1.0	5.8	19.5	62.7	134.2	204.0
2003.....	12.6	*	*	*	*	*	1.2	5.3	22.9	65.5	126.5	220.8
2002.....	12.9	*	*	*	*	*	1.1	5.3	21.5	64.0	141.3	226.3
2001.....	12.3	*	*	*	*	*	1.2	5.6	21.0	65.1	124.8	223.5
2000.....	12.0	*	*	*	*	*	1.3	5.1	23.4	62.0	124.8	211.6
1999.....	12.5	*	*	*	*	*	1.0	4.9	22.8	69.4	134.9	204.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)												
All races 3/, both sexes												
2004.....	5.7	*	*	*	0.1	0.2	0.4	0.9	2.6	9.6	42.9	164.9
2003.....	6.0	*	*	*	0.1	0.2	0.4	1.0	2.8	9.6	45.0	176.2
2002.....	6.1	*	*	*	0.1	0.2	0.4	0.9	2.5	9.8	46.3	186.0
2001.....	6.1	*	*	*	0.1	0.2	0.4	1.0	2.6	10.0	45.8	189.4
2000.....	5.9	*	*	*	0.1	0.2	0.4	1.0	2.5	10.3	44.5	187.6
1999.....	5.5	*	*	*	0.1	0.2	0.4	0.8	2.5	9.5	41.1	175.6
All races, male												
2004.....	5.9	*	*	*	0.1	0.2	0.4	1.2	3.2	12.9	58.6	237.2
2003.....	6.1	*	*	*	0.1	0.2	0.4	1.2	3.5	12.8	61.7	246.2
2002.....	6.3	*	*	*	0.1	0.2	0.5	1.1	3.1	13.0	63.8	268.5
2001.....	6.1	*	*	*	0.1	0.2	0.5	1.2	3.2	13.3	63.1	270.3
2000.....	6.1	*	*	*	0.1	0.2	0.5	1.2	3.0	14.4	62.3	276.0
1999.....	5.7	*	*	*	0.1	0.2	0.4	0.9	3.2	13.1	59.7	258.4
All races, female												
2004.....	5.5	*	*	*	*	0.1	0.3	0.7	1.9	6.8	32.3	132.4
2003.....	5.8	*	*	*	*	0.1	0.3	0.7	2.1	6.9	33.8	145.3
2002.....	5.9	*	*	*	*	0.2	0.3	0.6	2.0	7.1	34.7	150.2
2001.....	6.0	*	*	*	*	0.2	0.3	0.7	2.1	7.2	34.5	155.6
2000.....	5.8	*	*	*	*	0.1	0.3	0.7	1.9	7.0	33.0	151.6
1999.....	5.3	*	*	*	*	0.2	0.3	0.6	1.8	6.6	29.1	142.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)												
White, both sexes												
2004.....	6.3	*	*	*	0.1	0.2	0.3	0.9	2.4	9.4	43.2	167.8
2003.....	6.5	*	*	*	0.1	0.2	0.3	1.0	2.6	9.3	44.7	178.7
2002.....	6.7	*	*	*	0.1	0.2	0.4	0.8	2.4	9.5	46.5	189.5
2001.....	6.6	*	*	*	0.1	0.2	0.4	0.9	2.5	9.6	45.8	190.8
2000.....	6.5	*	*	*	0.1	0.1	0.4	0.9	2.3	9.8	44.4	191.0
1999.....	5.9	*	*	*	0.1	0.2	0.3	0.7	2.3	9.0	40.7	178.0
White, male												
2004.....	6.5	*	*	*	*	0.2	0.4	1.2	2.9	12.4	58.8	241.7
2003.....	6.6	*	*	*	0.1	0.2	0.4	1.2	3.2	12.4	61.1	250.8
2002.....	6.9	*	*	*	0.1	0.2	0.5	1.1	3.0	12.5	64.0	274.6
2001.....	6.7	*	*	*	*	0.2	0.5	1.2	3.1	12.9	63.3	275.4
2000.....	6.6	*	*	*	*	0.1	0.5	1.1	2.9	13.4	62.1	284.0
1999.....	6.1	*	*	*	*	0.2	0.4	0.8	2.9	12.2	58.7	263.6
White, female												
2004.....	6.1	*	*	*	*	0.1	0.3	0.6	1.9	6.8	32.6	134.5
2003.....	6.4	*	*	*	*	0.1	0.3	0.7	2.0	6.7	33.6	146.7
2002.....	6.6	*	*	*	*	0.2	0.3	0.6	1.9	6.9	34.8	152.5
2001.....	6.6	*	*	*	*	0.2	0.3	0.7	2.1	6.9	34.2	155.3
2000.....	6.3	*	*	*	*	0.2	0.3	0.7	1.8	6.8	32.8	153.2
1999.....	5.8	*	*	*	*	0.2	0.3	0.6	1.7	6.4	29.0	143.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)												
Black, both sexes												
2004.....	3.9	*	*	*	*	*	0.6	1.3	4.8	13.9	46.9	144.7
2003.....	4.2	*	*	*	*	*	0.7	1.3	5.1	13.7	54.4	163.4
2002.....	4.1	*	*	*	*	*	0.7	1.4	3.7	14.8	52.2	165.1
2001.....	4.3	*	*	*	*	*	0.6	1.6	4.0	15.1	53.2	185.5
2000.....	4.2	*	*	*	*	*	0.7	1.9	4.2	17.5	51.5	159.8
1999.....	4.2	*	*	*	*	*	0.7	1.5	4.4	16.2	52.2	167.1
Black, male												
2004.....	4.2	*	*	*	*	*	0.7	1.7	7.1	21.2	67.7	202.7
2003.....	4.3	*	*	*	*	*	*	1.6	7.1	19.6	77.8	212.7
2002.....	4.2	*	*	*	*	*	0.8	2.0	4.8	21.7	73.4	233.7
2001.....	4.2	*	*	*	*	*	0.7	1.9	5.3	20.4	75.3	243.9
2000.....	4.4	*	*	*	*	*	0.8	2.7	5.0	27.8	74.5	209.9
1999.....	4.5	*	*	*	*	*	0.9	1.8	6.4	25.0	80.9	228.5
Black, female												
2004.....	3.6	*	*	*	*	*	*	1.0	3.0	8.7	35.2	121.7
2003.....	4.1	*	*	*	*	*	0.8	1.0	3.6	9.6	41.1	144.2
2002.....	3.9	*	*	*	*	*	*	0.9	2.9	9.9	40.1	138.8
2001.....	4.4	*	*	*	*	*	*	1.4	2.9	11.3	40.8	163.8
2000.....	3.9	*	*	*	*	*	*	1.2	3.5	10.3	38.5	141.2
1999.....	3.8	*	*	*	*	*	*	1.2	2.8	10.0	36.1	144.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

All races 3/, both sexes

2004.....	9.3	6.5	0.7	0.2	0.3	0.4	1.2	3.2	9.5	29.9	74.2	139.4
2003.....	8.8	7.0	0.5	0.1	0.3	0.5	1.1	3.3	8.8	28.1	68.5	138.1
2002.....	8.7	6.9	0.5	0.2	0.3	0.5	1.2	3.2	9.2	28.7	68.0	133.5
2001.....	9.2	7.3	0.6	0.2	0.4	0.5	1.3	3.5	9.6	30.1	73.2	142.7
2000.....	9.1	7.2	0.6	0.2	0.3	0.6	1.4	3.5	9.9	29.6	72.0	141.7
1999.....	9.0	6.6	0.6	0.2	0.3	0.6	1.3	3.3	9.5	29.4	70.7	147.6

All races, male

2004.....	9.3	6.6	0.7	0.2	0.3	0.4	1.2	3.5	10.8	36.9	93.9	168.8
2003.....	8.7	8.0	0.6	0.2	0.4	0.6	1.1	3.5	10.2	34.1	85.3	165.0
2002.....	8.5	7.9	0.7	0.2	0.3	0.4	1.3	3.3	10.5	35.3	83.9	156.6
2001.....	8.9	9.0	0.6	0.2	0.4	0.5	1.4	3.8	10.9	36.5	89.3	169.1
2000.....	9.0	7.9	0.6	0.2	0.4	0.6	1.5	3.8	11.4	35.7	90.5	179.7
1999.....	8.9	7.5	0.7	0.2	0.3	0.6	1.4	3.5	10.6	36.2	89.9	182.3

All races, female

2004.....	9.3	6.4	0.6	0.1	0.3	0.4	1.2	3.0	8.4	24.1	61.0	126.1
2003.....	8.9	5.9	0.5	0.1	0.3	0.5	1.1	3.0	7.4	23.1	57.2	126.2
2002.....	8.8	5.8	0.4	0.2	0.3	0.5	1.1	3.1	8.0	23.1	57.5	123.4
2001.....	9.4	5.5	0.6	0.2	0.3	0.6	1.2	3.1	8.4	24.8	62.6	131.6
2000.....	9.2	6.4	0.6	0.2	0.3	0.6	1.2	3.2	8.6	24.7	59.9	126.2
1999.....	9.1	5.8	0.5	0.2	0.2	0.7	1.1	3.1	8.5	23.7	58.3	133.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

White, both sexes

2004.....	10.2	4.9	0.5	0.1	0.3	0.4	1.1	2.9	9.3	30.7	76.8	144.1
2003.....	9.6	5.4	0.4	0.1	0.3	0.5	0.9	2.9	8.6	29.0	70.8	143.2
2002.....	9.5	4.8	0.5	0.2	0.3	0.4	1.0	2.9	8.9	29.3	69.9	137.9
2001.....	10.0	5.0	0.5	0.2	0.3	0.4	1.1	3.0	9.3	30.6	75.4	146.0
2000.....	9.9	5.6	0.5	0.2	0.3	0.6	1.2	3.2	9.7	30.0	73.7	145.9
1999.....	9.7	4.9	0.5	0.2	0.3	0.6	1.1	3.0	9.2	29.6	72.2	150.7

White, male

2004.....	10.1	4.9	0.6	0.1	0.3	0.4	1.0	3.2	10.6	37.8	96.7	174.3
2003.....	9.4	5.8	0.4	0.2	0.3	0.5	1.0	3.3	9.9	35.1	87.6	170.5
2002.....	9.2	5.4	0.6	0.2	0.3	0.4	1.1	3.0	10.1	36.1	86.1	162.0
2001.....	9.7	5.9	0.5	0.2	0.3	0.4	1.2	3.4	10.6	37.4	92.0	173.5
2000.....	9.7	6.6	0.5	0.2	0.3	0.6	1.3	3.4	11.1	36.1	92.5	184.4
1999.....	9.6	5.6	0.6	0.2	0.4	0.6	1.2	3.2	10.3	36.8	92.0	185.7

White, female

2004.....	10.2	4.9	0.4	*	0.3	0.4	1.1	2.6	8.1	24.7	63.2	130.5
2003.....	9.8	5.0	0.4	*	0.3	0.4	0.9	2.6	7.2	23.8	59.4	131.0
2002.....	9.7	4.2	*	0.2	0.2	0.4	1.0	2.8	7.7	23.4	59.1	127.4
2001.....	10.2	4.0	0.6	0.2	0.3	0.5	0.9	2.6	8.2	24.9	64.3	134.4
2000.....	10.0	4.6	0.6	0.2	0.3	0.6	1.1	2.9	8.4	25.0	61.4	130.3
1999.....	9.8	4.3	0.5	0.2	0.2	0.5	0.9	2.8	8.2	23.6	59.3	136.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Black, both sexes

2004.....	6.2	14.8	1.2	*	0.4	0.9	2.6	6.2	13.4	27.5	53.2	92.8
2003.....	5.9	15.0	1.1	*	0.6	1.0	2.4	6.4	12.2	24.2	50.8	89.6
2002.....	6.0	16.6	1.1	0.4	0.7	0.9	2.3	6.1	13.6	26.5	51.5	90.3
2001.....	6.7	18.0	1.0	*	0.6	1.3	2.9	7.3	13.6	29.5	56.7	109.6
2000.....	6.5	16.7	0.8	*	0.7	1.1	2.7	6.8	13.4	30.5	58.0	98.2
1999.....	6.5	15.4	1.0	*	0.4	1.3	2.9	6.4	13.4	30.8	57.9	111.6

Black, male

2004.....	6.3	15.7	*	*	*	1.0	3.0	6.7	15.2	34.5	69.0	115.2
2003.....	6.0	18.4	*	*	0.8	1.1	2.2	6.3	15.1	30.8	68.5	113.6
2002.....	6.0	19.5	*	*	*	0.9	2.7	6.5	16.1	31.7	65.7	100.8
2001.....	6.4	23.1	*	*	*	1.3	2.6	7.6	15.3	32.9	69.4	129.3
2000.....	6.5	16.3	*	*	0.8	1.2	3.1	7.8	15.2	36.0	73.0	132.6
1999.....	6.4	17.6	*	*	*	1.0	2.9	7.1	15.5	35.1	72.6	145.3

Black, female

2004.....	6.1	13.9	*	*	*	0.8	2.2	5.8	11.9	22.4	44.2	83.9
2003.....	5.7	11.4	*	*	*	1.0	2.7	6.5	9.8	19.4	40.8	80.2
2002.....	6.0	13.6	*	*	0.7	0.8	2.0	5.8	11.5	22.9	43.5	86.3
2001.....	7.0	12.6	*	*	*	1.2	3.1	7.0	12.2	27.1	49.6	102.3
2000.....	6.4	17.2	*	*	*	1.0	2.2	5.9	11.9	26.6	49.6	85.4
1999.....	6.7	13.1	*	*	*	1.6	2.8	5.8	11.6	27.8	49.7	99.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Peptic ulcer (K25-K28)												
All races 3/, both sexes												
2004.....	1.3	*	*	*	*	*	0.2	0.6	1.3	3.2	8.9	24.4
2003.....	1.3	*	*	*	*	0.1	0.2	0.7	1.4	3.2	9.9	26.4
2002.....	1.4	*	*	*	*	0.1	0.3	0.7	1.5	3.6	10.1	28.3
2001.....	1.6	*	*	*	*	0.1	0.3	0.7	1.7	3.9	11.4	33.4
2000.....	1.6	*	*	*	*	*	0.3	0.8	1.8	4.1	11.4	35.6
1999.....	1.6	*	*	*	*	0.1	0.3	0.7	1.8	4.5	12.3	34.6
All races, male												
2004.....	1.2	*	*	*	*	*	0.3	0.9	1.6	3.6	10.2	24.6
2003.....	1.3	*	*	*	*	*	0.3	1.0	1.8	4.1	11.8	28.9
2002.....	1.4	*	*	*	*	*	0.3	0.9	1.9	4.6	12.2	31.6
2001.....	1.5	*	*	*	*	0.1	0.4	1.0	2.2	4.8	13.0	34.9
2000.....	1.6	*	*	*	*	*	0.4	1.2	2.4	5.3	13.7	40.7
1999.....	1.6	*	*	*	*	*	0.3	0.9	2.3	6.1	14.8	37.5
All races, female												
2004.....	1.3	*	*	*	*	*	0.1	0.4	1.0	2.8	8.0	24.3
2003.....	1.4	*	*	*	*	*	0.2	0.4	1.0	2.4	8.7	25.3
2002.....	1.5	*	*	*	*	*	0.2	0.5	1.2	2.9	8.7	26.9
2001.....	1.7	*	*	*	*	*	0.1	0.5	1.1	3.2	10.4	32.7
2000.....	1.6	*	*	*	*	*	0.1	0.5	1.3	3.0	10.0	33.5
1999.....	1.7	*	*	*	*	*	0.2	0.5	1.3	3.3	10.6	33.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Peptic ulcer (K25-K28)												
White, both sexes												
2004.....	1.3	*	*	*	*	*	0.2	0.6	1.2	3.1	8.9	25.1
2003.....	1.4	*	*	*	*	0.1	0.2	0.6	1.3	3.2	9.8	26.9
2002.....	1.5	*	*	*	*	*	0.2	0.6	1.4	3.6	10.3	29.1
2001.....	1.7	*	*	*	*	*	0.2	0.7	1.5	3.9	11.5	34.4
2000.....	1.7	*	*	*	*	*	0.2	0.7	1.8	4.0	11.5	36.3
1999.....	1.8	*	*	*	*	*	0.2	0.6	1.7	4.4	12.3	35.3
White, male												
2004.....	1.2	*	*	*	*	*	0.2	0.8	1.6	3.6	10.0	25.5
2003.....	1.4	*	*	*	*	*	0.2	0.8	1.6	4.1	11.4	29.0
2002.....	1.5	*	*	*	*	*	0.3	0.8	1.8	4.4	12.3	32.5
2001.....	1.5	*	*	*	*	*	0.3	0.8	2.0	4.5	12.9	35.7
2000.....	1.7	*	*	*	*	*	0.3	1.0	2.4	5.2	13.5	40.7
1999.....	1.7	*	*	*	*	*	0.2	0.8	2.2	5.9	14.9	38.4
White, female												
2004.....	1.5	*	*	*	*	*	0.1	0.4	1.0	2.7	8.2	25.0
2003.....	1.5	*	*	*	*	*	0.2	0.4	1.0	2.5	8.8	26.0
2002.....	1.6	*	*	*	*	*	0.1	0.5	1.1	2.9	9.0	27.6
2001.....	1.9	*	*	*	*	*	0.1	0.5	1.1	3.3	10.7	33.9
2000.....	1.8	*	*	*	*	*	0.1	0.4	1.2	3.0	10.2	34.5
1999.....	1.9	*	*	*	*	*	0.2	0.4	1.3	3.2	10.7	34.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Peptic ulcer (K25-K28)

Black, both sexes

2004.....	0.9	*	*	*	*	*	0.5	1.3	2.0	4.2	7.7	16.1
2003.....	1.0	*	*	*	*	*	0.4	1.5	2.2	3.2	11.0	16.9
2002.....	1.0	*	*	*	*	*	0.6	1.2	2.6	4.1	8.0	19.0
2001.....	1.1	*	*	*	*	*	0.6	1.3	2.7	4.4	9.1	20.9
2000.....	1.2	*	*	*	*	*	0.5	1.7	2.7	4.8	10.0	26.6
1999.....	1.2	*	*	*	*	*	0.8	1.5	2.8	5.7	10.9	25.5

Black, male

2004.....	1.0	*	*	*	*	*	*	1.8	2.9	4.6	11.1	*
2003.....	1.2	*	*	*	*	*	*	2.3	3.4	5.2	16.1	*
2002.....	1.2	*	*	*	*	*	0.8	1.9	3.8	5.9	11.5	*
2001.....	1.4	*	*	*	*	*	1.0	2.2	4.2	7.0	12.4	23.8
2000.....	1.3	*	*	*	*	*	0.8	2.3	3.7	7.0	13.9	31.1
1999.....	1.4	*	*	*	*	*	1.2	2.3	4.4	8.1	12.3	28.1

Black, female

2004.....	0.9	*	*	*	*	*	*	0.8	1.3	4.0	5.8	16.5
2003.....	0.8	*	*	*	*	*	*	0.8	*	*	8.1	15.8
2002.....	0.8	*	*	*	*	*	*	*	1.6	2.8	6.0	18.5
2001.....	0.9	*	*	*	*	*	*	*	1.5	2.7	7.3	19.8
2000.....	1.0	*	*	*	*	*	*	1.1	1.8	3.3	7.8	24.9
1999.....	1.1	*	*	*	*	*	*	*	*	3.9	10.2	24.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diseases of appendix (K35-K38)

All races 3/, both sexes

2004.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.1	1.7
2003.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.3	1.0	1.7
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.9	2.0
2001.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.7	2.1
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.0	1.6
1999.....	0.1	*	*	0.1	*	*	0.1	0.1	0.2	0.4	0.8	1.5

All races, male

2004.....	0.2	*	*	*	*	*	*	0.2	0.2	0.5	1.3	2.5
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.5	2.4
2002.....	0.2	*	*	*	*	*	*	0.2	0.4	0.7	1.3	2.2
2001.....	0.2	*	*	*	*	*	*	0.1	0.2	0.7	1.0	2.8
2000.....	0.2	*	*	*	*	*	*	0.2	0.4	0.5	1.2	2.0
1999.....	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.0	2.9

All races, female

2004.....	0.1	*	*	*	*	*	*	*	0.2	0.4	0.9	1.4
2003.....	0.1	*	*	*	*	*	*	0.1	*	*	0.7	1.3
2002.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	2.0
2001.....	0.1	*	*	*	*	*	*	*	*	0.3	0.5	1.8
2000.....	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.5
1999.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	0.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diseases of appendix (K35-K38)

White, both sexes

2004.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.0	1.8
2003.....	0.2	*	*	*	*	*	*	0.1	0.1	0.3	1.0	1.7
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.0	2.1
2001.....	0.1	*	*	*	*	*	*	0.1	0.1	0.5	0.7	2.1
2000.....	0.2	*	*	*	*	*	*	0.1	0.2	0.3	0.9	1.7
1999.....	0.1	*	*	*	*	*	*	0.1	0.1	0.4	0.8	1.5

White, male

2004.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.3	2.3
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	2.5
2002.....	0.2	*	*	*	*	*	*	0.2	0.4	0.6	1.4	2.0
2001.....	0.2	*	*	*	*	*	*	0.1	0.2	0.7	1.0	2.5
2000.....	0.2	*	*	*	*	*	*	*	0.3	0.4	1.2	2.0
1999.....	0.2	*	*	*	*	*	*	0.2	0.2	0.4	0.9	3.1

White, female

2004.....	0.1	*	*	*	*	*	*	*	*	0.4	0.8	1.6
2003.....	0.1	*	*	*	*	*	*	*	*	*	0.7	1.3
2002.....	0.2	*	*	*	*	*	*	*	*	0.3	0.7	2.1
2001.....	0.1	*	*	*	*	*	*	*	*	0.3	0.5	1.9
2000.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.6
1999.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	0.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Hernia (K40-K46)

All races 3/, both sexes

2004.....	0.5	1.0	*	*	*	*	0.1	0.2	0.4	1.3	3.9	11.0
2003.....	0.6	0.9	*	*	*	*	0.1	0.3	0.6	1.3	3.6	11.9
2002.....	0.6	0.7	*	*	*	*	0.1	0.2	0.5	1.2	3.8	12.8
2001.....	0.5	1.1	*	*	*	*	0.1	0.2	0.5	1.2	3.5	12.6
2000.....	0.5	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.8	13.1
1999.....	0.5	1.2	*	*	*	*	0.1	0.2	0.4	1.2	3.7	12.7

All races, male

2004.....	0.5	1.0	*	*	*	*	0.1	0.2	0.4	1.4	3.8	11.3
2003.....	0.5	1.0	*	*	*	*	*	0.3	0.5	1.3	3.9	12.3
2002.....	0.4	*	*	*	*	*	*	0.2	0.4	1.2	3.7	11.7
2001.....	0.4	1.4	*	*	*	*	*	0.2	0.5	1.1	3.5	12.0
2000.....	0.4	1.4	*	*	*	*	*	0.2	0.4	1.1	4.0	12.6
1999.....	0.4	1.3	*	*	*	*	0.1	0.2	0.5	1.1	4.0	13.2

All races, female

2004.....	0.6	1.0	*	*	*	*	*	0.2	0.5	1.2	4.0	10.8
2003.....	0.6	*	*	*	*	*	*	0.2	0.6	1.3	3.4	11.8
2002.....	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.9	13.3
2001.....	0.6	*	*	*	*	*	*	0.2	0.5	1.2	3.6	12.8
2000.....	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.6	13.3
1999.....	0.6	*	*	*	*	*	*	0.1	0.4	1.2	3.5	12.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hernia (K40-K46)												
White, both sexes												
2004.....	0.6	1.0	*	*	*	*	0.1	0.2	0.4	1.3	4.1	11.2
2003.....	0.6	0.9	*	*	*	*	0.1	0.3	0.5	1.3	3.6	12.3
2002.....	0.6	0.8	*	*	*	*	0.1	0.2	0.4	1.2	4.0	13.4
2001.....	0.6	1.0	*	*	*	*	0.1	0.2	0.5	1.2	3.5	12.8
2000.....	0.6	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.8	13.3
1999.....	0.6	1.1	*	*	*	*	0.1	0.1	0.5	1.1	3.7	13.0
White, male												
2004.....	0.5	*	*	*	*	*	*	0.2	0.4	1.4	3.9	11.4
2003.....	0.5	*	*	*	*	*	*	0.3	0.5	1.3	3.9	12.5
2002.....	0.5	*	*	*	*	*	*	0.2	0.4	1.2	3.9	11.9
2001.....	0.4	*	*	*	*	*	*	0.2	0.5	1.2	3.4	11.8
2000.....	0.5	1.5	*	*	*	*	*	0.2	0.4	1.1	4.1	12.7
1999.....	0.5	1.3	*	*	*	*	0.1	0.2	0.6	1.2	4.1	12.9
White, female												
2004.....	0.7	*	*	*	*	*	*	0.2	0.4	1.2	4.2	11.2
2003.....	0.7	*	*	*	*	*	*	0.2	0.6	1.3	3.4	12.2
2002.....	0.8	*	*	*	*	*	*	0.2	0.5	1.2	4.1	14.0
2001.....	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.6	13.2
2000.....	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.6	13.6
1999.....	0.7	*	*	*	*	*	*	*	0.4	1.1	3.5	13.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Chronic liver disease and cirrhosis (K70,K73-K74)

All races 3/, both sexes

2004.....	9.2	*	*	*	*	0.8	6.3	18.0	22.6	27.7	28.8	19.7
2003.....	9.5	*	*	*	*	0.9	6.8	18.3	23.0	29.5	30.0	20.1
2002.....	9.5	*	*	*	0.1	0.9	7.0	18.0	22.9	29.4	31.4	21.4
2001.....	9.5	*	*	*	0.1	1.0	7.4	18.5	22.7	30.0	30.2	22.2
2000.....	9.4	*	*	*	0.1	1.0	7.5	17.7	23.8	29.8	31.0	23.1
1999.....	9.4	*	*	*	0.1	1.0	7.3	17.4	23.7	30.6	31.9	23.2

All races, male

2004.....	12.1	*	*	*	*	1.1	8.7	26.3	32.8	37.0	36.7	25.5
2003.....	12.5	*	*	*	*	1.2	9.2	27.5	33.8	39.1	36.9	27.5
2002.....	12.3	*	*	*	*	1.2	9.3	26.9	33.1	38.5	38.4	28.4
2001.....	12.4	*	*	*	*	1.3	9.8	27.5	33.1	40.2	36.8	32.0
2000.....	12.5	*	*	*	*	1.4	10.3	26.9	34.4	39.9	38.4	34.3
1999.....	12.5	*	*	*	*	1.3	10.3	26.9	34.6	41.6	39.6	33.1

All races, female

2004.....	6.3	*	*	*	*	0.4	4.0	10.0	13.1	19.9	23.4	17.1
2003.....	6.5	*	*	*	*	0.5	4.4	9.4	13.1	21.4	25.4	16.8
2002.....	6.7	*	*	*	*	0.7	4.8	9.5	13.5	21.9	26.7	18.4
2001.....	6.7	*	*	*	*	0.7	5.0	9.8	13.1	21.5	25.9	18.1
2000.....	6.5	*	*	*	*	0.7	4.7	8.8	14.0	21.5	26.1	18.5
1999.....	6.4	*	*	*	*	0.8	4.4	8.3	13.7	21.6	26.9	19.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic liver disease and cirrhosis (K70,K73-K74)												
White, both sexes												
2004.....	9.9	*	*	*	*	0.8	6.5	18.4	22.7	28.9	30.2	20.3
2003.....	10.2	*	*	*	*	0.9	7.1	18.4	23.1	30.7	31.6	20.9
2002.....	10.1	*	*	*	0.1	0.9	7.3	18.1	23.1	30.7	32.8	22.3
2001.....	10.1	*	*	*	*	1.0	7.5	18.3	22.7	30.9	31.5	23.2
2000.....	10.0	*	*	*	0.1	1.0	7.4	17.5	23.8	30.6	32.3	23.9
1999.....	9.9	*	*	*	*	1.0	7.2	17.0	23.3	31.4	33.0	23.8
White, male												
2004.....	13.0	*	*	*	*	1.1	9.0	26.9	32.7	38.2	38.4	26.7
2003.....	13.4	*	*	*	*	1.3	9.6	27.8	33.8	40.5	38.5	28.4
2002.....	13.2	*	*	*	*	1.2	9.7	27.1	33.2	39.8	40.2	29.6
2001.....	13.1	*	*	*	*	1.3	9.9	27.3	32.8	41.1	38.4	33.5
2000.....	13.2	*	*	*	*	1.4	10.3	26.8	34.5	40.9	39.9	35.5
1999.....	13.1	*	*	*	*	1.3	10.3	26.4	34.1	42.4	40.8	34.2
White, female												
2004.....	6.8	*	*	*	*	0.5	4.0	10.1	13.1	21.0	24.6	17.4
2003.....	7.0	*	*	*	*	0.5	4.4	9.2	13.1	22.4	27.0	17.5
2002.....	7.2	*	*	*	*	0.6	4.8	9.1	13.5	23.0	27.9	19.1
2001.....	7.1	*	*	*	*	0.6	5.0	9.4	13.3	22.4	27.0	18.9
2000.....	6.8	*	*	*	*	0.6	4.5	8.2	13.7	21.9	27.2	19.2
1999.....	6.8	*	*	*	*	0.7	4.2	7.8	13.2	22.2	27.9	19.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic liver disease and cirrhosis (K70,K73-K74)												
Black, both sexes												
2004.....	6.6	*	*	*	*	0.4	5.4	18.3	25.3	21.6	14.4	12.1
2003.....	6.9	*	*	*	*	0.7	5.7	19.8	25.8	23.8	14.4	12.3
2002.....	6.9	*	*	*	*	0.9	5.5	20.8	25.0	21.6	18.5	12.2
2001.....	7.5	*	*	*	*	0.9	7.3	22.6	25.9	25.0	18.5	10.4
2000.....	7.5	*	*	*	*	1.0	7.8	22.1	25.9	26.2	18.0	13.8
1999.....	7.8	*	*	*	*	1.0	8.0	22.7	29.3	26.6	21.5	17.2
Black, male												
2004.....	9.1	*	*	*	*	*	7.4	27.1	39.1	33.8	20.6	*
2003.....	9.5	*	*	*	*	1.0	7.1	30.4	40.8	33.4	20.6	21.9
2002.....	9.0	*	*	*	*	1.0	6.7	30.0	37.8	32.4	22.4	*
2001.....	10.4	*	*	*	*	1.0	10.0	34.5	41.5	39.1	22.4	*
2000.....	9.9	*	*	*	*	1.0	10.5	32.7	38.0	37.7	24.5	*
1999.....	10.7	*	*	*	*	1.1	11.1	35.4	43.3	40.7	29.5	*
Black, female												
2004.....	4.3	*	*	*	*	*	3.7	10.8	14.2	12.9	10.9	11.4
2003.....	4.5	*	*	*	*	*	4.4	10.6	13.7	16.9	10.8	8.5
2002.....	4.9	*	*	*	*	0.8	4.4	12.9	14.7	14.0	16.2	10.7
2001.....	4.8	*	*	*	*	0.7	4.8	12.4	13.4	15.0	16.4	9.7
2000.....	5.2	*	*	*	*	1.0	5.5	13.1	16.2	18.1	14.3	11.6
1999.....	5.2	*	*	*	*	0.9	5.3	11.8	18.1	16.7	17.0	15.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcoholic liver disease (K70)												
All races 3/, both sexes												
2004.....	4.3	*	*	*	*	0.6	4.2	10.6	11.7	10.0	5.5	2.1
2003.....	4.3	*	*	*	*	0.6	4.4	10.5	11.6	9.8	5.7	1.8
2002.....	4.2	*	*	*	*	0.6	4.4	10.3	11.6	10.1	5.7	2.1
2001.....	4.3	*	*	*	*	0.7	4.8	10.7	11.5	10.1	5.7	2.4
2000.....	4.3	*	*	*	*	0.7	4.9	10.3	12.2	10.3	6.1	2.2
1999.....	4.3	*	*	*	*	0.6	4.8	10.4	12.0	10.5	6.5	2.8
All races, male												
2004.....	6.3	*	*	*	*	0.8	5.8	15.5	18.2	16.5	10.1	4.6
2003.....	6.4	*	*	*	*	0.9	6.0	16.1	18.3	16.1	10.1	3.6
2002.....	6.2	*	*	*	*	0.8	5.8	15.6	18.3	16.5	9.7	4.7
2001.....	6.4	*	*	*	*	0.9	6.5	16.1	18.4	16.6	10.1	5.5
2000.....	6.5	*	*	*	*	1.0	6.8	16.0	19.0	17.3	11.5	5.3
1999.....	6.5	*	*	*	*	0.8	6.8	16.3	19.2	18.1	11.2	6.1
All races, female												
2004.....	2.3	*	*	*	*	0.3	2.7	5.7	5.6	4.5	2.3	1.0
2003.....	2.2	*	*	*	*	0.3	2.9	5.1	5.4	4.5	2.7	1.0
2002.....	2.3	*	*	*	*	0.4	3.0	5.2	5.4	4.8	3.1	1.0
2001.....	2.3	*	*	*	*	0.4	3.1	5.5	5.1	4.7	2.8	1.0
2000.....	2.2	*	*	*	*	0.4	3.1	4.8	6.0	4.5	2.5	1.0
1999.....	2.1	*	*	*	*	0.5	2.9	4.7	5.4	4.3	3.4	1.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcoholic liver disease (K70)												
White, both sexes												
2004.....	4.6	*	*	*	*	0.6	4.4	11.0	11.7	10.3	5.7	2.2
2003.....	4.5	*	*	*	*	0.6	4.6	10.7	11.7	10.0	5.9	1.9
2002.....	4.4	*	*	*	*	0.6	4.6	10.5	11.7	10.4	5.8	2.1
2001.....	4.5	*	*	*	*	0.7	4.8	10.8	11.5	10.2	5.8	2.3
2000.....	4.5	*	*	*	*	0.7	4.9	10.3	12.2	10.5	6.1	2.3
1999.....	4.4	*	*	*	*	0.6	4.8	10.2	11.8	10.5	6.4	2.7
White, male												
2004.....	6.8	*	*	*	*	0.9	6.0	16.2	18.3	16.8	10.4	4.8
2003.....	6.8	*	*	*	*	0.9	6.2	16.4	18.5	16.4	10.4	3.7
2002.....	6.6	*	*	*	*	0.8	6.1	16.0	18.5	16.8	9.9	4.5
2001.....	6.7	*	*	*	*	0.9	6.5	16.3	18.4	16.6	10.4	5.4
2000.....	6.8	*	*	*	*	1.0	6.8	16.2	19.2	17.5	11.7	5.4
1999.....	6.7	*	*	*	*	0.8	6.7	16.2	19.0	17.9	11.1	5.8
White, female												
2004.....	2.4	*	*	*	*	0.3	2.8	5.8	5.6	4.7	2.4	1.0
2003.....	2.3	*	*	*	*	0.3	3.0	5.1	5.3	4.5	2.9	1.1
2002.....	2.3	*	*	*	*	0.3	3.0	5.1	5.2	5.0	3.1	1.1
2001.....	2.3	*	*	*	*	0.4	3.1	5.3	5.1	4.8	2.8	1.0
2000.....	2.1	*	*	*	*	0.3	3.0	4.4	5.6	4.6	2.4	1.0
1999.....	2.1	*	*	*	*	0.4	2.8	4.4	5.0	4.3	3.3	1.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcoholic liver disease (K70)												
Black, both sexes												
2004.....	3.0	*	*	*	*	*	2.9	9.0	12.5	8.9	3.1	*
2003.....	3.2	*	*	*	*	*	3.3	9.9	12.5	9.3	3.4	*
2002.....	3.2	*	*	*	*	0.5	2.9	10.0	12.3	8.6	5.3	*
2001.....	3.6	*	*	*	*	0.5	4.2	11.5	12.9	10.7	4.7	*
2000.....	3.8	*	*	*	*	0.6	5.0	12.0	13.3	10.3	5.5	*
1999.....	4.1	*	*	*	*	0.5	5.0	12.8	15.1	11.8	7.1	*
Black, male												
2004.....	4.4	*	*	*	*	*	3.9	12.9	20.3	16.9	*	*
2003.....	4.8	*	*	*	*	*	4.1	15.8	20.1	16.0	6.8	*
2002.....	4.4	*	*	*	*	*	3.8	14.7	18.7	15.0	9.2	*
2001.....	5.3	*	*	*	*	*	5.9	17.1	21.4	19.5	7.7	*
2000.....	5.3	*	*	*	*	*	6.9	17.4	19.4	18.1	10.0	*
1999.....	5.9	*	*	*	*	*	6.7	19.7	24.2	21.8	12.6	*
Black, female												
2004.....	1.8	*	*	*	*	*	1.9	5.6	6.2	3.3	*	*
2003.....	1.8	*	*	*	*	*	2.5	4.7	6.4	4.5	*	*
2002.....	2.0	*	*	*	*	*	2.2	5.9	7.1	4.1	*	*
2001.....	2.1	*	*	*	*	*	2.7	6.7	6.1	4.4	*	*
2000.....	2.4	*	*	*	*	*	3.3	7.3	8.4	4.7	*	*
1999.....	2.4	*	*	*	*	*	3.4	6.9	7.9	4.8	4.0	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)												
All races 3/, both sexes												
2004.....	4.9	*	*	*	*	0.2	2.1	7.5	10.9	17.8	23.3	17.6
2003.....	5.2	*	*	*	*	0.3	2.4	7.8	11.4	19.7	24.3	18.2
2002.....	5.2	*	*	*	0.1	0.3	2.6	7.7	11.3	19.3	25.7	19.3
2001.....	5.2	*	*	*	*	0.3	2.6	7.8	11.2	19.9	24.5	19.9
2000.....	5.1	*	*	*	0.1	0.3	2.5	7.3	11.6	19.5	24.9	20.9
1999.....	5.1	*	*	*	*	0.4	2.5	7.0	11.7	20.1	25.5	20.4
All races, male												
2004.....	5.8	*	*	*	*	0.3	2.9	10.8	14.6	20.5	26.6	21.0
2003.....	6.2	*	*	*	*	0.3	3.2	11.4	15.5	23.0	26.8	23.9
2002.....	6.1	*	*	*	*	0.4	3.5	11.2	14.8	22.0	28.7	23.7
2001.....	6.1	*	*	*	*	0.4	3.4	11.4	14.7	23.6	26.6	26.5
2000.....	6.0	*	*	*	*	0.4	3.5	10.9	15.4	22.7	26.9	29.0
1999.....	6.0	*	*	*	*	0.5	3.5	10.6	15.3	23.6	28.4	27.0
All races, female												
2004.....	4.1	*	*	*	*	0.1	1.3	4.2	7.5	15.4	21.1	16.1
2003.....	4.3	*	*	*	*	0.2	1.5	4.3	7.6	17.0	22.6	15.8
2002.....	4.5	*	*	*	*	0.3	1.8	4.3	8.1	17.1	23.7	17.4
2001.....	4.4	*	*	*	*	0.2	1.9	4.3	8.0	16.8	23.1	17.1
2000.....	4.3	*	*	*	*	0.3	1.6	4.0	8.0	16.9	23.6	17.5
1999.....	4.3	*	*	*	*	0.3	1.4	3.6	8.3	17.3	23.5	17.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)												
White, both sexes												
2004.....	5.3	*	*	*	*	0.2	2.1	7.5	10.9	18.6	24.6	18.1
2003.....	5.7	*	*	*	*	0.3	2.5	7.7	11.4	20.8	25.7	19.0
2002.....	5.7	*	*	*	0.1	0.3	2.7	7.5	11.4	20.3	27.0	20.2
2001.....	5.6	*	*	*	*	0.3	2.7	7.5	11.2	20.7	25.7	20.9
2000.....	5.5	*	*	*	*	0.3	2.5	7.2	11.6	20.1	26.1	21.6
1999.....	5.5	*	*	*	*	0.4	2.5	6.8	11.5	20.9	26.6	21.1
White, male												
2004.....	6.2	*	*	*	*	0.3	3.0	10.7	14.5	21.3	28.0	22.0
2003.....	6.6	*	*	*	*	0.3	3.4	11.4	15.3	24.1	28.1	24.7
2002.....	6.5	*	*	*	*	0.4	3.7	11.1	14.7	23.0	30.3	25.1
2001.....	6.4	*	*	*	*	0.4	3.4	11.0	14.4	24.4	28.0	28.1
2000.....	6.4	*	*	*	*	0.3	3.5	10.6	15.3	23.4	28.2	30.1
1999.....	6.4	*	*	*	*	0.5	3.5	10.2	15.1	24.4	29.7	28.4
White, female												
2004.....	4.5	*	*	*	*	0.1	1.2	4.2	7.5	16.4	22.2	16.4
2003.....	4.7	*	*	*	*	0.2	1.5	4.1	7.8	17.9	24.0	16.5
2002.....	4.9	*	*	*	*	0.3	1.8	4.0	8.3	18.0	24.8	18.1
2001.....	4.8	*	*	*	*	0.2	1.9	4.2	8.2	17.6	24.2	17.9
2000.....	4.7	*	*	*	*	0.2	1.5	3.8	8.1	17.3	24.8	18.2
1999.....	4.6	*	*	*	*	0.3	1.4	3.4	8.2	17.9	24.6	18.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other chronic liver disease and cirrhosis (K73-K74)

Black, both sexes

2004.....	3.5	*	*	*	*	*	2.5	9.3	12.8	12.6	11.3	11.3
2003.....	3.7	*	*	*	*	0.4	2.4	9.9	13.3	14.5	11.0	10.8
2002.....	3.7	*	*	*	*	0.5	2.5	10.8	12.7	13.0	13.2	9.8
2001.....	3.8	*	*	*	*	0.4	3.1	11.1	13.0	14.3	13.9	7.7
2000.....	3.7	*	*	*	*	0.4	2.8	10.1	12.6	16.0	12.5	12.2
1999.....	3.8	*	*	*	*	0.5	3.0	9.9	14.2	14.8	14.5	12.8

Black, male

2004.....	4.7	*	*	*	*	*	3.5	14.2	18.8	16.9	15.3	*
2003.....	4.7	*	*	*	*	*	3.0	14.6	20.8	17.3	13.8	*
2002.....	4.6	*	*	*	*	*	2.9	15.3	19.1	17.4	13.2	*
2001.....	5.1	*	*	*	*	*	4.1	17.4	20.1	19.5	14.7	*
2000.....	4.7	*	*	*	*	*	3.6	15.2	18.6	19.6	14.5	*
1999.....	4.8	*	*	*	*	*	4.4	15.8	19.1	19.0	16.9	*

Black, female

2004.....	2.5	*	*	*	*	*	1.7	5.1	8.0	9.6	9.0	11.0
2003.....	2.7	*	*	*	*	*	1.9	5.9	7.3	12.4	9.4	8.1
2002.....	2.9	*	*	*	*	*	2.2	7.0	7.6	9.9	13.2	9.9
2001.....	2.7	*	*	*	*	*	2.1	5.7	7.3	10.6	13.4	*
2000.....	2.8	*	*	*	*	*	2.2	5.8	7.8	13.4	11.4	11.2
1999.....	2.9	*	*	*	*	*	1.9	4.9	10.2	11.9	13.1	13.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

All races 3/, both sexes

2004.....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.5	7.8	23.8
2003.....	1.0	*	*	*	*	0.1	0.1	0.3	0.7	2.7	8.2	21.8
2002.....	1.0	*	*	*	*	0.1	0.1	0.3	0.8	2.8	7.8	22.9
2001.....	1.0	*	*	*	*	*	0.1	0.3	1.0	2.9	7.5	23.7
2000.....	1.0	*	*	*	*	*	0.1	0.3	0.8	2.4	7.5	24.8
1999.....	1.0	*	*	*	*	*	0.1	0.3	0.9	2.6	7.8	23.9

All races, male

2004.....	0.9	*	*	*	*	*	0.1	0.4	1.0	2.9	9.3	25.5
2003.....	0.9	*	*	*	*	*	0.1	0.3	0.8	3.2	10.1	22.7
2002.....	0.9	*	*	*	*	*	0.1	0.4	1.0	3.1	9.8	24.1
2001.....	1.0	*	*	*	*	*	0.1	0.4	1.2	3.2	9.3	26.9
2000.....	0.9	*	*	*	*	*	0.1	0.3	0.9	2.9	9.4	27.6
1999.....	0.9	*	*	*	*	*	0.1	0.3	1.1	3.0	9.3	27.7

All races, female

2004.....	1.2	*	*	*	*	*	0.1	0.3	0.7	2.1	6.8	23.0
2003.....	1.1	*	*	*	*	*	*	0.3	0.5	2.2	6.9	21.4
2002.....	1.1	*	*	*	*	*	0.1	0.2	0.7	2.5	6.4	22.4
2001.....	1.1	*	*	*	*	*	*	0.3	0.8	2.6	6.3	22.4
2000.....	1.1	*	*	*	*	*	0.1	0.3	0.6	2.1	6.2	23.7
1999.....	1.1	*	*	*	*	*	0.1	0.4	0.7	2.2	6.8	22.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

White, both sexes

2004.....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.3	7.7	24.2
2003.....	1.1	*	*	*	*	*	0.1	0.3	0.6	2.7	8.2	21.9
2002.....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.7	7.6	23.5
2001.....	1.1	*	*	*	*	*	0.1	0.3	0.9	2.7	7.4	23.9
2000.....	1.1	*	*	*	*	*	0.1	0.2	0.7	2.3	7.4	25.6
1999.....	1.1	*	*	*	*	*	0.1	0.3	0.7	2.5	7.8	24.4

White, male

2004.....	1.0	*	*	*	*	*	0.1	0.3	0.9	2.9	9.3	26.7
2003.....	1.0	*	*	*	*	*	*	0.3	0.7	3.2	10.0	23.0
2002.....	1.0	*	*	*	*	*	0.1	0.4	0.9	3.1	9.8	24.7
2001.....	1.0	*	*	*	*	*	0.1	0.4	1.1	3.1	9.3	26.7
2000.....	1.0	*	*	*	*	*	0.1	0.2	0.9	2.8	9.6	28.9
1999.....	1.0	*	*	*	*	*	0.1	0.3	0.9	3.0	9.6	28.1

White, female

2004.....	1.2	*	*	*	*	*	0.1	0.3	0.6	1.9	6.6	23.1
2003.....	1.2	*	*	*	*	*	*	0.2	0.5	2.2	6.9	21.4
2002.....	1.2	*	*	*	*	*	*	0.2	0.6	2.4	6.2	23.0
2001.....	1.2	*	*	*	*	*	*	0.3	0.7	2.4	6.1	22.8
2000.....	1.2	*	*	*	*	*	*	0.2	0.5	1.8	6.0	24.3
1999.....	1.2	*	*	*	*	*	*	0.3	0.6	2.1	6.7	22.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

Black, both sexes

2004.....	0.8	*	*	*	*	*	*	0.8	1.6	3.5	9.1	19.5
2003.....	0.7	*	*	*	*	*	*	0.7	1.2	2.9	7.9	20.7
2002.....	0.7	*	*	*	*	*	*	0.6	1.1	3.3	9.7	14.8
2001.....	0.8	*	*	*	*	*	*	0.7	1.7	4.7	8.3	19.7
2000.....	0.8	*	*	*	*	*	*	0.7	1.2	3.9	8.3	17.8
1999.....	0.7	*	*	*	*	*	0.4	0.5	1.5	3.6	7.2	15.3

Black, male

2004.....	0.7	*	*	*	*	*	*	1.0	1.6	3.6	8.9	*
2003.....	0.7	*	*	*	*	*	*	*	*	3.2	10.4	20.9
2002.....	0.6	*	*	*	*	*	*	*	*	*	10.9	*
2001.....	0.7	*	*	*	*	*	*	*	2.1	4.9	6.2	27.2
2000.....	0.6	*	*	*	*	*	*	*	*	3.7	8.2	*
1999.....	0.6	*	*	*	*	*	*	*	*	3.7	7.1	*

Black, female

2004.....	1.0	*	*	*	*	*	*	*	1.6	3.5	9.3	21.7
2003.....	0.8	*	*	*	*	*	*	0.8	*	2.7	6.5	20.7
2002.....	0.8	*	*	*	*	*	*	*	*	3.7	9.1	16.0
2001.....	0.9	*	*	*	*	*	*	*	*	4.5	9.4	16.8
2000.....	1.0	*	*	*	*	*	*	*	*	4.1	8.4	19.3
1999.....	0.8	*	*	*	*	*	*	*	*	3.5	7.2	15.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

All races 3/, both sexes

2004.....	14.5	4.3	*	0.1	0.2	0.6	1.8	5.0	13.6	38.6	108.4	286.6
2003.....	14.6	4.5	*	0.1	0.2	0.7	1.8	4.9	13.6	40.1	109.5	293.1
2002.....	14.2	4.3	*	0.1	0.2	0.7	1.7	4.7	13.0	39.2	109.1	288.6
2001.....	13.9	3.3	*	*	0.2	0.6	1.7	4.6	13.0	40.2	104.2	287.7
2000.....	13.2	4.3	*	0.1	0.2	0.6	1.6	4.4	12.8	38.0	100.8	277.8
1999.....	12.7	4.4	*	0.1	0.2	0.6	1.6	4.0	12.0	37.1	97.6	268.9

All races, male

2004.....	14.1	5.3	*	*	0.2	0.6	2.0	5.9	15.1	43.6	134.3	372.2
2003.....	14.3	5.2	*	*	0.3	0.7	2.1	6.0	15.4	45.3	136.8	387.6
2002.....	13.9	5.3	*	*	0.2	0.6	1.8	5.5	14.3	45.8	136.2	385.5
2001.....	13.5	3.7	*	*	0.2	0.6	1.8	5.5	14.1	46.1	130.0	393.7
2000.....	12.9	5.3	*	*	0.2	0.6	1.9	5.3	14.2	42.9	127.3	380.7
1999.....	12.4	4.6	*	*	0.2	0.6	1.8	4.7	13.4	42.3	125.3	371.1

All races, female

2004.....	14.8	3.2	*	*	0.2	0.7	1.6	4.1	12.2	34.3	90.9	248.1
2003.....	14.9	3.8	*	*	0.2	0.7	1.5	3.9	12.0	35.6	91.3	251.4
2002.....	14.5	3.2	*	*	0.2	0.7	1.5	4.0	11.8	33.7	91.1	246.6
2001.....	14.2	2.9	*	*	0.2	0.6	1.6	3.9	11.9	35.2	87.1	243.3
2000.....	13.6	3.2	*	*	0.2	0.6	1.4	3.5	11.5	34.0	83.5	235.9
1999.....	13.0	4.1	*	*	0.2	0.6	1.3	3.4	10.8	33.0	79.6	227.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)												
White, both sexes												
2004.....	14.1	3.1	*	*	0.2	0.4	1.3	3.4	10.3	32.9	101.5	278.2
2003.....	14.3	3.8	*	*	0.2	0.5	1.2	3.4	10.6	33.9	102.4	284.8
2002.....	13.9	3.6	*	*	0.1	0.5	1.1	3.2	9.9	33.4	102.2	278.8
2001.....	13.5	2.4	*	*	0.2	0.4	1.2	3.2	9.7	34.2	96.9	278.2
2000.....	12.9	2.9	*	0.1	0.2	0.4	1.1	3.0	9.7	31.9	94.0	268.7
1999.....	12.3	3.4	*	0.1	0.1	0.4	1.0	2.7	9.1	31.5	90.1	255.7
White, male												
2004.....	13.9	3.8	*	*	0.2	0.3	1.5	4.0	11.7	37.9	128.6	366.7
2003.....	14.0	4.5	*	*	0.2	0.5	1.4	4.1	11.9	39.0	130.1	383.6
2002.....	13.7	4.4	*	*	0.1	0.5	1.2	3.8	10.7	40.0	130.4	377.4
2001.....	13.3	2.8	*	*	0.2	0.3	1.4	3.8	10.9	40.4	123.5	388.9
2000.....	12.7	3.8	*	*	0.2	0.4	1.3	3.6	11.0	37.7	121.1	376.2
1999.....	12.1	4.1	*	*	*	0.4	1.2	3.1	10.6	36.9	117.2	360.8
White, female												
2004.....	14.4	2.3	*	*	0.2	0.5	1.1	2.9	9.0	28.6	83.1	238.3
2003.....	14.5	3.0	*	*	0.1	0.5	0.9	2.7	9.3	29.5	83.6	241.0
2002.....	14.1	2.7	*	*	0.1	0.5	1.0	2.7	9.1	27.8	83.3	235.9
2001.....	13.7	2.1	*	*	0.2	0.5	1.0	2.6	8.6	29.0	79.2	231.9
2000.....	13.0	2.0	*	*	0.2	0.4	0.9	2.4	8.4	27.1	76.2	225.0
1999.....	12.4	2.8	*	*	0.2	0.4	0.9	2.3	7.8	26.9	72.3	213.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)												
Black, both sexes												
2004.....	20.3	10.1	*	*	0.5	1.9	5.5	16.6	44.5	95.4	202.8	430.9
2003.....	20.6	9.1	*	*	0.5	2.2	6.2	16.4	42.8	103.3	207.0	440.5
2002.....	19.8	8.2	*	*	0.6	2.0	5.3	16.2	41.0	100.1	203.2	441.6
2001.....	19.6	8.1	*	*	0.4	1.8	5.3	15.8	41.6	102.5	202.1	429.6
2000.....	18.9	11.5	*	*	0.4	1.8	5.3	15.8	40.6	99.9	195.1	406.4
1999.....	18.6	9.6	*	*	0.5	1.9	5.4	15.2	37.2	95.4	195.2	433.3
Black, male												
2004.....	19.2	12.2	*	*	*	1.9	6.2	21.0	49.8	108.1	232.7	512.6
2003.....	20.0	9.5	*	*	0.6	2.4	7.1	21.2	51.5	120.5	246.0	518.2
2002.....	19.1	10.2	*	*	*	1.9	6.0	20.1	49.1	115.0	235.2	561.8
2001.....	18.0	8.7	*	*	*	2.2	5.2	19.3	45.4	114.0	233.4	511.7
2000.....	17.7	13.4	*	*	*	1.9	6.5	20.3	44.9	104.8	231.6	490.1
1999.....	17.5	8.1	*	*	*	1.8	6.5	18.9	40.4	102.9	250.0	526.1
Black, female												
2004.....	21.3	8.0	*	*	*	1.8	4.9	12.9	40.3	86.3	185.8	398.5
2003.....	21.1	8.7	*	*	*	1.9	5.4	12.2	35.8	91.0	185.0	410.4
2002.....	20.5	6.1	*	*	0.7	2.1	4.7	12.8	34.4	89.6	185.0	395.5
2001.....	21.0	7.6	*	*	*	1.4	5.5	12.7	38.5	94.4	184.6	399.1
2000.....	20.0	9.6	*	*	*	1.7	4.3	11.9	37.1	96.5	174.6	375.2
1999.....	19.5	11.1	*	*	*	1.9	4.5	12.0	34.6	90.1	164.5	398.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

All races 3/, both sexes

2004.....	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.8
2003.....	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.6
2002.....	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.9
2001.....	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.9
2000.....	0.1	*	*	*	*	*	*	*	*	0.2	0.4	1.2
1999.....	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.8

All races, male

2004.....	0.1	*	*	*	*	*	*	*	*	*	0.5	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	0.4	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.5	1.8
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.7	*

All races, female

2004.....	0.1	*	*	*	*	*	*	*	*	*	0.3	0.8
2003.....	0.1	*	*	*	*	*	*	*	*	*	0.4	0.6
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.4	0.9
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.4	0.7
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.3	0.9
1999.....	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

All races 3/, both sexes

2004.....	0.2	*	*	*	*	0.1	0.1	0.1	0.2	0.5	1.3	2.8
2003.....	0.2	*	*	*	*	*	0.0	0.1	0.2	0.6	1.3	2.8
2002.....	0.2	*	*	*	*	0.1	0.1	0.1	0.2	0.5	1.4	2.8
2001.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.2	2.8
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.4	2.8
1999.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	1.6	3.2

All races, male

2004.....	0.2	*	*	*	*	*	*	0.1	0.3	0.5	1.6	4.0
2003.....	0.2	*	*	*	*	*	*	0.2	0.2	0.6	1.6	4.1
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.8	3.8
2001.....	0.2	*	*	*	*	*	*	0.1	0.2	0.8	1.7	3.7
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.7	1.7	4.5
1999.....	0.2	*	*	*	*	*	*	0.1	0.3	0.5	2.2	5.0

All races, female

2004.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.1	2.2
2003.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.1	2.2
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.1	2.3
2001.....	0.2	*	*	*	*	*	*	*	*	0.5	0.8	2.4
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.2	2.1
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.3	2.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

White, both sexes

2004.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.2	2.7
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.2	2.6
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	2.6
2001.....	0.2	*	*	*	*	*	*	0.1	0.1	0.6	1.1	2.9
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	2.4
1999.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.6	2.9

White, male

2004.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.6	3.7
2003.....	0.2	*	*	*	*	*	*	0.1	*	0.5	1.4	3.8
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.7	3.8
2001.....	0.2	*	*	*	*	*	*	*	*	0.7	1.7	4.0
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.7	4.0
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.5	2.2	4.6

White, female

2004.....	0.2	*	*	*	*	*	*	*	*	0.4	1.0	2.3
2003.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.0	2.1
2002.....	0.2	*	*	*	*	*	*	*	*	0.4	1.0	2.0
2001.....	0.2	*	*	*	*	*	*	*	*	0.5	0.8	2.4
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.1	1.8
1999.....	0.2	*	*	*	*	*	*	*	*	0.5	1.2	2.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Renal failure (N17-N19)												
All races 3/, both sexes												
2004.....	14.2	4.0	*	*	0.2	0.6	1.7	4.8	13.4	37.9	106.6	282.9
2003.....	14.4	4.2	*	0.1	0.2	0.7	1.7	4.8	13.4	39.3	107.8	289.7
2002.....	13.9	4.1	*	*	0.2	0.6	1.6	4.6	12.7	38.6	107.2	284.8
2001.....	13.6	3.1	*	*	0.2	0.5	1.6	4.5	12.7	39.4	102.6	284.0
2000.....	13.0	4.0	*	*	0.2	0.6	1.6	4.2	12.5	37.2	98.9	273.7
1999.....	12.4	4.3	*	0.1	0.2	0.6	1.5	3.9	11.7	36.4	95.3	264.8
All races, male												
2004.....	13.8	5.1	*	*	0.2	0.5	2.0	5.7	14.8	42.9	132.1	367.4
2003.....	14.1	4.8	*	*	0.2	0.7	2.0	5.7	15.2	44.5	134.8	383.1
2002.....	13.7	5.1	*	*	0.2	0.6	1.8	5.4	14.0	45.0	134.0	380.7
2001.....	13.2	3.5	*	*	0.2	0.5	1.8	5.3	13.8	45.2	128.0	388.7
2000.....	12.6	5.1	*	*	0.2	0.6	1.8	5.1	13.8	42.1	125.1	374.2
1999.....	12.1	4.5	*	*	0.1	0.5	1.7	4.5	13.1	41.5	122.3	364.6
All races, female												
2004.....	14.6	3.0	*	*	0.2	0.6	1.5	3.9	12.0	33.7	89.5	244.9
2003.....	14.6	3.7	*	*	0.2	0.6	1.5	3.8	11.8	34.9	89.7	248.4
2002.....	14.2	3.0	*	*	0.2	0.7	1.4	3.8	11.6	33.2	89.5	243.2
2001.....	14.0	2.7	*	*	0.2	0.5	1.5	3.7	11.8	34.5	85.9	240.1
2000.....	13.3	2.8	*	*	0.2	0.5	1.3	3.4	11.2	33.3	81.8	232.8
1999.....	12.7	4.1	*	*	0.2	0.6	1.3	3.3	10.4	32.2	77.8	224.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Renal failure (N17-N19)												
White, both sexes												
2004.....	13.9	2.9	*	*	0.1	0.4	1.2	3.3	10.0	32.3	99.8	274.7
2003.....	14.0	3.5	*	*	0.2	0.5	1.1	3.3	10.3	33.2	100.7	281.5
2002.....	13.6	3.4	*	*	0.1	0.4	1.1	3.1	9.6	32.7	100.5	275.3
2001.....	13.2	2.2	*	*	0.2	0.4	1.1	3.1	9.5	33.4	95.3	274.6
2000.....	12.6	2.7	*	*	0.2	0.4	1.1	2.9	9.4	31.2	92.2	265.0
1999.....	12.0	3.3	*	*	0.1	0.4	1.0	2.5	8.9	30.8	87.9	251.9
White, male												
2004.....	13.6	3.6	*	*	0.2	0.3	1.4	3.9	11.4	37.3	126.5	362.3
2003.....	13.8	4.1	*	*	0.2	0.5	1.3	3.9	11.7	38.3	128.3	379.3
2002.....	13.4	4.2	*	*	*	0.4	1.2	3.6	10.4	39.2	128.3	372.7
2001.....	13.0	2.6	*	*	0.2	0.3	1.3	3.7	10.7	39.5	121.5	383.9
2000.....	12.4	3.6	*	*	0.2	0.4	1.2	3.4	10.6	36.9	118.9	370.4
1999.....	11.8	3.9	*	*	*	0.4	1.1	3.0	10.3	36.2	114.4	354.9
White, female												
2004.....	14.1	2.1	*	*	0.1	0.4	1.0	2.8	8.7	28.1	81.6	235.2
2003.....	14.2	2.8	*	*	*	0.5	0.9	2.7	9.0	28.8	82.1	238.2
2002.....	13.8	2.6	*	*	*	0.5	1.0	2.6	8.9	27.2	81.8	232.9
2001.....	13.4	1.8	*	*	0.2	0.4	0.9	2.5	8.4	28.3	78.0	228.8
2000.....	12.8	1.7	*	*	0.2	0.4	0.9	2.3	8.2	26.5	74.6	222.1
1999.....	12.2	2.8	*	*	0.1	0.4	0.8	2.2	7.6	26.2	70.6	210.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Renal failure (N17-N19)												
Black, both sexes												
2004.....	20.0	9.8	*	*	0.4	1.7	5.3	16.3	44.0	94.2	200.3	426.4
2003.....	20.3	8.9	*	*	0.5	2.1	6.1	16.1	42.5	102.0	204.4	436.4
2002.....	19.5	7.7	*	*	0.5	1.9	5.3	15.7	40.6	99.6	200.2	434.2
2001.....	19.3	8.1	*	*	0.4	1.7	5.2	15.4	41.1	101.3	200.0	425.6
2000.....	18.5	11.1	*	*	0.4	1.7	5.1	15.2	40.1	98.6	192.4	400.4
1999.....	18.2	9.6	*	*	0.4	1.8	5.2	14.8	36.4	94.1	191.8	425.3
Black, male												
2004.....	18.9	11.9	*	*	*	1.9	6.0	20.6	49.0	106.8	230.0	506.7
2003.....	19.7	9.2	*	*	*	2.4	7.0	20.8	51.2	118.8	241.7	513.0
2002.....	18.8	9.6	*	*	*	1.7	5.9	19.6	48.8	114.2	231.5	555.3
2001.....	17.7	8.7	*	*	*	2.1	5.0	18.8	44.7	112.4	231.6	508.3
2000.....	17.3	13.4	*	*	*	1.8	6.3	19.5	44.2	103.3	229.2	480.9
1999.....	17.1	8.1	*	*	*	1.7	6.2	18.5	39.7	101.5	245.7	513.2
Black, female												
2004.....	21.0	7.7	*	*	*	1.6	4.7	12.6	39.9	85.3	183.5	394.6
2003.....	20.9	8.7	*	*	*	1.8	5.3	12.1	35.5	90.0	183.2	406.7
2002.....	20.2	*	*	*	*	2.0	4.7	12.3	34.1	89.2	182.6	387.7
2001.....	20.7	7.6	*	*	*	1.3	5.4	12.6	38.2	93.5	182.3	394.9
2000.....	19.6	8.6	*	*	*	1.6	4.0	11.5	36.7	95.3	171.7	370.4
1999.....	19.1	11.1	*	*	*	1.9	4.3	11.7	33.8	88.9	161.6	392.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Other disorders of kidney (N25,N27)

Black, both sexes

Table with 13 columns (years 2004-1999) and 13 rows (ages 1-85+). All cells contain an asterisk (*).

Black, male

Table with 13 columns (years 2004-1999) and 13 rows (ages 1-85+). All cells contain an asterisk (*).

Black, female

Table with 13 columns (years 2004-1999) and 13 rows (ages 1-85+). All cells contain an asterisk (*).

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Infections of kidney (N10-N12,N13.6,N15.1)

All races 3/, both sexes

2004.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.8	5.2
2003.....	0.3	*	*	*	*	*	0.1	0.2	0.2	0.6	1.9	5.9
2002.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.9	6.3
2001.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.5	2.0	6.0
2000.....	0.3	*	*	*	*	*	0.0	0.1	0.2	0.6	2.0	7.5
1999.....	0.3	*	*	*	*	*	0.1	0.2	0.2	0.7	2.0	7.5

All races, male

2004.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.5	4.5
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.4	5.4
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.4	5.3
2001.....	0.2	*	*	*	*	*	*	*	*	0.5	1.5	4.3
2000.....	0.2	*	*	*	*	*	*	*	*	0.3	1.5	5.6
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.7	6.8

All races, female

2004.....	0.4	*	*	*	*	*	0.1	0.2	0.2	0.7	2.1	5.5
2003.....	0.4	*	*	*	*	*	0.1	0.2	0.3	0.7	2.2	6.1
2002.....	0.4	*	*	*	*	*	*	0.1	0.3	0.6	2.2	6.7
2001.....	0.4	*	*	*	*	*	0.1	0.2	0.3	0.5	2.3	6.7
2000.....	0.4	*	*	*	*	*	*	0.1	0.3	0.8	2.3	8.3
1999.....	0.4	*	*	*	*	*	0.1	0.2	0.3	0.9	2.1	7.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Infections of kidney (N10-N12,N13.6,N15.1)

White, both sexes

2004.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.8	5.2
2003.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.9	6.0
2002.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.9	6.4
2001.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.5	1.9	6.1
2000.....	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.9	7.6
1999.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.7	2.0	7.7

White, male

2004.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.5	4.7
2003.....	0.2	*	*	*	*	*	*	0.1	*	0.4	1.3	5.8
2002.....	0.2	*	*	*	*	*	*	*	*	0.5	1.4	5.4
2001.....	0.2	*	*	*	*	*	*	*	*	0.4	1.4	4.3
2000.....	0.2	*	*	*	*	*	*	*	*	*	1.5	5.9
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.8	7.2

White, female

2004.....	0.4	*	*	*	*	*	0.1	0.1	0.2	0.8	2.0	5.5
2003.....	0.4	*	*	*	*	*	*	0.2	0.3	0.8	2.2	6.0
2002.....	0.4	*	*	*	*	*	*	*	0.2	0.7	2.3	6.8
2001.....	0.4	*	*	*	*	*	*	0.2	0.3	0.5	2.3	6.9
2000.....	0.4	*	*	*	*	*	*	*	0.3	0.7	2.2	8.3
1999.....	0.4	*	*	*	*	*	*	0.2	0.2	0.9	2.2	7.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Pregnancy, childbirth and the puerperium (000-099)

All races 3/, both sexes

2004.....	0.2	*	0.4	0.7	0.4	0.2	*	*	*	*
2003.....	0.2	*	0.3	0.6	0.4	0.1	*	*	*	*
2002.....	0.1	*	0.2	0.4	0.2	*	*	*	*	*
2001.....	0.1	*	0.3	0.5	0.2	*	*	*	*	*
2000.....	0.1	*	0.2	0.5	0.3	*	*	*	*	*
1999.....	0.1	*	0.3	0.5	0.3	*	*	*	*	*

All races, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2004.....	0.5	*	0.8	1.5	0.7	0.4	*	*	*	*
2003.....	0.4	*	0.6	1.2	0.7	0.1	*	*	*	*
2002.....	0.3	*	0.5	0.9	0.5	*	*	*	*	*
2001.....	0.3	*	0.6	1.0	0.5	*	*	*	*	*
2000.....	0.3	*	0.5	0.9	0.5	*	*	*	*	*
1999.....	0.3	*	0.5	0.9	0.5	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Pregnancy, childbirth and the puerperium (000-099)

White, both sexes												
2004	0.2	*	0.3	0.5	0.3	0.2	*	*	*	*
2003	0.1	*	0.2	0.4	0.3	0.1	*	*	*	*
2002	0.1	*	0.1	0.3	0.2	*	*	*	*	*
2001	0.1	*	0.2	0.4	0.1	*	*	*	*	*
2000	0.1	*	0.2	0.3	0.2	*	*	*	*	*
1999	0.1	*	0.1	0.3	0.2	*	*	*	*	*
White, male												
2004
2003
2002
2001
2000
1999
White, female												
2004	0.3	*	0.6	1.1	0.6	0.3	*	*	*	*
2003	0.3	*	0.4	0.9	0.5	0.1	*	*	*	*
2002	0.2	*	0.3	0.6	0.3	*	*	*	*	*
2001	0.2	*	0.4	0.7	0.3	*	*	*	*	*
2000	0.2	*	0.4	0.7	0.4	*	*	*	*	*
1999	0.2	*	0.3	0.6	0.4	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Pregnancy, childbirth and the puerperium (000-099)

Black, both sexes

2004.....	0.7	*	1.1	2.0	0.8	0.5	*	*	*	*
2003.....	0.5	*	0.7	1.5	1.0	*	*	*	*	*
2002.....	0.4	*	0.7	1.2	0.7	*	*	*	*	*
2001.....	0.4	*	0.7	1.2	0.7	*	*	*	*	*
2000.....	0.4	*	0.6	1.2	0.7	*	*	*	*	*
1999.....	0.4	*	0.8	1.3	0.7	*	*	*	*	*

Black, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

Black, female

2004.....	1.2	*	2.3	3.8	1.5	0.9	*	*	*	*
2003.....	1.0	*	1.4	2.9	1.9	*	*	*	*	*
2002.....	0.8	*	1.5	2.3	1.4	*	*	*	*	*
2001.....	0.8	*	1.5	2.3	1.4	*	*	*	*	*
2000.....	0.7	*	1.2	2.3	1.2	*	*	*	*	*
1999.....	0.9	*	1.6	2.5	1.4	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other complications of pregnancy, childbirth and the puerperium (O10-O99)

All races 3/, both sexes

2004.....	0.2	*	0.4	0.7	0.3	0.2	*	*	*	*
2003.....	0.2	*	0.3	0.5	0.4	0.1	*	*	*	*
2002.....	0.1	*	0.2	0.4	0.2	*	*	*	*	*
2001.....	0.1	*	0.2	0.4	0.2	*	*	*	*	*
2000.....	0.1	*	0.2	0.4	0.2	*	*	*	*	*
1999.....	0.1	*	0.2	0.4	0.3	*	*	*	*	*

All races, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races, female

2004.....	0.4	*	0.8	1.4	0.7	0.4	*	*	*	*
2003.....	0.3	*	0.5	1.1	0.7	0.1	*	*	*	*
2002.....	0.2	*	0.4	0.8	0.4	*	*	*	*	*
2001.....	0.3	*	0.5	0.9	0.4	*	*	*	*	*
2000.....	0.3	*	0.5	0.8	0.5	*	*	*	*	*
1999.....	0.3	*	0.5	0.8	0.5	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other complications of pregnancy, childbirth and the puerperium (O10-O99)

White, both sexes

2004.....	0.2	*	0.3	0.5	0.3	0.2	*	*	*	*
2003.....	0.1	*	0.2	0.4	0.3	0.1	*	*	*	*
2002.....	0.1	*	0.1	0.3	0.2	*	*	*	*	*
2001.....	0.1	*	0.2	0.3	0.1	*	*	*	*	*
2000.....	0.1	*	0.2	0.3	0.2	*	*	*	*	*
1999.....	0.1	*	0.1	0.3	0.2	*	*	*	*	*

White, male

2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

White, female

2004.....	0.3	*	0.5	1.1	0.6	0.3	*	*	*	*
2003.....	0.3	*	0.4	0.8	0.5	0.1	*	*	*	*
2002.....	0.2	*	0.3	0.6	0.3	*	*	*	*	*
2001.....	0.2	*	0.4	0.7	0.3	*	*	*	*	*
2000.....	0.2	*	0.4	0.6	0.4	*	*	*	*	*
1999.....	0.2	*	0.3	0.6	0.4	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other complications of pregnancy, childbirth and the puerperium (O10-O99)

Black, both sexes

2004	0.6	*	1.0	1.9	0.7	0.5	*	*	*	*
2003	0.5	*	0.6	1.4	0.9	*	*	*	*	*
2002	0.4	*	0.7	1.1	0.6	*	*	*	*	*
2001	0.4	*	0.6	1.1	0.7	*	*	*	*	*
2000	0.3	*	0.5	1.0	0.6	*	*	*	*	*
1999	0.4	*	0.7	1.1	0.7	*	*	*	*	*

Black, male

2004
2003
2002
2001
2000
1999

Black, female

2004	1.1	*	2.1	3.6	1.3	0.9	*	*	*	*
2003	0.9	*	1.1	2.7	1.8	*	*	*	*	*
2002	0.7	*	1.4	2.2	1.2	*	*	*	*	*
2001	0.7	*	1.3	2.1	1.3	*	*	*	*	*
2000	0.6	*	1.0	2.0	1.1	*	*	*	*	*
1999	0.8	*	1.5	2.2	1.3	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races 3/, both sexes

2004.....	3.6	137.9	3.6	1.0	1.2	1.0	1.1	1.7	2.2	2.3	3.6	6.1
2003.....	3.6	140.4	3.4	0.9	1.1	1.1	1.3	1.6	2.4	2.3	3.6	6.8
2002.....	3.7	139.4	3.4	1.0	1.2	1.2	1.3	1.8	2.4	2.4	3.8	6.6
2001.....	3.7	136.7	3.6	0.9	1.3	1.2	1.3	1.6	2.3	2.4	3.7	7.2
2000.....	3.8	150.9	3.2	1.0	1.1	1.2	1.3	1.6	2.3	2.5	3.9	7.5
1999.....	3.7	144.2	3.6	1.0	1.1	1.2	1.2	1.8	2.3	2.5	4.0	7.8

All races, male

2004.....	3.8	139.0	3.9	1.0	1.3	1.2	1.3	1.8	2.3	2.5	3.7	6.8
2003.....	3.8	147.6	3.4	0.9	1.3	1.2	1.4	1.7	2.5	2.4	3.5	6.7
2002.....	3.9	142.6	3.5	1.0	1.4	1.4	1.3	1.9	2.5	2.6	4.0	7.0
2001.....	3.9	143.1	3.6	1.0	1.5	1.2	1.4	1.7	2.3	2.7	4.1	7.7
2000.....	4.0	155.4	3.3	1.0	1.3	1.4	1.4	1.9	2.3	2.7	4.0	10.6
1999.....	3.9	147.9	4.0	1.0	1.3	1.2	1.3	1.9	2.5	2.5	4.2	8.5

All races, female

2004.....	3.4	136.8	3.2	0.9	1.0	0.9	1.0	1.5	2.1	2.2	3.6	5.8
2003.....	3.4	132.8	3.5	0.9	0.9	1.0	1.2	1.5	2.3	2.2	3.7	6.8
2002.....	3.5	136.1	3.3	1.0	1.1	1.0	1.2	1.6	2.3	2.3	3.8	6.5
2001.....	3.4	129.9	3.7	0.8	1.0	1.1	1.2	1.6	2.3	2.2	3.5	7.0
2000.....	3.5	146.2	3.1	1.0	0.9	1.0	1.1	1.4	2.4	2.4	3.9	6.3
1999.....	3.5	140.3	3.2	1.1	0.9	1.1	1.1	1.6	2.2	2.6	3.9	7.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

White, both sexes

2004.....	3.5	134.5	3.2	0.9	1.1	1.1	1.1	1.7	2.3	2.5	3.7	6.3
2003.....	3.5	137.6	3.2	0.9	1.1	1.1	1.3	1.7	2.5	2.3	3.7	6.9
2002.....	3.6	138.3	3.3	1.0	1.2	1.2	1.3	1.8	2.4	2.5	3.9	6.9
2001.....	3.6	135.0	3.2	0.9	1.2	1.2	1.3	1.7	2.4	2.5	3.8	7.4
2000.....	3.6	147.1	2.9	0.9	1.1	1.2	1.2	1.7	2.4	2.5	4.1	7.8
1999.....	3.6	141.3	3.2	0.9	1.1	1.2	1.2	1.8	2.4	2.6	4.0	8.0

White, male

2004.....	3.6	134.3	3.5	1.0	1.2	1.2	1.3	1.9	2.4	2.6	3.7	7.2
2003.....	3.7	143.2	3.1	0.9	1.2	1.2	1.4	1.7	2.6	2.3	3.4	6.9
2002.....	3.8	143.3	3.4	1.0	1.3	1.4	1.3	1.9	2.5	2.6	3.9	7.3
2001.....	3.8	143.0	3.2	1.0	1.4	1.3	1.4	1.7	2.4	2.9	4.2	7.5
2000.....	3.9	151.9	2.9	0.9	1.3	1.4	1.4	1.9	2.4	2.6	4.3	11.1
1999.....	3.8	146.5	3.5	0.9	1.3	1.2	1.3	1.9	2.6	2.5	4.1	8.6

White, female

2004.....	3.4	134.7	2.9	0.8	1.0	1.0	1.0	1.6	2.2	2.3	3.6	6.0
2003.....	3.4	131.8	3.2	0.9	0.9	1.0	1.2	1.6	2.4	2.2	3.9	6.9
2002.....	3.5	133.1	3.1	1.0	1.1	1.0	1.2	1.7	2.3	2.4	3.9	6.7
2001.....	3.4	126.7	3.3	0.7	1.0	1.1	1.2	1.6	2.5	2.2	3.6	7.3
2000.....	3.5	142.0	2.9	0.9	0.9	1.0	1.1	1.5	2.4	2.5	4.0	6.5
1999.....	3.5	135.7	2.9	1.0	0.9	1.2	1.2	1.7	2.3	2.6	4.0	7.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Black, both sexes

2004.....	4.5	162.9	5.1	1.3	1.7	1.1	1.2	1.7	1.6	1.6	3.9	*
2003.....	4.5	157.5	4.7	1.1	1.5	1.4	1.2	1.6	2.2	2.3	3.1	5.8
2002.....	4.7	155.2	4.5	1.2	1.6	1.4	1.6	1.9	2.6	2.4	3.3	*
2001.....	4.6	158.4	5.3	1.1	1.9	1.3	1.6	1.7	1.9	2.2	3.0	6.1
2000.....	4.9	179.2	5.0	1.3	1.5	1.4	1.6	1.6	2.4	2.6	2.7	*
1999.....	4.8	169.3	5.5	1.6	1.4	1.1	1.3	1.9	1.9	2.8	4.2	7.7

Black, male

2004.....	5.1	169.9	5.8	1.1	2.2	1.4	1.6	1.8	1.9	*	*	*
2003.....	5.1	170.7	4.8	1.1	1.7	1.5	1.2	1.9	2.7	3.5	*	*
2002.....	5.1	151.7	4.6	1.4	1.9	1.7	1.7	2.3	3.4	3.0	*	*
2001.....	5.0	160.8	5.5	1.0	2.4	1.2	1.7	1.7	2.0	*	*	*
2000.....	5.4	181.4	5.5	1.5	2.0	1.8	1.9	2.1	*	3.8	*	*
1999.....	5.2	172.7	6.2	1.6	1.8	1.2	1.6	2.3	2.1	*	*	*

Black, female

2004.....	4.0	155.6	4.3	1.4	1.2	0.8	0.9	1.6	1.4	*	3.3	*
2003.....	3.9	143.8	4.6	1.0	1.2	1.3	1.2	1.4	1.8	*	*	*
2002.....	4.3	158.9	4.4	1.0	1.3	1.1	1.5	1.5	2.0	*	*	*
2001.....	4.3	155.9	5.1	1.2	1.4	1.3	1.4	1.7	1.8	2.5	*	*
2000.....	4.4	176.9	4.6	1.2	1.1	1.1	1.4	1.1	3.0	*	*	*
1999.....	4.3	165.8	4.8	1.6	1.1	1.0	1.2	1.5	1.8	3.1	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races 3/, both sexes

2004.....	10.3	82.4	1.2	0.3	1.6	2.7	4.3	5.8	7.0	12.6	39.8	221.5
2003.....	10.8	82.9	1.6	0.4	2.0	3.3	5.4	7.0	7.4	13.8	38.9	224.1
2002.....	10.4	85.7	1.6	0.3	1.8	3.3	5.3	6.7	7.5	13.1	38.8	208.0
2001.....	11.4	86.5	1.8	0.4	2.4	3.9	6.9	8.9	9.1	15.1	39.8	206.4
2000.....	11.3	95.6	1.6	0.4	2.2	3.9	6.6	8.3	9.3	16.5	40.7	208.1
1999.....	9.6	93.2	1.5	0.3	1.6	2.7	4.6	5.9	7.8	14.6	36.1	187.7

All races, male

2004.....	9.5	96.9	1.2	0.3	2.1	3.6	5.6	7.6	9.1	15.0	41.8	183.6
2003.....	10.2	96.5	1.8	0.4	2.7	4.5	6.9	9.2	9.6	17.1	41.9	177.9
2002.....	10.1	98.5	1.8	0.4	2.4	4.5	6.9	8.8	10.0	15.9	43.8	175.5
2001.....	11.3	98.9	2.1	0.5	3.2	5.2	9.0	11.9	12.0	18.0	43.8	175.0
2000.....	11.5	110.6	1.7	0.5	2.9	5.3	8.9	11.5	12.3	20.6	46.0	182.7
1999.....	9.4	106.6	1.7	0.4	2.1	3.6	6.0	8.1	10.4	18.2	40.1	166.3

All races, female

2004.....	11.0	67.3	1.1	0.3	1.0	1.8	3.0	4.0	5.1	10.6	38.4	238.5
2003.....	11.4	68.6	1.5	0.4	1.4	2.1	3.8	5.0	5.3	10.9	36.9	244.6
2002.....	10.7	72.3	1.4	0.3	1.2	2.1	3.8	4.7	5.3	10.8	35.4	222.1
2001.....	11.4	73.6	1.5	0.4	1.5	2.5	4.9	6.1	6.3	12.7	37.2	219.5
2000.....	11.2	79.9	1.5	0.4	1.5	2.5	4.4	5.3	6.5	13.2	37.2	218.5
1999.....	9.7	79.2	1.3	0.3	1.0	1.9	3.2	3.8	5.5	11.7	33.5	196.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
White, both sexes												
2004.....	10.5	65.8	1.0	0.2	1.6	2.7	4.2	5.5	6.5	12.1	39.5	224.4
2003.....	11.1	68.1	1.4	0.4	2.0	3.2	5.1	6.4	6.9	13.2	39.0	228.3
2002.....	10.5	71.0	1.3	0.3	1.8	3.2	5.1	5.9	7.0	12.7	38.6	211.6
2001.....	11.6	70.9	1.5	0.4	2.4	3.8	6.8	8.6	8.5	14.4	39.6	210.0
2000.....	11.4	77.2	1.5	0.4	2.1	3.7	6.5	7.8	8.6	15.6	40.2	212.2
1999.....	9.4	74.7	1.3	0.3	1.4	2.5	4.1	5.1	7.1	13.9	35.1	190.2
White, male												
2004.....	9.4	76.9	1.1	0.3	2.1	3.6	5.5	7.1	8.3	14.4	41.4	186.0
2003.....	10.1	82.4	1.4	0.4	2.7	4.4	6.6	8.4	8.8	16.3	41.7	180.0
2002.....	9.8	82.5	1.4	0.3	2.4	4.3	6.5	7.7	9.2	15.3	42.9	178.2
2001.....	11.2	82.0	1.7	0.4	3.2	5.2	8.9	11.4	11.0	17.2	43.9	177.0
2000.....	11.3	89.4	1.7	0.4	2.9	5.1	8.9	10.7	11.2	19.2	45.5	186.8
1999.....	9.0	86.4	1.6	0.3	2.0	3.2	5.4	6.9	9.3	17.3	38.7	167.3
White, female												
2004.....	11.6	54.2	1.0	0.2	1.0	1.7	2.8	3.9	4.8	10.1	38.2	241.7
2003.....	12.0	53.3	1.3	0.4	1.3	2.0	3.7	4.4	5.1	10.5	37.2	249.7
2002.....	11.2	59.0	1.2	0.2	1.1	2.0	3.6	4.2	5.0	10.5	35.7	226.1
2001.....	11.9	59.3	1.4	0.4	1.5	2.4	4.8	5.8	6.2	12.1	36.7	223.8
2000.....	11.5	64.3	1.3	0.3	1.3	2.3	4.1	4.9	6.1	12.7	36.7	222.5
1999.....	9.9	62.4	1.1	0.3	0.8	1.7	2.7	3.4	5.0	11.2	32.8	199.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Black, both sexes

2004.....	11.6	171.7	1.9	0.5	1.8	4.0	6.5	9.7	13.3	20.4	53.0	220.8
2003.....	12.3	159.3	3.1	0.8	2.4	5.0	8.5	13.4	14.2	23.4	45.3	210.9
2002.....	12.3	162.3	2.6	0.7	2.5	5.1	8.5	13.7	13.7	20.1	49.6	196.0
2001.....	12.6	169.5	2.8	0.6	2.7	5.0	9.0	13.9	14.8	24.4	49.5	191.6
2000.....	13.7	197.1	2.3	0.7	3.2	5.8	9.2	14.7	17.4	28.9	55.5	186.4
1999.....	12.7	192.8	2.7	0.5	2.6	4.8	9.0	13.2	15.9	24.5	54.1	182.4

Black, male

2004.....	12.6	204.3	2.1	*	2.3	5.2	8.0	13.8	19.0	24.8	59.0	181.8
2003.....	13.5	170.7	3.4	0.8	3.0	6.7	11.1	17.6	20.3	31.4	52.5	193.9
2002.....	14.1	183.0	3.5	0.8	3.0	6.8	11.3	19.4	19.7	26.4	68.3	181.2
2001.....	13.9	187.4	3.5	0.8	3.7	6.0	11.1	18.0	23.3	30.4	51.8	184.9
2000.....	15.5	227.1	2.3	0.8	3.7	7.7	10.8	20.6	25.2	39.9	65.1	161.5
1999.....	14.6	217.6	2.8	0.6	3.5	6.8	11.5	19.4	23.1	32.5	66.1	184.0

Black, female

2004.....	10.8	137.7	1.7	0.6	1.3	3.0	5.1	6.2	8.8	17.3	49.6	236.3
2003.....	11.3	147.5	2.9	0.7	1.9	3.3	6.1	9.8	9.2	17.7	41.3	217.5
2002.....	10.6	140.8	1.8	0.7	2.0	3.6	6.0	8.8	8.9	15.7	39.0	201.7
2001.....	11.4	150.6	2.0	*	1.7	4.1	7.2	10.3	7.9	20.2	48.3	194.1
2000.....	12.0	166.0	2.3	*	2.6	4.1	7.7	9.7	11.0	21.2	50.1	195.7
1999.....	10.9	167.1	2.7	*	1.8	3.1	6.8	7.8	10.1	18.9	47.3	181.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All races 3/, both sexes

2004.....	68.8	29.2	3.8	2.2	4.7	8.1	18.3	38.9	67.0	141.4	409.6	1,461.6
2003.....	69.3	29.7	4.0	2.3	4.9	8.4	18.9	38.9	67.1	144.4	411.5	1,508.4
2002.....	67.5	28.4	4.2	2.1	4.6	7.9	18.9	37.8	65.3	140.6	405.3	1,493.0
2001.....	65.5	28.0	4.2	2.1	4.5	7.8	18.0	36.2	63.4	140.9	395.3	1,476.7
2000.....	63.4	29.0	4.0	2.1	4.5	7.8	18.0	35.4	62.4	136.0	388.5	1,445.7
1999.....	61.5	30.2	4.2	2.3	4.6	7.6	17.7	33.9	60.5	136.6	373.0	1,410.7

All races, male

2004.....	57.2	33.8	4.0	2.3	5.4	9.2	21.5	47.8	77.1	151.9	410.5	1,236.6
2003.....	57.3	34.8	4.2	2.6	5.6	9.3	22.3	47.2	76.5	155.6	415.1	1,258.1
2002.....	55.8	33.4	4.6	2.3	5.1	9.1	22.4	46.3	74.5	152.6	406.7	1,262.9
2001.....	53.8	31.4	4.6	2.2	5.2	8.6	21.3	43.7	71.2	153.4	397.4	1,271.6
2000.....	52.3	32.3	4.3	2.3	5.3	9.0	21.4	42.8	70.7	148.4	391.5	1,266.2
1999.....	51.0	35.0	4.5	2.3	5.3	8.8	21.6	40.5	69.0	148.7	384.2	1,225.4

All races, female

2004.....	79.9	24.5	3.6	2.0	3.9	7.0	15.1	30.4	57.6	132.5	408.9	1,562.8
2003.....	81.0	24.5	3.8	1.9	4.2	7.4	15.6	31.0	58.4	135.0	409.1	1,619.1
2002.....	78.8	23.0	3.9	1.9	4.0	6.6	15.5	29.6	56.8	130.6	404.4	1,592.8
2001.....	76.7	24.5	3.8	2.0	3.8	6.9	14.6	29.1	56.2	130.5	394.0	1,562.6
2000.....	74.0	25.6	3.6	2.0	3.7	6.5	14.6	28.2	54.8	125.8	386.5	1,518.8
1999.....	71.6	25.3	3.9	2.2	3.8	6.4	13.9	27.4	52.7	126.7	365.7	1,485.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year	2/	years	years	years	years	years	years	years	years	and over

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

White, both sexes

2004.....	73.3	25.4	3.5	2.1	4.4	7.1	16.9	36.2	63.8	138.4	410.8	1,486.9
2003.....	73.9	25.5	3.7	2.2	4.7	7.5	17.4	35.9	63.8	142.0	412.1	1,534.2
2002.....	71.8	25.2	3.9	2.1	4.2	7.1	17.3	34.8	62.0	137.8	405.2	1,517.6
2001.....	69.4	23.7	3.9	2.1	4.2	6.9	16.0	32.8	59.7	138.1	395.1	1,498.0
2000.....	67.0	24.9	3.5	2.0	4.3	6.9	16.0	32.0	58.6	132.7	386.9	1,464.6
1999.....	65.0	25.2	4.0	2.1	4.3	7.0	15.5	30.3	56.9	132.6	371.6	1,434.0

White, male

2004.....	59.9	29.6	3.7	2.3	5.2	8.4	20.6	45.1	73.3	147.6	407.8	1,251.8
2003.....	59.9	31.5	3.8	2.5	5.7	8.8	21.0	44.0	72.4	151.4	411.4	1,279.1
2002.....	58.2	30.8	4.1	2.2	5.0	8.4	21.3	43.2	70.5	148.1	403.4	1,274.7
2001.....	55.8	25.9	4.4	2.1	5.1	7.8	19.5	39.9	66.8	149.4	394.8	1,282.9
2000.....	54.1	26.8	3.9	2.2	5.2	8.3	19.5	39.3	66.0	143.4	387.1	1,274.8
1999.....	52.7	29.2	4.3	2.1	5.1	8.4	19.6	36.7	64.9	143.7	379.3	1,231.6

White, female

2004.....	86.4	21.0	3.3	1.9	3.5	5.8	13.1	27.4	54.8	130.6	412.9	1,592.9
2003.....	87.6	19.3	3.7	1.8	3.7	6.2	13.6	27.9	55.5	133.9	412.6	1,647.2
2002.....	85.2	19.3	3.6	1.9	3.4	5.6	13.3	26.6	54.0	129.0	406.3	1,623.1
2001.....	82.7	21.4	3.4	2.0	3.4	5.8	12.6	25.8	53.0	128.6	395.2	1,588.0
2000.....	79.6	22.9	3.2	1.8	3.4	5.4	12.5	24.7	51.6	123.6	386.7	1,541.7
1999.....	77.0	20.9	3.7	2.1	3.4	5.5	11.4	24.0	49.5	123.3	366.5	1,515.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Black, both sexes

2004.....	60.9	49.3	5.3	2.8	6.7	15.2	30.9	67.9	110.8	200.4	482.4	1,405.8
2003.....	60.9	51.8	5.3	2.7	6.5	15.1	33.0	69.6	111.4	196.9	480.2	1,435.7
2002.....	59.7	45.2	5.9	2.6	6.8	14.5	33.7	68.4	108.1	194.2	477.7	1,425.4
2001.....	59.1	51.3	5.4	2.4	6.5	14.1	33.9	70.4	107.2	196.2	470.9	1,418.0
2000.....	58.1	49.6	5.9	2.8	6.5	15.0	34.2	69.3	107.4	191.9	467.7	1,388.9
1999.....	56.2	56.7	5.3	3.0	6.7	12.9	35.3	67.8	103.4	199.8	440.9	1,286.2

Black, male

2004.....	56.1	55.7	5.7	2.5	6.9	16.0	32.2	79.7	132.4	234.3	540.1	1,291.5
2003.....	55.7	55.3	6.1	2.9	5.9	15.0	35.4	82.2	132.3	233.6	544.4	1,224.1
2002.....	54.8	48.8	6.4	2.8	6.7	15.4	35.3	81.5	129.1	229.4	522.1	1,319.8
2001.....	54.6	58.4	5.1	2.4	6.6	14.9	38.2	84.2	126.4	229.8	524.1	1,313.9
2000.....	54.3	59.8	6.1	2.5	7.0	15.6	38.7	81.1	130.2	228.4	523.5	1,325.1
1999.....	52.8	64.7	5.4	3.5	7.1	13.4	39.3	80.3	121.7	226.2	505.1	1,291.3

Black, female

2004.....	65.4	42.6	4.9	3.0	6.6	14.6	29.8	57.8	93.3	176.2	449.9	1,451.1
2003.....	65.6	48.2	4.5	2.6	7.1	15.2	30.9	58.8	94.6	170.7	444.0	1,517.9
2002.....	64.1	41.5	5.4	2.3	6.9	13.7	32.2	57.2	91.3	169.2	452.5	1,465.9
2001.....	63.1	43.8	5.6	2.5	6.3	13.3	30.0	58.5	91.8	172.6	441.0	1,456.6
2000.....	61.6	39.0	5.7	3.0	6.0	14.4	30.2	59.1	89.1	166.3	436.2	1,412.7
1999.....	59.3	48.3	5.1	2.5	6.2	12.5	31.9	57.0	88.7	181.2	405.0	1,284.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All races 3/, both sexes

2004.....	38.1	25.8	10.3	6.5	37.0	32.6	37.3	40.7	33.2	44.0	103.7	276.7
2003.....	37.6	23.6	10.9	6.4	37.1	31.5	37.8	38.8	32.9	44.1	101.9	278.9
2002.....	37.0	23.5	10.5	6.6	38.0	31.5	37.2	36.6	31.4	44.2	101.3	275.4
2001.....	35.7	24.2	11.2	6.9	36.1	29.9	35.4	34.1	30.3	42.8	100.9	276.4
2000.....	34.8	23.1	11.9	7.3	36.0	29.5	34.1	32.6	30.9	41.9	95.1	273.5
1999.....	35.1	22.3	12.4	7.6	35.3	29.6	33.8	31.8	30.6	44.6	100.5	282.4

All races, male

2004.....	49.8	29.0	11.8	8.0	53.3	48.5	53.0	58.6	46.9	59.2	134.7	345.5
2003.....	49.3	26.4	13.0	7.9	53.3	47.1	54.2	55.9	47.1	58.9	132.9	352.5
2002.....	48.9	26.8	12.9	8.2	54.9	47.7	53.7	53.3	45.2	60.6	131.6	346.6
2001.....	47.2	26.2	13.3	8.6	52.9	45.4	51.8	50.6	43.0	57.3	133.4	358.5
2000.....	46.2	25.7	14.0	8.8	52.1	45.0	51.0	49.0	44.1	55.9	124.3	359.7
1999.....	46.4	25.4	14.4	9.4	50.7	45.0	50.2	47.9	44.0	60.2	132.4	375.4

All races, female

2004.....	26.8	22.5	8.7	5.1	19.9	16.0	21.7	23.4	20.5	31.1	82.9	245.8
2003.....	26.2	20.7	8.7	4.8	19.9	15.3	21.4	22.4	19.7	31.6	81.1	246.4
2002.....	25.6	20.0	8.1	5.0	20.1	14.9	20.8	20.5	18.6	30.6	81.2	244.6
2001.....	24.5	22.1	8.9	5.2	18.4	14.0	19.2	18.0	18.5	30.8	79.6	242.0
2000.....	23.8	20.5	9.7	5.6	19.1	13.7	17.5	16.8	18.8	30.3	76.1	238.4
1999.....	24.1	19.0	10.3	5.6	19.2	14.0	17.6	16.3	18.3	31.8	79.7	244.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
White, both sexes												
2004.....	40.2	22.2	10.0	6.2	40.3	34.2	38.9	40.7	32.6	43.9	106.3	289.9
2003.....	39.5	20.0	10.6	6.1	40.1	33.0	39.0	38.5	32.3	43.8	104.0	291.1
2002.....	38.7	18.6	9.9	6.2	40.8	32.8	38.1	36.2	30.7	43.9	103.6	286.1
2001.....	37.0	20.6	11.0	6.4	38.2	30.9	35.8	33.1	29.4	42.4	102.7	285.5
2000.....	35.9	19.1	11.1	6.9	38.3	29.9	34.3	31.3	30.2	41.1	96.5	282.8
1999.....	36.0	18.9	11.5	7.1	37.2	30.0	33.8	30.5	29.2	43.5	100.9	290.4
White, male												
2004.....	52.0	25.5	11.5	7.5	57.8	50.5	54.8	58.0	45.1	58.8	137.0	359.4
2003.....	51.3	22.5	12.8	7.6	57.4	49.3	55.6	54.9	45.8	58.3	134.4	365.7
2002.....	50.4	22.3	11.8	7.6	58.7	49.3	54.7	52.3	43.5	59.2	133.0	356.6
2001.....	48.4	22.1	13.3	7.8	55.8	46.7	52.3	48.6	41.1	56.3	134.3	368.1
2000.....	47.0	21.7	13.2	8.4	55.0	45.4	51.0	46.7	42.1	54.2	124.3	369.8
1999.....	46.9	21.8	13.2	8.8	52.8	45.4	50.0	45.4	41.5	58.0	131.8	383.6
White, female												
2004.....	28.8	18.8	8.4	4.7	21.5	17.0	22.6	23.6	20.7	31.2	85.4	258.6
2003.....	28.0	17.4	8.3	4.7	21.7	15.9	22.1	22.3	19.5	31.4	83.5	258.0
2002.....	27.3	14.7	7.8	4.7	21.8	15.5	21.3	20.3	18.6	30.9	83.9	255.4
2001.....	25.9	18.9	8.5	5.0	19.5	14.4	19.2	17.8	18.3	30.7	81.6	251.0
2000.....	25.1	16.4	9.0	5.4	20.4	13.8	17.4	16.2	18.9	30.1	78.1	247.4
1999.....	25.3	15.8	9.6	5.4	20.5	14.1	17.4	15.8	17.7	31.3	80.6	253.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
Black, both sexes												
2004.....	32.8	45.8	12.5	8.5	25.8	31.4	36.7	49.8	43.7	48.2	87.1	152.3
2003.....	32.4	43.5	13.4	7.7	25.6	29.9	38.6	48.4	42.3	50.4	82.7	160.2
2002.....	33.1	48.9	14.1	9.1	27.8	31.4	39.1	46.8	41.4	49.3	83.6	173.4
2001.....	33.5	43.1	13.6	9.4	28.7	30.2	40.0	47.7	42.0	51.0	88.5	183.6
2000.....	33.5	44.5	16.2	9.3	28.7	32.5	39.7	47.0	40.8	52.5	84.9	181.1
1999.....	35.2	41.6	17.7	10.5	29.5	32.2	40.0	47.8	44.7	57.8	103.3	207.6
Black, male												
2004.....	46.7	48.3	14.2	10.3	37.8	50.3	54.1	76.7	70.5	73.7	132.0	213.6
2003.....	46.1	47.6	15.7	9.7	38.6	46.7	57.8	74.5	67.9	74.3	123.3	225.2
2002.....	47.9	51.1	18.5	11.6	41.8	50.6	58.4	72.2	68.7	76.9	129.7	260.5
2001.....	48.2	48.5	15.4	12.8	43.2	47.4	59.9	77.2	67.6	74.9	136.0	255.3
2000.....	49.0	47.7	18.8	12.0	43.0	51.8	62.0	75.4	68.8	79.4	132.6	275.6
1999.....	51.0	45.9	21.0	13.8	44.9	51.4	62.0	78.6	71.1	88.2	150.0	316.4
Black, female												
2004.....	20.1	43.2	10.6	6.6	13.6	14.1	21.3	26.7	22.2	30.0	61.7	128.0
2003.....	19.9	39.3	11.0	5.7	12.5	14.5	21.6	26.1	21.7	33.4	59.8	134.9
2002.....	19.7	46.6	9.6	6.6	13.8	14.1	22.0	25.1	19.5	29.7	57.5	140.1
2001.....	20.1	37.5	11.9	6.0	14.2	14.7	22.4	22.3	21.4	34.2	61.8	157.0
2000.....	19.5	41.3	13.5	6.5	14.6	15.0	19.9	22.6	18.3	33.5	58.0	145.9
1999.....	20.8	37.3	14.3	7.1	14.3	14.9	20.6	21.4	23.5	36.4	77.1	166.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Transport accidents (V01-V99,Y85)												
All races 3/, both sexes												
2004.....	16.4	3.6	4.1	4.3	27.3	18.8	16.4	16.7	15.5	17.5	25.5	27.2
2003.....	16.5	3.7	4.1	4.3	27.7	18.3	17.1	16.6	15.9	17.5	26.2	29.7
2002.....	16.8	3.1	4.1	4.2	29.1	18.7	17.1	16.4	15.7	18.3	27.0	28.8
2001.....	16.6	3.7	4.4	4.4	27.9	18.8	17.4	16.0	15.3	17.9	27.9	28.9
2000.....	16.6	4.6	4.5	4.7	28.1	18.6	17.0	15.8	16.0	18.0	27.0	31.5
1999.....	16.6	5.0	4.6	4.8	27.6	18.5	16.8	15.4	16.0	19.4	28.4	32.0
All races, male												
2004.....	23.0	3.9	4.4	5.0	37.8	27.7	24.0	24.7	22.2	23.7	36.5	46.5
2003.....	23.3	4.0	4.5	5.2	38.6	27.3	24.8	24.3	23.0	23.7	36.6	52.0
2002.....	23.7	3.5	4.8	4.9	40.7	28.1	24.6	24.0	22.5	25.3	37.7	53.0
2001.....	23.5	3.3	5.0	5.3	39.6	28.3	25.1	23.3	21.6	24.3	39.8	52.7
2000.....	23.3	4.8	5.0	5.4	39.3	27.7	24.8	23.0	22.5	24.3	38.1	59.8
1999.....	23.1	5.2	5.1	5.8	38.1	27.2	24.0	22.4	22.5	26.1	40.7	63.5
All races, female												
2004.....	9.9	3.2	3.8	3.6	16.1	9.5	8.8	9.0	9.3	12.3	18.1	18.5
2003.....	10.0	3.4	3.7	3.4	16.3	9.0	9.4	9.1	9.4	12.4	19.3	19.9
2002.....	10.1	2.8	3.4	3.5	16.9	9.2	9.7	9.0	9.3	12.5	19.8	18.3
2001.....	10.0	4.2	3.8	3.5	15.6	9.2	9.8	8.9	9.4	12.7	20.1	18.9
2000.....	10.2	4.3	4.0	3.9	16.4	9.3	9.3	8.9	10.0	12.8	19.8	19.9
1999.....	10.4	4.8	4.2	3.9	16.6	9.6	9.6	8.7	10.1	14.0	20.4	19.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Transport accidents (V01-V99,Y85)												
White, both sexes												
2004.....	16.9	3.3	4.0	4.4	29.5	19.1	16.7	16.7	15.5	17.6	25.9	28.4
2003.....	17.0	3.6	4.0	4.4	29.8	18.7	17.4	16.6	15.7	17.5	26.5	30.5
2002.....	17.3	2.7	3.9	4.2	31.3	19.1	17.3	16.4	15.6	18.4	27.5	29.6
2001.....	17.0	3.7	4.3	4.3	29.5	19.2	17.6	15.8	15.1	17.9	28.5	29.9
2000.....	17.0	4.1	4.3	4.7	30.0	18.5	17.1	15.6	16.0	17.9	27.5	32.2
1999.....	17.0	4.6	4.1	4.8	29.3	18.7	16.8	15.2	15.7	19.4	28.6	32.3
White, male												
2004.....	23.7	3.3	4.2	5.1	40.8	27.8	24.4	24.4	22.0	23.6	36.7	48.6
2003.....	23.9	4.0	4.5	5.2	41.1	27.8	25.0	24.1	22.7	23.7	36.7	52.8
2002.....	24.2	3.1	4.5	5.0	43.4	28.3	24.7	24.0	22.3	24.9	37.7	53.8
2001.....	23.9	3.2	4.9	5.1	41.7	28.7	25.1	22.9	21.2	24.0	40.3	54.1
2000.....	23.6	4.5	4.7	5.5	41.6	27.4	24.7	22.6	22.0	23.7	38.1	61.6
1999.....	23.4	5.2	4.4	5.7	40.1	27.3	23.8	21.8	21.9	25.8	40.6	63.7
White, female												
2004.....	10.3	3.2	3.8	3.6	17.4	9.9	8.9	9.0	9.5	12.4	18.5	19.3
2003.....	10.3	3.1	3.4	3.5	17.7	9.1	9.6	9.1	9.1	12.2	19.7	20.6
2002.....	10.5	2.2	3.4	3.4	18.3	9.4	9.8	8.9	9.4	12.8	20.6	19.1
2001.....	10.3	4.1	3.6	3.5	16.5	9.3	10.0	8.8	9.4	12.7	20.7	19.7
2000.....	10.5	3.7	3.8	4.0	17.7	9.3	9.4	8.7	10.3	12.9	20.5	20.3
1999.....	10.7	3.9	3.8	3.9	17.8	9.8	9.6	8.6	10.0	14.0	20.7	19.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Transport accidents (V01-V99,Y85)												
Black, both sexes												
2004.....	15.2	4.8	5.1	4.4	19.6	20.8	17.1	19.5	17.3	16.4	21.6	14.7
2003.....	15.1	4.2	5.0	4.1	19.7	19.3	18.3	18.1	18.5	18.2	22.4	21.0
2002.....	15.4	5.3	5.0	4.6	21.2	20.4	18.2	17.8	17.1	17.1	21.7	18.4
2001.....	15.9	*	5.2	5.3	22.2	19.9	18.7	18.8	18.1	18.8	21.2	17.5
2000.....	16.0	6.5	5.6	4.7	21.7	21.4	18.6	19.3	17.4	19.6	20.0	21.6
1999.....	16.3	7.3	7.1	5.4	22.0	19.9	19.0	19.0	18.3	19.3	26.0	27.7
Black, male												
2004.....	22.8	5.9	5.4	4.8	27.9	33.8	26.8	30.9	27.7	26.7	38.4	28.8
2003.....	22.6	*	5.1	5.1	29.3	30.6	28.4	29.0	29.1	26.6	37.5	42.7
2002.....	23.1	*	6.4	5.3	30.5	33.1	27.5	27.8	28.3	28.5	40.4	46.1
2001.....	23.8	*	5.7	6.9	32.5	31.6	28.9	30.2	29.6	27.4	36.2	35.2
2000.....	24.1	6.7	6.2	5.7	31.6	34.4	29.7	29.6	29.4	32.2	36.3	45.0
1999.....	24.2	*	8.0	6.8	32.0	31.8	29.0	30.3	28.9	29.1	43.4	68.0
Black, female												
2004.....	8.2	*	4.8	3.9	11.1	8.9	8.6	9.6	9.0	9.1	12.1	9.1
2003.....	8.3	*	5.0	3.1	9.8	9.0	9.4	8.8	9.9	12.2	13.9	12.6
2002.....	8.3	*	3.4	3.9	11.8	9.0	9.9	9.2	8.1	9.0	11.0	*
2001.....	8.7	*	4.7	3.7	11.9	9.4	9.7	9.1	8.9	12.7	12.7	10.9
2000.....	8.5	*	5.0	3.6	11.9	9.7	8.7	10.5	7.6	10.8	10.7	12.9
1999.....	9.2	8.7	6.2	4.0	12.2	9.0	10.1	9.4	9.7	12.4	16.4	12.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

All races 3/, both sexes

2004.....	15.3	3.5	4.0	4.1	26.3	17.6	15.1	15.1	14.1	16.1	24.5	26.1
2003.....	15.4	3.6	3.9	4.0	26.6	17.1	15.7	14.9	14.2	16.2	24.9	28.8
2002.....	15.7	3.0	3.9	3.9	28.2	17.8	15.8	14.8	14.1	17.0	25.7	28.0
2001.....	15.4	3.6	4.1	4.1	26.8	17.5	15.7	14.2	13.7	16.3	26.3	27.8
2000.....	15.4	4.4	4.2	4.3	26.9	17.3	15.3	14.2	14.4	16.5	25.7	30.4
1999.....	15.2	4.8	4.2	4.3	26.2	16.9	14.9	13.6	14.2	17.8	26.5	30.4

All races, male

2004.....	21.3	3.8	4.2	4.7	36.4	25.8	21.9	21.9	19.8	21.3	34.7	44.2
2003.....	21.4	3.9	4.4	4.8	36.9	25.5	22.5	21.5	20.0	21.3	34.3	50.0
2002.....	21.9	3.3	4.5	4.6	39.3	26.5	22.3	21.3	19.9	22.7	35.3	51.7
2001.....	21.5	3.2	4.5	4.9	38.0	26.2	22.4	20.5	19.0	21.5	37.3	51.1
2000.....	21.3	4.6	4.7	5.0	37.4	25.5	22.0	20.4	19.8	21.7	35.6	57.5
1999.....	20.9	5.0	4.7	5.1	36.0	24.7	21.1	19.4	19.3	23.3	38.0	59.8

All races, female

2004.....	9.5	3.2	3.7	3.4	15.7	9.1	8.3	8.5	8.8	11.7	17.7	17.9
2003.....	9.5	3.3	3.5	3.2	15.8	8.5	8.9	8.5	8.8	11.9	18.7	19.5
2002.....	9.8	2.8	3.3	3.3	16.6	8.8	9.3	8.6	8.9	12.3	19.3	17.7
2001.....	9.5	4.0	3.6	3.3	15.1	8.7	9.1	8.1	8.8	12.0	19.2	18.0
2000.....	9.7	4.2	3.8	3.6	15.9	8.8	8.8	8.2	9.5	12.3	19.2	19.3
1999.....	9.7	4.7	3.8	3.6	15.9	8.9	8.8	8.0	9.5	13.3	19.1	18.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

White, both sexes

2004.....	15.8	3.2	3.9	4.1	28.5	17.8	15.3	15.0	14.1	16.2	24.9	27.3
2003.....	15.8	3.5	3.8	4.1	28.6	17.5	15.9	14.9	13.9	16.1	25.2	29.5
2002.....	16.2	2.6	3.7	3.9	30.3	18.0	15.9	14.8	14.1	17.1	26.3	28.8
2001.....	15.7	3.5	3.9	4.0	28.4	17.8	15.9	14.1	13.6	16.3	27.0	28.8
2000.....	15.7	3.9	4.0	4.3	28.7	17.1	15.4	14.0	14.3	16.4	26.1	31.1
1999.....	15.5	4.5	3.8	4.3	27.8	17.0	14.9	13.3	13.9	17.7	26.8	30.7

White, male

2004.....	21.9	3.3	4.0	4.8	39.4	25.8	22.2	21.7	19.5	21.1	34.9	46.3
2003.....	22.0	3.9	4.3	4.9	39.2	25.9	22.6	21.3	19.6	21.3	34.4	50.7
2002.....	22.4	2.9	4.2	4.6	41.9	26.6	22.3	21.2	19.6	22.4	35.5	52.4
2001.....	21.9	3.1	4.4	4.7	40.0	26.4	22.4	20.1	18.6	21.3	37.8	52.6
2000.....	21.6	4.2	4.5	5.0	39.6	25.1	21.8	20.0	19.2	21.1	35.6	59.2
1999.....	21.1	5.0	4.1	5.0	37.8	24.6	20.9	18.8	18.7	23.0	37.9	60.3

White, female

2004.....	9.8	3.2	3.7	3.4	16.8	9.4	8.4	8.4	9.0	11.9	18.1	18.7
2003.....	9.8	3.0	3.2	3.2	17.2	8.6	9.0	8.5	8.6	11.7	19.0	20.1
2002.....	10.1	2.2	3.2	3.2	17.9	9.0	9.4	8.5	8.9	12.6	20.1	18.5
2001.....	9.8	3.8	3.4	3.2	16.0	8.8	9.3	8.1	8.8	12.0	19.8	18.9
2000.....	10.0	3.5	3.6	3.7	17.1	8.9	8.9	8.0	9.8	12.4	19.8	19.7
1999.....	10.0	3.9	3.5	3.6	17.1	9.1	8.8	7.9	9.3	13.3	19.6	18.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Black, both sexes

2004.....	14.3	4.7	4.9	4.2	18.7	19.8	16.0	17.8	15.7	15.5	20.2	14.7
2003.....	14.2	4.2	4.9	3.9	18.8	18.4	17.1	16.5	17.0	17.0	21.0	20.7
2002.....	14.5	5.2	4.8	4.2	20.6	19.7	16.9	16.1	15.6	15.9	20.0	18.1
2001.....	14.8	*	5.0	4.9	21.5	18.9	17.2	16.9	16.4	17.3	19.7	16.6
2000.....	15.0	6.5	5.4	4.4	20.9	20.4	17.1	17.9	16.1	18.2	18.9	20.3
1999.....	15.1	7.1	6.4	5.0	21.1	18.6	17.4	17.1	16.5	17.9	24.1	25.5

Black, male

2004.....	21.2	*	5.2	4.6	26.4	31.8	24.7	28.0	24.7	24.9	35.9	28.8
2003.....	21.1	*	5.0	4.7	27.9	29.0	26.4	25.9	26.3	24.2	35.0	41.7
2002.....	21.5	*	6.3	4.9	29.6	31.7	25.3	24.7	25.2	25.5	36.1	45.0
2001.....	22.0	*	5.3	6.3	31.4	29.9	26.1	26.9	26.3	24.2	34.1	35.2
2000.....	22.5	6.7	5.8	5.4	30.2	32.6	27.2	27.1	27.1	29.4	34.5	43.8
1999.....	22.2	*	7.0	6.0	30.5	29.7	26.3	27.0	25.5	26.5	41.2	60.9

Black, female

2004.....	7.9	*	4.7	3.8	10.9	8.7	8.3	9.0	8.5	8.7	11.3	9.1
2003.....	8.0	*	4.8	3.1	9.6	8.7	8.9	8.4	9.6	11.9	13.1	12.6
2002.....	8.0	*	3.4	3.5	11.6	8.8	9.4	8.8	7.9	9.0	10.9	*
2001.....	8.2	*	4.6	3.3	11.7	8.9	9.2	8.3	8.3	12.5	11.6	9.7
2000.....	8.2	*	4.9	3.5	11.7	9.4	8.2	10.0	7.3	10.3	10.1	11.6
1999.....	8.7	8.7	5.8	3.9	11.8	8.6	9.6	8.7	9.2	11.9	14.5	12.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All Under 1-4 5-14 15-24 25-34 35-44 45-54 55-64 65-74 75-84 85 years
ages 1/ 1 year 2/ years years years years years years years years years and over

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All races 3/, both sexes

2004.....	0.4	*	*	0.1	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.6
2003.....	0.5	*	*	0.1	0.6	0.5	0.5	0.6	0.5	0.4	0.6	0.5
2002.....	0.4	*	*	0.2	0.4	0.3	0.5	0.5	0.4	0.4	0.5	*
2001.....	0.5	*	0.2	0.2	0.5	0.4	0.6	0.6	0.5	0.6	0.8	0.7
2000.....	0.5	*	0.2	0.2	0.6	0.5	0.7	0.6	0.5	0.5	0.7	0.8
1999.....	0.8	*	0.3	0.3	0.9	0.8	0.9	0.8	0.8	0.8	1.3	1.3

All races, male

2004.....	0.6	*	*	0.2	0.7	0.7	0.8	0.9	0.7	0.7	0.9	*
2003.....	0.7	*	*	0.2	1.0	0.8	0.8	1.0	0.8	0.8	1.0	*
2002.....	0.6	*	*	0.2	0.6	0.5	0.8	0.9	0.7	0.7	1.0	*
2001.....	0.8	*	0.3	0.2	0.8	0.8	1.0	1.0	0.9	1.0	1.1	*
2000.....	0.8	*	*	0.3	1.0	0.8	1.1	0.9	0.8	0.9	1.2	*
1999.....	1.1	*	0.3	0.4	1.4	1.2	1.4	1.3	1.2	1.2	1.8	2.9

All races, female

2004.....	0.2	*	*	*	0.2	0.2	0.2	0.2	0.2	0.3	*	*
2003.....	0.2	*	*	0.1	0.3	0.2	0.2	0.2	0.2	*	0.4	*
2002.....	0.1	*	*	0.1	0.1	0.2	0.2	0.1	*	*	*	*
2001.....	0.2	*	*	0.2	0.2	0.1	0.2	0.3	0.2	0.3	0.6	*
2000.....	0.3	*	*	0.2	0.3	0.2	0.3	0.4	0.2	0.2	0.5	*
1999.....	0.4	*	0.3	0.2	0.4	0.4	0.4	0.4	0.4	0.5	1.0	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

All races 3/, both sexes

2004.....	0.7	*	*	0.1	0.5	0.8	0.8	1.1	1.0	0.9	0.5	0.5
2003.....	0.7	*	*	0.2	0.5	0.6	0.9	1.1	1.2	0.9	0.7	*
2002.....	0.7	*	*	0.1	0.5	0.6	0.9	1.0	1.1	0.9	0.8	0.4
2001.....	0.7	*	0.2	0.1	0.5	0.9	1.0	1.1	1.0	1.0	0.8	0.5
2000.....	0.7	*	*	0.1	0.6	0.8	0.9	1.0	1.1	1.0	0.6	*
1999.....	0.7	*	*	0.2	0.5	0.8	0.9	1.0	1.1	0.9	0.5	*

All races, male

2004.....	1.1	*	*	0.1	0.7	1.2	1.4	1.8	1.7	1.7	0.9	*
2003.....	1.1	*	*	0.2	0.8	0.9	1.5	1.8	2.2	1.6	1.3	*
2002.....	1.1	*	*	0.1	0.8	1.1	1.5	1.8	1.9	1.9	1.5	*
2001.....	1.2	*	*	0.2	0.9	1.4	1.6	1.9	1.8	1.7	1.4	*
2000.....	1.2	*	*	0.2	0.9	1.3	1.6	1.7	1.8	1.7	1.3	*
1999.....	1.1	*	*	0.3	0.8	1.4	1.4	1.7	2.0	1.6	0.9	*

All races, female

2004.....	0.3	*	*	*	0.3	0.3	0.3	0.4	0.3	0.2	*	*
2003.....	0.2	*	*	0.1	0.2	0.3	0.3	0.4	0.3	0.3	0.3	*
2002.....	0.2	*	*	0.1	0.2	0.2	0.3	0.3	0.3	*	0.3	*
2001.....	0.3	*	*	0.1	0.2	0.3	0.4	0.4	0.3	0.3	0.4	*
2000.....	0.2	*	*	*	0.2	0.2	0.2	0.3	0.3	0.3	*	*
1999.....	0.3	*	*	0.1	0.2	0.3	0.4	0.3	0.3	0.2	0.3	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/	1-4 2/	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------	--------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

White, both sexes

2004.....	0.7	*	*	0.1	0.6	0.8	0.9	1.2	1.0	0.9	0.5	0.6
2003.....	0.7	*	*	0.2	0.5	0.7	1.0	1.2	1.3	0.9	0.7	*
2002.....	0.7	*	*	0.1	0.5	0.7	0.9	1.1	1.2	0.9	0.7	*
2001.....	0.8	*	0.2	0.1	0.6	0.9	1.1	1.2	1.1	1.0	0.8	*
2000.....	0.7	*	*	0.1	0.6	0.9	1.0	1.0	1.1	1.0	0.7	*
1999.....	0.7	*	*	0.2	0.6	0.9	1.0	1.1	1.1	0.9	0.5	*

White, male

2004.....	1.2	*	*	0.1	0.8	1.3	1.4	2.0	1.8	1.8	1.0	*
2003.....	1.2	*	*	0.2	0.8	1.0	1.6	2.0	2.3	1.7	1.3	*
2002.....	1.2	*	*	0.1	0.8	1.2	1.6	1.9	2.0	1.9	1.4	*
2001.....	1.3	*	*	0.2	0.9	1.5	1.8	1.9	1.8	1.8	1.4	*
2000.....	1.3	*	*	0.2	1.0	1.4	1.8	1.8	1.9	1.8	1.3	*
1999.....	1.2	*	*	0.3	0.9	1.5	1.5	1.8	2.0	1.6	0.9	*

White, female

2004.....	0.3	*	*	*	0.3	0.3	0.4	0.4	0.3	*	*	*
2003.....	0.3	*	*	0.2	0.2	0.3	0.4	0.4	0.3	0.3	*	*
2002.....	0.2	*	*	*	0.2	0.2	0.3	0.3	0.4	*	0.3	*
2001.....	0.3	*	*	*	0.3	0.4	0.5	0.4	0.4	0.3	0.4	*
2000.....	0.2	*	*	*	0.3	0.2	0.3	0.3	0.4	0.3	*	*
1999.....	0.3	*	*	0.1	0.3	0.4	0.4	0.3	0.3	0.3	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)												
All races 3/, both sexes												
2004.....	21.8	22.2	6.1	2.2	9.8	13.8	20.9	24.0	17.7	26.5	78.2	249.5
2003.....	21.0	19.9	6.8	2.1	9.3	13.2	20.7	22.3	16.9	26.6	75.6	249.2
2002.....	20.2	20.3	6.4	2.4	8.9	12.7	20.1	20.2	15.7	25.9	74.4	246.7
2001.....	19.0	20.5	6.8	2.5	8.2	11.1	18.0	18.1	15.0	24.9	73.0	247.5
2000.....	18.2	18.6	7.4	2.6	7.9	11.0	17.2	16.8	14.9	23.8	68.1	242.0
1999.....	18.4	17.3	7.7	2.7	7.7	11.1	17.0	16.4	14.6	25.1	72.1	250.3
All races, male												
2004.....	26.8	25.1	7.4	3.0	15.4	20.8	28.9	34.0	24.6	35.5	98.2	299.0
2003.....	26.0	22.4	8.5	2.8	14.7	19.9	29.4	31.6	24.1	35.3	96.3	300.5
2002.....	25.2	23.3	8.1	3.2	14.2	19.6	29.1	29.3	22.7	35.3	93.9	293.6
2001.....	23.8	22.9	8.3	3.3	13.3	17.1	26.8	27.3	21.4	33.0	93.6	305.8
2000.....	22.9	20.9	9.0	3.4	12.8	17.3	26.2	26.0	21.6	31.5	86.2	299.8
1999.....	23.3	20.2	9.3	3.7	12.5	17.7	26.2	25.6	21.5	34.1	91.7	311.9
All races, female												
2004.....	16.9	19.3	4.8	1.5	3.8	6.5	12.9	14.4	11.2	18.9	64.8	227.3
2003.....	16.2	17.3	5.0	1.4	3.6	6.3	12.0	13.3	10.3	19.3	61.8	226.5
2002.....	15.4	17.2	4.7	1.5	3.2	5.7	11.1	11.5	9.2	18.1	61.4	226.3
2001.....	14.5	17.9	5.1	1.6	2.8	4.9	9.4	9.2	9.1	18.1	59.5	223.1
2000.....	13.6	16.2	5.7	1.7	2.7	4.5	8.2	7.9	8.8	17.5	56.3	218.4
1999.....	13.7	14.2	6.1	1.7	2.6	4.4	8.0	7.6	8.2	17.8	59.4	225.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)												
White, both sexes												
2004.....	23.3	19.0	6.0	1.8	10.8	15.1	22.1	24.1	17.0	26.3	80.4	261.5
2003.....	22.5	16.4	6.6	1.8	10.3	14.3	21.6	21.9	16.6	26.3	77.5	260.6
2002.....	21.4	15.9	5.9	1.9	9.6	13.7	20.8	19.8	15.1	25.5	76.1	256.5
2001.....	20.0	16.9	6.7	2.1	8.7	11.7	18.2	17.3	14.3	24.6	74.2	255.7
2000.....	18.9	15.1	6.9	2.2	8.2	11.4	17.2	15.8	14.2	23.2	69.0	250.6
1999.....	19.0	14.3	7.3	2.3	7.9	11.3	17.0	15.3	13.5	24.1	72.3	258.1
White, male												
2004.....	28.3	22.1	7.3	2.4	17.0	22.6	30.4	33.6	23.1	35.2	100.3	310.8
2003.....	27.4	18.6	8.3	2.3	16.3	21.5	30.6	30.7	23.1	34.6	97.8	312.9
2002.....	26.2	19.2	7.4	2.6	15.3	20.9	29.9	28.3	21.3	34.3	95.2	302.8
2001.....	24.5	18.9	8.4	2.7	14.1	18.0	27.2	25.7	19.9	32.3	94.0	314.0
2000.....	23.4	17.3	8.4	2.9	13.4	18.0	26.2	24.1	20.2	30.4	86.2	308.2
1999.....	23.5	16.5	8.8	3.0	12.8	18.1	26.1	23.5	19.6	32.3	91.3	319.8
White, female												
2004.....	18.5	15.6	4.6	1.2	4.2	7.2	13.6	14.6	11.3	18.7	66.9	239.3
2003.....	17.7	14.2	4.9	1.1	3.9	6.8	12.4	13.2	10.4	19.2	63.8	237.4
2002.....	16.8	12.5	4.4	1.2	3.5	6.1	11.5	11.3	9.2	18.0	63.3	236.3
2001.....	15.6	14.8	4.8	1.5	3.0	5.0	9.2	9.0	9.0	18.0	60.9	231.2
2000.....	14.6	12.8	5.2	1.4	2.7	4.5	8.0	7.5	8.6	17.2	57.7	227.2
1999.....	14.6	11.8	5.8	1.5	2.7	4.3	7.8	7.2	7.7	17.3	59.9	233.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)												
Black, both sexes												
2004.....	17.6	41.0	7.4	4.1	6.2	10.6	19.6	30.4	26.4	31.7	65.4	137.6
2003.....	17.2	39.3	8.4	3.6	5.9	10.6	20.3	30.3	23.9	32.2	60.3	139.2
2002.....	17.8	43.6	9.2	4.5	6.6	11.0	20.9	29.0	24.3	32.2	61.9	155.0
2001.....	17.6	40.2	8.4	4.1	6.5	10.3	21.3	28.9	23.8	32.3	67.3	166.1
2000.....	17.6	38.0	10.6	4.6	7.0	11.0	21.1	27.7	23.5	32.8	64.9	159.5
1999.....	18.9	34.4	10.6	5.1	7.5	12.3	21.0	28.7	26.4	38.5	77.2	179.8
Black, male												
2004.....	23.9	42.4	8.9	5.4	9.9	16.5	27.3	45.7	42.8	47.0	93.5	184.8
2003.....	23.5	44.0	10.6	4.6	9.2	16.2	29.5	45.5	38.7	47.7	85.7	182.5
2002.....	24.8	45.9	12.1	6.3	11.2	17.5	30.9	44.4	40.4	48.4	89.2	214.4
2001.....	24.4	46.4	9.7	5.8	10.7	15.8	31.0	47.1	37.9	47.5	99.8	220.1
2000.....	24.9	40.9	12.6	6.3	11.4	17.4	32.3	45.8	39.3	47.3	96.3	230.6
1999.....	26.8	40.0	13.0	7.0	12.9	19.5	32.9	48.3	42.2	59.1	106.7	248.4
Black, female												
2004.....	11.9	39.5	5.9	2.7	2.5	5.2	12.7	17.1	13.2	20.9	49.6	118.9
2003.....	11.6	34.3	6.0	2.6	2.6	5.5	12.2	17.3	11.9	21.2	45.9	122.3
2002.....	11.4	41.2	6.2	2.7	2.0	5.1	12.0	15.9	11.4	20.7	46.4	132.3
2001.....	11.5	33.7	7.1	2.4	2.4	5.3	12.6	13.2	12.5	21.6	49.1	146.1
2000.....	11.0	35.1	8.5	2.9	2.7	5.3	11.3	12.1	10.6	22.7	47.2	133.1
1999.....	11.7	28.5	8.1	3.1	2.0	5.9	10.5	12.0	13.8	24.0	60.8	154.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Falls (W00-W19)												
All races 3/, both sexes												
2004.....	6.4	0.6	0.3	0.1	0.6	0.8	1.5	2.8	4.8	12.2	43.8	143.3
2003.....	5.9	*	0.3	0.1	0.6	0.7	1.4	2.6	4.4	11.2	40.8	135.9
2002.....	5.6	*	0.2	0.1	0.6	0.8	1.5	2.5	4.1	10.8	38.3	130.4
2001.....	5.3	0.6	0.2	0.2	0.6	0.9	1.4	2.6	4.0	10.0	35.3	121.5
2000.....	4.7	*	0.2	0.1	0.6	0.8	1.3	2.3	3.9	9.0	31.1	112.6
1999.....	4.7	*	0.4	0.1	0.6	0.9	1.4	2.3	3.7	9.0	31.5	110.2
All races, male												
2004.....	6.8	*	0.4	0.1	1.0	1.3	2.4	4.5	7.1	16.5	54.8	174.4
2003.....	6.2	*	0.5	0.1	0.9	1.2	2.4	4.0	6.2	14.5	51.6	165.1
2002.....	6.0	*	0.4	0.1	1.0	1.3	2.3	3.9	6.0	14.5	48.1	158.9
2001.....	5.8	*	0.3	0.2	1.1	1.5	2.4	4.3	5.9	13.1	45.7	155.7
2000.....	5.2	*	0.3	0.1	1.0	1.4	2.2	3.7	5.8	11.9	39.9	145.4
1999.....	5.2	*	0.5	0.2	1.1	1.5	2.4	3.7	5.7	12.4	40.5	142.5
All races, female												
2004.....	6.0	*	*	*	0.2	0.2	0.6	1.3	2.7	8.6	36.4	129.2
2003.....	5.6	*	*	*	0.2	0.2	0.5	1.2	2.7	8.4	33.5	122.9
2002.....	5.3	*	*	*	0.2	0.2	0.6	1.2	2.3	7.7	31.8	118.1
2001.....	4.8	*	*	0.1	0.1	0.2	0.5	1.0	2.2	7.4	28.5	107.2
2000.....	4.3	*	*	*	0.2	0.2	0.5	0.9	2.1	6.6	25.3	99.2
1999.....	4.3	*	*	*	0.2	0.2	0.5	0.9	1.9	6.3	25.7	97.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Falls (W00-W19)												
White, both sexes												
2004.....	7.3	*	0.3	0.1	0.7	0.9	1.6	2.9	4.8	12.7	46.4	152.6
2003.....	6.7	*	0.3	0.1	0.6	0.8	1.5	2.5	4.4	11.5	42.8	144.4
2002.....	6.4	*	0.2	0.1	0.7	0.9	1.5	2.6	4.1	11.1	40.4	138.0
2001.....	6.0	*	0.2	0.2	0.8	0.9	1.5	2.6	4.0	10.4	37.0	128.0
2000.....	5.3	*	0.2	0.1	0.7	0.8	1.4	2.3	3.9	9.2	32.6	119.1
1999.....	5.2	*	0.4	0.1	0.7	0.9	1.5	2.2	3.5	9.1	32.9	115.9
White, male												
2004.....	7.6	*	0.4	*	1.1	1.4	2.5	4.5	6.9	17.1	57.6	183.7
2003.....	6.9	*	0.5	0.2	1.0	1.3	2.4	3.9	6.2	14.7	53.8	175.1
2002.....	6.6	*	*	0.1	1.1	1.5	2.4	3.9	6.0	14.8	50.2	166.4
2001.....	6.4	*	0.3	0.2	1.3	1.6	2.5	4.2	5.8	13.5	47.4	162.3
2000.....	5.7	*	*	0.2	1.2	1.5	2.2	3.7	5.7	12.1	41.4	152.8
1999.....	5.6	*	0.5	0.2	1.2	1.5	2.4	3.5	5.4	12.3	41.9	149.1
White, female												
2004.....	7.0	*	*	*	0.2	0.2	0.6	1.3	2.8	8.9	38.7	138.6
2003.....	6.5	*	*	*	0.2	0.3	0.5	1.1	2.8	8.8	35.4	130.8
2002.....	6.2	*	*	*	0.2	0.2	0.7	1.2	2.4	8.0	33.9	125.6
2001.....	5.5	*	*	*	0.1	0.2	0.4	1.1	2.2	7.7	30.2	113.6
2000.....	5.0	*	*	*	0.2	0.2	0.5	0.9	2.1	6.7	26.8	105.4
1999.....	4.9	*	*	*	0.2	0.2	0.6	0.8	1.8	6.4	27.1	102.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Falls (W00-W19)												
Black, both sexes												
2004.....	2.6	*	*	*	*	0.7	1.4	3.0	5.6	10.1	20.3	46.5
2003.....	2.4	*	*	*	*	0.5	1.3	3.3	4.8	8.9	19.4	44.1
2002.....	2.2	*	*	*	*	0.4	1.3	2.9	4.2	8.2	16.1	49.3
2001.....	2.3	*	*	*	*	0.5	1.6	2.9	4.4	8.7	18.5	44.5
2000.....	2.1	*	*	*	*	0.4	1.3	2.7	4.7	8.3	14.9	38.8
1999.....	2.4	*	*	*	0.4	0.8	1.3	3.4	5.5	9.2	17.1	45.3
Black, male												
2004.....	3.4	*	*	*	*	1.1	2.1	4.9	10.1	14.4	29.2	71.5
2003.....	3.0	*	*	*	*	0.8	2.2	5.2	7.1	12.5	24.8	56.3
2002.....	3.0	*	*	*	*	0.8	2.2	4.8	6.7	11.8	24.4	77.2
2001.....	3.1	*	*	*	*	0.8	2.6	5.4	7.0	12.6	28.5	68.1
2000.....	2.8	*	*	*	*	*	2.1	4.7	8.2	11.4	21.2	61.1
1999.....	3.3	*	*	*	*	1.3	2.5	5.8	8.8	14.7	24.3	58.6
Black, female												
2004.....	1.8	*	*	*	*	*	0.8	1.3	2.0	7.0	15.3	36.6
2003.....	1.9	*	*	*	*	*	*	1.6	2.9	6.3	16.3	39.3
2002.....	1.5	*	*	*	*	*	*	1.2	2.2	5.6	11.4	38.6
2001.....	1.6	*	*	*	*	*	*	*	2.3	5.9	12.9	35.8
2000.....	1.4	*	*	*	*	*	*	1.0	2.0	6.1	11.4	30.5
1999.....	1.7	*	*	*	*	*	*	1.3	2.9	5.3	13.1	40.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental discharge of firearms (W32-W34)

All races 3/, both sexes

2004.....	0.2	*	*	0.1	0.4	0.2	0.2	0.3	0.1	0.2	0.2	*
2003.....	0.3	*	*	0.1	0.5	0.3	0.3	0.2	0.2	0.2	0.3	*
2002.....	0.3	*	*	0.1	0.5	0.4	0.3	0.2	0.2	0.2	0.3	*
2001.....	0.3	*	*	0.1	0.5	0.3	0.3	0.3	0.3	0.2	0.3	*
2000.....	0.3	*	*	0.2	0.5	0.3	0.3	0.2	0.2	0.2	0.2	*
1999.....	0.3	*	*	0.2	0.6	0.4	0.3	0.3	0.2	0.2	0.2	*

All races, male

2004.....	0.4	*	*	0.2	0.7	0.5	0.4	0.4	0.2	0.4	0.4	*
2003.....	0.5	*	*	0.2	0.9	0.5	0.5	0.4	0.4	0.4	0.6	*
2002.....	0.5	*	*	0.2	0.9	0.6	0.4	0.4	0.4	0.4	0.5	*
2001.....	0.5	*	*	0.2	0.9	0.5	0.5	0.5	0.5	0.3	0.7	*
2000.....	0.5	*	*	0.3	0.9	0.5	0.6	0.4	0.4	0.4	0.5	*
1999.....	0.5	*	*	0.3	1.2	0.6	0.5	0.4	0.4	0.4	0.4	*

All races, female

2004.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000.....	0.1	*	*	*	*	0.1	0.1	*	*	*	*	*
1999.....	0.1	*	*	*	0.1	0.1	0.1	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental discharge of firearms (W32-W34)

White, both sexes

2004.....	0.2	*	*	0.1	0.4	0.2	0.2	0.3	0.1	0.2	0.2	*
2003.....	0.2	*	*	0.1	0.5	0.3	0.3	0.2	0.2	0.2	0.3	*
2002.....	0.3	*	*	0.1	0.4	0.3	0.3	0.2	0.2	0.2	0.2	*
2001.....	0.3	*	*	0.1	0.5	0.3	0.3	0.3	0.3	0.2	0.3	*
2000.....	0.3	*	*	0.2	0.5	0.3	0.3	0.2	0.2	0.2	0.2	*
1999.....	0.3	*	*	0.2	0.6	0.4	0.3	0.3	0.2	0.2	0.2	*

White, male

2004.....	0.4	*	*	0.2	0.6	0.4	0.4	0.5	0.2	0.4	*	*
2003.....	0.4	*	*	0.2	0.8	0.5	0.5	0.4	0.5	0.4	0.6	*
2002.....	0.4	*	*	0.2	0.8	0.5	0.4	0.4	0.4	0.4	0.5	*
2001.....	0.5	*	*	0.3	0.9	0.5	0.6	0.5	0.5	0.3	0.7	*
2000.....	0.5	*	*	0.3	0.8	0.5	0.6	0.4	0.4	0.4	0.5	*
1999.....	0.5	*	*	0.3	1.0	0.6	0.5	0.4	0.4	0.4	*	*

White, female

2004.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental drowning and submersion (W65-W74)												
All races 3/, both sexes												
2004.....	1.1	1.5	2.7	0.7	1.4	1.0	1.0	1.1	0.9	1.0	1.2	1.6
2003.....	1.1	1.4	2.9	0.7	1.4	0.9	1.0	1.0	0.9	1.1	1.3	1.8
2002.....	1.2	1.6	2.9	0.8	1.5	1.1	1.2	0.9	0.8	0.9	1.3	1.7
2001.....	1.2	1.7	3.0	0.8	1.5	0.9	1.0	0.9	0.8	0.9	1.2	1.7
2000.....	1.2	2.0	3.2	0.9	1.6	1.1	1.1	0.9	0.9	1.0	1.3	1.5
1999.....	1.3	1.8	3.2	0.9	1.7	1.1	1.1	1.0	0.9	1.0	1.3	1.6
All races, male												
2004.....	1.8	1.8	3.2	0.9	2.4	1.6	1.7	1.8	1.5	1.7	2.2	2.9
2003.....	1.8	1.7	3.7	0.9	2.5	1.6	1.8	1.6	1.6	1.7	2.0	3.3
2002.....	1.9	1.9	3.9	1.2	2.8	1.9	2.0	1.5	1.3	1.6	2.1	3.6
2001.....	1.8	1.5	3.7	1.1	2.7	1.6	1.7	1.6	1.3	1.5	2.0	3.5
2000.....	2.0	2.2	4.0	1.3	2.9	1.8	1.8	1.6	1.5	1.6	1.8	2.8
1999.....	2.0	2.0	4.0	1.3	3.0	2.0	1.8	1.7	1.5	1.6	2.3	2.3
All races, female												
2004.....	0.5	1.2	2.1	0.4	0.3	0.3	0.3	0.4	0.4	0.5	0.6	1.0
2003.....	0.5	1.2	2.0	0.3	0.2	0.2	0.3	0.4	0.4	0.6	0.8	1.1
2002.....	0.5	1.2	1.9	0.4	0.3	0.3	0.4	0.4	0.4	0.4	0.8	0.9
2001.....	0.5	1.9	2.2	0.5	0.3	0.2	0.4	0.3	0.4	0.3	0.7	1.0
2000.....	0.5	1.7	2.4	0.5	0.3	0.3	0.3	0.3	0.3	0.5	0.9	1.0
1999.....	0.5	1.6	2.4	0.5	0.3	0.2	0.3	0.3	0.4	0.6	0.6	1.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental drowning and submersion (W65-W74)												
White, both sexes												
2004.....	1.1	1.6	3.0	0.4	1.2	0.9	1.0	1.1	0.9	1.0	1.2	1.5
2003.....	1.1	1.4	3.1	0.5	1.3	0.9	1.0	1.0	0.9	1.1	1.3	1.6
2002.....	1.1	1.6	3.0	0.5	1.4	1.0	1.1	0.9	0.8	0.9	1.3	1.6
2001.....	1.1	1.7	3.3	0.5	1.3	0.9	1.0	0.9	0.8	0.9	1.2	1.7
2000.....	1.2	1.6	3.5	0.7	1.5	1.0	1.0	0.9	0.9	1.0	1.2	1.5
1999.....	1.2	1.5	3.5	0.6	1.5	1.0	1.1	1.0	0.9	1.0	1.3	1.6
White, male												
2004.....	1.7	1.8	3.5	0.6	2.2	1.6	1.6	1.7	1.4	1.6	2.1	2.8
2003.....	1.8	1.5	4.0	0.7	2.3	1.5	1.7	1.5	1.5	1.8	2.1	3.3
2002.....	1.8	1.9	4.0	0.7	2.5	1.8	1.9	1.4	1.2	1.5	2.1	3.7
2001.....	1.7	1.6	4.2	0.7	2.4	1.4	1.6	1.5	1.2	1.5	2.0	3.5
2000.....	1.9	1.8	4.4	0.9	2.6	1.7	1.7	1.5	1.6	1.6	1.8	2.7
1999.....	1.9	1.7	4.3	0.8	2.7	1.8	1.8	1.7	1.4	1.4	2.2	2.4
White, female												
2004.....	0.5	1.4	2.4	0.3	0.2	0.3	0.3	0.4	0.4	0.4	0.5	0.9
2003.....	0.5	1.3	2.2	0.3	0.2	0.2	0.3	0.4	0.4	0.6	0.7	0.9
2002.....	0.4	*	2.0	0.3	0.2	0.3	0.4	0.3	0.4	0.3	0.7	0.7
2001.....	0.5	1.8	2.4	0.4	0.3	0.3	0.4	0.3	0.4	0.3	0.7	1.0
2000.....	0.5	*	2.5	0.4	0.3	0.2	0.3	0.3	0.3	0.5	0.8	1.0
1999.....	0.5	1.4	2.6	0.3	0.3	0.2	0.4	0.3	0.4	0.5	0.6	1.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)												
All races 3/, both sexes												
2004.....	1.1	0.6	1.4	0.6	0.4	0.6	0.7	1.2	1.4	1.9	3.5	5.6
2003.....	1.2	0.7	1.4	0.5	0.5	0.6	0.9	1.3	1.4	2.3	3.6	5.4
2002.....	1.1	0.9	1.4	0.6	0.5	0.7	0.8	1.1	1.3	2.1	3.4	4.8
2001.....	1.2	1.2	1.5	0.6	0.5	0.6	1.0	1.1	1.5	2.2	3.4	5.7
2000.....	1.2	1.0	1.9	0.6	0.5	0.6	0.9	1.2	1.5	2.2	3.8	6.2
1999.....	1.2	1.1	2.0	0.6	0.5	0.7	0.9	1.0	1.3	2.3	4.0	5.9
All races, male												
2004.....	1.3	*	1.5	0.7	0.5	0.8	0.9	1.6	1.8	2.5	4.5	7.2
2003.....	1.4	*	1.6	0.5	0.6	0.8	1.2	1.6	1.9	2.9	4.4	7.4
2002.....	1.4	1.0	1.6	0.7	0.6	0.8	1.1	1.5	1.7	2.9	4.5	7.0
2001.....	1.4	1.5	1.8	0.7	0.7	0.8	1.4	1.5	1.9	2.8	4.1	8.3
2000.....	1.5	1.0	2.2	0.7	0.6	0.8	1.2	1.6	2.1	2.8	5.0	9.4
1999.....	1.5	1.1	2.2	0.7	0.7	1.0	1.3	1.4	1.7	2.9	5.1	9.3
All races, female												
2004.....	0.9	*	1.3	0.5	0.4	0.4	0.5	0.8	1.1	1.4	2.9	4.9
2003.....	1.0	1.1	1.2	0.5	0.4	0.5	0.6	0.9	1.0	1.8	3.0	4.5
2002.....	0.8	*	1.2	0.5	0.3	0.5	0.5	0.8	0.9	1.4	2.6	3.9
2001.....	0.9	1.0	1.1	0.5	0.4	0.4	0.6	0.7	1.1	1.8	2.9	4.6
2000.....	0.9	*	1.5	0.6	0.4	0.4	0.6	0.7	1.0	1.7	3.0	4.9
1999.....	0.9	1.1	1.7	0.6	0.4	0.4	0.5	0.6	1.0	1.7	3.2	4.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)												
White, both sexes												
2004.....	1.0	*	1.1	0.4	0.4	0.6	0.7	1.0	1.2	1.6	3.1	4.9
2003.....	1.1	0.7	1.2	0.4	0.5	0.6	0.9	1.2	1.3	2.0	3.1	4.7
2002.....	1.0	*	1.0	0.4	0.4	0.6	0.8	1.0	1.1	1.8	3.0	4.0
2001.....	1.0	1.1	1.2	0.5	0.5	0.6	0.9	0.9	1.3	1.9	2.8	4.8
2000.....	1.0	0.8	1.3	0.5	0.5	0.5	0.8	1.0	1.3	1.8	3.1	5.1
1999.....	1.0	0.9	1.5	0.5	0.5	0.7	0.8	0.9	1.2	1.8	3.3	4.7
White, male												
2004.....	1.2	*	1.2	0.5	0.5	0.8	0.9	1.4	1.6	2.1	3.9	6.5
2003.....	1.3	*	1.4	0.4	0.6	0.7	1.2	1.5	1.7	2.6	3.6	6.7
2002.....	1.2	*	1.1	0.6	0.6	0.8	1.1	1.3	1.5	2.5	4.1	5.8
2001.....	1.3	*	1.6	0.6	0.7	0.7	1.2	1.2	1.6	2.2	3.4	7.0
2000.....	1.3	*	1.6	0.5	0.6	0.7	1.1	1.5	1.7	2.3	4.1	8.2
1999.....	1.3	*	1.7	0.5	0.7	1.0	1.2	1.2	1.5	2.3	4.2	8.0
White, female												
2004.....	0.8	*	1.0	0.4	0.4	0.4	0.5	0.7	0.9	1.2	2.5	4.2
2003.....	0.9	*	1.0	0.4	0.4	0.5	0.6	0.8	0.9	1.5	2.7	3.8
2002.....	0.7	*	1.0	0.3	0.3	0.4	0.5	0.7	0.7	1.3	2.2	3.2
2001.....	0.8	*	0.8	0.4	0.3	0.4	0.5	0.6	0.9	1.6	2.5	3.8
2000.....	0.8	*	1.1	0.5	0.3	0.4	0.5	0.6	0.8	1.5	2.4	3.9
1999.....	0.8	*	1.3	0.5	0.3	0.4	0.5	0.5	0.8	1.4	2.6	3.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)												
Black, both sexes												
2004.....	2.1	*	2.8	1.6	0.6	1.0	1.2	2.4	3.4	4.8	9.7	14.9
2003.....	2.0	*	3.0	1.4	0.6	0.9	1.2	2.6	2.9	5.3	10.2	14.9
2002.....	2.1	*	3.5	1.4	0.8	1.1	1.0	2.1	3.3	5.0	8.6	15.7
2001.....	2.3	*	3.3	1.3	0.7	1.2	1.8	2.4	3.5	6.1	9.7	17.8
2000.....	2.5	*	4.8	1.5	0.7	1.2	1.7	2.4	4.2	5.9	13.2	20.3
1999.....	2.5	*	4.5	1.5	0.7	1.0	1.8	2.1	3.4	6.9	13.2	21.7
Black, male												
2004.....	2.6	*	2.9	1.9	0.7	1.3	1.7	3.8	4.4	7.5	13.4	*
2003.....	2.4	*	3.2	1.5	*	1.1	1.4	3.3	4.3	7.5	16.1	*
2002.....	2.6	*	4.1	1.6	1.0	1.4	1.7	3.0	4.7	7.8	10.9	25.7
2001.....	2.9	*	3.6	1.6	0.8	1.4	2.7	3.9	5.0	9.0	12.9	27.2
2000.....	3.2	*	5.6	1.7	0.9	1.5	2.4	3.5	6.3	8.8	18.5	27.7
1999.....	3.0	*	5.0	1.7	0.8	1.2	2.7	3.2	4.1	10.1	18.1	29.3
Black, female												
2004.....	1.6	*	2.7	1.2	*	*	0.7	1.2	2.6	3.0	7.6	13.4
2003.....	1.7	*	2.7	1.3	0.6	*	1.0	2.0	1.7	3.7	6.9	13.4
2002.....	1.6	*	2.8	1.3	*	0.9	*	1.4	2.2	3.0	7.3	11.9
2001.....	1.7	*	2.9	1.0	*	1.1	1.0	1.1	2.4	4.1	7.9	14.3
2000.....	2.0	*	4.1	1.3	*	0.9	1.1	1.4	2.4	3.9	10.2	17.6
1999.....	2.0	*	3.9	1.3	*	0.8	1.0	1.3	2.7	4.6	10.5	18.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)												
All races 3/, both sexes												
2004.....	7.1	*	*	0.1	5.4	9.1	14.6	14.5	5.4	2.3	2.4	3.4
2003.....	6.7	0.5	0.3	0.1	4.9	8.6	14.0	13.3	4.9	2.2	2.2	3.5
2002.....	6.1	0.6	0.2	0.1	4.1	7.8	13.4	11.7	4.1	2.1	2.3	3.9
2001.....	4.9	*	0.2	0.1	3.4	6.3	11.2	9.1	3.2	1.6	2.2	3.5
2000.....	4.5	*	0.2	0.1	3.0	6.0	10.3	8.1	2.8	1.5	2.0	4.3
1999.....	4.4	*	0.2	0.1	2.5	5.9	10.1	7.8	2.8	1.6	2.2	3.4
All races, male												
2004.....	9.6	*	*	0.1	8.2	13.1	19.0	19.1	6.6	2.6	2.7	4.2
2003.....	9.2	*	0.3	0.1	7.4	12.4	18.8	18.0	6.3	2.7	2.8	4.0
2002.....	8.5	1.0	*	0.1	6.3	11.5	18.4	15.9	5.4	2.5	2.7	5.1
2001.....	7.1	*	0.3	0.1	5.3	9.3	15.7	13.0	4.1	2.0	2.7	3.8
2000.....	6.6	*	0.3	0.1	4.6	9.0	15.1	12.0	3.4	1.8	2.1	6.2
1999.....	6.5	*	*	0.1	3.8	9.0	14.9	11.7	3.9	2.1	2.4	4.4
All races, female												
2004.....	4.7	*	*	0.1	2.4	4.9	10.3	10.0	4.3	2.0	2.2	3.0
2003.....	4.3	*	0.3	0.2	2.2	4.7	9.3	8.8	3.7	1.8	1.8	3.3
2002.....	3.7	*	*	*	1.9	4.0	8.4	7.6	3.0	1.7	2.1	3.4
2001.....	2.9	*	*	0.1	1.4	3.3	6.7	5.2	2.3	1.2	1.9	3.3
2000.....	2.5	*	*	*	1.3	2.8	5.6	4.4	2.3	1.3	1.9	3.6
1999.....	2.3	*	*	*	1.1	2.6	5.3	4.0	1.8	1.3	2.0	3.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)												
White, both sexes												
2004.....	7.6	*	*	0.1	6.4	10.3	15.7	14.6	5.0	2.2	2.4	3.4
2003.....	7.1	*	0.3	0.1	5.8	9.7	14.8	13.3	4.8	2.0	2.3	3.5
2002.....	6.3	*	*	0.1	4.9	8.7	14.0	11.3	3.8	2.1	2.3	3.9
2001.....	5.0	*	0.2	0.1	3.9	6.9	11.4	8.4	2.8	1.5	2.1	3.3
2000.....	4.6	*	*	0.1	3.4	6.4	10.4	7.5	2.6	1.4	2.1	4.1
1999.....	4.3	*	0.2	0.1	2.8	6.1	10.0	7.1	2.5	1.5	2.2	3.3
White, male												
2004.....	10.2	*	*	0.2	9.7	14.9	20.3	18.9	5.7	2.4	2.5	4.3
2003.....	9.7	*	*	0.1	8.8	13.9	19.8	17.6	5.8	2.2	2.8	3.7
2002.....	8.8	*	*	0.2	7.4	12.7	19.2	15.2	4.5	2.3	2.6	5.2
2001.....	7.2	*	*	*	6.0	10.2	16.2	11.9	3.4	1.9	2.5	3.7
2000.....	6.7	*	*	*	5.3	9.7	15.2	10.9	2.9	1.6	2.1	5.9
1999.....	6.4	*	*	*	4.2	9.5	14.9	10.5	3.2	1.9	2.5	4.5
White, female												
2004.....	5.0	*	*	*	2.8	5.5	11.0	10.3	4.4	2.0	2.2	3.0
2003.....	4.5	*	*	*	2.6	5.2	9.8	9.0	3.8	1.8	1.9	3.4
2002.....	3.9	*	*	*	2.2	4.5	8.7	7.5	3.0	1.8	2.1	3.4
2001.....	2.9	*	*	0.1	1.6	3.5	6.6	5.0	2.3	1.2	1.8	3.1
2000.....	2.5	*	*	*	1.3	3.0	5.5	4.1	2.3	1.3	2.0	3.4
1999.....	2.3	*	*	*	1.2	2.6	5.1	3.8	1.8	1.2	2.1	2.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Black, both sexes

2004.....	6.5	*	*	*	1.8	5.3	12.8	18.8	9.8	3.7	3.5	*
2003.....	6.3	*	*	*	1.4	5.7	13.5	17.9	7.6	4.1	2.0	*
2002.....	6.3	*	*	*	1.4	5.6	13.8	17.9	8.4	2.7	2.8	*
2001.....	5.8	*	*	*	1.6	5.2	13.1	16.7	6.5	2.5	3.6	6.4
2000.....	5.5	*	*	*	1.3	5.6	13.0	15.5	5.5	2.6	*	8.1
1999.....	5.7	*	*	*	1.6	6.1	13.1	15.5	6.0	3.0	*	*

Black, male

2004.....	8.9	*	*	*	2.7	7.6	16.4	27.1	16.2	6.2	5.6	*
2003.....	8.8	*	*	*	2.0	8.0	18.6	26.2	12.8	7.6	*	*
2002.....	9.1	*	*	*	2.2	8.5	19.3	26.5	15.0	5.0	*	*
2001.....	8.4	*	*	*	2.3	7.5	17.9	26.6	11.3	3.9	5.9	*
2000.....	8.2	*	*	*	1.7	8.4	18.7	24.9	9.5	4.5	*	*
1999.....	8.7	*	*	*	2.7	9.1	19.4	25.9	11.0	4.4	*	*

Black, female

2004.....	4.3	*	*	*	1.0	3.3	9.7	11.8	4.7	*	*	*
2003.....	3.9	*	*	*	0.7	3.6	8.9	10.8	3.3	*	*	*
2002.....	3.7	*	*	*	0.7	3.0	8.9	10.5	3.2	*	*	*
2001.....	3.5	*	*	*	0.9	3.1	9.0	8.2	2.6	*	*	*
2000.....	3.1	*	*	*	1.0	3.0	7.8	7.4	2.3	*	*	*
1999.....	3.0	*	*	*	*	3.4	7.6	6.5	2.0	2.1	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All races 3/, both sexes

2004.....	5.8	19.2	1.6	0.6	1.5	2.1	2.9	4.2	5.0	8.8	27.0	95.5
2003.....	5.9	16.9	1.8	0.5	1.6	2.0	3.0	3.9	5.1	9.6	27.5	102.5
2002.....	6.0	16.8	1.6	0.7	1.6	2.1	3.0	3.7	5.2	9.8	28.8	105.6
2001.....	6.2	16.6	1.8	0.6	1.6	2.0	3.1	4.2	5.3	10.0	30.5	115.0
2000.....	6.2	15.0	1.7	0.6	1.7	2.2	3.2	4.0	5.6	9.9	29.8	117.1
1999.....	6.6	13.8	1.9	0.8	1.7	2.2	3.2	4.1	5.6	11.0	32.9	128.9

All races, male

2004.....	6.8	21.7	2.0	0.9	2.6	3.4	4.6	6.6	7.5	11.8	33.6	109.8
2003.....	6.9	19.5	2.2	0.8	2.6	3.4	4.8	6.0	7.8	13.1	34.9	120.2
2002.....	6.9	18.8	2.0	0.9	2.6	3.4	4.8	6.0	7.9	13.4	36.0	118.6
2001.....	7.2	18.9	2.1	0.9	2.6	3.4	5.0	6.6	7.7	13.2	38.4	134.0
2000.....	7.2	16.8	2.1	0.9	2.8	3.8	5.4	6.6	8.3	13.0	36.9	135.2
1999.....	7.6	16.2	2.3	1.1	2.9	3.7	5.3	6.6	8.3	14.8	40.9	152.7

All races, female

2004.....	4.8	16.5	1.1	0.3	0.4	0.7	1.2	1.8	2.7	6.4	22.6	89.1
2003.....	4.9	14.1	1.3	0.3	0.5	0.7	1.2	1.9	2.6	6.7	22.6	94.7
2002.....	5.0	14.6	1.2	0.4	0.5	0.6	1.2	1.5	2.7	6.8	24.0	100.0
2001.....	5.3	14.2	1.5	0.4	0.6	0.6	1.1	1.9	3.1	7.3	25.4	107.0
2000.....	5.2	13.2	1.4	0.4	0.5	0.6	1.1	1.5	3.0	7.4	25.1	109.8
1999.....	5.6	11.3	1.5	0.5	0.5	0.7	1.2	1.7	3.1	8.0	27.7	119.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

White, both sexes

2004.....	6.2	16.0	1.5	0.6	1.7	2.2	3.0	4.3	4.9	8.7	27.2	98.8
2003.....	6.3	13.6	1.7	0.5	1.7	2.2	3.1	3.8	4.9	9.4	27.8	106.3
2002.....	6.3	12.9	1.4	0.7	1.7	2.2	3.1	3.7	5.1	9.4	28.8	108.8
2001.....	6.6	13.4	1.7	0.6	1.8	2.1	3.1	4.1	5.1	9.7	30.6	117.7
2000.....	6.6	12.2	1.6	0.7	1.7	2.4	3.2	3.8	5.4	9.5	29.8	120.5
1999.....	6.9	11.2	1.8	0.8	1.9	2.3	3.3	3.9	5.2	10.6	32.5	132.4

White, male

2004.....	7.2	19.0	2.0	0.9	2.8	3.6	4.7	6.7	7.3	11.6	33.7	112.9
2003.....	7.3	15.7	2.2	0.8	2.8	3.6	4.9	5.8	7.4	12.9	34.9	123.9
2002.....	7.3	15.3	1.8	0.9	2.9	3.7	5.0	6.0	7.6	12.7	35.7	121.2
2001.....	7.5	15.2	2.0	0.9	2.9	3.5	5.1	6.4	7.4	12.9	38.0	137.0
2000.....	7.4	14.0	2.0	0.9	2.9	4.0	5.4	6.2	7.9	12.5	36.2	137.6
1999.....	7.8	12.9	2.1	1.1	3.1	3.8	5.4	6.2	7.7	14.0	40.1	155.2

White, female

2004.....	5.2	12.8	0.9	0.3	0.5	0.7	1.3	1.9	2.7	6.2	22.8	92.5
2003.....	5.3	11.4	1.2	0.3	0.5	0.7	1.2	1.8	2.5	6.5	23.0	98.5
2002.....	5.4	10.5	1.1	0.4	0.5	0.7	1.1	1.5	2.6	6.5	24.2	103.4
2001.....	5.8	11.5	1.4	0.4	0.6	0.6	1.1	1.9	3.0	7.1	25.7	109.7
2000.....	5.7	10.3	1.2	0.4	0.5	0.6	1.0	1.5	3.0	7.1	25.6	113.6
1999.....	6.1	9.5	1.4	0.5	0.6	0.7	1.2	1.6	2.8	7.7	27.4	123.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Black, both sexes

2004.....	4.8	37.1	1.8	0.6	1.0	1.9	2.9	4.6	6.5	11.7	30.6	70.2
2003.....	4.9	34.9	2.0	0.5	1.3	1.8	3.2	5.4	7.3	12.8	27.6	73.2
2002.....	5.2	37.5	2.4	0.8	1.1	1.7	3.1	4.7	7.3	15.2	32.3	83.1
2001.....	5.4	34.2	2.5	0.6	1.0	1.9	3.2	5.6	7.9	14.1	33.8	96.1
2000.....	5.4	30.6	2.3	0.7	1.5	2.2	3.5	5.6	8.2	14.9	33.7	90.1
1999.....	6.1	27.9	2.5	0.7	1.3	2.4	3.4	6.6	10.0	17.9	42.9	105.5

Black, male

2004.....	6.1	38.3	1.7	1.0	1.5	3.3	4.7	7.5	10.3	16.4	42.3	87.4
2003.....	6.4	40.5	2.8	0.7	1.9	2.9	5.1	8.5	11.9	18.3	39.8	91.8
2002.....	6.6	38.9	3.3	1.1	1.7	2.9	4.7	7.4	12.1	21.6	45.6	106.1
2001.....	6.8	40.4	2.8	0.9	1.5	3.2	5.1	9.0	11.9	19.8	48.8	114.6
2000.....	7.3	32.0	2.5	1.0	2.5	3.7	6.0	10.1	13.8	20.3	53.0	124.5
1999.....	8.0	33.2	3.4	0.9	2.3	4.0	5.7	11.1	15.7	26.9	57.2	153.5

Black, female

2004.....	3.6	35.8	2.0	*	*	*	1.2	2.2	3.5	8.4	23.9	63.4
2003.....	3.6	29.1	*	*	0.7	0.7	1.5	2.8	3.6	8.8	20.7	66.0
2002.....	4.0	36.0	*	*	*	*	1.7	2.3	3.5	10.7	24.7	74.3
2001.....	4.1	27.7	2.2	*	*	0.7	1.6	2.7	4.6	10.0	25.3	89.3
2000.....	3.8	29.1	2.0	*	*	0.9	1.3	1.8	3.6	11.1	22.8	77.3
1999.....	4.4	22.5	1.6	*	*	1.0	1.4	2.7	5.4	11.6	34.9	87.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
All races 3/, both sexes												
2004.....	11.0	0.7	10.3	12.7	15.0	16.6	13.8	12.3	16.3	16.4
2003.....	10.8	0.6	9.7	12.7	14.9	15.9	13.8	12.7	16.4	16.9
2002.....	11.0	0.6	9.9	12.6	15.3	15.7	13.6	13.5	17.7	18.0
2001 4/.....	10.8	0.7	9.9	12.8	14.7	15.2	13.1	13.3	17.4	17.5
2000.....	10.4	0.7	10.2	12.0	14.5	14.4	12.1	12.5	17.6	19.6
1999.....	10.5	0.6	10.1	12.7	14.3	13.9	12.2	13.4	18.1	19.3
All races, male												
2004.....	17.7	0.9	16.8	20.4	23.0	24.8	22.1	22.6	34.8	45.0
2003.....	17.6	0.9	16.0	20.6	23.2	24.4	22.3	23.4	35.1	47.8
2002.....	17.9	0.9	16.5	20.5	23.7	24.4	22.2	24.7	38.1	50.7
2001 4/.....	17.6	1.0	16.6	21.0	23.1	23.4	21.1	24.6	37.8	51.1
2000.....	17.1	1.2	17.1	19.6	22.8	22.4	19.4	22.7	38.6	57.5
1999.....	17.1	0.9	16.8	20.7	22.4	21.5	19.8	24.7	38.8	57.1
All races, female												
2004.....	4.6	0.5	3.6	4.7	7.1	8.6	6.1	3.8	3.9	3.6
2003.....	4.3	0.3	3.0	4.6	6.6	7.7	5.9	3.8	4.0	3.3
2002.....	4.3	0.3	2.9	4.6	6.9	7.4	5.7	4.1	4.2	3.8
2001 4/.....	4.1	0.3	2.9	4.4	6.4	7.2	5.7	3.9	4.0	3.4
2000.....	4.0	0.3	3.0	4.3	6.4	6.7	5.4	4.0	4.0	4.2
1999.....	4.0	0.3	3.0	4.6	6.4	6.6	5.2	4.1	4.7	4.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
White, both sexes												
2004.....	12.3	0.7	11.0	13.6	16.9	18.6	15.1	13.4	17.5	17.5
2003.....	12.1	0.6	10.2	13.7	16.6	17.9	15.2	13.8	17.6	18.1
2002.....	12.2	0.7	10.7	13.6	17.0	17.6	15.0	14.7	19.0	19.1
2001 4/.....	11.9	0.7	10.6	13.8	16.4	16.9	14.3	14.3	18.6	18.6
2000.....	11.5	0.8	10.7	12.8	16.1	15.9	13.3	13.5	18.8	21.0
1999.....	11.5	0.6	10.5	13.5	15.8	15.3	13.2	14.4	19.3	20.7
White, male												
2004.....	19.6	0.9	17.9	21.8	25.6	27.6	23.9	24.2	37.1	48.4
2003.....	19.5	0.9	16.9	21.8	25.7	27.3	24.3	25.2	37.5	51.4
2002.....	19.9	1.0	17.7	21.6	26.2	27.0	24.3	26.8	40.6	53.9
2001 4/.....	19.5	1.0	17.6	22.4	25.4	26.0	22.8	26.3	40.2	55.0
2000.....	18.8	1.2	17.9	20.7	24.9	24.5	21.2	24.3	41.1	61.6
1999.....	18.7	1.0	17.5	21.8	24.4	23.4	21.2	26.4	41.2	61.5
White, female												
2004.....	5.1	0.5	3.8	5.0	8.1	9.8	6.8	4.1	4.1	3.6
2003.....	4.7	0.3	3.1	5.1	7.5	8.7	6.5	4.1	4.2	3.3
2002.....	4.8	0.4	3.1	5.2	7.8	8.4	6.2	4.4	4.4	4.0
2001 4/.....	4.6	0.3	3.1	4.9	7.3	8.0	6.4	4.2	4.3	3.4
2000.....	4.4	0.3	3.1	4.6	7.2	7.5	6.0	4.4	4.1	4.5
1999.....	4.5	0.3	3.2	4.9	7.2	7.4	5.7	4.4	4.9	4.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

Black, both sexes

2004.....	5.2	0.7	7.2	9.4	6.6	5.8	5.7	4.4	6.0	*
2003.....	5.1	0.6	7.1	9.3	7.2	5.6	5.0	4.5	4.5	*
2002.....	5.1	0.5	6.5	9.3	7.6	5.9	4.8	4.9	5.5	5.9
2001 4/.....	5.3	0.6	7.1	9.1	7.3	6.2	5.2	5.5	5.7	*
2000.....	5.4	0.6	8.1	8.7	7.6	6.5	4.2	5.1	5.8	*
1999.....	5.4	0.4	8.1	9.2	7.6	5.7	4.9	5.8	5.7	*

Black, male

2004.....	9.0	0.9	12.2	16.2	11.3	9.7	10.8	9.8	15.0	*
2003.....	8.8	1.0	12.1	16.5	12.2	9.1	8.7	8.3	11.3	*
2002.....	9.1	0.8	11.3	17.3	13.0	10.2	8.4	9.7	13.8	*
2001 4/.....	9.2	0.9	13.0	16.5	12.3	9.9	9.4	10.7	13.5	*
2000.....	9.4	1.0	14.2	15.5	13.2	11.2	7.5	11.1	12.1	*
1999.....	9.6	0.6	14.3	16.8	13.3	10.0	9.1	11.8	13.8	*

Black, female

2004.....	1.8	*	2.2	3.2	2.5	2.5	1.6	*	*	*
2003.....	1.8	*	2.0	2.7	2.9	2.6	2.0	*	*	*
2002.....	1.5	*	1.7	2.0	2.9	2.2	1.9	*	*	*
2001 4/.....	1.7	*	1.3	2.4	2.9	3.0	1.8	*	*	*
2000.....	1.7	*	2.2	2.6	2.7	2.5	1.6	*	*	*
1999.....	1.6	*	1.9	2.3	2.6	2.0	1.6	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
All races 3/, both sexes												
2004.....	5.7	0.1	5.0	5.7	6.5	8.0	8.0	8.8	12.2	11.1
2003.....	5.8	0.2	5.0	6.0	6.6	8.0	8.3	9.3	12.4	11.9
2002.....	5.9	0.2	5.1	6.0	7.0	7.8	8.4	9.7	13.2	12.0
2001.....	5.9	0.2	5.3	6.5	6.7	7.7	8.2	9.6	13.1	12.2
2000.....	5.9	0.3	5.8	6.1	7.1	7.6	7.6	9.2	13.0	13.2
1999.....	5.9	0.3	6.0	6.5	6.9	7.3	7.8	9.7	13.1	12.6
All races, male												
2004.....	10.0	0.2	8.8	9.7	10.8	13.5	14.2	17.5	28.3	34.4
2003.....	10.4	0.3	8.9	10.4	11.2	13.8	14.9	18.4	28.7	36.6
2002.....	10.6	0.3	9.1	10.3	11.9	13.6	15.1	19.5	31.0	37.8
2001.....	10.6	0.3	9.4	11.2	11.4	13.3	14.6	19.2	30.7	38.9
2000.....	10.5	0.4	10.2	10.6	11.8	13.0	13.5	18.4	30.8	42.9
1999.....	10.6	0.4	10.5	11.2	11.5	12.5	14.0	19.5	31.1	41.6
All races, female												
2004.....	1.5	*	1.1	1.5	2.2	2.8	2.2	1.6	1.4	*
2003.....	1.4	*	1.0	1.4	2.1	2.5	2.2	1.6	1.5	0.9
2002.....	1.4	*	1.0	1.6	2.2	2.2	2.2	1.6	1.4	0.8
2001.....	1.5	0.1	1.0	1.6	2.1	2.4	2.4	1.7	1.5	1.0
2000.....	1.5	0.1	1.2	1.7	2.4	2.3	2.1	1.7	1.4	1.1
1999.....	1.5	0.1	1.2	1.8	2.3	2.3	2.1	1.7	1.4	0.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
White, both sexes												
2004.....	6.5	0.2	5.4	6.1	7.4	9.2	9.0	9.7	13.3	12.1
2003.....	6.6	0.2	5.3	6.4	7.6	9.2	9.3	10.3	13.4	12.9
2002.....	6.7	0.2	5.5	6.5	7.9	8.9	9.4	10.8	14.2	12.9
2001.....	6.7	0.2	5.6	7.1	7.5	8.7	9.2	10.5	14.1	13.2
2000.....	6.6	0.3	6.1	6.6	7.9	8.5	8.3	10.1	14.1	14.3
1999.....	6.6	0.3	6.2	7.0	7.7	8.2	8.6	10.6	14.1	13.6
White, male												
2004.....	11.3	0.3	9.4	10.3	12.1	15.3	15.8	19.1	30.4	37.5
2003.....	11.7	0.3	9.4	10.9	12.6	15.6	16.5	20.2	30.9	39.9
2002.....	11.9	0.4	9.6	10.8	13.3	15.2	16.7	21.4	33.3	40.5
2001.....	11.8	0.4	9.7	12.0	12.6	14.9	16.1	20.9	32.9	42.1
2000.....	11.6	0.5	10.6	11.2	13.1	14.4	14.7	19.8	33.3	46.5
1999.....	11.7	0.4	10.8	11.8	12.6	13.9	15.2	21.0	33.3	44.9
White, female												
2004.....	1.7	*	1.1	1.7	2.6	3.2	2.5	1.8	1.6	*
2003.....	1.6	*	1.0	1.7	2.4	2.9	2.5	1.8	1.6	0.9
2002.....	1.6	*	1.0	1.9	2.5	2.6	2.5	1.7	1.5	0.9
2001.....	1.7	*	1.1	1.8	2.4	2.7	2.7	1.8	1.6	1.1
2000.....	1.7	*	1.2	1.8	2.7	2.6	2.4	1.9	1.5	1.2
1999.....	1.7	0.1	1.3	2.0	2.6	2.6	2.4	1.9	1.5	1.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Black, both sexes												
2004.....	2.6	*	4.0	4.8	2.7	2.4	3.0	3.5	4.8	*
2003.....	2.6	*	4.3	5.1	2.6	2.5	2.7	3.3	4.0	*
2002.....	2.8	*	4.1	5.0	3.5	2.9	2.8	3.3	4.4	*
2001.....	2.9	*	4.9	4.9	3.4	2.8	2.8	3.7	5.2	*
2000.....	3.0	*	5.2	5.0	3.4	3.3	2.9	3.7	3.5	*
1999.....	3.1	*	5.6	5.0	3.5	3.0	3.4	4.0	4.7	*
Black, male												
2004.....	4.9	*	7.1	9.0	5.2	4.4	5.8	8.3	12.8	*
2003.....	5.0	*	7.8	10.0	4.8	4.7	5.1	6.6	9.9	*
2002.....	5.4	*	7.5	10.0	6.3	5.5	5.5	7.1	11.5	*
2001.....	5.5	*	9.2	9.4	6.3	4.8	5.7	7.7	12.4	*
2000.....	5.5	*	9.6	9.3	6.2	5.8	5.7	8.8	8.2	*
1999.....	5.9	*	10.3	9.6	6.7	5.6	7.0	8.7	12.0	*
Black, female												
2004.....	0.5	*	0.9	0.9	*	*	*	*	*	*
2003.....	0.5	*	0.7	*	0.7	*	*	*	*	*
2002.....	0.5	*	0.7	*	0.9	*	*	*	*	*
2001.....	0.6	*	*	0.9	0.9	1.0	*	*	*	*
2000.....	0.7	*	0.9	1.1	0.8	1.1	*	*	*	*
1999.....	0.6	*	1.0	0.8	0.7	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

All races 3/, both sexes

2004.....	5.3	0.6	5.3	7.0	8.5	8.5	5.8	3.5	4.1	5.4
2003.....	5.0	0.4	4.6	6.7	8.3	7.8	5.5	3.5	4.0	5.1
2002.....	5.0	0.4	4.7	6.6	8.2	7.9	5.2	3.8	4.5	6.0
2001 4/.....	4.8	0.5	4.6	6.3	8.0	7.4	4.9	3.7	4.3	5.3
2000.....	4.5	0.5	4.4	5.9	7.5	6.8	4.6	3.2	4.6	6.5
1999.....	4.5	0.3	4.1	6.2	7.5	6.6	4.4	3.7	5.0	6.7

All races, male

2004.....	7.6	0.7	8.0	10.6	12.2	11.3	7.8	5.1	6.5	10.7
2003.....	7.3	0.6	7.1	10.2	12.1	10.6	7.3	5.0	6.4	11.1
2002.....	7.3	0.6	7.4	10.2	11.8	10.8	7.1	5.2	7.1	12.9
2001 4/.....	7.1	0.7	7.2	9.8	11.7	10.2	6.5	5.4	7.1	12.2
2000.....	6.6	0.7	6.9	9.0	10.9	9.4	6.0	4.3	7.8	14.6
1999.....	6.6	0.5	6.3	9.5	10.9	8.9	5.8	5.3	7.7	15.5

All races, female

2004.....	3.1	0.4	2.5	3.2	4.9	5.9	3.9	2.2	2.5	3.0
2003.....	2.8	0.2	2.1	3.2	4.5	5.2	3.7	2.2	2.5	2.4
2002.....	2.9	0.2	2.0	3.0	4.7	5.2	3.5	2.5	2.8	3.0
2001 4/.....	2.6	0.2	1.9	2.8	4.3	4.8	3.4	2.3	2.5	2.4
2000.....	2.5	0.2	1.8	2.7	4.0	4.4	3.3	2.3	2.6	3.2
1999.....	2.5	0.1	1.8	2.8	4.1	4.3	3.0	2.4	3.3	3.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
White, both sexes												
2004.....	5.8	0.5	5.6	7.5	9.5	9.4	6.2	3.6	4.2	5.5
2003.....	5.5	0.4	4.9	7.3	9.1	8.7	5.9	3.6	4.2	5.2
2002.....	5.5	0.4	5.2	7.1	9.1	8.8	5.6	3.9	4.7	6.2
2001 4/.....	5.3	0.5	5.0	6.8	8.8	8.2	5.2	3.8	4.5	5.5
2000.....	4.9	0.5	4.6	6.2	8.1	7.5	5.0	3.4	4.6	6.7
1999.....	4.9	0.4	4.3	6.5	8.2	7.2	4.6	3.8	5.2	7.1
White, male												
2004.....	8.3	0.6	8.4	11.4	13.5	12.3	8.2	5.2	6.6	10.9
2003.....	7.9	0.6	7.5	10.9	13.1	11.7	7.7	5.0	6.6	11.5
2002.....	8.0	0.6	8.1	10.7	12.9	11.8	7.6	5.4	7.4	13.4
2001 4/.....	7.7	0.7	7.9	10.4	12.8	11.0	6.8	5.4	7.3	12.9
2000.....	7.1	0.7	7.3	9.5	11.8	10.0	6.5	4.5	7.8	15.2
1999.....	7.0	0.6	6.6	9.9	11.8	9.6	6.1	5.4	7.9	16.6
White, female												
2004.....	3.4	0.4	2.6	3.2	5.5	6.6	4.3	2.4	2.5	3.0
2003.....	3.1	0.2	2.1	3.4	5.1	5.8	4.1	2.3	2.6	2.4
2002.....	3.2	0.3	2.1	3.3	5.3	5.8	3.7	2.6	2.9	3.1
2001 4/.....	2.9	0.2	2.0	3.0	4.9	5.3	3.7	2.4	2.7	2.3
2000.....	2.8	0.2	1.9	2.8	4.5	4.9	3.6	2.5	2.6	3.3
1999.....	2.8	0.1	1.9	2.9	4.6	4.8	3.3	2.5	3.4	3.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

Black, both sexes												
2004.....	2.6	0.6	3.2	4.6	3.9	3.5	2.7	*	*	*
2003.....	2.5	0.5	2.8	4.2	4.6	3.1	2.2	1.2	*	*
2002.....	2.3	0.4	2.4	4.2	4.2	3.0	2.0	1.5	*	*
2001 4/.....	2.3	0.4	2.3	4.2	3.9	3.4	2.4	1.7	*	*
2000.....	2.4	0.5	2.9	3.7	4.2	3.2	1.3	1.4	2.3	*
1999.....	2.3	0.4	2.5	4.2	4.1	2.7	1.5	1.9	*	*
Black, male												
2004.....	4.1	0.7	5.1	7.2	6.1	5.3	5.0	*	*	*
2003.....	3.8	0.9	4.3	6.5	7.3	4.4	3.6	*	*	*
2002.....	3.7	0.6	3.8	7.3	6.7	4.7	2.9	*	*	*
2001 4/.....	3.7	0.6	3.8	7.1	6.1	5.0	3.7	3.0	*	*
2000.....	3.9	0.9	4.6	6.3	6.9	5.4	1.8	*	*	*
1999.....	3.7	*	4.0	7.2	6.6	4.4	2.2	3.1	*	*
Black, female												
2004.....	1.3	*	1.3	2.3	2.0	1.9	*	*	*	*
2003.....	1.3	*	1.3	2.0	2.2	2.1	*	*	*	*
2002.....	1.1	*	1.0	1.5	1.9	1.6	1.4	*	*	*
2001 4/.....	1.1	*	0.7	1.5	2.0	2.0	*	*	*	*
2000.....	1.0	*	1.3	1.5	1.9	1.4	*	*	*	*
1999.....	1.0	*	0.9	1.6	1.9	1.2	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
All races 3/, both sexes												
2004.....	5.9	8.0	2.4	0.8	12.2	11.2	6.8	4.8	3.0	2.4	2.2	2.1
2003.....	6.1	8.5	2.4	0.8	13.0	11.3	7.0	4.9	2.8	2.4	2.5	2.2
2002.....	6.1	7.5	2.7	0.9	12.9	11.2	7.2	4.8	3.2	2.3	2.3	2.1
2001 4/.....	7.1	8.2	2.7	0.8	13.3	13.1	9.5	6.3	4.0	2.9	2.5	2.4
2000.....	6.0	9.2	2.3	0.9	12.6	10.4	7.1	4.7	3.0	2.4	2.4	2.4
1999.....	6.1	8.7	2.5	1.1	12.9	10.5	7.1	4.6	3.0	2.6	2.5	2.4
All races, male												
2004.....	9.4	8.0	2.5	1.0	20.4	18.5	10.1	7.3	4.6	3.6	2.6	2.7
2003.....	9.7	10.0	2.5	1.0	21.8	18.5	10.4	7.4	4.2	3.4	3.3	3.2
2002.....	9.6	7.9	2.9	0.9	21.5	18.2	10.7	7.1	4.8	3.3	3.1	3.0
2001 4/.....	11.1	9.5	3.0	0.9	22.2	20.8	13.9	9.5	5.9	4.0	3.1	3.2
2000.....	9.3	10.4	2.5	1.1	20.9	16.7	10.3	6.9	4.6	3.4	3.2	3.3
1999.....	9.3	9.6	2.5	1.2	21.1	16.5	10.3	6.8	4.7	3.7	3.4	3.4
All races, female												
2004.....	2.5	7.9	2.2	0.6	3.5	3.7	3.4	2.4	1.6	1.4	1.9	1.9
2003.....	2.6	6.9	2.3	0.5	3.7	3.9	3.6	2.5	1.5	1.5	1.9	1.8
2002.....	2.7	7.1	2.5	0.8	3.8	4.2	3.8	2.6	1.6	1.4	1.8	1.7
2001 4/.....	3.3	6.9	2.4	0.7	3.9	5.3	5.1	3.2	2.2	2.0	2.1	2.0
2000.....	2.8	7.9	2.1	0.7	3.9	4.1	4.0	2.5	1.6	1.6	2.0	2.0
1999.....	2.9	7.8	2.4	0.9	4.3	4.4	4.0	2.4	1.5	1.7	1.9	2.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

White, both sexes

2004.....	3.6	6.5	1.6	0.6	6.5	6.1	4.5	3.4	2.3	2.0	1.9	1.9
2003.....	3.7	6.9	1.7	0.5	6.7	6.1	4.6	3.5	2.2	1.9	2.1	2.0
2002.....	3.7	5.4	2.0	0.6	6.7	5.9	4.8	3.3	2.6	1.9	1.9	1.8
2001 4/.....	4.9	6.2	1.9	0.7	7.2	8.3	7.3	5.0	3.4	2.5	2.1	2.2
2000.....	3.6	6.6	1.7	0.7	6.4	5.7	4.7	3.3	2.4	1.9	2.0	2.2
1999.....	3.8	6.7	1.7	0.8	6.8	6.0	4.8	3.3	2.4	2.2	2.0	2.1

White, male

2004.....	5.3	6.9	1.8	0.7	10.2	9.3	6.3	4.9	3.4	2.8	2.1	2.3
2003.....	5.4	8.1	1.5	0.6	10.6	9.2	6.3	5.0	3.1	2.7	2.8	3.0
2002.....	5.4	6.2	2.1	0.6	10.6	8.9	6.8	4.6	3.7	2.6	2.4	2.7
2001 4/.....	7.2	7.3	2.2	0.7	11.2	12.3	10.7	7.4	4.9	3.2	2.6	2.8
2000.....	5.2	8.2	1.9	0.9	9.9	8.4	6.5	4.6	3.5	2.4	2.5	2.8
1999.....	5.5	7.9	1.8	0.9	10.3	8.8	6.7	4.8	3.6	2.9	2.7	2.7

White, female

2004.....	1.9	6.2	1.4	0.5	2.5	2.7	2.6	1.9	1.4	1.4	1.7	1.8
2003.....	2.0	5.7	1.8	0.3	2.5	2.8	2.8	2.1	1.4	1.3	1.7	1.6
2002.....	2.0	4.6	1.8	0.6	2.5	2.8	2.8	2.1	1.5	1.3	1.6	1.5
2001 4/.....	2.6	5.1	1.6	0.6	3.0	4.1	3.8	2.7	2.0	1.8	1.8	2.0
2000.....	2.1	5.0	1.5	0.6	2.7	2.9	2.8	2.1	1.4	1.5	1.8	1.9
1999.....	2.1	5.4	1.7	0.7	3.0	3.1	2.9	1.8	1.3	1.6	1.6	1.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

Black, both sexes

2004.....	21.1	15.6	6.3	1.9	43.5	44.2	22.6	15.3	8.8	6.0	6.3	*
2003.....	22.0	16.1	6.1	2.2	47.6	45.2	23.4	15.4	8.1	6.2	6.6	*
2002.....	22.0	17.3	6.3	2.0	46.8	45.3	23.7	15.5	8.3	5.7	7.4	*
2001 4/.....	22.1	18.9	6.8	1.4	47.3	43.7	25.0	16.1	8.9	7.1	6.7	*
2000.....	21.5	22.8	5.8	1.8	47.8	40.8	23.9	14.8	8.3	7.6	7.0	*
1999.....	21.1	19.5	6.1	2.2	47.7	39.3	23.2	14.0	8.2	6.3	7.4	*

Black, male

2004.....	37.1	14.2	6.3	2.4	77.6	81.6	37.9	26.0	15.1	11.5	10.3	*
2003.....	38.9	17.8	7.4	2.9	84.6	82.5	40.2	26.3	14.8	11.1	11.6	*
2002.....	38.4	16.3	7.1	2.5	83.1	82.2	38.8	27.1	15.4	10.6	13.8	*
2001 4/.....	38.3	21.0	8.0	2.0	85.7	78.8	38.5	26.9	15.0	12.6	10.0	*
2000.....	37.2	23.3	5.9	2.1	85.3	73.9	38.5	26.0	14.7	13.9	11.8	*
1999.....	36.1	19.5	5.9	2.7	84.7	69.2	37.2	23.4	15.1	11.8	12.0	*

Black, female

2004.....	6.4	17.0	6.3	1.5	8.8	9.8	9.1	6.0	3.7	2.1	4.1	*
2003.....	6.6	14.2	4.8	1.5	10.1	11.1	8.5	6.0	2.7	2.7	3.8	*
2002.....	7.0	18.5	5.5	1.5	10.3	11.9	10.4	5.6	2.5	2.3	3.7	*
2001 4/.....	7.4	16.7	5.6	0.8	8.9	12.1	13.0	6.8	3.9	3.3	4.8	*
2000.....	7.2	22.2	5.7	1.6	10.7	10.9	11.0	5.3	3.3	3.1	4.3	*
1999.....	7.6	19.5	6.3	1.7	11.2	12.4	10.8	5.9	2.6	2.5	4.8	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
All races 3/, both sexes												
2004.....	4.0	*	0.2	0.5	9.9	8.8	4.3	2.6	1.6	1.0	0.8	0.6
2003.....	4.1	*	0.3	0.5	10.7	8.9	4.4	2.7	1.4	0.9	0.7	0.5
2002.....	4.1	*	0.3	0.5	10.6	8.7	4.5	2.6	1.5	0.9	0.9	*
2001.....	4.0	*	0.4	0.4	10.5	8.4	4.4	2.4	1.4	1.0	0.8	*
2000.....	3.8	*	0.2	0.5	10.1	7.7	4.2	2.5	1.5	1.0	0.7	0.7
1999.....	3.9	*	0.3	0.5	10.4	7.6	4.1	2.4	1.6	1.0	0.8	0.6
All races, male												
2004.....	6.9	*	*	0.6	17.3	15.3	7.1	4.0	2.5	1.6	0.8	*
2003.....	7.1	*	0.2	0.6	18.8	15.5	6.9	4.3	2.2	1.4	0.7	*
2002.....	7.0	*	0.4	0.6	18.5	14.9	7.3	3.9	2.4	1.2	1.3	*
2001.....	6.8	*	0.3	0.6	18.5	14.3	6.8	3.8	2.2	1.5	1.1	*
2000.....	6.5	*	*	0.7	17.7	13.2	6.6	3.9	2.3	1.3	0.8	*
1999.....	6.5	*	0.3	0.7	18.0	12.9	6.3	3.8	2.6	1.5	1.0	*
All races, female												
2004.....	1.1	*	*	0.3	2.0	2.0	1.5	1.1	0.8	0.5	0.8	*
2003.....	1.2	*	0.3	0.3	2.2	2.1	1.8	1.2	0.7	0.5	0.7	*
2002.....	1.3	*	*	0.4	2.4	2.3	1.9	1.3	0.8	0.6	0.7	*
2001.....	1.3	*	0.4	0.3	2.1	2.3	2.0	1.0	0.8	0.7	0.6	*
2000.....	1.3	*	*	0.2	2.2	2.1	1.8	1.2	0.8	0.8	0.6	*
1999.....	1.3	*	0.3	0.3	2.5	2.2	2.0	1.2	0.6	0.7	0.6	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
White, both sexes												
2004.....	2.1	*	*	0.3	4.7	4.3	2.7	1.8	1.3	0.9	0.7	0.5
2003.....	2.2	*	0.2	0.3	5.0	4.2	2.7	1.9	1.1	0.7	0.6	0.5
2002.....	2.2	*	0.3	0.3	5.0	4.1	2.8	1.8	1.2	0.7	0.8	*
2001.....	2.2	*	0.3	0.4	5.0	4.2	2.9	1.7	1.2	0.9	0.7	*
2000.....	2.1	*	*	0.4	4.6	3.7	2.7	1.8	1.3	0.8	0.6	0.6
1999.....	2.2	*	0.2	0.4	4.9	3.8	2.7	1.7	1.2	0.9	0.7	0.6
White, male												
2004.....	3.5	*	*	0.4	7.8	7.0	4.1	2.6	1.8	1.2	0.7	*
2003.....	3.5	*	*	0.4	8.5	6.9	3.9	2.8	1.6	1.1	0.5	*
2002.....	3.5	*	0.3	0.4	8.4	6.6	4.2	2.5	1.8	0.9	1.0	*
2001.....	3.5	*	*	0.5	8.3	6.7	4.3	2.6	1.6	1.1	0.8	*
2000.....	3.2	*	*	0.5	7.7	5.9	3.9	2.6	1.8	1.0	0.6	*
1999.....	3.4	*	*	0.5	8.0	6.1	4.0	2.6	1.9	1.2	0.7	*
White, female												
2004.....	0.9	*	*	0.2	1.3	1.3	1.2	0.9	0.7	0.6	0.7	*
2003.....	0.9	*	*	0.1	1.3	1.4	1.5	1.0	0.7	0.4	0.6	*
2002.....	1.0	*	*	0.3	1.4	1.5	1.4	1.1	0.7	0.6	0.6	*
2001.....	1.0	*	*	0.3	1.5	1.6	1.5	0.9	0.8	0.7	0.6	*
2000.....	1.0	*	*	0.2	1.4	1.5	1.4	1.0	0.8	0.7	0.6	*
1999.....	0.9	*	*	0.3	1.5	1.5	1.5	0.9	0.6	0.7	0.6	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Black, both sexes												
2004.....	16.1	*	0.8	1.2	38.8	37.7	15.6	8.4	4.8	2.4	*	*
2003.....	16.8	*	*	1.4	41.9	38.8	15.9	8.7	3.9	2.5	*	*
2002.....	16.7	*	*	1.2	41.8	38.2	16.3	8.5	4.1	2.4	2.2	*
2001.....	15.8	*	*	0.8	41.8	35.1	15.0	7.5	3.7	2.7	*	*
2000.....	15.6	*	*	1.0	41.2	33.2	15.0	8.4	3.5	2.7	*	*
1999.....	15.4	*	*	1.2	42.0	31.4	14.1	8.0	4.4	2.4	2.3	*
Black, male												
2004.....	30.3	*	*	1.8	71.3	72.2	28.9	15.2	8.6	5.4	*	*
2003.....	31.6	*	*	1.9	76.8	74.1	29.1	15.9	7.5	4.6	*	*
2002.....	31.0	*	*	1.6	76.1	72.8	29.2	15.1	7.6	4.7	*	*
2001.....	29.8	*	*	1.3	78.2	66.9	26.0	13.7	7.1	5.6	*	*
2000.....	29.2	*	*	1.5	76.5	63.3	26.7	15.1	6.3	5.4	*	*
1999.....	28.4	*	*	1.8	76.9	59.1	24.0	13.9	8.9	4.6	*	*
Black, female												
2004.....	3.0	*	*	*	5.8	6.1	3.8	2.5	1.7	*	*	*
2003.....	3.3	*	*	0.8	6.5	6.6	4.2	2.5	*	*	*	*
2002.....	3.6	*	*	0.8	7.3	6.9	4.8	2.8	1.4	*	*	*
2001.....	3.1	*	*	*	5.4	6.3	5.4	2.2	*	*	*	*
2000.....	3.2	*	*	*	6.4	6.1	4.6	2.6	*	*	*	*
1999.....	3.7	*	*	0.6	7.6	6.5	5.3	3.0	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

All races 3/, both sexes

2004.....	2.0	7.8	2.1	0.4	2.3	2.5	2.5	2.3	1.4	1.4	1.5	1.5
2003.....	2.0	8.3	2.1	0.3	2.3	2.4	2.6	2.2	1.4	1.5	1.8	1.7
2002.....	2.0	7.3	2.4	0.4	2.2	2.6	2.7	2.2	1.6	1.4	1.4	1.7
2001 4/.....	3.1	8.0	2.3	0.4	2.7	4.8	5.1	3.9	2.6	1.9	1.7	2.0
2000.....	2.1	8.9	2.1	0.4	2.5	2.7	2.9	2.1	1.5	1.4	1.8	1.7
1999.....	2.2	8.5	2.1	0.5	2.5	2.9	3.0	2.1	1.5	1.5	1.7	1.8

All races, male

2004.....	2.5	7.9	2.3	0.3	3.1	3.2	3.0	3.3	2.1	1.9	1.9	1.7
2003.....	2.6	9.7	2.3	0.4	3.1	3.0	3.5	3.1	2.1	2.0	2.6	2.5
2002.....	2.6	7.8	2.6	0.3	3.0	3.2	3.4	3.2	2.4	2.1	1.9	2.3
2001 4/.....	4.3	9.2	2.7	0.3	3.6	6.5	7.1	5.7	3.8	2.5	2.0	2.7
2000.....	2.8	10.0	2.3	0.4	3.3	3.5	3.7	3.0	2.3	2.1	2.4	2.5
1999.....	2.8	9.4	2.1	0.5	3.1	3.6	4.0	3.1	2.2	2.2	2.4	2.6

All races, female

2004.....	1.4	7.7	2.0	0.4	1.5	1.7	1.9	1.3	0.8	0.9	1.2	1.5
2003.....	1.4	6.9	2.0	0.3	1.6	1.8	1.8	1.3	0.8	1.0	1.2	1.4
2002.....	1.4	6.8	2.2	0.4	1.4	1.9	1.9	1.3	0.9	0.8	1.1	1.4
2001 4/.....	2.0	6.7	2.0	0.4	1.8	3.1	3.1	2.1	1.5	1.3	1.5	1.7
2000.....	1.5	7.7	2.0	0.5	1.7	1.9	2.1	1.3	0.8	0.9	1.4	1.4
1999.....	1.6	7.6	2.1	0.6	1.8	2.2	2.0	1.2	0.8	1.0	1.3	1.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

White, both sexes

2004.....	1.5	6.5	1.5	0.3	1.8	1.8	1.8	1.6	1.1	1.1	1.2	1.4
2003.....	1.5	6.9	1.5	0.2	1.7	1.8	1.9	1.6	1.1	1.2	1.5	1.6
2002.....	1.5	5.3	1.7	0.3	1.7	1.8	2.0	1.6	1.3	1.2	1.1	1.4
2001 4/.....	2.6	6.0	1.6	0.3	2.2	4.1	4.4	3.3	2.3	1.6	1.4	1.8
2000.....	1.5	6.5	1.5	0.4	1.8	2.0	2.0	1.5	1.1	1.0	1.5	1.5
1999.....	1.6	6.6	1.5	0.4	1.9	2.1	2.0	1.6	1.2	1.3	1.4	1.5

White, male

2004.....	1.9	6.8	1.7	0.3	2.4	2.2	2.2	2.3	1.6	1.5	1.4	1.5
2003.....	1.9	8.0	1.4	0.3	2.1	2.3	2.4	2.2	1.5	1.5	2.2	2.3
2002.....	1.9	6.2	1.8	0.2	2.2	2.3	2.5	2.1	1.9	1.8	1.3	2.0
2001 4/.....	3.7	7.0	1.9	0.3	2.8	5.6	6.4	4.8	3.3	2.1	1.8	2.3
2000.....	2.0	8.1	1.7	0.4	2.3	2.5	2.6	2.0	1.7	1.4	1.9	2.1
1999.....	2.1	7.7	1.5	0.4	2.3	2.7	2.7	2.3	1.7	1.7	2.0	1.9

White, female

2004.....	1.1	6.1	1.3	0.3	1.2	1.4	1.4	1.0	0.6	0.8	1.0	1.3
2003.....	1.1	5.6	1.6	0.2	1.2	1.4	1.4	1.0	0.7	0.9	1.1	1.2
2002.....	1.1	4.5	1.6	0.4	1.1	1.3	1.4	1.0	0.8	0.7	1.0	1.2
2001 4/.....	1.6	4.9	1.4	0.4	1.5	2.5	2.3	1.8	1.3	1.1	1.2	1.6
2000.....	1.1	4.8	1.3	0.4	1.3	1.4	1.4	1.1	0.6	0.7	1.2	1.3
1999.....	1.2	5.4	1.5	0.5	1.4	1.6	1.4	0.9	0.7	0.9	1.0	1.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Black, both sexes

2004.....	5.0	15.1	5.4	0.7	4.7	6.4	7.0	6.9	4.0	3.6	4.9	*
2003.....	5.2	15.2	5.3	0.8	5.7	6.4	7.5	6.8	4.1	3.7	4.9	*
2002.....	5.3	16.5	5.8	0.8	5.0	7.1	7.4	7.1	4.1	3.4	5.2	*
2001 4/.....	6.3	18.6	6.1	0.5	5.5	8.7	9.9	8.5	5.2	4.4	4.8	*
2000.....	5.9	21.6	5.5	0.8	6.5	7.6	8.9	6.5	4.9	4.8	5.1	*
1999.....	5.7	19.2	5.4	1.0	5.7	7.9	9.1	6.0	3.8	4.0	5.1	*

Black, male

2004.....	6.8	13.9	5.5	0.6	6.4	9.5	9.0	10.8	6.5	6.1	8.4	*
2003.....	7.4	16.1	6.5	0.9	7.7	8.4	11.1	10.5	7.3	6.5	8.7	*
2002.....	7.3	15.7	6.4	0.9	7.0	9.4	9.6	12.0	7.8	5.9	9.2	*
2001 4/.....	8.5	20.7	7.3	0.7	7.5	11.8	12.6	13.2	8.0	7.0	*	*
2000.....	8.0	21.1	5.5	*	8.8	10.6	11.7	10.8	8.3	8.5	8.2	*
1999.....	7.7	19.2	5.4	0.9	7.8	10.1	13.2	9.6	6.2	7.2	7.1	*

Black, female

2004.....	3.4	16.4	5.4	0.9	3.0	3.6	5.3	3.5	2.0	*	*	*
2003.....	3.3	14.2	4.1	0.7	3.6	4.6	4.2	3.6	1.6	*	*	*
2002.....	3.4	17.3	5.1	0.7	3.0	5.0	5.6	2.8	*	*	*	*
2001 4/.....	4.3	16.4	4.8	*	3.5	5.8	7.6	4.6	2.9	2.7	4.3	*
2000.....	4.0	22.2	5.5	1.1	4.3	4.8	6.4	2.7	2.1	2.3	3.4	*
1999.....	3.9	19.1	5.4	1.1	3.6	5.9	5.5	2.9	1.9	*	4.0	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All races 3/, both sexes												
2004.....	1.7	2.6	0.3	0.2	1.2	2.0	3.1	3.1	1.4	0.8	0.9	1.6
2003.....	1.7	1.9	0.2	0.1	1.4	2.1	3.6	3.1	1.3	0.7	0.7	1.1
2002.....	1.7	2.5	0.4	0.2	1.2	2.1	3.4	2.8	1.2	0.8	0.7	1.4
2001.....	1.5	2.1	0.3	0.1	1.0	1.8	3.1	2.5	0.9	0.5	0.9	1.0
2000.....	1.4	1.7	0.3	0.1	0.9	1.7	2.8	2.2	0.9	0.6	0.6	1.3
1999.....	1.4	1.4	0.3	0.1	1.1	1.8	2.9	2.2	0.9	0.6	0.8	1.4
All races, male												
2004.....	2.2	3.1	0.3	0.2	1.8	2.7	4.0	4.0	1.7	1.1	1.0	1.9
2003.....	2.3	2.4	*	0.2	1.9	2.9	4.6	4.0	1.7	0.8	1.0	1.5
2002.....	2.2	2.5	0.4	0.2	1.7	2.9	4.4	3.5	1.6	1.0	0.9	2.2
2001.....	2.0	2.4	*	0.2	1.5	2.6	4.2	3.4	1.2	0.8	1.1	1.5
2000.....	1.9	1.4	0.3	0.2	1.4	2.4	3.9	3.1	1.3	0.8	1.0	1.7
1999.....	2.0	1.7	0.3	0.2	1.7	2.6	4.0	3.0	1.3	0.9	1.1	2.1
All races, female												
2004.....	1.2	2.0	0.3	0.1	0.6	1.3	2.3	2.3	1.1	0.6	0.9	1.6
2003.....	1.2	1.3	*	*	0.8	1.3	2.6	2.2	1.0	0.7	0.6	0.9
2002.....	1.2	2.5	0.4	0.1	0.7	1.2	2.4	2.2	0.8	0.6	0.6	1.1
2001.....	0.9	1.7	0.3	*	0.5	1.1	2.1	1.6	0.6	0.3	0.7	0.8
2000.....	0.8	2.0	*	*	0.4	1.0	1.8	1.4	0.6	0.3	0.4	1.2
1999.....	0.9	1.2	0.3	*	0.5	1.1	1.8	1.4	0.6	0.4	0.6	1.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

White, both sexes												
2004.....	1.7	2.1	0.2	0.1	1.3	2.1	3.3	3.2	1.4	0.8	0.9	1.6
2003.....	1.8	1.7	0.2	0.1	1.5	2.3	3.7	3.1	1.3	0.6	0.7	1.2
2002.....	1.7	1.8	0.4	0.2	1.3	2.1	3.4	2.8	1.2	0.7	0.7	1.4
2001.....	1.5	1.4	0.3	0.1	1.1	1.9	3.1	2.5	0.9	0.5	0.9	1.0
2000.....	1.3	1.4	0.2	0.1	1.0	1.7	2.8	2.1	0.9	0.5	0.6	1.3
1999.....	1.4	1.2	0.2	0.1	1.1	1.9	2.8	2.0	0.9	0.6	0.8	1.4
White, male												
2004.....	2.2	2.7	*	0.1	1.9	2.9	4.0	3.9	1.6	1.0	1.0	1.9
2003.....	2.3	1.9	*	0.2	2.1	3.0	4.7	3.9	1.5	0.7	1.0	1.5
2002.....	2.2	1.8	0.4	0.2	1.9	2.9	4.4	3.3	1.5	0.9	0.9	2.3
2001.....	2.0	1.7	*	0.2	1.6	2.7	4.0	3.2	1.1	0.7	1.2	*
2000.....	1.8	*	*	0.2	1.4	2.4	3.8	2.8	1.1	0.7	1.0	*
1999.....	1.9	1.4	*	0.2	1.8	2.5	3.8	2.6	1.1	0.8	1.1	2.1
White, female												
2004.....	1.3	1.5	*	*	0.7	1.3	2.5	2.5	1.2	0.6	0.9	1.5
2003.....	1.3	1.4	*	*	0.9	1.5	2.7	2.4	1.0	0.6	0.6	1.0
2002.....	1.2	1.8	0.4	0.1	0.7	1.3	2.5	2.2	0.9	0.6	0.6	1.1
2001.....	1.0	*	*	*	0.6	1.2	2.2	1.7	0.6	0.4	0.6	0.8
2000.....	0.8	1.6	*	*	0.5	1.0	1.7	1.4	0.6	0.4	0.4	1.1
1999.....	0.9	*	*	*	0.4	1.2	1.8	1.4	0.7	0.4	0.6	1.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Black, both sexes

2004.....	1.8	4.8	*	0.4	0.9	1.8	3.3	3.9	1.7	1.3	*	*
2003.....	1.8	3.2	*	*	0.8	1.8	3.8	4.0	1.9	1.2	*	*
2002.....	2.0	5.5	*	*	0.9	2.3	4.2	4.2	1.5	*	*	*
2001.....	1.7	5.5	*	*	0.7	1.7	3.9	3.8	1.3	*	*	*
2000.....	1.8	*	*	0.3	0.9	1.9	4.1	3.7	1.8	*	*	*
1999.....	1.9	*	*	*	1.2	2.2	4.2	4.1	1.2	*	*	*

Black, male

2004.....	2.6	*	*	*	1.3	2.8	4.6	5.8	2.9	*	*	*
2003.....	2.7	*	*	*	1.4	2.9	5.3	6.4	3.2	*	*	*
2002.....	2.9	*	*	*	1.4	3.5	6.0	6.4	2.5	*	*	*
2001.....	2.7	6.0	*	*	1.3	2.6	6.2	6.3	2.4	*	*	*
2000.....	2.8	*	*	*	1.6	3.1	6.0	5.9	3.0	*	*	*
1999.....	3.0	*	*	*	1.9	3.6	6.4	7.2	2.5	*	*	*

Black, female

2004.....	1.1	*	*	*	*	0.9	2.2	2.2	*	*	*	*
2003.....	0.9	*	*	*	*	*	2.4	1.9	*	*	*	*
2002.....	1.2	*	*	*	*	1.2	2.6	2.4	*	*	*	*
2001.....	0.9	*	*	*	*	0.9	1.9	1.6	*	*	*	*
2000.....	0.9	*	*	*	*	0.9	2.4	1.8	*	*	*	*
1999.....	0.9	*	*	*	*	1.0	2.3	1.6	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All races 3/, both sexes

2004.....	1.6	2.6	0.3	0.1	1.1	1.9	3.0	3.1	1.3	0.8	0.9	1.6
2003.....	1.7	1.9	0.2	0.1	1.2	2.0	3.5	3.0	1.2	0.7	0.7	1.1
2002.....	1.6	2.5	0.4	0.2	1.0	2.0	3.3	2.8	1.2	0.7	0.7	1.4
2001.....	1.4	2.1	0.3	0.1	0.9	1.7	3.0	2.5	0.8	0.5	0.8	1.0
2000.....	1.3	1.7	0.3	0.1	0.8	1.6	2.8	2.2	0.9	0.5	0.6	1.3
1999.....	1.3	1.4	0.3	0.1	0.8	1.7	2.7	2.1	0.9	0.5	0.7	1.3

All races, male

2004.....	2.1	3.1	0.3	0.2	1.5	2.6	3.8	3.8	1.6	0.9	0.9	1.8
2003.....	2.2	2.4	*	0.1	1.7	2.7	4.5	3.9	1.5	0.7	0.8	1.4
2002.....	2.1	2.5	0.3	0.2	1.4	2.7	4.2	3.4	1.5	0.9	0.8	2.2
2001.....	1.9	2.4	*	0.2	1.3	2.4	4.0	3.3	1.1	0.7	1.0	*
2000.....	1.8	1.4	0.3	0.2	1.1	2.2	3.8	3.0	1.2	0.7	0.9	*
1999.....	1.8	1.7	*	0.1	1.2	2.4	3.8	2.8	1.2	0.7	0.9	1.8

All races, female

2004.....	1.2	2.0	0.3	*	0.6	1.2	2.3	2.3	1.1	0.6	0.8	1.5
2003.....	1.2	1.3	*	*	0.7	1.3	2.5	2.2	1.0	0.7	0.6	0.9
2002.....	1.1	2.5	0.4	0.1	0.6	1.2	2.3	2.1	0.8	0.5	0.6	1.1
2001.....	0.9	1.7	0.3	*	0.5	1.1	2.0	1.6	0.6	0.3	0.6	0.8
2000.....	0.8	2.0	*	*	0.4	0.9	1.7	1.4	0.6	0.3	0.4	1.2
1999.....	0.8	1.2	0.3	*	0.4	1.1	1.7	1.4	0.6	0.4	0.5	1.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

White, both sexes

2004.....	1.7	2.1	0.2	0.1	1.2	2.1	3.2	3.1	1.3	0.7	0.8	1.6
2003.....	1.7	1.7	0.2	0.1	1.4	2.2	3.6	3.0	1.2	0.6	0.7	1.2
2002.....	1.6	1.8	0.4	0.1	1.1	2.0	3.3	2.7	1.2	0.6	0.7	1.4
2001.....	1.4	1.4	0.3	0.1	1.0	1.8	3.0	2.4	0.8	0.5	0.8	1.0
2000.....	1.3	1.4	0.2	0.1	0.8	1.6	2.7	2.1	0.8	0.5	0.6	1.2
1999.....	1.3	1.2	0.2	0.1	0.8	1.8	2.6	1.9	0.9	0.5	0.7	1.3

White, male

2004.....	2.1	2.7	*	0.1	1.7	2.8	3.9	3.7	1.5	0.8	0.9	1.8
2003.....	2.2	1.9	*	0.1	1.9	2.9	4.6	3.8	1.3	0.6	0.8	*
2002.....	2.0	1.8	0.4	0.1	1.5	2.8	4.2	3.2	1.5	0.7	0.8	2.3
2001.....	1.9	1.7	*	0.2	1.4	2.5	3.9	3.1	1.0	0.6	1.0	*
2000.....	1.7	*	*	0.2	1.2	2.2	3.7	2.8	1.0	0.6	0.9	*
1999.....	1.7	1.4	*	0.1	1.3	2.3	3.5	2.5	1.1	0.7	0.9	*

White, female

2004.....	1.2	1.5	*	*	0.6	1.3	2.4	2.4	1.1	0.6	0.8	1.5
2003.....	1.3	1.4	*	*	0.8	1.5	2.6	2.3	1.0	0.6	0.6	1.0
2002.....	1.2	1.8	0.4	0.1	0.7	1.3	2.4	2.2	0.9	0.6	0.6	1.1
2001.....	1.0	*	*	*	0.6	1.1	2.1	1.7	0.6	0.3	0.6	0.8
2000.....	0.8	1.6	*	*	0.5	1.0	1.7	1.4	0.6	0.3	0.4	1.1
1999.....	0.8	*	*	*	0.4	1.2	1.7	1.4	0.7	0.4	0.6	1.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Black, both sexes

2004.....	1.7	4.8	*	0.3	0.6	1.6	3.3	3.8	1.7	1.2	*	*
2003.....	1.7	3.2	*	*	0.5	1.6	3.7	3.9	1.9	1.2	*	*
2002.....	1.9	5.5	*	*	0.6	2.1	4.1	4.2	1.4	*	*	*
2001.....	1.7	5.5	*	*	0.5	1.6	3.9	3.7	1.1	*	*	*
2000.....	1.7	*	*	*	0.5	1.7	3.9	3.6	1.6	*	*	*
1999.....	1.7	*	*	*	0.6	2.0	4.1	4.0	1.1	*	*	*

Black, male

2004.....	2.4	*	*	*	1.0	2.5	4.5	5.7	2.9	*	*	*
2003.....	2.6	*	*	*	0.8	2.7	5.2	6.3	3.2	*	*	*
2002.....	2.7	*	*	*	0.8	3.1	5.8	6.3	2.5	*	*	*
2001.....	2.5	6.0	*	*	0.8	2.4	6.2	6.1	2.1	*	*	*
2000.....	2.5	*	*	*	1.0	2.6	5.8	5.8	2.8	*	*	*
1999.....	2.7	*	*	*	0.9	3.1	6.2	6.9	2.5	*	*	*

Black, female

2004.....	1.1	*	*	*	*	0.9	2.2	2.2	*	*	*	*
2003.....	0.9	*	*	*	*	*	2.3	1.9	*	*	*	*
2002.....	1.2	*	*	*	*	1.2	2.6	2.4	*	*	*	*
2001.....	0.8	*	*	*	*	0.9	1.8	1.6	*	*	*	*
2000.....	0.9	*	*	*	*	0.8	2.3	1.8	*	*	*	*
1999.....	0.9	*	*	*	*	0.9	2.2	1.6	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Complications of medical and surgical care (Y40-Y84,Y88)

All races 3/, both sexes

2004.....	1.0	*	*	*	0.1	0.2	0.4	0.7	1.4	3.0	6.1	10.6
2003.....	1.0	*	0.2	*	0.1	0.2	0.4	0.7	1.4	2.9	6.2	11.2
2002.....	1.0	*	*	0.1	0.1	0.2	0.4	0.7	1.3	3.1	6.2	11.3
2001.....	1.1	0.6	0.2	*	0.1	0.2	0.4	0.7	1.5	3.1	6.9	12.8
2000.....	1.1	*	0.1	0.1	0.1	0.2	0.3	0.8	1.5	3.5	6.7	13.6
1999.....	1.0	1.4	0.2	*	0.1	0.2	0.3	0.6	1.3	3.2	6.6	12.7

All races, male

2004.....	0.9	*	*	*	0.1	0.2	0.3	0.7	1.6	3.3	7.4	11.9
2003.....	0.9	*	*	*	0.1	0.2	0.4	0.7	1.5	3.2	6.9	12.9
2002.....	0.9	*	*	0.1	*	0.2	0.4	0.6	1.3	3.1	6.3	12.3
2001.....	1.0	*	*	*	*	0.2	0.4	0.7	1.6	3.2	7.7	14.5
2000.....	1.0	*	*	*	*	0.2	0.3	0.8	1.6	4.0	8.2	16.0
1999.....	1.0	1.6	*	*	0.1	0.2	0.3	0.5	1.4	3.4	7.8	16.5

All races, female

2004.....	1.0	*	*	*	0.1	0.2	0.4	0.7	1.2	2.8	5.2	9.9
2003.....	1.0	*	*	*	*	0.2	0.4	0.6	1.3	2.7	5.7	10.5
2002.....	1.1	*	*	*	*	0.2	0.4	0.7	1.3	3.0	6.2	10.9
2001.....	1.1	*	*	*	0.1	0.2	0.4	0.7	1.4	3.0	6.3	12.1
2000.....	1.1	*	*	*	*	0.2	0.3	0.8	1.5	3.1	5.7	12.7
1999.....	1.1	1.2	*	*	*	0.2	0.3	0.6	1.2	3.0	5.9	11.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)												
White, both sexes												
2004.....	1.0	*	*	*	0.1	0.2	0.3	0.7	1.3	2.9	5.9	10.4
2003.....	1.0	*	*	*	0.1	0.2	0.4	0.6	1.2	2.8	6.1	11.0
2002.....	1.0	*	*	0.1	0.1	0.2	0.3	0.6	1.2	3.0	6.1	11.2
2001.....	1.1	*	*	*	0.1	0.2	0.3	0.6	1.4	2.8	6.8	12.1
2000.....	1.1	*	*	0.1	0.1	0.2	0.3	0.7	1.4	3.3	6.4	13.1
1999.....	1.0	1.2	0.2	*	0.1	0.2	0.3	0.5	1.2	3.0	6.3	12.1
White, male												
2004.....	1.0	*	*	*	*	0.2	0.3	0.7	1.5	3.3	7.3	11.8
2003.....	1.0	*	*	*	0.1	0.2	0.4	0.6	1.4	3.1	7.0	12.8
2002.....	0.9	*	*	*	*	0.2	0.4	0.6	1.2	2.9	6.3	11.8
2001.....	0.9	*	*	*	*	0.2	0.3	0.6	1.4	2.8	7.7	13.6
2000.....	1.1	*	*	*	*	0.2	0.3	0.7	1.4	3.7	8.2	15.8
1999.....	1.0	1.4	*	*	*	0.2	0.3	0.5	1.3	3.2	7.5	15.6
White, female												
2004.....	1.0	*	*	*	*	0.2	0.3	0.6	1.1	2.5	4.9	9.8
2003.....	1.0	*	*	*	*	0.2	0.3	0.5	1.1	2.5	5.4	10.2
2002.....	1.1	*	*	*	*	0.1	0.3	0.5	1.2	3.0	6.0	10.9
2001.....	1.2	*	*	*	*	0.2	0.3	0.6	1.3	2.7	6.2	11.5
2000.....	1.1	*	*	*	*	0.2	0.3	0.7	1.3	3.0	5.3	12.0
1999.....	1.1	*	*	*	*	0.1	0.3	0.6	1.1	2.9	5.6	10.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)												
Black, both sexes												
2004.....	1.1	*	*	*	*	*	0.8	1.2	2.6	5.0	9.3	13.8
2003.....	1.1	*	*	*	*	*	0.6	1.4	2.7	4.6	8.8	15.2
2002.....	1.2	*	*	*	*	*	0.8	1.6	2.7	4.6	8.3	14.0
2001.....	1.4	*	*	*	*	0.4	0.9	1.6	3.4	6.3	9.2	23.3
2000.....	1.3	*	*	*	*	0.4	0.5	1.4	3.5	6.1	10.5	21.0
1999.....	1.2	*	*	*	*	*	0.6	1.3	2.6	5.2	10.6	20.1
Black, male												
2004.....	0.9	*	*	*	*	*	*	1.0	2.9	4.6	10.0	*
2003.....	1.0	*	*	*	*	*	*	1.8	2.8	3.9	7.6	*
2002.....	1.0	*	*	*	*	*	*	1.3	2.5	5.3	7.2	*
2001.....	1.4	*	*	*	*	*	1.0	1.8	4.2	7.8	9.1	29.5
2000.....	1.1	*	*	*	*	*	*	1.3	3.1	7.6	10.3	*
1999.....	1.1	*	*	*	*	*	*	1.4	2.8	5.9	12.3	23.4
Black, female												
2004.....	1.3	*	*	*	*	*	1.0	1.4	2.3	5.2	9.0	13.4
2003.....	1.3	*	*	*	*	*	0.7	1.1	2.5	5.0	9.4	15.4
2002.....	1.3	*	*	*	*	*	1.1	1.8	2.7	4.1	8.9	11.5
2001.....	1.4	*	*	*	*	*	0.7	1.3	2.7	5.3	9.3	21.0
2000.....	1.5	*	*	*	*	*	*	1.5	3.8	5.0	10.6	21.9
1999.....	1.3	*	*	*	*	*	*	1.2	2.4	4.8	9.6	18.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 5/

All races 3/, both sexes

2004.....	7.2	*	*	*	0.3	1.5	7.6	17.3	18.7	15.7	9.3	4.3
2003.....	7.1	*	*	*	0.3	1.5	8.1	17.3	18.5	15.1	9.2	4.1
2002.....	7.0	*	*	*	0.3	1.5	8.1	16.8	18.4	15.5	9.3	4.4
2001.....	7.1	*	*	*	0.3	1.6	8.3	17.2	18.2	15.5	9.6	5.0
2000.....	7.0	*	*	*	0.2	1.6	8.5	16.3	18.7	15.8	9.9	5.4
1999.....	7.0	*	*	*	0.3	1.6	8.5	16.4	18.7	15.9	10.6	5.5

All races, male

2004.....	11.0	*	*	*	0.5	2.3	11.0	26.4	30.2	26.5	17.1	9.0
2003.....	10.9	*	*	*	0.5	2.3	11.4	27.0	29.8	25.4	16.8	7.8
2002.....	10.8	*	*	*	0.4	2.3	11.5	26.3	30.0	26.0	16.6	9.7
2001.....	10.8	*	*	*	0.4	2.3	11.9	26.6	29.7	26.3	17.2	11.6
2000.....	10.9	*	*	*	0.4	2.5	12.5	25.9	30.0	27.1	18.8	11.9
1999.....	10.9	*	*	*	0.4	2.3	12.6	26.2	30.5	27.4	19.4	11.8

All races, female

2004.....	3.5	*	*	*	0.2	0.7	4.2	8.6	8.1	6.6	3.9	2.3
2003.....	3.4	*	*	*	0.1	0.6	4.7	7.9	8.1	6.5	4.1	2.5
2002.....	3.4	*	*	*	*	0.7	4.7	7.7	7.6	6.8	4.5	2.1
2001.....	3.4	*	*	*	*	0.9	4.8	8.1	7.5	6.6	4.6	2.3
2000.....	3.2	*	*	*	*	0.7	4.5	7.0	8.3	6.6	4.1	2.7
1999.....	3.2	*	*	*	0.1	0.9	4.5	7.0	7.7	6.5	4.8	3.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 5/

White, both sexes												
2004.....	7.5	*	*	*	0.3	1.5	7.9	17.8	18.5	15.4	9.4	4.5
2003.....	7.4	*	*	*	0.3	1.5	8.2	17.4	18.2	15.2	9.3	4.2
2002.....	7.2	*	*	*	0.3	1.5	8.3	16.9	17.8	15.4	9.3	4.3
2001.....	7.2	*	*	*	0.2	1.6	8.3	16.8	17.6	15.3	9.6	4.8
2000.....	7.1	*	*	*	0.2	1.6	8.4	15.8	18.0	15.6	9.8	5.3
1999.....	7.0	*	*	*	0.3	1.6	8.2	15.7	17.7	15.3	10.3	5.3
White, male												
2004.....	11.5	*	*	*	0.5	2.3	11.4	26.9	29.5	25.6	17.3	9.3
2003.....	11.3	*	*	*	0.5	2.4	11.8	26.8	29.0	25.4	16.8	7.8
2002.....	11.1	*	*	*	0.4	2.3	11.8	26.4	29.0	25.4	16.5	9.3
2001.....	11.0	*	*	*	0.4	2.3	11.7	26.0	28.4	25.4	17.2	11.1
2000.....	11.0	*	*	*	0.4	2.5	12.3	25.3	28.8	26.4	18.6	11.6
1999.....	10.9	*	*	*	0.4	2.3	12.2	25.2	28.9	26.0	18.9	11.4
White, female												
2004.....	3.6	*	*	*	0.1	0.7	4.3	8.7	8.0	6.7	4.0	2.3
2003.....	3.5	*	*	*	*	0.7	4.7	8.0	8.0	6.5	4.3	2.5
2002.....	3.4	*	*	*	*	0.7	4.7	7.6	7.3	7.0	4.5	2.2
2001.....	3.4	*	*	*	*	0.8	4.8	7.6	7.4	6.7	4.6	2.2
2000.....	3.2	*	*	*	*	0.7	4.4	6.5	7.8	6.5	4.1	2.7
1999.....	3.1	*	*	*	*	0.9	4.1	6.4	7.2	6.5	4.6	2.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 5/

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Black, both sexes												
2004.....	6.1	*	*	*	*	1.0	5.7	17.1	23.5	20.1	7.6	*
2003.....	6.3	*	*	*	*	1.1	6.8	18.4	24.2	16.9	8.9	*
2002.....	6.4	*	*	*	*	1.3	6.7	18.5	25.3	18.3	11.2	*
2001.....	7.3	*	*	*	*	1.3	8.7	22.5	25.9	20.5	11.1	*
2000.....	7.4	*	*	*	*	1.7	9.2	22.7	26.4	21.0	11.5	6.3
1999.....	7.8	*	*	*	*	1.3	9.9	23.4	29.1	23.2	14.6	7.7
Black, male												
2004.....	9.7	*	*	*	*	1.6	8.8	27.0	41.5	39.5	14.5	*
2003.....	10.0	*	*	*	*	1.8	9.6	31.2	42.2	30.7	18.9	*
2002.....	10.0	*	*	*	*	2.0	9.4	29.3	43.5	35.8	21.8	*
2001.....	11.6	*	*	*	*	1.9	13.2	35.6	46.3	40.0	21.8	*
2000.....	11.4	*	*	*	*	2.5	13.6	36.0	43.8	39.9	24.2	*
1999.....	12.2	*	*	*	*	1.8	14.4	37.7	51.0	45.3	28.3	*
Black, female												
2004.....	2.8	*	*	*	*	*	2.9	8.6	9.0	6.2	3.8	*
2003.....	2.9	*	*	*	*	*	4.3	7.4	9.8	7.1	3.2	*
2002.....	3.2	*	*	*	*	*	4.4	9.3	10.7	5.8	5.2	*
2001.....	3.5	*	*	*	*	0.8	4.6	11.1	9.4	6.8	5.1	*
2000.....	3.7	*	*	*	*	1.0	5.2	11.3	12.3	7.7	4.3	*
1999.....	3.9	*	*	*	*	0.9	5.9	11.2	11.6	7.7	6.9	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/

All races 3/, both sexes

2004.....	10.5	0.7	0.2	0.2	7.0	12.6	20.9	21.7	9.1	4.3	4.8	6.3
2003.....	9.9	0.6	0.2	0.1	6.4	12.2	20.6	20.0	8.0	4.1	4.2	6.0
2002.....	9.0	0.7	0.2	0.1	5.5	11.1	19.7	17.9	6.9	3.6	3.8	5.7
2001.....	7.6	0.5	0.2	0.1	4.5	9.4	17.0	14.8	5.4	3.0	3.5	5.1
2000.....	7.0	*	*	0.1	4.0	8.8	16.0	13.2	4.9	2.6	3.5	5.7
1999.....	6.9	0.6	0.2	0.1	3.5	8.9	15.7	12.6	4.9	3.0	3.8	4.8

All races, male

2004.....	13.4	*	*	0.1	10.2	17.5	26.0	27.0	10.7	4.7	5.3	8.6
2003.....	12.9	*	0.3	0.1	9.2	16.6	26.5	25.7	9.4	4.6	4.9	7.8
2002.....	11.8	1.0	*	0.1	7.8	15.5	25.6	23.0	8.1	3.7	4.2	7.3
2001.....	10.2	*	0.3	0.1	6.6	13.2	22.7	19.7	6.3	3.5	3.8	5.7
2000.....	9.5	*	*	0.1	5.8	12.5	21.8	18.2	5.4	2.8	3.8	7.5
1999.....	9.4	*	*	*	5.0	12.7	21.9	17.5	6.0	3.4	3.8	6.2

All races, female

2004.....	7.6	*	*	0.2	3.6	7.7	15.8	16.5	7.7	3.9	4.4	5.2
2003.....	7.0	*	*	0.1	3.3	7.6	14.8	14.6	6.7	3.7	3.7	5.2
2002.....	6.3	*	0.3	*	3.0	6.6	13.9	13.1	5.7	3.4	3.5	4.9
2001.....	5.1	*	*	0.1	2.4	5.6	11.5	10.0	4.6	2.6	3.3	4.9
2000.....	4.6	*	*	*	2.1	5.1	10.2	8.5	4.4	2.5	3.2	4.9
1999.....	4.4	*	*	*	2.0	5.0	9.7	8.0	3.8	2.7	3.8	4.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All Under 1-4 5-14 15-24 25-34 35-44 45-54 55-64 65-74 75-84 85 years
ages 1/ 1 year 2/ years years years years years years years years years and over

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/

White, both sexes

2004.....	11.1	*	*	0.2	8.3	14.3	22.4	21.7	8.8	4.3	4.9	6.4
2003.....	10.4	*	0.2	0.1	7.6	13.6	21.7	20.0	7.8	4.1	4.5	6.3
2002.....	9.4	*	*	0.1	6.4	12.3	20.6	17.5	6.6	3.7	3.9	5.8
2001.....	7.8	*	*	0.1	5.2	10.3	17.5	14.0	5.1	3.0	3.6	5.1
2000.....	7.1	*	*	0.1	4.6	9.5	16.1	12.4	4.8	2.7	3.7	5.6
1999.....	6.9	*	*	0.1	3.9	9.4	15.6	11.7	4.6	3.0	4.0	4.8

White, male

2004.....	14.1	*	*	0.1	12.1	19.7	27.7	26.5	9.5	4.4	5.3	8.7
2003.....	13.5	*	*	*	11.0	18.6	27.8	25.1	8.7	4.3	5.2	7.8
2002.....	12.2	*	*	0.1	9.1	17.1	26.6	21.9	7.2	3.6	4.2	7.6
2001.....	10.4	*	*	*	7.6	14.4	23.2	18.1	5.5	3.4	3.8	5.6
2000.....	9.6	*	*	0.1	6.7	13.3	21.9	16.5	4.9	2.7	4.0	7.2
1999.....	9.3	*	*	*	5.5	13.3	21.7	15.6	5.2	3.3	3.9	6.4

White, female

2004.....	8.2	*	*	0.2	4.2	8.6	17.0	17.0	8.1	4.3	4.7	5.3
2003.....	7.4	*	*	*	3.9	8.3	15.6	15.0	7.0	3.9	4.0	5.6
2002.....	6.7	*	*	*	3.5	7.3	14.5	13.2	6.0	3.7	3.8	4.9
2001.....	5.4	*	*	*	2.7	6.0	11.7	10.1	4.8	2.7	3.5	4.9
2000.....	4.7	*	*	*	2.3	5.5	10.2	8.4	4.7	2.7	3.4	4.9
1999.....	4.5	*	*	*	2.2	5.2	9.5	7.9	4.1	2.8	4.1	4.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

All Under 1-4 5-14 15-24 25-34 35-44 45-54 55-64 65-74 75-84 85 years
ages 1/ 1 year 2/ years years years years years years years years years and over

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/

Black, both sexes

2004.....	9.4	*	*	*	2.2	7.9	18.7	28.1	14.4	4.6	3.9	*
2003.....	9.2	*	*	*	1.8	8.1	20.8	26.9	11.7	5.0	2.2	*
2002.....	9.2	*	*	*	2.1	8.5	20.7	26.8	11.5	3.1	2.7	*
2001.....	8.5	*	*	*	1.9	7.7	20.0	25.2	8.8	3.2	2.6	6.1
2000.....	8.3	*	*	*	2.0	8.3	20.1	23.8	7.5	3.0	2.4	7.5
1999.....	8.6	*	*	*	2.1	9.1	20.9	23.7	8.3	3.5	2.2	*

Black, male

2004.....	12.8	*	*	*	3.2	11.1	23.8	39.1	24.1	8.5	7.0	*
2003.....	12.7	*	*	*	2.5	10.7	27.7	38.9	19.0	8.6	*	*
2002.....	12.8	*	*	*	2.9	12.0	27.7	39.2	19.9	5.4	*	*
2001.....	12.2	*	*	*	2.9	10.8	27.0	39.3	15.1	5.4	*	*
2000.....	12.0	*	*	*	2.7	12.0	28.2	37.6	12.4	5.3	*	*
1999.....	12.7	*	*	*	3.2	13.3	30.1	38.2	15.2	5.4	*	*

Black, female

2004.....	6.3	*	*	*	1.3	5.0	14.2	18.6	6.6	*	*	*
2003.....	6.1	*	*	*	1.1	5.8	14.6	16.6	5.8	2.4	*	*
2002.....	5.8	*	*	*	1.3	5.3	14.5	16.1	4.8	*	*	*
2001.....	5.1	*	*	*	0.9	4.8	13.8	13.0	3.8	*	*	*
2000.....	4.9	*	*	*	1.3	4.9	13.0	11.9	3.5	*	*	*
1999.....	4.8	*	*	*	1.0	5.2	12.9	11.3	2.8	2.1	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/

All races 3/, both sexes

2004.....	10.1	*	0.3	0.7	15.7	15.0	11.3	11.0	9.8	10.2	13.3	11.9
2003.....	10.4	*	0.3	0.8	16.6	15.5	11.5	11.2	10.1	10.4	13.5	12.5
2002.....	10.5	*	0.4	0.8	16.7	15.4	12.1	10.8	10.2	10.9	14.4	12.5
2001.....	10.4	*	0.5	0.8	16.7	15.5	11.7	10.5	10.1	10.9	14.3	12.8
2000.....	10.2	*	0.3	0.9	16.8	14.5	11.9	10.5	9.4	10.6	13.9	14.2
1999.....	10.3	*	0.4	1.0	17.6	14.9	11.6	10.2	9.7	11.0	14.2	13.5

All races, male

2004.....	17.6	*	0.4	1.1	27.5	26.1	18.7	18.3	17.1	19.7	29.8	35.9
2003.....	18.3	*	0.3	1.2	29.2	27.1	19.1	18.8	17.7	20.3	30.2	37.8
2002.....	18.4	*	0.5	1.2	29.3	26.5	20.1	18.2	18.0	21.3	32.9	38.9
2001.....	18.2	*	0.5	1.2	29.6	26.7	19.2	17.8	17.4	21.1	32.8	40.2
2000.....	17.8	*	0.4	1.4	29.4	24.9	19.4	17.6	16.3	20.3	32.2	44.7
1999.....	18.1	*	0.5	1.5	30.6	25.5	18.9	17.0	17.1	21.5	32.8	43.3

All races, female

2004.....	2.7	*	0.3	0.3	3.2	3.7	3.9	4.0	3.1	2.2	2.3	1.0
2003.....	2.7	*	0.3	0.3	3.3	3.6	4.0	3.8	2.9	2.2	2.2	1.3
2002.....	2.8	*	0.3	0.5	3.5	4.0	4.2	3.6	3.1	2.3	2.1	1.1
2001.....	2.8	*	0.4	0.4	3.2	4.0	4.3	3.5	3.3	2.4	2.2	1.3
2000.....	2.8	*	*	0.4	3.5	4.0	4.4	3.6	3.0	2.5	2.0	1.7
1999.....	2.9	*	0.4	0.5	3.9	4.1	4.4	3.6	2.9	2.4	2.1	1.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All	Under	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 years
	ages 1/	1 year 2/	years	years	years	years	years	years	years	years	years	and over

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/

White, both sexes

2004.....	9.0	*	*	0.6	10.7	10.9	10.6	11.4	10.4	11.0	14.3	12.8
2003.....	9.2	*	0.2	0.6	11.1	11.2	10.8	11.5	10.8	11.3	14.4	13.5
2002.....	9.3	*	0.3	0.7	11.3	11.2	11.3	11.1	10.9	11.8	15.3	13.4
2001.....	9.4	*	0.4	0.8	11.4	11.9	11.1	10.9	10.7	11.7	15.2	13.8
2000.....	9.1	*	0.2	0.9	11.5	10.9	11.2	10.6	9.9	11.2	15.0	15.2
1999.....	9.2	*	0.3	0.9	12.0	11.5	11.0	10.3	10.1	11.8	15.1	14.5

White, male

2004.....	15.5	*	*	0.9	18.4	18.2	17.1	18.7	17.9	20.9	31.8	38.9
2003.....	16.0	*	*	1.0	19.2	18.8	17.5	19.2	18.9	21.9	32.3	41.0
2002.....	16.1	*	0.4	0.9	19.4	18.5	18.5	18.4	19.0	22.9	34.8	41.7
2001.....	16.2	*	0.4	1.1	19.5	19.9	18.0	18.3	18.3	22.4	34.7	43.3
2000.....	15.6	*	*	1.3	19.6	18.1	17.9	17.7	17.0	21.4	34.5	48.4
1999.....	15.9	*	*	1.3	20.5	19.2	17.6	17.2	17.6	22.8	34.7	46.8

White, female

2004.....	2.7	*	*	0.3	2.6	3.2	4.0	4.3	3.3	2.5	2.4	1.1
2003.....	2.6	*	*	0.2	2.5	3.1	4.0	4.0	3.2	2.3	2.3	1.3
2002.....	2.7	*	*	0.4	2.6	3.5	4.1	3.8	3.3	2.4	2.2	1.2
2001.....	2.7	*	*	0.4	2.7	3.6	4.1	3.7	3.6	2.6	2.3	1.4
2000.....	2.7	*	*	0.3	2.8	3.5	4.3	3.7	3.2	2.8	2.1	1.8
1999.....	2.8	*	*	0.5	3.0	3.6	4.3	3.6	3.1	2.6	2.2	1.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/

Black, both sexes

2004.....	19.3	*	1.2	1.5	44.1	43.8	18.8	11.1	8.0	6.2	6.6	*
2003.....	20.1	*	0.9	1.7	47.7	45.4	19.0	11.4	6.8	6.0	6.1	*
2002.....	20.2	*	*	1.5	47.7	44.7	20.5	11.8	7.2	5.8	7.1	*
2001.....	19.3	*	0.9	1.2	48.1	40.7	19.0	10.6	6.9	6.6	7.5	*
2000.....	19.3	*	*	1.4	48.1	39.4	19.2	12.1	6.7	6.9	5.5	*
1999.....	19.4	*	1.1	1.6	49.9	37.7	18.3	11.5	8.2	6.6	7.5	*

Black, male

2004.....	36.4	*	*	2.3	80.7	83.6	35.1	20.2	14.8	14.4	15.9	*
2003.....	37.8	*	*	2.5	87.6	87.2	34.8	21.1	12.9	11.7	13.8	*
2002.....	37.8	*	*	2.1	87.1	85.6	36.9	21.4	13.5	12.1	17.5	*
2001.....	36.4	*	*	1.8	90.3	77.7	33.2	19.1	13.7	13.8	18.0	*
2000.....	36.1	*	*	2.2	89.3	74.8	34.3	21.7	12.6	15.1	12.1	*
1999.....	36.0	*	*	2.3	91.4	71.3	32.2	20.4	16.5	14.0	18.1	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2004

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. Multiple-race data was reported for California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and for California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these States were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/

Black, female

2004.....	3.7	*	*	0.7	6.9	7.1	4.3	3.3	2.6	*	*	*
2003.....	3.9	*	*	0.9	7.4	7.3	5.0	3.1	2.0	2.0	*	*
2002.....	4.2	*	*	0.9	8.1	7.5	5.9	3.5	2.1	*	*	*
2001.....	3.8	*	*	*	6.1	7.3	6.5	3.2	1.5	*	*	*
2000.....	4.0	*	*	*	7.6	7.4	5.7	3.8	1.9	*	*	*
1999.....	4.4	*	*	0.8	9.0	7.4	6.1	3.8	1.6	*	*	*

* Figure does not meet standards of reliability or precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

1/ Figures for age not stated included in "All ages" but not distributed among age groups.

2/ Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).

3/ Includes races other than white and black.

4/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

5/ Included in selected categories.

Source: CDC/NCHS, National Vital Statistics System, Mortality.