

**Table 14. Tuberculosis Cases and Percentages by Reason Tuberculosis Therapy Stopped: United States, 1993–2005**

Year	Total Cases <sup>1</sup>	Completed Therapy		Moved		Lost		Refused		Died <sup>2</sup>		Unknown <sup>3</sup>	
	No.	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
1993	23741	18042	(76.0)	1120	(4.7)	1087	(4.6)	223	(0.9)	3052	(12.9)	217	(0.9)
1994	23051	17763	(77.1)	1194	(5.2)	739	(3.2)	183	(0.8)	2743	(11.9)	429	(1.9)
1995	21712	17285	(79.6)	969	(4.5)	566	(2.6)	156	(0.7)	2391	(11.0)	345	(1.6)
1996	20297	16510	(81.3)	783	(3.9)	520	(2.6)	156	(0.8)	1992	(9.8)	336	(1.7)
1997	18930	15654	(82.7)	667	(3.5)	435	(2.3)	119	(0.6)	1755	(9.3)	300	(1.6)
1998	17584	14763	(84.0)	534	(3.0)	400	(2.3)	104	(0.6)	1578	(9.0)	205	(1.2)
1999	16863	14219	(84.3)	456	(2.7)	356	(2.1)	104	(0.6)	1436	(8.5)	292	(1.7)
2000	15786	13399	(84.9)	408	(2.6)	392	(2.5)	112	(0.7)	1294	(8.2)	181	(1.1)
2001	15410	13195	(85.6)	375	(2.4)	364	(2.4)	98	(0.6)	1115	(7.2)	263	(1.7)
2002	14559	12430	(85.4)	336	(2.3)	364	(2.5)	86	(0.6)	1071	(7.4)	272	(1.9)
2003	14381	12391	(86.2)	312	(2.2)	359	(2.5)	85	(0.6)	989	(6.9)	245	(1.7)
2004	14078	11998	(85.2)	341	(2.4)	331	(2.4)	82	(0.6)	960	(6.8)	366	(2.6)
2005	13662	11629	(85.1)	336	(2.5)	317	(2.3)	88	(0.6)	974	(7.1)	318	(2.3)

<sup>1</sup>Includes all cases in persons reported as alive at diagnosis and taking one or more TB drugs.

<sup>2</sup>Died = died of any cause (not only TB).

<sup>3</sup>Includes cases in persons reporting reason therapy stopped = Other, Missing, or Unknown.

**Note:** Data for all years are updated through April 23, 2008.

Data complete through 2005 only. See Technical Notes (page 9) for details.