

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

SLAC-PUB-2089
March 1978
(T)

Cont - 780145 -- 1 2

REMARKS ON THE TOPOLOGY OF GAUGE FIELDS*

Y. Nambu[†]

Stanford Linear Accelerator Center
Stanford University, Stanford, California 94305

In contrast to the short distance behavior of quantum chromodynamics (QCD), which is within the scope of perturbation theory and thus can be subjected to quantitative tests, the large distance or strong coupling regime of QCD is not well understood yet even qualitatively. An overriding problem of interest in quark confinement, and various theoretical schemes have been put forward to show that the quarks can indeed be confined. These schemes, though varying from one to another in detail, rely on the idea that the growing coupling constant at large distances plays a key role. It is not clear yet, however, whether confinement is a natural consequence of QCD alone, or it requires some independent and extraneous assumptions.

(Introductory remarks made at Orbis Scientiae Conference,
15-20 January 1978, Coral Gables, Florida.)

*Work supported in part by the Department of Energy and the Guggenheim Fellowship.

†On leave from the University of Chicago.

Another important aspect of gauge fields is the existence of topologically nontrivial configurations such as vortices, monopoles and instantons. They may be regarded as natural consequences of gauge physics. Most likely these configurations also are crucial to the understanding of confinement as is claimed by a number of recent papers. For this reason I have planned this session to be organized mainly around the topological problems.

My own remarks on these questions will be very brief:

1. Action versus Free Action

There is a formal analogy between statistical mechanics and the Feynman formulation of quantum field theory, which seems to become especially relevant in gauge theories. This analogy was emphasized by the Princeton group¹ who made use of the concept of entropy. Let us write down the Feynman integral (in the Euclidean form) for a gauge field

$$Z = \int \exp \left[- \frac{1}{4g^2} \int F_{\mu\nu} F_{\mu\nu} d^4x \right] D(A_\mu) \quad (1)$$

where the coupling constant does not enter the definition of $F_{\mu\nu}$. The functional measure is not clearly defined, but we intend to integrate over the A_μ without imposing special gauge conditions. For a given value of the action, there is a corresponding phase space volume which we write symbolically as $\exp S$:

$$\begin{aligned} Z &= \int \exp \left[- \frac{1}{g^2} I + S(I) \right] dI \\ &= \exp \left[- \Theta / g^2 \right], \\ \Theta &\approx (I - g^2 S)_{\min}. \end{aligned} \quad (2)$$

MASTER
MASTER

EB

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

Let us call Θ free action in contrast to the action I . In principle, we should minimize Θ rather than I , and this difference could become important for large g^2 , which plays the role of temperature.

This statement takes on a real significance when the topology of gauge fields is considered. Suppose we start from the lowest value of I , i.e. the pure gauges: $F_{\mu\nu} = 0$, $I = 0$. This part of phase space can be parametrized by a unitary matrix field u ,

$$A_\mu = iu^\dagger \partial_\mu u, \quad u^\dagger u = 1 \quad (2a)$$

The field u , however, may be topologically non-trivial. For example, choose

$$u = \tau_\mu x_\mu / |x|, \quad \tau_\mu = (\vec{\tau}, i) \quad (3)$$

in the case of $SU(2)$ theory. Then u becomes singular at the origin, and $F_{\mu\nu}$ and I cannot be identically zero since the Pontrjagin index

$$\begin{aligned} \frac{1}{16\pi^2} \text{Tr} \int F_{\mu\nu} \tilde{F}_{\mu\nu} d^4\chi &= \int Q_\mu dS_\mu, \\ Q_\mu &= -\frac{1}{3} \text{Tr} u^\dagger \partial_\nu u \partial_\lambda u^\dagger \partial_\rho u \epsilon^{\mu\nu\lambda\rho} \end{aligned} \quad (4)$$

is computed to be nonzero. Thus we might say that pure gauges are not a well defined concept, or the entropy cannot be defined for given I , at least not for $I = 0$. There seems to be a sort of uncertainty principle between I and S due to topology. Clearly this is related to the Gribov problem² and Singer's observation³ of its generality. At any rate we are forced to enlarge pure gauges to their neighborhood which I will call almost pure gauges. It is conceivable that for large enough g^2 the gain in S by including various topologies can outweigh the cost of larger I , thus actually lowering the free action. Then the topology-averaged field configurations which minimize the free action may

deviate substantially from the naive classical configurations which minimize only I .

2. A class of almost pure gauges.⁴

It is both natural and convenient to consider the following class of almost pure gauges

$$A_\mu = ifu^\dagger \partial_\mu u \quad (5)$$

where f is a scalar function which vanishes at the singularities of u . The topological characterization of a singularity will not be altered by this if $f \rightarrow 1$ in a region surrounding it. From Eq. (5) we obtain

$$F_{\mu\nu} = f(1-f) \partial_\mu u^\dagger \partial_\nu u + \partial_\mu f u^\dagger \partial_\nu u - (\mu \leftrightarrow \nu). \quad (6)$$

It is not surprising that this class of configurations cover various known examples of nontrivial topology such as instanton, meron, monopole, and string.

A more interesting point is that we can generalize Eq. (5) further as

$$A_\mu = i \sum_i f_i u_i^\dagger \partial_\mu u_i. \quad (7)$$

The many-instanton solution of 't Hooft (in the singular gauge) is indeed of this form. Furthermore, Eq. (7) can be written as a London relation⁵

$$A_\mu = \lambda J_\mu = \lambda \sum_{i=1}^N J_{i\mu},$$

$$J_{i\mu} = ih_i u_i^\dagger D_\mu u_i$$

where

$$D_\mu u = \partial_\mu u + iu A_\mu, \quad (8)$$

f_i and h_i are related by

$$f_i = \lambda h_i / (1 + \lambda \sum_j h_j) . \quad (9)$$

Then a simple superposition principle holds for the currents $J_{i\mu}$ to generate many instantons, and the total current is conserved: $D_\mu J_\mu = 0$. It might also be instructive to observe that Eq. (8) can be further simplified if J_μ is defined in terms of a $2N \times 2$ rectangular matrix U :

$$U = \sum \oplus h_i^{\frac{1}{2}} u_i \quad (\text{assuming } h_i \geq 0), \quad U^\dagger U = (\sum h_i) 1 ,$$

$$J_\mu = \frac{i}{2} [U^\dagger (D_\mu U) - (D_\mu U^\dagger) U] . \quad (10)$$

U has a gauged $SU(2)$ symmetry acting from the right, and a global $SU(2) \times S_N$ symmetry acting from the left which may be generalized to $SU(2N)$. These two are independent, like color and flavor.

3. A remark concerning the Wilson criteria.

The Wilson criterion is widely used to test the confinement property of a theory. In the context of functional integration, one evaluates

$$\langle W \rangle = \sum W_i \rho_i ,$$

$$W = \text{Tr} \exp [i \oint A_\mu dx_\mu] \quad (11)$$

where ρ_i is the weight of a configuration i . In general Eq. (11) is expected to yield an asymptotic form

$$\langle W \rangle \sim a \exp [-\lambda L] + b \exp [-\mu L^n] + \dots, \quad n > 1 \quad (12)$$

where L is the linear dimension of the Wilson loop. What I would like to emphasize is that to prove confinement, one must first show the absence of configurations contributing to the first term rather than the presence of

configurations contributing to the second term. I suppose that the former task is more demanding and difficult than the latter.

REFERENCES

1. C. Callan et al., Phys. Lett 66B, 375 (1977).
2. V. M. Gribov, Lectures at the 12th Winter School of the Leningrad Physics Institute (1977).
3. I. M. Singer, Berkeley preprint (Math. Department), 1977.
4. See also Y. Nambu, Cal Tech preprint CALT 68-634, 1977.
5. Y. Nambu, Annals of the New York Academy of Sciences 294 (Five Decades of Weak Interactions) p. 74 (1977).