

Mean serum total cholesterol of adults 20 years and older, 1999-2002

S U D A A N

Software for the Statistical Analysis of Correlated Data
 Copyright Research Triangle Institute February

2005

Release 9.0.1

Number of observations read : 19759 Weighted count :278652243
 Observations in subpopulation : 9471 Weighted count:198419259
 Denominator degrees of freedom : 29

Date: 01-31-2007 Page : 1
 Variance Estimation Method: Taylor Series (WR) Table : 1
 For Subpopulation: RIDAGEYR >= 20
 Prevalence of SPs with measured high blood pressure : NHANES 1999-2002
 by: Variable, Gender, AGE GROUP, Race Ethnicity.

for: Variable = HBP: High BP, Gender = Total.

AGE GROUP	Race Ethnicity	Sample Size	Percent	SE

Total				
	Total	8960	29.1630	0.9817
	NH White	4439	29.6593	1.1555
	NH Black	1677	36.9562	1.5130
	Mexican American	2087	17.1412	1.2180
	Other	757	25.8660	2.2956
20-39				
	Total	3175	7.1830	0.8315
	NH White	1408	6.3050	0.8653
	NH Black	602	11.9044	1.3178
	Mexican American	825	5.0434	0.9021
	Other	340	8.8807	2.3944
40-59				
	Total	2692	29.7023	1.3820
	NH White	1287	27.4901	1.7061
	NH Black	555	46.4504	2.1302
	Mexican American	620	26.0275	1.9296
	Other	230	30.7528	4.3219
60+				
	Total	3093	66.7150	1.2708

NH White	1744	65.3819	1.5779
NH Black	520	81.4957	1.6193
Mexican American	642	62.0765	1.4276
Other	187	67.3857	5.0892

Date: 01-31-2007

Page : 2

Variance Estimation Method: Taylor Series (WR)

Table : 1

For Subpopulation: RIDAGEYR >= 20

Prevalence of SPs with measured high blood pressure : NHANES 1999-2002

by: Variable, Gender, AGE GROUP, Race Ethnicity.

for: Variable = HBP: High BP, Gender = Male.

AGE GROUP	Sample		
Race Ethnicity	Size	Percent	SE
Total			
Total	4228	27.2721	1.2213
NH White	2143	27.7521	1.2987
NH Black	800	35.3589	1.5435
Mexican American	957	17.2746	1.6922
Other	328	23.0655	2.8433
20-39			
Total	1352	10.1333	1.3483
NH White	595	9.3704	1.4354
NH Black	265	15.4747	1.4174
Mexican American	350	6.8777	1.6786
Other	142	12.1588	3.8178
40-59			
Total	1343	28.3341	1.9214
NH White	668	26.8355	2.2446
NH Black	283	41.5988	3.3480
Mexican American	292	26.6817	2.5467
Other	100	27.3367	5.6711
60+			
Total	1533	59.4742	1.7058
NH White	880	58.2755	1.9405
NH Black	252	77.9679	1.8563
Mexican American	315	57.2239	2.3411
Other	86	55.0099	9.0828

Date: 01-31-2007

Page : 3

Variance Estimation Method: Taylor Series (WR)

Table : 1

For Subpopulation: RIDAGEYR >= 20

Prevalence of SPs with measured high blood pressure : NHANES 1999-2002

by: Variable, Gender, AGE GROUP, Race Ethnicity.

for: Variable = HBP: High BP, Gender = Female.

AGE GROUP Race Ethnicity	Sample Size	Percent	SE

Total			
Total	4732	30.9091	1.0690
NH White	2296	31.4636	1.3439
NH Black	877	38.2763	2.1365
Mexican American	1130	17.0021	1.5080
Other	429	28.1374	3.1949
20-39			
Total	1823	4.2766	0.6670
NH White	813	3.2251	0.6012
NH Black	337	8.9185	2.1830
Mexican American	475	3.0391	0.8401
Other	198	5.7442	2.7978
40-59			
Total	1349	31.0173	1.5526
NH White	619	28.1486	1.9675
NH Black	272	50.7866	3.3318
Mexican American	328	25.3509	2.4511
Other	130	33.2027	4.6175
60+			
Total	1560	72.2896	1.5304
NH White	864	71.0079	1.9252
NH Black	268	83.8735	2.0034
Mexican American	327	65.9937	1.8603
Other	101	75.2974	4.8470
