The Case of the Cryptic Clustered Cryptococci

(Outbreak of Cryptococcus neoformans var. gattii on Vancouver Island , British Columbia)

Mike Starr, Murray Fyfe, Marc Romney, Bill Black, Pamela Kibsey, Laura MacDougall, Sunny Mak, Marty Pearce, Judith Isaac-Renton, Louise Stein, Craig Stephen, David Patrick University of British Columbia Centre for Disease Control, Vancouver, CANADA




Cryptococcus

- *Encapsulated yeast-like fungus
- *World-wide distribution
- 37 species
- C neoformans is the only species that is pathogenic


C neoformans var. neoformans


- *90% of recognised C neoformans disease
- Usually seen as opportunistic pathogen in patients with AIDS
- Most common cause of fungal meningitis
- Pigeon or other avian dung reservoir



C neoformans var. gattii (CNVG)


- Less common pathogen
- Immunocompetent hosts
- Large mass lesions in lung &/or brain
- Geographically restricted
- Specific ecological association with Eucalyptus camaldulensis







Vancouver Island Background Information



- Mean Temp: 2.3°C Jan, 17.6°C Aug
- Precip: 857-3295mm/yr
- ❖ Population: 703,052
- Median age: 40.2yrs (37.2yrs for BC)
- Over 65 yo: 16.3% (12.8% for BC)
- Aboriginal population: 39,010 (5.5%)


Vancouver Island Cluster

- January 1, 1999 March 1, 2002:41 confirmed cases of Cryptococcal Disease
- Mainly immunocompetent adults
- Over the same period, vets reported
 ↑ cryptococcal cases in domestic pets
- Initial testing suggested isolates to be CNVG

CNVG on Vancouver Island

- Incidence pre-1999: ??
- Incidence since 1999:
 - * 1999 = 8.5 per million
 - * 2000 = 26 per million
 - * 2001 = 24 per million
- (Incidence in Aust: ~1/million/year)

No. of Cases of Cryptococcus by date of initial presentation



Investigation

- Chart Reviews
- Case Interviews
- Case Control Study
- Veterinary Study
- Molecular typing
- Serology
- Environmental Studies

Patient Characteristics

- Mean age 61 years (range 19-88)
- *61% male
- 64% immunocompetent
- 36 cases involved lungs (88%)
- Meningitis in 10 (24%)
- 6 deaths 2 attributable

Preliminary findings

- Possible exposures
 - Wooded area (70%)
 - Gardening (64%)
 - Fertilizer/compost (60%)
- Unlikely exposures
 - Animal exposures farms, pets, birds
 - * Outdoor leisure activities
 - No ubiquitous exposure to eucalyptus, fig, almond trees

Preliminary findings

- Possible predisposer
 - ♦ Smoking (68%)
 - Immunocomp (36%)
 - Steroids (25%)
 - Smoking or Immunocomp (89%)

Distribution of Human and Animal Cryptococcal Cases in Southwestern British Columbia, 1999-2002*



Environmental Sampling


- Samples taken from tree holes, rotting wood in "hot spots"
- Garry Oak and Arbutus
- GPS readings
- Bird seed agar
- No cryptococci to date

Laboratory Typing

- 21 C neoformans isolates from humans typed
 - **♦ 15 CNVG**
 - ◆ 1 CNVN
 - 5 not able to be typed
- 4 animal isolates typed all CNVG

Other typing

- Molecular typing
 - * PCR fingerprinting
 - * AFLP
- Immunohistochemistry
- Type-specific serology
 - Immunoblotting of sera against CNVG protein extracts
 - * IgG reactive to CNVG protein detected



Conclusions

- CNVG usu seen in tropical/subtropical climate
- Incidence of CNVG in endemic areas like Australia is approx 1/million/year
- Estimated incidence on Vancouver Island since 1999 - 24/million/year
- Concurrent outbreak amongst animals
- This outbreak may in fact represent the emergence of a new disease

Acknowledgements

- Sarah Kidd
- Weiland Meyer
- Jim Kronstad
- *Richard Stanwick
- Linda Poffenroth
- Fred Rockwell
- *Brian Emerson
- Karen Bartlett

- Sultana Mithani
- * Amelia Trinidad
- Sally Lester
- Josh Waddington
- Peter Phillips
- Tania Sorrell
- David Ellis
- Bryan Speed