

Detection of CD8a in Frozen Mouse Tissue

Reagent and Antibody Information

[1X Wash Buffer](#)
[3% Hydrogen Peroxide](#)
[1% BSA Diluent](#)
[DAB Chromagen](#)
[Hematoxylin](#)
[Rapid Fixx](#)

Blocking Serum: Normal Goat Serum
Jackson ImmunoResearch Laboratories, Inc.
West Grove, PA 19390
www.jacksonimmuno.com
1-800-367-5296
Catalog # 005-000-121

Avidin / Biotin Blocking Kit
Vector Laboratories, Inc.
Burlingame, CA 94010
www.vectorlabs.com
1-800-227-6666
Catalog # SP-2001

Primary Antibody: Rat Anti-Mouse CD8a Monoclonal Antibody
BD Biosciences
San Jose, CA 95131
www.bdbioscience.com
1-877-232-8995
Catalog # 550281

Negative Control Serum: Purified Rat IgG2a
BD Biosciences
San Jose, CA 95131
www.bdpharma.com
1-877-232-8995
Catalog # 559073

Secondary Antibody: Biotin Polyclonal Goat Anti-Rat Ig (Multiple Adsorbed)
BD Biosciences
San Jose, CA 95131
www.bdpharma.com
1-877-232-8995
Catalog #559286

Label Complex: Peroxidase-Conjugated Streptavidin SS Label

Biogenex Laboratories

San Ramon, CA 94583

www.biogenex.com

1-800-421-4149

Catalog # HK330-9K

Staining Procedure

Positive Control Tissue: Spleen – most T-cell

Stain Localization: Membrane and cytoplasmic

1. Cut each frozen section at 6µm and mount on a positively charged slide.
Immediately fix the section in Rapid Fix Solution for 7 seconds.
Rinse the slide thoroughly in tap water to remove excess fixative and then place in 1X Wash Buffer.
Once all the slides have undergone this process, proceed to step 2.
2. Rinse the slides in 2 changes of 1X Wash Buffer for 5 minutes each.
3. Quench endogenous peroxidase by placing the slides in 0.3% hydrogen peroxide for 30 minutes.
4. Rinse the slides in 2 changes of 1X Wash Buffer for 5 minutes each.
5. Block with 5% Normal Goat Serum for 20 minutes at room temperature.
Lot # _____ Date Reconstituted _____

DO NOT RINSE SLIDES. CONTINUE TO AVIDIN-BIOTIN BLOCK.

6. Avidin / Biotin Blocking Kit
Lot # _____ Exp. Date _____ New Kit: yes / no
Apply avidin block - 15 minutes at room temperature.
Quick rinse in 1X Wash Buffer.
Apply biotin block - 15 minutes at room temperature.

DO NOT RINSE SLIDES WITH BUFFER BEFORE ADDING PRIMARY ANTIBODY.
ONLY WIPE EXCESS BUFFER.

7. Apply primary antibody at a 1:30 dilution and incubate for one hour at room temperature.
Lot # _____ Exp. Date _____

For negative control slides, dilute the protein concentration of the purified Rat IgG2a to match that of the primary antibody, if necessary. Make a 1:30 dilution from this normalized serum and apply to the slides. Incubate for one hour at room temperature.

Lot # _____ Date Reconstituted _____

8. Rinse the slides in 2 changes of 1X Wash Buffer for 5 minutes each.

9. Apply the goat anti-rat Ig secondary antibody at a 1:200 dilution and incubate for 30 minutes at room temperature.

Lot # _____ Exp. Date _____

10. Rinse the slides in 2 changes of 1X Wash Buffer for 5 minutes each.

11. Apply the Streptavidin SS Label and incubate for 30 minutes at room temperature.

Lot # _____ Exp. Date _____

12. Rinse the slides in 2 changes of 1X Wash Buffer for 5 minutes each.

13. Apply the DAB chromagen and incubate in the dark for 6 minutes at room temperature.

(Add 1 drop of DAB per ml of substrate)

Lot # _____ Exp. Date _____ New Kit: yes / no

14. Rinse the slides in tap water 3 minutes.

15. Counterstain with Harris Hematoxylin for 30 seconds.

16. Rinse the slides in tap water until water is clear.

17. Gently agitate slides in 1X Wash Buffer until they turn blue.

18. Dehydrate through the following solutions:

95% Ethanol	1 time	3 minutes
100% Ethanol	3 times	3 minutes
Xylene	2 times	5 minutes

19. Coverslip

Updated 09/27/05