

Estimated Pregnancy Rates for the United States, 1990–2000: An Update

by Stephanie J. Ventura, M.A.; Joyce C. Abma, Ph.D.; William D. Mosher, Ph.D., Division of Vital Statistics; and Stanley Henshaw, Ph.D., The Alan Guttmacher Institute

Abstract

Objectives—This report presents detailed pregnancy rates for 1990–2000, updating a national series of rates extending since 1976.

Methods—Tabular data on pregnancy rates by age, race, and Hispanic origin, and by marital status are presented and briefly described.

Results—In 2000 an estimated 6,401,000 pregnancies resulted in 4.06 million live births, 1.31 million induced abortions, and 1.03 million fetal losses. The 2000 pregnancy rate of 104.0 pregnancies per 1,000 women aged 15–44 years is 10 percent lower than the 1990 peak of 115.6. The teenage pregnancy rate dropped 27 percent during 1990–2000, reaching an historic low of 84.5 pregnancies per 1,000 women aged 15–19 years. Rates for younger teenagers declined relatively more than for older teenagers.

Introduction

This report presents pregnancy rates for 1990–2000, by age, race, Hispanic origin, and by marital status. The data in this report, together with previously published pregnancy rates for earlier years, provide a consistent series of rates for the United States for 1976–2000 (1). This update contains two new elements: new data on pregnancy rates for the year 2000 and revised rates for teenagers and marital status groups for 1991–99. The revised rates for 1991–99 use newly released population estimates that are consistent with the April 1, 2000, census (2). Rates for the 1990s that were published in 2003 (3) were based on 2000 census level population estimates prepared by the National Center for Health Statistics (2,4). Rates initially issued for 1991–97 were based on postcensal population estimates consistent with the 1990 census (1,5).

Rates are presented by pregnancy outcome, age, race, and Hispanic origin, and by marital status, race, and Hispanic origin (tables 1–4). Details of sources and methods of estimation are presented in earlier reports (1,3,5–7).

Overview of findings

An estimated 6,401,000 pregnancies ended in 2000, about 6 percent fewer than the 1990 peak (6,778,000) for the period 1976–2000. The 2000 total included 4.06 million live births, 1.31 million induced abortions, and 1.03 million fetal losses. The estimated pregnancy rate for U.S. women was 104.0 pregnancies per 1,000 women aged 15–44 years, 10 percent below the 1990 peak (115.6 per 1,000).

The pregnancy rate for teenagers fell 27 percent overall during 1990–2000, to 84.5 pregnancies per 1,000 women aged 15–19 years. This rate was the lowest reported since 1976. The overall decline is reflected in significant reductions in birth and abortion rates. Pregnancy rates fell more for 15–17-year-old teenagers (33 percent) than for 18–19 year olds (20 percent). Rates declined about one-third each for white and black non-Hispanic teenagers and by 15 percent for Hispanic teenagers (table 1).

During the 1990–2000 decade, pregnancy rates declined about 8 percent for married women and 12 percent for unmarried women (table 4). The birth rate for married women declined about 6 percent, while the abortion rate fell one-fourth. The birth rate for unmarried women changed little, but the abortion rate for unmarried women also declined about one-fourth.

Findings from Cycle 6 of the National Survey of Family Growth (available later in 2004) are expected to help explain the trends and variations in pregnancy rates and the factors that account for them.

Acknowledgments

This report was prepared in the Division of Vital Statistics under the general direction of Charles J. Rothwell, Director of the Division of Vital Statistics. Yashodhara Patel provided table review. The report was edited by Demarius V. Miller and typeset by Jacqueline M. Davis of the Office of Information Services, Information Design and Publishing Staff.

References

1. Ventura SJ, Mosher WD, Curtin SC, Abma JC, Henshaw S. Trends in pregnancies and pregnancy rates by outcome: Estimates for the United States, 1976–96. National Center for Health Statistics. Vital Health Stat 21(56). 2000.
2. National Center for Health Statistics. U.S. census populations with bridged race categories. Available at: <http://www.cdc.gov/nchs/about/major/dvs/popbridge/popbridge.htm>.
3. Ventura SJ, Abma JC, Mosher WD, Henshaw S. Revised pregnancy rates, 1990–97, and new rates for 1998–99: United States. National vital statistics reports; vol 52 no 7. Hyattsville, Maryland: National Center for Health Statistics. 2003.
4. Hamilton BE, Sutton PD, Ventura SJ. Revised birth and fertility rates for the 1990s and new rates for Hispanic populations, 2000 and 2001: United States. National vital statistics reports; vol 51 no 12. Hyattsville, Maryland: National Center for Health Statistics. 2003.
5. Ventura SJ, Mosher WD, Curtin SC, Abma JC, Henshaw S. Trends in pregnancy rates for the United States, 1976–97: An update. National vital statistics reports; vol 49 no 4. Hyattsville, Maryland: National Center for Health Statistics. 2001.
6. Elam-Evans LD, Strauss LT, Herndon J, et al. Abortion surveillance—United States, 2000. In: MMWR 52 (SS-12). 2003.
7. Henshaw S. The Alan Guttmacher Institute. Unpublished tabulations, 2004.

List of Detailed Tables

| | |
|---|---|
| 1. Pregnancy, live birth, induced abortion, and fetal loss rates by age, race, and Hispanic origin of woman: United States, 1990–2000 | 3 |
| 2. Number and percent distribution of pregnancies by outcome of pregnancy according to age, race, and Hispanic origin of woman: United States, 2000 | 7 |
| 3. Estimates of lifetime pregnancy, live birth, and induced abortion rates by race and Hispanic origin of woman: United States, 1990 and 2000 | 8 |
| 4. Pregnancy, live birth, and induced abortion rates by marital status and race and Hispanic origin of woman: United States, 1990–2000 | 9 |

Table 1. Pregnancy, live birth, induced abortion, and fetal loss rates by age, race, and Hispanic origin of woman: United States, 1990–2000

[Rates are pregnancy outcomes per 1,000 women in specified group, estimated as of April 1 for 1990 and 2000 and as of July 1 for all other years]

| Pregnancy outcome, race, and Hispanic origin and year | Age of woman | | | | | | | | | | | |
|--|--------------------|--------------------------------|-------------|----------------|----------------|-------|-------|----------------|----------------|----------------|----------------|-----------------------------|
| | Total ¹ | Under 15 years ² | 15–19 years | | | | | 20–24 years | 25–29 years | 30–34 years | 35–39 years | 40–44 years ³ |
| | | | Total | 15–17 years | 18–19 years | | | | | | | |
| All races ⁴ | | | | | | | | | | | | |
| All pregnancies: | | | | | | | | | | | | |
| 2000 | 104.0 | 2.1 | 84.5 | 53.5 | 129.9 | 178.2 | 169.3 | 132.2 | 66.4 | 15.2 | | |
| 1999 | 102.1 | 2.2 | 86.7 | 56.0 | 131.8 | 176.2 | 166.6 | 126.6 | 63.8 | 14.4 | | |
| 1998 | 102.1 | 2.3 | 89.9 | 59.6 | 135.4 | 177.3 | 165.4 | 123.9 | 62.4 | 14.2 | | |
| 1997 | 101.6 | 2.5 | 92.6 | 62.5 | 139.4 | 177.0 | 163.2 | 121.0 | 60.5 | 13.7 | | |
| 1996 | 102.7 | 2.7 | 97.0 | 66.6 | 144.0 | 178.8 | 163.8 | 119.8 | 59.7 | 13.4 | | |
| 1995 | 103.4 | 3.0 | 101.1 | 70.7 | 148.2 | 178.2 | 163.4 | 118.4 | 58.3 | 12.9 | | |
| 1994 | 106.1 | 3.2 | 106.1 | 74.5 | 154.3 | 183.2 | 166.7 | 118.1 | 57.7 | 12.7 | | |
| 1993 | 108.7 | 3.3 | 109.2 | 76.0 | 158.7 | 188.7 | 170.4 | 118.0 | 56.9 | 12.2 | | |
| 1992 | 111.0 | 3.4 | 112.0 | 76.8 | 163.7 | 192.6 | 173.8 | 117.9 | 56.6 | 11.9 | | |
| 1991 | 112.5 | 3.3 | 116.0 | 79.5 | 166.5 | 195.0 | 175.5 | 117.5 | 56.0 | 11.1 | | |
| 1990 | 115.6 | 3.5 | 116.3 | 80.3 | 162.4 | 196.7 | 179.6 | 120.2 | 56.1 | 11.3 | | |
| Live births: | | | | | | | | | | | | |
| 2000 | 65.9 | 0.9 | 47.7 | 26.9 | 78.1 | 109.7 | 113.5 | 91.2 | 39.7 | 8.4 | | |
| 1999 | 64.4 | 0.9 | 48.8 | 28.2 | 79.0 | 107.9 | 111.2 | 87.1 | 37.8 | 7.8 | | |
| 1998 | 64.3 | 1.0 | 50.3 | 29.9 | 80.9 | 108.4 | 110.2 | 85.2 | 36.9 | 7.7 | | |
| 1997 | 63.6 | 1.1 | 51.3 | 31.4 | 82.1 | 107.3 | 108.3 | 83.0 | 35.7 | 7.4 | | |
| 1996 | 64.1 | 1.2 | 53.5 | 33.3 | 84.7 | 107.8 | 108.6 | 82.1 | 34.9 | 7.1 | | |
| 1995 | 64.6 | 1.3 | 56.0 | 35.5 | 87.7 | 107.5 | 108.8 | 81.1 | 34.0 | 6.8 | | |
| 1994 | 65.9 | 1.4 | 58.2 | 37.2 | 90.3 | 109.2 | 111.0 | 80.4 | 33.4 | 6.6 | | |
| 1993 | 67.0 | 1.4 | 59.0 | 37.4 | 91.1 | 111.3 | 113.2 | 79.9 | 32.7 | 6.3 | | |
| 1992 | 68.4 | 1.4 | 60.3 | 37.5 | 93.7 | 113.7 | 115.7 | 79.6 | 32.3 | 6.1 | | |
| 1991 | 69.3 | 1.4 | 61.8 | 38.5 | 94.0 | 115.3 | 117.2 | 79.2 | 31.9 | 5.7 | | |
| 1990 | 70.9 | 1.4 | 59.9 | 37.5 | 88.6 | 116.5 | 120.2 | 80.8 | 31.7 | 5.6 | | |
| Induced abortions: | | | | | | | | | | | | |
| 2000 | 21.3 | 0.9 | 24.0 | 14.5 | 37.7 | 46.3 | 31.6 | 18.7 | 9.7 | 3.2 | | |
| 1999 | 21.4 | 0.9 | 24.7 | 15.2 | 38.6 | 46.4 | 31.7 | 18.3 | 9.7 | 3.2 | | |
| 1998 | 21.5 | 1.0 | 25.8 | 16.4 | 39.9 | 47.0 | 31.7 | 17.9 | 9.5 | 3.2 | | |
| 1997 | 21.9 | 1.0 | 27.1 | 17.2 | 42.6 | 48.1 | 31.9 | 17.7 | 9.5 | 3.1 | | |
| 1996 | 22.4 | 1.1 | 28.6 | 18.6 | 44.0 | 49.3 | 32.1 | 17.7 | 9.7 | 3.2 | | |
| 1995 | 22.5 | 1.2 | 29.4 | 19.5 | 44.8 | 49.1 | 31.5 | 17.5 | 9.7 | 3.2 | | |
| 1994 | 23.7 | 1.3 | 31.6 | 21.0 | 47.9 | 51.9 | 32.1 | 18.1 | 9.9 | 3.2 | | |
| 1993 | 25.0 | 1.4 | 33.9 | 22.2 | 51.3 | 54.9 | 33.2 | 18.6 | 10.2 | 3.2 | | |
| 1992 | 25.7 | 1.4 | 35.2 | 22.9 | 53.3 | 55.9 | 33.5 | 18.9 | 10.3 | 3.2 | | |
| 1991 | 26.2 | 1.4 | 37.4 | 24.2 | 55.7 | 56.4 | 33.4 | 19.0 | 10.4 | 3.0 | | |
| 1990 | 27.4 | 1.5 | 40.3 | 26.5 | 57.9 | 56.7 | 33.9 | 19.7 | 10.8 | 3.2 | | |
| Fetal losses ⁵ : | | | | | | | | | | | | |
| 2000 | 16.7 | 0.3 | 12.8 | 12.0 | 14.1 | 22.2 | 24.2 | 22.3 | 17.0 | 3.6 | | |
| 1999 | 16.3 | 0.4 | 13.2 | 12.6 | 14.2 | 21.8 | 23.7 | 21.3 | 16.3 | 3.3 | | |
| 1998 | 16.3 | 0.4 | 13.8 | 13.3 | 14.5 | 21.9 | 23.5 | 20.8 | 15.9 | 3.3 | | |
| 1997 | 16.1 | 0.4 | 14.2 | 13.9 | 14.7 | 21.7 | 23.0 | 20.3 | 15.3 | 3.2 | | |
| 1996 | 16.2 | 0.5 | 14.9 | 14.7 | 15.2 | 21.7 | 23.1 | 20.0 | 15.0 | 3.1 | | |
| 1995 | 16.3 | 0.5 | 15.7 | 15.7 | 15.8 | 21.7 | 23.1 | 19.8 | 14.6 | 2.9 | | |
| 1994 | 16.5 | 0.5 | 16.3 | 16.3 | 16.2 | 22.1 | 23.6 | 19.6 | 14.4 | 2.8 | | |
| 1993 | 16.6 | 0.5 | 16.3 | 16.3 | 16.3 | 22.5 | 24.1 | 19.5 | 14.0 | 2.7 | | |
| 1992 | 16.9 | 0.5 | 16.5 | 16.3 | 16.8 | 23.0 | 24.6 | 19.4 | 13.9 | 2.6 | | |
| 1991 | 17.0 | 0.5 | 16.8 | 16.7 | 16.8 | 23.3 | 24.9 | 19.3 | 13.7 | 2.4 | | |
| 1990 | 17.2 | 0.5 | 16.1 | 16.2 | 15.9 | 23.5 | 25.5 | 19.7 | 13.6 | 2.4 | | |

See footnotes at end of table.

Table 1. Pregnancy, live birth, induced abortion, and fetal loss rates by age, race, and Hispanic origin of woman: United States, 1990–2000—Con.

[Rates are pregnancy outcomes per 1,000 women in specified group, estimated as of April 1 for 1990 and 2000 and as of July 1 for all other years]

| Pregnancy outcome, race, and Hispanic origin and year | Age of woman | | | | | | | | | |
|--|--------------------|--------------------------------|-------------|----------------|----------------|----------------|----------------|----------------|----------------|-----------------------------|
| | Total ¹ | Under 15 years ² | 15–19 years | | | 20–24 years | 25–29 years | 30–34 years | 35–39 years | 40–44 years ³ |
| | | | Total | 15–17 years | 18–19 years | | | | | |
| Non-Hispanic white | | | | | | | | | | |
| All pregnancies: | | | | | | | | | | |
| 2000 | 85.2 | 0.8 | 56.9 | 32.5 | 92.8 | 136.1 | 147.9 | 125.0 | 62.2 | 13.3 |
| 1999 | 84.5 | 0.9 | 59.7 | 35.4 | 95.6 | 135.8 | 147.3 | 120.6 | 60.0 | 12.8 |
| 1998 | 84.9 | 1.0 | 62.6 | 38.6 | 99.1 | 137.8 | 146.8 | 117.9 | 58.8 | 12.5 |
| 1997 | 84.5 | 1.1 | 65.4 | 41.2 | 103.0 | 138.1 | 144.5 | 114.6 | 56.7 | 12.0 |
| 1996 | 85.3 | 1.1 | 68.3 | 43.9 | 106.2 | 139.2 | 145.1 | 112.9 | 55.7 | 11.6 |
| 1995 | 86.3 | 1.2 | 71.8 | 46.8 | 110.8 | 140.4 | 145.8 | 111.5 | 54.1 | 11.2 |
| 1994 | 87.7 | 1.3 | 74.6 | 48.7 | 114.4 | 142.9 | 147.7 | 110.3 | 53.1 | 10.9 |
| 1993 | 89.7 | 1.3 | 77.1 | 49.9 | 117.8 | 147.5 | 150.7 | 109.3 | 51.9 | 10.4 |
| 1992 | 91.6 | 1.4 | 79.4 | 50.5 | 122.2 | 150.8 | 154.0 | 108.8 | 51.2 | 10.0 |
| 1991 | 94.0 | 1.4 | 84.9 | 54.2 | 127.3 | 155.6 | 157.2 | 108.8 | 51.1 | 9.2 |
| 1990 | 97.9 | 1.5 | 87.7 | 56.5 | 126.8 | 160.1 | 162.3 | 111.5 | 51.1 | 9.4 |
| Live births: | | | | | | | | | | |
| 2000 | 58.5 | 0.3 | 32.6 | 15.8 | 57.5 | 91.2 | 109.4 | 93.2 | 38.8 | 7.7 |
| 1999 | 57.7 | 0.3 | 34.1 | 17.1 | 59.4 | 90.6 | 108.6 | 89.5 | 37.3 | 7.3 |
| 1998 | 57.6 | 0.3 | 35.3 | 18.3 | 60.9 | 91.2 | 107.4 | 87.2 | 36.4 | 7.1 |
| 1997 | 56.8 | 0.4 | 36.0 | 19.3 | 62.1 | 90.0 | 104.8 | 84.3 | 34.8 | 6.8 |
| 1996 | 57.1 | 0.4 | 37.6 | 20.6 | 64.0 | 90.1 | 104.9 | 82.8 | 33.9 | 6.5 |
| 1995 | 57.5 | 0.4 | 39.3 | 22.0 | 65.2 | 90.2 | 105.1 | 81.5 | 32.8 | 6.1 |
| 1994 | 58.2 | 0.5 | 40.4 | 22.7 | 67.6 | 90.9 | 106.6 | 80.2 | 32.0 | 5.9 |
| 1993 | 58.9 | 0.5 | 40.7 | 22.7 | 67.7 | 92.2 | 108.2 | 79.0 | 31.0 | 5.6 |
| 1992 | 60.0 | 0.5 | 41.6 | 22.7 | 69.8 | 93.8 | 110.7 | 78.3 | 30.4 | 5.3 |
| 1991 | 60.9 | 0.5 | 43.3 | 23.6 | 70.6 | 95.7 | 112.2 | 77.7 | 30.2 | 4.8 |
| 1990 | 62.8 | 0.5 | 42.6 | 23.3 | 66.9 | 97.9 | 115.3 | 79.2 | 29.9 | 4.8 |
| Induced abortions: | | | | | | | | | | |
| 2000 | 11.7 | 0.3 | 14.8 | 8.5 | 24.1 | 26.0 | 16.2 | 10.2 | 6.0 | 2.2 |
| 1999 | 11.9 | 0.4 | 15.5 | 9.4 | 24.6 | 26.3 | 16.6 | 10.2 | 6.1 | 2.2 |
| 1998 | 12.5 | 0.4 | 16.9 | 10.7 | 26.3 | 27.7 | 17.5 | 10.4 | 6.3 | 2.2 |
| 1997 | 13.2 | 0.5 | 18.6 | 11.8 | 29.1 | 29.4 | 18.3 | 10.7 | 6.4 | 2.2 |
| 1996 | 13.6 | 0.5 | 19.3 | 12.6 | 29.8 | 30.3 | 18.8 | 10.8 | 6.7 | 2.3 |
| 1995 | 14.2 | 0.6 | 20.5 | 13.4 | 31.6 | 31.6 | 19.3 | 11.1 | 6.7 | 2.3 |
| 1994 | 14.8 | 0.6 | 21.8 | 14.1 | 33.7 | 33.2 | 19.2 | 11.4 | 6.7 | 2.3 |
| 1993 | 16.1 | 0.6 | 24.0 | 15.4 | 36.9 | 36.2 | 20.4 | 11.9 | 7.1 | 2.3 |
| 1992 | 16.7 | 0.7 | 25.3 | 16.0 | 38.9 | 37.5 | 20.8 | 12.2 | 7.2 | 2.3 |
| 1991 | 18.1 | 0.7 | 28.7 | 18.3 | 43.0 | 40.0 | 22.2 | 13.0 | 7.5 | 2.2 |
| 1990 | 19.7 | 0.8 | 32.5 | 21.1 | 46.8 | 41.9 | 23.4 | 13.8 | 7.9 | 2.4 |
| Fetal losses ⁵ : | | | | | | | | | | |
| 2000 | 15.0 | 0.2 | 9.4 | 8.2 | 11.2 | 18.9 | 22.3 | 21.7 | 17.3 | 3.4 |
| 1999 | 14.8 | 0.2 | 10.0 | 8.9 | 11.6 | 18.8 | 22.2 | 20.9 | 16.6 | 3.3 |
| 1998 | 14.8 | 0.2 | 10.5 | 9.6 | 11.9 | 18.9 | 21.9 | 20.3 | 16.2 | 3.2 |
| 1997 | 14.5 | 0.2 | 10.9 | 10.1 | 12.1 | 18.7 | 21.4 | 19.6 | 15.5 | 3.0 |
| 1996 | 14.6 | 0.2 | 11.4 | 10.7 | 12.5 | 18.7 | 21.4 | 19.3 | 15.1 | 2.9 |
| 1995 | 14.6 | 0.2 | 12.0 | 11.4 | 12.9 | 18.7 | 21.4 | 19.0 | 14.6 | 2.7 |
| 1994 | 14.7 | 0.2 | 12.4 | 11.9 | 13.2 | 18.9 | 21.8 | 18.7 | 14.3 | 2.6 |
| 1993 | 14.7 | 0.2 | 12.4 | 11.8 | 13.2 | 19.1 | 22.1 | 18.4 | 13.8 | 2.5 |
| 1992 | 14.9 | 0.2 | 12.5 | 11.8 | 13.6 | 19.5 | 22.6 | 18.2 | 13.6 | 2.4 |
| 1991 | 15.0 | 0.2 | 12.9 | 12.3 | 13.7 | 19.9 | 22.9 | 18.1 | 13.5 | 2.2 |
| 1990 | 15.3 | 0.2 | 12.5 | 12.1 | 13.0 | 20.3 | 23.5 | 18.5 | 13.3 | 2.1 |

See footnotes at end of table.

Table 1. Pregnancy, live birth, induced abortion, and fetal loss rates by age, race, and Hispanic origin of woman: United States, 1990–2000—Con.

[Rates are pregnancy outcomes per 1,000 women in specified group, estimated as of April 1 for 1990 and 2000 and as of July 1 for all other years]

| Pregnancy outcome, race, and Hispanic origin and year | Age of woman | | | | | | | | | |
|--|--------------------|--------------------------------|-------------|----------------|----------------|----------------|----------------|----------------|----------------|-----------------------------|
| | Total ¹ | Under 15 years ² | 15–19 years | | | 20–24 years | 25–29 years | 30–34 years | 35–39 years | 40–44 years ³ |
| | | | Total | 15–17 years | 18–19 years | | | | | |
| Non-Hispanic black | | | | | | | | | | |
| All pregnancies: | | | | | | | | | | |
| 2000 | 148.7 | 5.9 | 151.0 | 100.7 | 224.5 | 304.6 | 218.3 | 143.9 | 67.3 | 17.1 |
| 1999 | 146.5 | 6.2 | 153.8 | 102.2 | 229.3 | 300.9 | 215.0 | 138.6 | 65.3 | 15.8 |
| 1998 | 147.1 | 6.8 | 159.9 | 109.9 | 233.8 | 301.0 | 213.8 | 137.4 | 64.2 | 15.9 |
| 1997 | 147.4 | 7.3 | 166.0 | 116.3 | 242.9 | 298.6 | 210.6 | 135.8 | 63.2 | 15.6 |
| 1996 | 148.9 | 8.3 | 173.2 | 124.4 | 249.3 | 299.0 | 209.7 | 134.7 | 62.4 | 15.2 |
| 1995 | 149.7 | 9.2 | 180.1 | 133.6 | 253.1 | 293.6 | 205.8 | 134.6 | 61.6 | 14.8 |
| 1994 | 160.1 | 10.5 | 197.0 | 146.5 | 275.6 | 315.2 | 215.9 | 138.6 | 62.8 | 14.8 |
| 1993 | 169.0 | 10.7 | 208.3 | 155.2 | 288.1 | 333.6 | 225.2 | 143.3 | 64.1 | 14.8 |
| 1992 | 173.6 | 11.2 | 214.8 | 158.4 | 297.1 | 341.5 | 229.3 | 143.6 | 63.9 | 14.3 |
| 1991 | 176.2 | 11.3 | 220.2 | 163.4 | 298.2 | 344.3 | 229.3 | 143.3 | 62.9 | 14.2 |
| 1990 | 180.2 | 11.8 | 221.3 | 165.0 | 295.3 | 342.5 | 233.2 | 145.9 | 63.3 | 14.4 |
| Live births: | | | | | | | | | | |
| 2000 | 71.4 | 2.4 | 79.2 | 50.1 | 121.9 | 145.4 | 102.8 | 66.5 | 31.8 | 7.5 |
| 1999 | 69.9 | 2.6 | 81.0 | 51.7 | 123.9 | 142.1 | 99.8 | 63.9 | 30.6 | 6.8 |
| 1998 | 70.9 | 2.9 | 85.7 | 56.8 | 128.2 | 142.5 | 99.9 | 64.4 | 30.4 | 6.9 |
| 1997 | 70.3 | 3.2 | 88.3 | 60.7 | 131.0 | 138.8 | 97.2 | 63.6 | 29.6 | 6.8 |
| 1996 | 70.7 | 3.6 | 91.9 | 64.8 | 134.1 | 137.0 | 96.7 | 63.2 | 29.1 | 6.4 |
| 1995 | 72.8 | 4.2 | 97.2 | 70.4 | 139.2 | 137.8 | 98.5 | 64.4 | 28.8 | 6.3 |
| 1994 | 77.5 | 4.6 | 105.7 | 77.0 | 150.4 | 146.8 | 104.1 | 66.3 | 29.1 | 6.2 |
| 1993 | 81.5 | 4.6 | 110.5 | 81.1 | 154.6 | 154.5 | 109.2 | 68.1 | 29.4 | 6.1 |
| 1992 | 84.5 | 4.8 | 114.7 | 82.9 | 161.1 | 160.8 | 112.8 | 68.4 | 29.1 | 5.8 |
| 1991 | 87.0 | 4.9 | 118.2 | 86.1 | 162.2 | 164.8 | 115.1 | 68.9 | 28.7 | 5.7 |
| 1990 | 89.0 | 5.0 | 116.2 | 84.9 | 157.5 | 165.2 | 118.3 | 70.2 | 28.6 | 5.8 |
| Induced abortions: | | | | | | | | | | |
| 2000 | 57.4 | 2.9 | 57.4 | 37.3 | 86.8 | 120.2 | 89.9 | 50.7 | 23.0 | 6.6 |
| 1999 | 57.2 | 3.0 | 58.1 | 36.7 | 89.4 | 120.8 | 90.3 | 49.0 | 22.7 | 6.3 |
| 1998 | 56.5 | 3.2 | 58.6 | 38.0 | 89.0 | 120.3 | 89.0 | 47.0 | 21.9 | 6.2 |
| 1997 | 57.5 | 3.2 | 61.3 | 39.5 | 95.0 | 122.7 | 89.3 | 46.6 | 22.0 | 6.2 |
| 1996 | 58.6 | 3.8 | 64.1 | 42.4 | 97.9 | 125.3 | 88.9 | 46.1 | 21.9 | 6.4 |
| 1995 | 56.7 | 3.9 | 64.6 | 44.6 | 96.0 | 118.9 | 82.8 | 44.3 | 21.4 | 6.1 |
| 1994 | 61.1 | 4.6 | 71.2 | 49.1 | 105.8 | 129.2 | 85.9 | 45.7 | 22.3 | 6.2 |
| 1993 | 65.0 | 4.8 | 76.9 | 52.6 | 113.5 | 137.7 | 88.8 | 47.9 | 23.1 | 6.3 |
| 1992 | 65.9 | 5.1 | 78.6 | 53.5 | 115.2 | 137.6 | 88.4 | 47.7 | 23.4 | 6.2 |
| 1991 | 65.5 | 5.1 | 80.0 | 54.5 | 115.0 | 135.4 | 85.4 | 46.8 | 22.9 | 6.2 |
| 1990 | 67.0 | 5.4 | 83.5 | 57.7 | 117.4 | 133.1 | 85.4 | 47.5 | 23.5 | 6.4 |
| Fetal losses ⁵ : | | | | | | | | | | |
| 2000 | 19.9 | 0.6 | 14.3 | 13.3 | 15.8 | 38.9 | 25.6 | 26.7 | 12.5 | 3.0 |
| 1999 | 19.4 | 0.7 | 14.7 | 13.7 | 16.0 | 38.0 | 24.9 | 25.7 | 12.0 | 2.7 |
| 1998 | 19.7 | 0.8 | 15.7 | 15.1 | 16.6 | 38.2 | 24.9 | 25.9 | 11.9 | 2.7 |
| 1997 | 19.6 | 0.9 | 16.4 | 16.1 | 16.9 | 37.2 | 24.2 | 25.6 | 11.6 | 2.7 |
| 1996 | 19.7 | 1.0 | 17.3 | 17.2 | 17.3 | 36.7 | 24.1 | 25.4 | 11.4 | 2.5 |
| 1995 | 20.2 | 1.1 | 18.4 | 18.7 | 18.0 | 36.9 | 24.5 | 25.9 | 11.3 | 2.5 |
| 1994 | 21.5 | 1.2 | 20.1 | 20.4 | 19.4 | 39.3 | 25.9 | 26.7 | 11.4 | 2.4 |
| 1993 | 22.5 | 1.2 | 20.9 | 21.5 | 20.0 | 41.4 | 27.2 | 27.4 | 11.6 | 2.4 |
| 1992 | 23.2 | 1.3 | 21.5 | 22.0 | 20.8 | 43.1 | 28.1 | 27.5 | 11.4 | 2.3 |
| 1991 | 23.7 | 1.3 | 22.1 | 22.8 | 21.0 | 44.1 | 28.7 | 27.7 | 11.3 | 2.3 |
| 1990 | 24.2 | 1.3 | 21.6 | 22.5 | 20.4 | 44.2 | 29.5 | 28.2 | 11.2 | 2.3 |

See footnotes at end of table.

Table 1. Pregnancy, live birth, induced abortion, and fetal loss rates by age, race, and Hispanic origin of woman: United States, 1990–2000—Con.

[Rates are pregnancy outcomes per 1,000 women in specified group, estimated as of April 1 for 1990 and 2000 and as of July 1 for all other years]

| Pregnancy outcome, race, and Hispanic origin and year | Age of woman | | | | | | | | | |
|--|--------------------|--------------------------------|-------------|----------------|----------------|----------------|----------------|----------------|----------------|-----------------------------|
| | Total ¹ | Under 15 years ² | 15–19 years | | | 20–24 years | 25–29 years | 30–34 years | 35–39 years | 40–44 years ³ |
| | | | Total | 15–17 years | 18–19 years | | | | | |
| Hispanic ⁶ | | | | | | | | | | |
| All pregnancies: | | | | | | | | | | |
| 2000 | 145.4 | 3.0 | 132.0 | 83.1 | 201.7 | 236.8 | 211.4 | 143.2 | 81.4 | 22.5 |
| 1999 | 142.7 | 3.2 | 133.1 | 86.5 | 199.9 | 234.0 | 207.4 | 137.2 | 78.7 | 21.1 |
| 1998 | 143.0 | 3.4 | 136.0 | 89.6 | 204.7 | 236.1 | 207.9 | 134.5 | 77.0 | 21.3 |
| 1997 | 143.3 | 3.6 | 137.0 | 91.4 | 205.6 | 238.5 | 208.8 | 132.5 | 76.4 | 21.2 |
| 1996 | 148.4 | 4.0 | 145.9 | 97.7 | 217.9 | 248.8 | 213.5 | 134.9 | 77.7 | 21.0 |
| 1995 | 150.1 | 4.2 | 151.4 | 103.0 | 223.3 | 251.9 | 212.3 | 134.3 | 77.6 | 21.2 |
| 1994 | 155.4 | 4.4 | 156.9 | 107.5 | 229.7 | 262.2 | 218.1 | 137.4 | 78.6 | 21.8 |
| 1993 | 159.4 | 4.4 | 158.2 | 105.2 | 236.6 | 269.1 | 223.3 | 140.8 | 79.4 | 21.5 |
| 1992 | 165.1 | 4.4 | 161.9 | 107.1 | 242.4 | 278.3 | 228.9 | 145.0 | 83.0 | 22.5 |
| 1991 | 163.0 | 4.2 | 161.4 | 104.7 | 242.7 | 271.2 | 226.7 | 142.2 | 80.2 | 21.3 |
| 1990 | 163.2 | 4.0 | 155.8 | 101.0 | 231.4 | 263.0 | 229.2 | 146.4 | 81.0 | 21.7 |
| Live births: | | | | | | | | | | |
| 2000 | 95.9 | 1.7 | 87.3 | 55.5 | 132.6 | 161.3 | 139.9 | 97.1 | 46.6 | 12.0 |
| 1999 | 93.0 | 1.9 | 86.8 | 56.9 | 129.5 | 157.3 | 135.8 | 92.3 | 44.5 | 11.1 |
| 1998 | 93.2 | 1.9 | 87.9 | 58.5 | 131.5 | 159.3 | 136.1 | 90.5 | 43.4 | 11.3 |
| 1997 | 94.2 | 2.1 | 89.6 | 61.1 | 132.4 | 162.6 | 137.5 | 89.6 | 43.4 | 11.2 |
| 1996 | 97.5 | 2.4 | 94.6 | 64.2 | 140.0 | 170.2 | 140.7 | 91.3 | 43.9 | 11.2 |
| 1995 | 98.8 | 2.6 | 99.3 | 68.3 | 145.4 | 171.9 | 140.4 | 90.5 | 43.7 | 11.2 |
| 1994 | 100.7 | 2.6 | 101.3 | 69.9 | 147.5 | 175.7 | 142.4 | 91.1 | 43.4 | 11.1 |
| 1993 | 103.3 | 2.6 | 101.8 | 68.5 | 151.1 | 180.0 | 146.0 | 93.2 | 44.1 | 11.1 |
| 1992 | 106.1 | 2.5 | 103.3 | 68.9 | 153.8 | 185.1 | 148.7 | 94.7 | 45.3 | 11.4 |
| 1991 | 106.9 | 2.4 | 104.6 | 69.2 | 155.5 | 184.6 | 149.9 | 95.0 | 44.7 | 11.1 |
| 1990 | 107.6 | 2.4 | 100.2 | 65.8 | 147.6 | 180.8 | 152.8 | 98.1 | 45.2 | 11.4 |
| Induced abortions: | | | | | | | | | | |
| 2000 | 30.6 | 1.0 | 30.3 | 18.4 | 47.3 | 58.6 | 40.5 | 25.2 | 13.8 | 5.0 |
| 1999 | 31.4 | 1.0 | 32.1 | 20.2 | 49.1 | 60.3 | 41.5 | 25.2 | 14.1 | 5.0 |
| 1998 | 31.6 | 1.1 | 33.7 | 21.5 | 51.6 | 60.1 | 41.6 | 24.6 | 14.0 | 5.0 |
| 1997 | 30.7 | 1.1 | 32.7 | 20.3 | 51.4 | 58.8 | 40.8 | 23.7 | 13.4 | 4.9 |
| 1996 | 31.9 | 1.2 | 35.7 | 22.9 | 54.9 | 60.9 | 41.6 | 24.1 | 14.0 | 4.8 |
| 1995 | 32.2 | 1.2 | 35.8 | 23.5 | 54.0 | 62.1 | 40.8 | 24.4 | 14.3 | 5.0 |
| 1994 | 35.4 | 1.4 | 39.0 | 26.1 | 58.0 | 68.2 | 44.1 | 26.8 | 15.6 | 5.7 |
| 1993 | 36.4 | 1.4 | 39.7 | 25.5 | 60.7 | 70.4 | 45.0 | 27.6 | 15.4 | 5.4 |
| 1992 | 38.8 | 1.5 | 41.6 | 26.8 | 63.3 | 73.9 | 47.3 | 29.9 | 17.2 | 5.9 |
| 1991 | 35.8 | 1.4 | 39.6 | 24.2 | 61.7 | 67.4 | 43.6 | 26.8 | 15.4 | 5.2 |
| 1990 | 35.1 | 1.1 | 39.1 | 24.3 | 59.5 | 63.4 | 42.6 | 27.2 | 15.4 | 5.2 |
| Fetal losses ⁵ : | | | | | | | | | | |
| 2000 | 18.9 | 0.3 | 14.4 | 9.1 | 21.8 | 16.8 | 31.0 | 20.8 | 21.0 | 5.4 |
| 1999 | 18.3 | 0.3 | 14.3 | 9.4 | 21.3 | 16.4 | 30.1 | 19.8 | 20.1 | 5.0 |
| 1998 | 18.2 | 0.3 | 14.4 | 9.6 | 21.6 | 16.6 | 30.2 | 19.4 | 19.6 | 5.1 |
| 1997 | 18.4 | 0.3 | 14.7 | 10.0 | 21.8 | 17.0 | 30.5 | 19.2 | 19.6 | 5.0 |
| 1996 | 18.9 | 0.4 | 15.6 | 10.6 | 23.0 | 17.8 | 31.2 | 19.6 | 19.8 | 5.0 |
| 1995 | 19.1 | 0.4 | 16.3 | 11.2 | 23.9 | 17.9 | 31.1 | 19.4 | 19.7 | 5.0 |
| 1994 | 19.4 | 0.4 | 16.7 | 11.5 | 24.2 | 18.3 | 31.6 | 19.5 | 19.5 | 5.0 |
| 1993 | 19.8 | 0.4 | 16.7 | 11.3 | 24.8 | 18.8 | 32.4 | 20.0 | 19.9 | 5.0 |
| 1992 | 20.3 | 0.4 | 17.0 | 11.3 | 25.3 | 19.3 | 33.0 | 20.3 | 20.4 | 5.1 |
| 1991 | 20.3 | 0.4 | 17.2 | 11.4 | 25.5 | 19.3 | 33.2 | 20.4 | 20.1 | 5.0 |
| 1990 | 20.5 | 0.4 | 16.5 | 10.8 | 24.3 | 18.9 | 33.9 | 21.0 | 20.4 | 5.1 |

¹Rates computed by relating the number of events to women of all ages to women aged 15–44 years.²Rates computed by relating the number of events to women under age 15 years to women aged 10–14 years.³Rates computed by relating the number of events to women aged 40 years and over to women aged 40–44 years.⁴Includes races other than white and black and origin not stated.⁵Spontaneous fetal losses from recognized pregnancies of all gestational periods as estimated from reports from women in the 1995 National Survey of Family Growth conducted by the National Center for Health Statistics. The rate of fetal loss depends on the degree to which losses at very early gestations are detected. See "Revised Pregnancy Rates, 1990–97, and New Rates for 1998–99: United States," for more information.⁶Includes all persons of Hispanic origin of any race.NOTES: Rates for 1990–99 for ages 15–17 and 18–19 years have been revised and may differ from rates previously published (4). For information on sources and methods, see "Revised Pregnancy Rates, 1990–97, and New Rates for 1998–99: United States." The revised populations are available on the Internet at: <http://www.cdc.gov/nchs>.

Table 2. Number and percent distribution of pregnancies by outcome of pregnancy according to age, race, and Hispanic origin of woman: United States, 2000

[Numbers in thousands]

| Pregnancy outcome and race and Hispanic origin | Total | Under 15 years | 15–19 years | | | | | | | |
|--|-------|----------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------------|
| | | | Total | 15–17 years | 18–19 years | 20–24 years | 25–29 years | 30–34 years | 35–39 years | 40 years and over |
| All races ¹ | | | | | | | | | | |
| Number of pregnancies | | | | | | | | | | |
| All pregnancies | 6,401 | 21 | 831 | 312 | 519 | 1,653 | 1,622 | 1,347 | 756 | 172 |
| Live births | 4,059 | 9 | 469 | 157 | 312 | 1,018 | 1,088 | 929 | 452 | 95 |
| Induced abortions | 1,313 | 9 | 235 | 85 | 151 | 430 | 303 | 190 | 110 | 37 |
| Fetal losses ² | 1,029 | 3 | 126 | 70 | 56 | 206 | 232 | 228 | 194 | 40 |
| Non-Hispanic white | | | | | | | | | | |
| All pregnancies | 3,497 | 5 | 361 | 123 | 238 | 794 | 894 | 842 | 492 | 109 |
| Live births | 2,400 | 2 | 207 | 60 | 148 | 532 | 661 | 627 | 308 | 63 |
| Induced abortions | 479 | 2 | 94 | 32 | 62 | 151 | 98 | 68 | 48 | 18 |
| Fetal losses ² | 618 | 1 | 60 | 31 | 29 | 110 | 135 | 146 | 137 | 28 |
| Non-Hispanic black | | | | | | | | | | |
| All pregnancies | 1,265 | 9 | 222 | 88 | 134 | 415 | 294 | 199 | 101 | 25 |
| Live births | 607 | 4 | 116 | 44 | 73 | 198 | 138 | 92 | 48 | 11 |
| Induced abortions | 488 | 5 | 84 | 33 | 52 | 164 | 121 | 70 | 35 | 10 |
| Fetal losses ² | 170 | 1 | 21 | 12 | 9 | 53 | 34 | 37 | 19 | 4 |
| Hispanic ³ | | | | | | | | | | |
| All pregnancies | 1,237 | 5 | 196 | 72 | 123 | 363 | 330 | 209 | 110 | 25 |
| Live births | 816 | 3 | 129 | 48 | 81 | 248 | 218 | 141 | 63 | 14 |
| Induced abortions | 261 | 2 | 45 | 16 | 29 | 90 | 63 | 37 | 19 | 6 |
| Fetal losses ² | 161 | – | 21 | 8 | 13 | 26 | 48 | 30 | 28 | 6 |
| All races ¹ | | | | | | | | | | |
| Percent distribution | | | | | | | | | | |
| All pregnancies | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 |
| Live births | 63.4 | 41.4 | 56.5 | 50.4 | 60.1 | 61.6 | 67.0 | 69.0 | 59.8 | 55.1 |
| Induced abortions | 20.5 | 41.6 | 28.3 | 27.2 | 29.1 | 26.0 | 18.7 | 14.1 | 14.5 | 21.3 |
| Fetal losses ² | 16.1 | 16.9 | 15.2 | 22.5 | 10.8 | 12.4 | 14.3 | 16.9 | 25.7 | 23.6 |
| Non-Hispanic white | | | | | | | | | | |
| All pregnancies | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 |
| Live births | 68.6 | 37.3 | 57.4 | 48.5 | 62.0 | 67.0 | 74.0 | 74.5 | 62.5 | 57.9 |
| Induced abortions | 13.7 | 43.2 | 26.0 | 26.2 | 26.0 | 19.1 | 10.9 | 8.1 | 9.7 | 16.3 |
| Fetal losses ² | 17.7 | 19.4 | 16.6 | 25.3 | 12.1 | 13.9 | 15.1 | 17.4 | 27.9 | 25.8 |
| Non-Hispanic black | | | | | | | | | | |
| All pregnancies | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 |
| Live births | 48.0 | 40.2 | 52.5 | 49.7 | 54.3 | 47.7 | 47.1 | 46.2 | 47.3 | 44.1 |
| Induced abortions | 38.6 | 49.1 | 38.1 | 37.1 | 38.7 | 39.5 | 41.2 | 35.2 | 34.1 | 38.6 |
| Fetal losses ² | 13.4 | 10.7 | 9.5 | 13.2 | 7.0 | 12.8 | 11.7 | 18.6 | 18.6 | 17.3 |
| Hispanic ³ | | | | | | | | | | |
| All pregnancies | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 |
| Live births | 65.9 | 57.6 | 66.1 | 66.8 | 65.7 | 68.1 | 66.2 | 67.8 | 57.3 | 53.5 |
| Induced abortions | 21.1 | 33.0 | 23.0 | 22.2 | 23.5 | 24.7 | 19.1 | 17.6 | 16.9 | 22.4 |
| Fetal losses ² | 13.0 | 9.5 | 10.9 | 11.0 | 10.8 | 7.1 | 14.7 | 14.5 | 25.8 | 24.1 |

– Quantity less than 500.

¹Includes races other than white and black and origin not stated.²Spontaneous fetal losses from recognized pregnancies of all gestational periods as estimated from reports from women in the 1995 National Survey of Family Growth conducted by the National Center for Health Statistics. The rate of fetal loss depends on the degree to which losses at very early gestations are detected. See "Revised Pregnancy Rates, 1990–97, and New Rates for 1998–99: United States."³Includes all persons of Hispanic origin of any race.

NOTE: Due to rounding, figures may not add to totals. Percent distributions based on unrounded frequencies.

Table 3. Estimates of lifetime pregnancy, live birth, and induced abortion rates by race and Hispanic origin of woman: United States, 1990 and 2000

[Rates are sums of rates for 5-year age groups (in table 1) multiplied by 5 and divided by 1,000]

| Race and Hispanic origin | Pregnancies ¹ | | Live births | | Induced abortions | |
|---------------------------------|--------------------------|------|-------------|------|-------------------|------|
| | 2000 | 1990 | 2000 | 1990 | 2000 | 1990 |
| Total ² | 3.2 | 3.4 | 2.1 | 2.1 | 0.7 | 0.8 |
| Non-Hispanic white | 2.7 | 2.9 | 1.9 | 1.8 | 0.4 | 0.6 |
| Non-Hispanic black | 4.5 | 5.2 | 2.2 | 2.5 | 1.8 | 1.9 |
| Hispanic ³ | 4.2 | 4.5 | 2.7 | 3.0 | 0.9 | 1.0 |

¹Includes estimates of fetal losses not shown separately.²Includes races other than white and black.³Includes all persons of Hispanic origin of any race.

NOTE: Total pregnancy rates are estimates of the number of pregnancies a woman would have if she experiences the age-specific pregnancy rates observed in a given year.

Table 4. Pregnancy, live birth, and induced abortion rates by marital status and race and Hispanic origin of woman: United States, 1990–2000

[Rates per 1,000 women aged 15–44 years in specified group; populations estimated as of July 1 for each year]

| Race and Hispanic origin and year | Married | | | Unmarried | | |
|-----------------------------------|------------------------------|------------|------------------|------------------------------|------------|------------------|
| | All pregnancies ¹ | Live birth | Induced abortion | All pregnancies ¹ | Live birth | Induced abortion |
| All races | | | | | | |
| 2000 | 116.5 | 87.4 | 7.9 | 89.6 | 44.1 | 34.9 |
| 1999 | 113.9 | 84.8 | 8.1 | 89.2 | 43.3 | 35.1 |
| 1998 | 113.0 | 84.2 | 7.9 | 90.0 | 43.3 | 35.8 |
| 1997 | 111.6 | 82.7 | 8.0 | 90.7 | 42.9 | 36.9 |
| 1996 | 111.6 | 82.3 | 8.5 | 92.9 | 43.8 | 37.9 |
| 1995 | 111.7 | 82.6 | 8.4 | 94.0 | 44.3 | 38.5 |
| 1994 | 112.5 | 82.9 | 8.9 | 98.7 | 46.2 | 40.8 |
| 1993 | 116.8 | 86.1 | 9.5 | 99.2 | 44.8 | 43.1 |
| 1992 | 119.8 | 88.5 | 9.7 | 100.6 | 44.9 | 44.6 |
| 1991 | 121.6 | 89.6 | 10.1 | 101.7 | 45.0 | 45.6 |
| 1990 | 126.3 | 93.2 | 10.6 | 102.3 | 43.8 | 47.7 |
| White total | | | | | | |
| 2000 | 116.1 | 88.7 | 5.9 | 73.0 | 38.2 | 25.5 |
| 1999 | 113.7 | 86.4 | 6.0 | 72.7 | 37.4 | 26.0 |
| 1998 | 113.1 | 85.8 | 6.0 | 73.2 | 36.9 | 27.1 |
| 1997 | 111.5 | 84.2 | 6.1 | 73.4 | 36.3 | 28.0 |
| 1996 | 111.5 | 84.1 | 6.4 | 75.2 | 37.0 | 28.9 |
| 1995 | 111.7 | 84.2 | 6.5 | 76.2 | 37.0 | 30.0 |
| 1994 | 112.0 | 84.3 | 6.8 | 79.0 | 37.8 | 31.8 |
| 1993 | 115.6 | 87.0 | 7.3 | 78.3 | 35.6 | 33.9 |
| 1992 | 118.4 | 89.1 | 7.5 | 79.1 | 35.0 | 35.6 |
| 1991 | 120.0 | 90.3 | 7.9 | 80.0 | 34.5 | 37.1 |
| 1990 | 125.1 | 94.1 | 8.5 | 80.2 | 32.9 | 39.4 |
| Non-Hispanic white | | | | | | |
| 2000 | 111.5 | 85.0 | 4.7 | 55.0 | 28.0 | 19.8 |
| 1999 | 108.9 | 82.9 | 4.7 | 55.5 | 27.9 | 20.4 |
| 1998 | 107.9 | 82.1 | 4.8 | 56.9 | 27.9 | 21.8 |
| 1997 | 105.7 | 80.2 | 5.0 | 58.0 | 27.5 | 23.4 |
| 1996 | 105.2 | 79.6 | 5.3 | 59.7 | 28.2 | 24.4 |
| 1995 | 105.6 | 79.9 | 5.4 | 61.0 | 28.1 | 25.8 |
| 1994 | 106.3 | 80.5 | 5.5 | 63.1 | 28.4 | 27.5 |
| Black total | | | | | | |
| 2000 | 106.9 | 68.6 | 20.3 | 160.2 | 70.5 | 71.1 |
| 1999 | 104.8 | 65.9 | 21.0 | 159.6 | 69.7 | 70.9 |
| 1998 | 102.8 | 65.0 | 20.4 | 161.6 | 71.6 | 70.8 |
| 1997 | 102.2 | 63.9 | 20.7 | 163.5 | 71.5 | 72.3 |
| 1996 | 101.8 | 62.2 | 22.5 | 166.0 | 72.8 | 73.2 |
| 1995 | 102.8 | 64.1 | 21.1 | 166.0 | 74.5 | 71.1 |
| 1994 | 107.7 | 66.1 | 23.5 | 179.6 | 80.9 | 76.6 |
| 1993 | 119.1 | 73.0 | 26.2 | 186.8 | 82.9 | 81.2 |
| Hispanic² | | | | | | |
| 2000 | 130.4 | 101.8 | 11.6 | 153.7 | 87.3 | 51.9 |
| 1999 | 130.5 | 100.0 | 12.4 | 153.8 | 84.9 | 53.6 |
| 1998 | 133.7 | 102.3 | 12.4 | 151.6 | 82.8 | 53.4 |
| 1997 | 136.5 | 103.7 | 12.6 | 151.2 | 83.2 | 51.8 |
| 1996 | 141.3 | 107.2 | 13.3 | 156.7 | 86.1 | 53.8 |
| 1995 | 141.5 | 107.1 | 13.8 | 160.3 | 88.8 | 54.4 |
| 1994 | 140.8 | 104.7 | 15.9 | 171.7 | 95.8 | 57.5 |
| 1993 | 151.1 | 113.0 | 16.4 | 169.6 | 91.4 | 60.6 |

¹Includes pregnancies ending in fetal loss, not shown separately.²Includes all persons of Hispanic origin of any race.

NOTES: Marital status is the woman's marital status as of the date the pregnancy ended, rather than at the time of conception. Figures for married women include pregnancies for separated women. Rates for 1990–99 have been revised and may differ from rates previously published (4). For information on data sources and methods, see "Revised Pregnancy Rates, 1990–97, and New Rates for 1998–99: United States."

Contents

Abstract 1
Introduction 1
Overview of findings. 1
References 2
List of Detailed Tables 2

Suggested citation

Ventura SJ, Abma JC, Mosher WD, Henshaw S. Estimated pregnancy rates for the United States, 1990–2000: An Update. National vital statistics reports; vol 52 no 23. Hyattsville, Maryland: National Center for Health Statistics. 2004.

National Center for Health Statistics

Director, Edward J. Sondik, Ph.D.
Deputy Director, Jack R. Anderson

Division of Vital Statistics

Director, Charles J. Rothwell

To receive this publication regularly, contact the National Center for Health Statistics by calling 1-866-441-6247. E-mail: nchsquery@cdc.gov
Internet: www.cdc.gov/nchs

Copyright information

All material appearing in this report is in the public domain and may be reproduced or copied without permission; citation as to source, however, is appreciated.

**U.S. DEPARTMENT OF
HEALTH & HUMAN SERVICES**

Centers for Disease Control and Prevention
National Center for Health Statistics
3311 Toledo Road
Hyattsville, Maryland 20782

DHHS Publication No. (PHS) 2004–1120
PRS 04-0392 (6/2004)

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE, \$300

| |
|--|
| FIRST CLASS POSTAGE & FEES PAID CDC/NCHS PERMIT NO. G-284 |
|--|